



Momentum Distribution Properties of Ultracold Hard-Core Anyons in One Dimension

Tim Skaras, Li Yang, Shah Saad Alam, and Han Pu

Dept. of Physics and Astronomy, Rice University, Houston, Texas, USA

Motivation

Anyons: mostly theoretical but can model real systems

Anyons:


- Are theoretical particles
- Used in 2D systems to realize topological quantum computers
- Help us understand real 1D systems of spinful particles

Background

Anyon is "mixture" between boson and fermion

Bosons vs. Fermions:

- Two types of particles in nature: bosons and fermions
- Different because they have different spin
- Bosons: spin quantum number is an integer (0, 1, 2, ...)
- Fermions: spin quantum number is odd multiple of $1/2$ ($1/2, 3/2, \dots$)




Boson Ex: Photon



Fermion Ex: Electron


Bosons & Fermions behave differently:

- Fermions follow Pauli Exclusion Principle (PEP), bosons do not
- PEP: two fermions cannot have same quantum state or position



Anyons:

- Not experimentally observed like bosons & fermions
- Described by anyon parameter κ between 0 and 2
- Can be bosons, fermions, or something in between as κ varies
- Need not follow PEP, unless $\kappa = 1$



Methods

Hard core gives particle fermi-like properties

What is a hard core?


- Anyons differ from fermions — e.g., only the latter obeys PEP
- Particles with hard cores cannot occupy same position
- So, hard-core anyons (HCA) are similar to fermions

known as "Fermionization"

The system we study comprises:

- Two spinless, hard-core anyons in harmonic potential: $V(x) = \frac{1}{2}m\omega^2 x^2$
- Ultracold HCA: this means HCA are in lowest energy configuration


Note: all the following equations use natural units — i.e., $\hbar = m = \omega = 1$



Anyon-fermion mapping:

- Transforms fermi wave function into HCA wave function
- Multiplies fermi w.f. by exponential function

Anyon-fermion mapping lets us study HCA



Momentum distribution $n_\kappa(p)$:

- Gives number density of particles in momentum space
- Calculated using the one-body density matrix $\rho(x, x')$

Results

HCA momentum distribution decays $\mathcal{O}(p^{-4})$


For large momentum p :

- Fermi momentum distribution $n_F(p)$ decays to zero exponentially
- HCA momentum distribution has power law decay

We show using asymptotic analysis:


$$\lim_{p \rightarrow \infty} = \left(\frac{2}{\pi} \right)^{3/2} \frac{\cos^2(\frac{\pi\kappa}{2})}{p^4} = \frac{C_\kappa}{p^4}$$

Numerically calculate C_κ to confirm our analytic result:



Expanding HCA mimic fermi momentum distribution

Time Evolution of HCA Momentum Distribution ($\kappa = 1/2$)




What happens if the potential is turned off? (i.e., $\omega = 1 \rightarrow \omega = 0$)

- We show numerically that $\lim_{t \rightarrow \infty} n_\kappa(p; t) = n_F(p)$
- As HCA gas expands, its momentum dist. becomes the fermi dist.
- This phenomenon is known as "dynamical fermionization"
- We show analytically: $C_\kappa(t) = \left(\frac{2}{\pi} \right)^{3/2} \cos^2\left(\frac{\pi\kappa}{2}\right) \frac{1}{(1+t^2)^{3/2}}$

Conclusions

HCA undergo dynamical fermionization

- Other papers have shown that dynamical fermionization occurs in systems of hard-core bosons
- We show that it also occurs in systems of hard-core anyons
- The momentum tail coefficient for hard-core anyons shares same time dependence of hard-core bosons



Future Work

Still need to study case where particles have spin

- We have studied the two hard-core anyon case
- We are trying to answer the same questions for the case where N hard-core anyons are confined in a harmonic trap
 - Less explored area because expressions for the momentum distribution grow more complex with more particles
- So far we have only studied spinless hard-core anyons, which means the spin of each particle is constrained and cannot change
 - When the spin of each particle is allowed to change, the dynamics are more sophisticated

References

1. Girardeau, M. "Relationship between Systems of Impenetrable Bosons and Fermions in One Dimension." *Journal of Mathematical Physics* 1, no. 6 (1960): 516-23. doi:10.1063/1.1703687.
2. Girardeau, M. D., E. M. Wright, and J. M. Triscari. "Ground-state Properties of a One-dimensional System of Hard-core Bosons in a Harmonic Trap." *Physical Review A* 63, no. 3 (2001). doi:10.1103/physreva.63.033601.
3. Girardeau, M. D. "Anyon-Fermion Mapping and Applications to Ultracold Gases in Tight Waveguides." *Physical Review Letters* 97, no. 10 (2006). doi:10.1103/physrevlett.97.100402.
4. Light Bulb by Schmidt Serje from the Noun Project
5. Minguzzi, A., P. Vignolo, and M.P. Tosi. "High-momentum Tail in the Tonks Gas under Harmonic Confinement." *Physics Letters A* 294, no. 3-4 (2002): 222-26. doi:10.1016/s0375-9601(02)00042-7.
6. Minguzzi, A., and D. M. Gangardt. "Exact Coherent States of a Harmonically Confined Tonks-Girardeau Gas." *Physical Review Letters* 94, no. 24 (2005). doi:10.1103/physrevlett.94.240404.
7. Photon by Anthony Ledoux from the Noun Project
8. quantum computing by Andrew Forrester from the Noun Project
9. spinning by Andrejs Kirma from the Noun Project

We would like to thank the Smalley-Curl Institute for funding this research