

**THE UNIVERSITY OF CALGARY**  
**FACULTY OF SCIENCE**  
**CHEMISTRY 209**  
**Winter 2013**  
**Final Exam**

Date: April 19<sup>th</sup>, 2013

Time Allotted: 3 hours

Please make sure you fill in your NAME, student ID and Lecture Section and encode all information on the optical score sheet.

Failure to do so will result in the loss of at least 2 marks.

Only Casio FX260 or similar non-programmable scientific calculators are permitted.

A Chemical Data and Formula Sheet is attached, which can be detached.

FIRST NAME: \_\_\_\_\_ LAST NAME: \_\_\_\_\_

ID: \_\_\_\_\_

Lecture Section: Please check one

- |                          |     |    |          |                   |
|--------------------------|-----|----|----------|-------------------|
| <input type="checkbox"/> | L01 | TR | 12:30 pm | Dr. Masood Parvez |
| <input type="checkbox"/> | L02 | TR | 8:00 am  | Dr. Scott Hinman  |

This exam consists of 14 pages, 21 multiple choice questions worth two marks each (total 42 marks), and 4 written answer questions (total 30 marks) for a total of 72 marks. Make sure to count the number of pages (should be 14 pages excluding data sheet) and LET THE INVIGILATOR KNOW IMMEDIATELY IF YOU ARE MISSING A PAGE.

All questions must be answered to earn full marks. The answers to the multiple choice questions must be entered on the optical score sheet using pencil. Answers to the written answer questions should be written in the exam booklet in ink.


**AT THE END OF THE EXAMINATION, HAND IN THE WHOLE EXAM BOOKLET AND THE OPTICAL SCORE SHEET**

Do not write in spaces in the table below

Q 22	Q 23	Q24	Q25

**SECTION I - Machine graded section (Total value 42 marks)**  
**Select the best answer and encode it on the Optical Score Sheet**

1. What is the mass percent of each element in sulfuric acid,  $\text{H}_2\text{SO}_4$ ?
- a. 2.055% H, 32.70% S, 65.25% O  
b. 1.028% H, 32.69% S, 66.28% O  
c. 28.57% H, 14.29% S, 57.17% O  
d. 1.028% H, 33.72% S, 65.25% O  
e. 2.016% H, 32.07% S, 65.91% O
2. A student analyzed a first-order reaction and obtained the graph below. Unfortunately, the student forgot to label the axes. What are the correct labels for the  $x$  and  $y$  axes?




- a.  $x$  axis = time,  $y$  axis =  $\ln[A]$   
b.  $x$  axis = time,  $y$  axis =  $[A]$   
c.  $x$  axis =  $\ln[\text{time}]$ ,  $y$  axis =  $[A]$   
d.  $x$  axis = time,  $y$  axis =  $1/[A]$   
e.  $x$  axis =  $1/\text{time}$ ,  $y$  axis =  $1/[A]$


3. According to the collision theory, which one of the following does NOT affect the rate of a reaction:

- a. temperature
- b. the speed of colliding particles
- c.  $\Delta H$  of the overall reaction
- d. the orientation of colliding particles
- e. the frequency with which particles collide

4. Nitrogen dioxide reacts with carbon monoxide to produce nitrogen monoxide and carbon dioxide.



A proposed mechanism for this reaction is:



What is a rate law that is consistent with the proposed mechanism?

- a.  $\text{rate} = k [\text{NO}_2]^2 [\text{CO}] [\text{NO}]^{-1}$
- b.  $\text{rate} = k [\text{NO}_2]^2 [\text{CO}]$
- c.  $\text{rate} = k [\text{NO}_2] [\text{CO}]$
- d.  $\text{rate} = k [\text{NO}_3] [\text{CO}]^2$
- e.  $\text{rate} = k [\text{NO}_2]^2$

5. At 25 °C, the equilibrium constant for the autoionization of water is  $K_w = 1.0 \times 10^{-14}$ . At 50 °C,  $K_w = 5.5 \times 10^{-14}$ . Determine  $\Delta H$  for the autoionization of water.

- a.  $-55 \text{ kJ mol}^{-1}$
- b.  $710 \text{ J mol}^{-1}$
- c.  $55 \text{ kJ mol}^{-1}$
- d.  $750 \text{ J mol}^{-1}$
- e.  $-14 \text{ kJ mol}^{-1}$

6. At 500 °C,  $K_c$  for the reaction:  $N_2 + 3H_2 \rightleftharpoons 2NH_3$  is  $K_c = 6.0 \times 10^{-2}$ . The numerical value of  $K_p$  at this temperature is:
- a.  $1.5 \times 10^{-5}$
  - b.  $2.5 \times 10^{-5}$
  - c.  $1.1 \times 10^{-9}$
  - d.  $2.0 \times 10^{-5}$
  - e.  $1.5 \times 10^{-9}$
7. An evacuated 2.0 L flask was filled with 0.10 moles  $SO_3(g)$  and 0.20 moles of  $NO(g)$ . The following reaction was allowed to come to equilibrium:  $SO_3(g) + NO(g) \rightleftharpoons SO_2(g) + NO_2(g)$   
If  $K_c$  for this reaction is 0.50, the equilibrium concentration of  $SO_2(g)$  is:
- a. 0.056 M
  - b. 0.021 M
  - c. 0.047 M
  - d. 0.035 M
  - e. 0.028 M
8. Colgate PreviDent™ toothpaste contains 1.0 mass percent NaF. Given that HF is a weak acid with  $K_a = 6.8 \times 10^{-4}$ , what mass of PreviDent™ toothpaste would have to be dissolved in pure water to give a total volume of 1.00 L of a solution with pH = 7.9?
- The logo for Colgate PreviDent toothpaste, featuring the word "Colgate" in white on a red background and "PreviDent" in white on a teal background.
- a. 210 g
  - b. 2.0 g
  - c. 4.9 g
  - d. 82 g
  - e. 180 g





9. In which of the following solutions would you expect LiF to be the most soluble given that  $K_{sp}$  for LiF is  $1.7 \times 10^{-3}$ ?
- a. 100.0 mL of pure water
  - b. 100.0 mL of 1.0 M HCl
  - c. 100.0 mL of 1.0 M  $\text{CH}_3\text{CO}_2\text{H}$
  - d. 100.0 mL of 1.0 M  $\text{NH}_3$
  - e. 100.0 mL of 1.0 M NaOH
10. What is the resulting pH when 50.00 mL of 0.10 M HCl is added to a buffer prepared by combining 0.100 moles of acetic acid and 0.110 moles of sodium acetate with 500 mL of pure water?  $K_a$  for acetic acid is  $1.8 \times 10^{-5}$ .
- a. 4.74
  - b. 4.83
  - c. 4.60
  - d. 5.12
  - e. 4.42

Topics in this question were not covered this year.

12.  $\text{CaSO}_4$  has  $K_{\text{sp}} = 2.4 \times 10^{-5}$  while  $\text{BaSO}_4$  has  $K_{\text{sp}} = 1.1 \times 10^{-10}$ . Determine the equilibrium constant,  $K_{\text{c}}$ , for the reaction:
- $$\text{CaSO}_4(s) + \text{Ba}^{2+}(aq) \rightleftharpoons \text{Ca}^{2+}(aq) + \text{BaSO}_4(s)$$

- a.  $2.2 \times 10^5$
- b.  $2.4 \times 10^{-5}$
- c.  $1.1 \times 10^{-10}$
- d.  $4.4 \times 10^{-6}$
- e.  $2.64 \times 10^{-15}$

**Standard potentials at 25 °C which may be used in Questions 13 to 15 and 24**




13. Which species is the strongest oxidizing agent under standard conditions?


- a.  $\text{Pb}(s)$
- b.  $\text{IO}_4^-$
- c.  $\text{Pb}^{2+}$
- d.  $\text{IO}_3^-$
- e.  $\text{H}_2\text{O}$

14. Determine the equilibrium constant for the reaction between  $\text{IO}_4^-$  and  $\text{Fe}^{2+}$ .


- a.  $4.32 \times 10^{27}$
- b.  $1.00 \times 10^{12}$
- c.  $0.57 \times 10^{13}$
- d.  $1.00 \times 10^6$
- e.  $5.76 \times 10^{19}$

15. What is the potential of the following cell?



- a. 1.26 V
  - b. 1.52 V
  - c. 1.49 V
  - d. 1.54 V
  - e. 1.23 V
- 



19. An element with the outermost electron configuration



could be\_\_\_\_\_.


- a.  $^{82}\text{Pb}$
- b.  $^{40}\text{Zr}$
- c.  $^{42}\text{Mo}$
- d.  $^{34}\text{Se}$
- e.  $^{19}\text{K}$





**SECTION II: Fill-in-the-blanks and long answer questions. (Total value 30 marks)**Write your answers in ink in the space provided.**22. Fill-in-the-blanks.****10 marks**


- a. Nitroglycerine decomposes violently according to the chemical equation below. What mass of carbon dioxide gas is produced from the decomposition of 5.00 g  $\text{C}_3\text{H}_5(\text{NO}_3)_3$ ?



- b. If 5.15 g  $\text{Fe}(\text{NO}_3)_3$  is dissolved in enough water to make exactly 150.0 mL of solution, what is the molar concentration of nitrate ion?

- c. The rate constant at 366 K for a first-order reaction is  $7.7 \times 10^{-3} \text{ s}^{-1}$  and the activation energy is 15.9 kJ/mol. What is the value of the frequency factor,  $A$ , in the Arrhenius equation?

- e. Assign formal charges to all atoms of the thiocyanate ion,  $\text{SCN}^-$ . Identify the most favored structure.



**8 marks**

b. After the addition of 10 mL of titrant

c. At the equivalence point

d. After the addition of 40 mL of titrant

**24. Long answer question (For full marks, you must show all work).****6 marks**

The standard electrode potential for the  $\text{Pb(s)}/\text{Pb}^{2+}$  redox couple is  $-0.126\text{ V}$ . Sketch the cell in which this potential was measured.

**Identify:**

- a. All concentrations/pressures
- b. The cathode and the anode
- c. The direction in which cations move in the salt bridge
- d. The direction in which electrons would move if a light bulb were connected across the cell.

25. For the species given below, draw Lewis (2D) and VSEPR (3D) structures and identify electron pair geometries, molecular geometries, bond angles around the central atom and molecular polarity. Enter your answers in the appropriate boxes:

<b>6 Marks</b>
----------------

Species	$\text{NCl}_3$	$\text{XeOF}_4$	$\text{SF}_4$
Lewis structure (2D) (Show all valence electrons)			
VSEPR structure (3D)			
Electro pair geometry			
Molecular Geometry			
Bond angles			
Polar or Non-polar			

\*\*\*\*\* END OF WRITTEN ANSWER SECTION \*\*\*\*\*

# Data Sheet – CHEM 209

## Periodic Table

Periodic Table																		18 8A
1 1A <div>H 1.008</div>	2 2A											13 3A	14 4A	15 5A	16 6A	17 7A	2 He 4.003	
3 Li 6.941	4 Be 9.012											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18	
11 Na 22.99	12 Mg 24.31	3	4	5	6	7	8	9	10	11	12	13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95	
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.88	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.38	31 Ga 69.72	32 Ge 72.59	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80	
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc (98)	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3	
55 Cs 132.9	56 Ba 137.3	57* La 138.9	72 Hf 178.5	73 Ta 180.9	74 W 183.9	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po (209)	85 At (210)	86 Rn (222)	
87 Fr (223)	88 Ra 226.0	89** Ac (227)	104 Rf (261)	105 Ha (262)	106 Sg (263)	107 Ns (262)	108 Hs (265)	109 Mt (266)	110 Uun (269)	111 Uuu (272)								

### Lanthanides \*

58 <b>Ce</b> 140.1	59 <b>Pr</b> 140.9	60 <b>Nd</b> 144.2	61 <b>Pm</b> (145)	62 <b>Sm</b> 150.4	63 <b>Eu</b> 152.0	64 <b>Gd</b> 157.3	65 <b>Tb</b> 158.9	66 <b>Dy</b> 162.5	67 <b>Ho</b> 164.9	68 <b>Er</b> 167.3	69 <b>Tm</b> 168.9	70 <b>Yb</b> 173.0	71 <b>Lu</b> 175.0
90 <b>Th</b> 232.0	91 <b>Pa</b> 231.0	92 <b>U</b> 238.0	93 <b>Np</b> 237.0	94 <b>Pu</b> (244)	95 <b>Am</b> (243)	96 <b>Cm</b> (247)	97 <b>Bk</b> (247)	98 <b>Cf</b> (251)	99 <b>Es</b> (252)	100 <b>Fm</b> (257)	101 <b>Md</b> (258)	102 <b>No</b> (259)	103 <b>Lr</b> (260)

### Actinides \*\*

Strong Acids: HCl, HBr, HI, HNO<sub>3</sub>, H<sub>2</sub>SO<sub>4</sub>, HClO<sub>4</sub>

Strong Bases: Hydroxides of Group 1A (Li to Cs) and Group 2A (Ca, Sr, Ba)

#### Constants:

Gas constant,  $R = 0.08205 \text{ L atm mol}^{-1} \text{ K}^{-1}$   
 $= 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$

Avogadro's number:  $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

Faraday:  $F = 96,485 \text{ C / mol electrons}$

Planck's constant  $h = 6.626 \times 10^{-34} \text{ Js}$

Speed of light,  $c = 2.998 \times 10^8 \text{ m/s}$

Rydberg constant,  $R = 1.09678 \times 10^{-7} \text{ m}^{-1}$

Factoring constant,  $R_H = Rhc = 2.18 \times 10^{-18} \text{ J}$

#### Conversion factors:

$1 \text{ J} = 1 \text{ kg m}^2 \text{ s}^{-2}$        $1 \text{ Pa} = 1 \text{ kg m s}^{-2}$

$T \text{ K} = T \text{ }^\circ\text{C} + 273.15$

$1 \text{ L atm} = 101.3 \text{ J}$

$1 \text{ atm} = 760.0 \text{ torr} = 101.3 \text{ kPa} = 760.0 \text{ mm Hg} = 1.013 \text{ bar}$

$1 \text{ L} = 10^{-3} \text{ m}^3$

$1 \text{ C} = 1 \text{ J / V}$        $1 \text{ A} = 1 \text{ C s}^{-1}$

STP conditions:  $0 \text{ }^\circ\text{C}$ ,  $1 \text{ atm}$

Electrochemical standard state:  $1 \text{ atm}$ ,  $1 \text{ M}$ ,  $25 \text{ }^\circ\text{C}$

$$[A]_t = -kt + [A]_0 \quad \ln [A]_t = -kt + \ln [A]_0 \quad \frac{1}{[A]_t} = kt + \frac{1}{[A]_0} \quad \ln \left( \frac{[A]_0}{[A]_t} \right) = kt$$

$$t_{1/2} = \frac{[A]_0}{2k} \quad t_{1/2} = \frac{0.693}{k} \quad t_{1/2} = \frac{1}{k[A]_0} \quad k = A e^{\frac{-E_a}{RT}} \quad \ln \left( \frac{K_2}{K_1} \right) = \frac{\Delta H}{R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right)$$

$$\ln \left( \frac{k_2}{k_1} \right) = \frac{E_a}{R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right) \quad PV = nRT \quad K_p = K_c(RT)^{\Delta n} \quad ax^2 + bx + c = 0$$

$$\text{pH} = -\log[\text{H}^+] \quad K_w = K_a K_b \quad K_{sp} = I / K_d \quad K_f = I / K_d \quad x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\text{pH} = \text{p}K_a + \log \left( \frac{[\text{cong. base}]}{[\text{cong. acid}]} \right) \quad \text{or} \quad \text{pOH} = \text{p}K_b + \log \left( \frac{[\text{cong. acid}]}{[\text{cong. base}]} \right)$$

$$E^\circ = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}} \quad E = E^\circ - \frac{0.0592}{n_e} \log Q \quad E^\circ = \frac{0.0592}{n_e} \log K \quad \text{or} \quad nFE^\circ = RT \ln K$$

$$q = It \quad q = n_e F \quad c = \lambda \nu \quad E = h\nu \quad E = mc^2 \quad \frac{1}{\lambda} = R \left( \frac{1}{n_1^2} - \frac{1}{n_2^2} \right) \quad \Delta E = -R_H \left( \frac{Z^2}{n_f^2} - \frac{Z^2}{n_i^2} \right)$$

$$E = -R_H \left( \frac{Z}{n} \right)^2 \quad \text{or} \quad E_n = -\frac{Rhc}{n^2} \quad \text{for single electron species}$$