

Electricity and Magnetism

- Physics 259 – L02
 - Lecture 19



UNIVERSITY OF
CALGARY

Chapter 23.3-4



Last time

- Chapter 23.2



This time



- Chapter 23. and 23.4



23-5: Electric field of a plane of charge

Nonconduction infinite sheet

$$\Phi_e = \oint \vec{E} \cdot d\vec{A} = \frac{Q_{in}}{\epsilon_0}$$



$$E_{plane} = \frac{\sigma}{2\epsilon_0}$$


Q2) Two very thin infinite sheets are uniformly charged with surface charge densities -2η and $+5\eta$ as indicated in the figure. What is the magnitude and direction of the electric field at point P located between the sheets? (note the direction of $+x$ in the figure)

a) $-3\eta/2\epsilon_0$

b) $+3\eta/2\epsilon_0$

c) $-7\eta/2\epsilon_0$

d) $+7\eta/2\epsilon_0$



$$E = \frac{\sigma}{2\epsilon_0}$$






$$\vec{E}_1 = -\frac{2\sigma}{2\epsilon_0} \hat{i}$$

$$\vec{E}_2 = -\frac{5\sigma}{2\epsilon_0} \hat{i}$$

$$\vec{E}_{net} = -\frac{7\sigma}{2\epsilon_0} \hat{i}$$

Two conducting Plates



$$E = \frac{2\sigma_1}{\epsilon_0} = \frac{\sigma}{\epsilon_0}$$


TopHat Question 1

$$Q = \lambda L$$

4 surfaces are coaxial with an infinitely long line of charge with a uniform linear charge density $= \lambda$. Choose all the surfaces through which $\Phi_E = \lambda L / \epsilon_0$

$$\Phi_E = \frac{q_{\text{enc}}}{\epsilon_0} = \frac{\lambda L}{\epsilon_0}$$

$$\Phi_E = \frac{q_{\text{enc}}}{\epsilon_0} = \frac{\lambda L}{\epsilon_0}$$



- A) I only
- B) I and II only
- C) I and III only
- ☒ D) I, II, and III only
- E) All four.


$$Q = \lambda L$$

$$\Phi_E = \frac{q}{\epsilon_0} = \frac{\lambda L}{\epsilon_0}$$

$$\Phi_E = \oint \vec{E} \cdot d\vec{A} = \frac{q_{\text{enc}}}{\epsilon_0}$$


$$\Phi_E = \frac{q_{\text{enc}}}{\epsilon_0}$$

$$\sigma = \frac{Q}{A} \rightarrow Q = \sigma A$$




$$\Phi_e = \frac{q_{enc}}{\epsilon_0} = \frac{\sigma A}{\epsilon_0}$$


What is the field inside the slab?



The slab has thickness L , we have to choose a Gaussian surface with the same symmetries as the slab: choose a box whose centre coincides with the centre of the slab.



$$\rho = \frac{Q}{V}$$



$$\oint \vec{E} \cdot d\vec{A} = \int_{\text{top}} \vec{E} \cdot d\vec{A} + \int_{\text{bottom}} \vec{E} \cdot d\vec{A} = \frac{q_{\text{enc}}}{\epsilon_0}$$

$$E \int_{\text{top}} dA + E \int_{\text{bottom}} dA = \frac{q_{\text{enc}}}{\epsilon_0}$$

$$\rho = \frac{q}{V}$$

$$q_{\text{enc}} = \rho V$$

$$V = A \times 2h$$

$$\rightarrow \underline{2EA = \frac{q_{\text{enc}}}{\epsilon_0}}$$

$$q_{\text{enc}} = \rho V = \underline{\underline{\rho A(2h)}}$$

$$\rightarrow \cancel{2EA} = \frac{\cancel{\rho A(2h)}}{\epsilon_0}$$

$$\boxed{E = \frac{\rho h}{\epsilon_0}} \quad \leftarrow$$

What about cylinder?


Homework 😊

Read appendix 1-chapter 23 posted on D2L.

TopHat Question

2

Two infinite planes are uniformly charged with the same charge per unit area σ (or η in your textbook). If one plane only were present, the E-field magnitude due to the one plane would be E . With both planes in place, the E-field magnitude in region B has magnitude:




A: zero

B: E

C: $2E$

D: depends on exact position.




$$E = \frac{\sigma}{2\epsilon_0}$$

Conductor
insulator

Conductor
insulator

Two infinite planes are uniformly charged with the same charge per unit area σ (~~or η in your textbook~~). If one plane only were present, the E-field magnitude due to the **one** plane would be E . With both planes in place, the E-field magnitude in region B has magnitude:



A: zero

B: E


C: $2E$

D: depends on exact position.

TopHat Question 3

Negative charges are uniformly distributed on the surface of an **insulating sphere** with charge density σ . Two additional point charges $-q$ are placed outside of the spherical charge distribution.

What is the magnitude of the E field at point P?



A: 0

B: non-zero

C: Not enough info given


Negative charges are uniformly distributed on the surface of an insulating sphere with charge density σ . Two additional point charges $-q$ are placed outside of the spherical charge distribution.

What is the magnitude of the E field at point P?

A: 0

B: non-zero


C: Not enough info given



The two outside charges $-q$ break the spherical symmetry.


If the 2 outside charges were **removed**, the E field at point P would indeed be ZERO.

In order to find the E field at point P, we draw a Gaussian surface that goes through point P. If the 2 outside charges were removed (*just for a moment*), the spherical charge distribution would insure that the E field = 0 at point P (also anywhere inside the spherical charge distribution). THEN, once we **re-insert** the 2 outside charges $-q$, they would produce their own E field at point P...therefore resulting in a net E field at point P.



TopHat Question 4

What is the charge of the insulating wire (the wire is reshaped to form a rectangle) with charge density $-\lambda$?



- A) $L^2\lambda$
- B) $-L^2\lambda$
- C) $4L\lambda$
- D) $-4L\lambda$

1

2

3

4


5

6

7

Field of a line charge


Consider an infinitely long, positively charged rod of linear charge density λ . How large is the flux through side A of the box? Suppose the values for l , a and λ are given.



Field of a line charge


- Consider an infinitely long, positively charged rod of linear charge density λ . How large is the flux through side A of the box? Suppose the values for l , a and λ are given.
- Gauss' law tells us that the total electric flux only depends on the enclosed charge – not the shape of the (closed) Gaussian surface:

$$\Phi_{\text{tot}} = Q_{\text{encl}}/\epsilon_0 = \lambda l/\epsilon_0$$




Field of a line charge

- Consider an infinitely long, positively charged rod of linear charge density λ . How large is the flux through side A of the box? Suppose the values for l , a and λ are given.
- Gauss' law tells us that the total electric flux only depends on the enclosed charge – not the shape of the (closed) Gaussian surface:
$$\Phi_{\text{tot}} = Q_{\text{encl}}/\epsilon_0 = \lambda l/\epsilon_0$$
- The total flux must be equally partitioned into flux through the four surfaces whose area vectors are parallel to the electric field.
Hence, $\Phi_A = \lambda l/4\epsilon_0$




Practice question:

Find the field at point p at the centre of a ring of charge composed of two oppositely charged half rings.



Find \vec{E} at P .
Radius, R

1. Cut the distribution into a bunch of tiny pieces each with charge dQ
2. Look for a symmetry \rightarrow can use $1/4$ circle (arc)
Find \vec{E} and multiply it by 4.
3. Calculate the magnitude of E-field due to ARBITRARY piece of charge dq .



4. Decompose field into components,

$$\frac{dE_x}{dE} = \sin \theta \quad \frac{dE_y}{dE} = \cos \theta$$

$$dE_x = dE \sin \theta$$

$$= \frac{1}{4\pi\epsilon_0} \frac{dq}{R^2} \cdot \sin \theta$$

$$dE_y = \frac{1}{4\pi\epsilon_0} \frac{dq}{R^2} \cos \theta$$

\Rightarrow not needed
due to symmetry

5. For each non-zero component, sum up all pieces dQ by integrating over the whole charge distribution

$$\int dE_x = \frac{1}{4\pi\epsilon_0} \int \frac{dq}{R^2} \sin \theta$$

6. Express dQ in terms of a variable to be integrated over using linear / surface / volume density

$$\int dE_x = \frac{1}{4\pi\epsilon_0} \int \frac{\lambda dL}{R^2} \cdot \sin\theta = \frac{1}{4\pi\epsilon_0} \int \frac{\lambda \cdot \cancel{R} \sin\theta d\theta}{\cancel{R^2}}$$

$$E_x = \frac{\lambda}{4\pi\epsilon_0 R} \int_0^{\pi/2} \sin\theta d\theta$$

$dL = R d\theta$
 $\pi/2$ - quarter of the circle

$$\int \sin x dx = -\cos x$$

$$E_x = \frac{\lambda}{4\pi\epsilon_0 R} [-\cos\theta]_0^{\pi/2} = \frac{\lambda}{4\pi\epsilon_0 R} [-\overset{0}{\cos(\frac{\pi}{2})} - (-\overset{1}{\cos 0^\circ})]$$



$$= \frac{\lambda}{4\pi\epsilon_0 R}$$

Total electric field

$$E_{\text{net}} = 4E_x = \frac{\cancel{4}\lambda}{4\pi\epsilon_0 R} = \frac{\lambda}{4\pi\epsilon_0 R^2}$$

$$\vec{E}_{\text{net}} = \frac{\lambda}{4\pi\epsilon_0 R} (-\hat{i})$$

Superposition





$$E_{\text{total}} = E_1 + E_2$$

Exercise: Coaxial Cable

Study appendix 1-Chapter 23

Assume there is a charge $+Q$ on the centre core and $-Q$ on the **metallic shield**. (Ignore the dielectric insulator and plastic jacket.)

Find the electric field outside the metallic shield (E_2) and just outside the central core (E_1).



This section we talked about:
Chapter 23 & Midterm Review

See you on Friday

