

Electricity and Magnetism

- Physics 259 – L02
 - Lecture 5

Section 21.1-3



Last time

- Charges and Force Between Charges
- Conductors and Insulators
- Van De Graaff Generator Experiment
- Solve Class Activity Question
- Coulomb's Law



This time

- Examples for Coulomb's law

Coulomb's Law

$$F_{1\text{ on } 2} = F_{2\text{ on } 1} = K \frac{|q_1||q_2|}{r^2}$$

K = electrostatic constant

$$K = 8.99 \times 10^9 \frac{N \cdot m^2}{C^2}$$

$$F_{1\text{ on } 2} = F_{2\text{ on } 1} = \frac{1}{4\pi\epsilon_0} \frac{|q_1||q_2|}{r^2}$$

ϵ_0 = permittivity of free space

$$\epsilon_0 = \frac{1}{4\pi K} = 8.85 \times 10^{-12} \frac{C^2}{N \cdot m^2}$$

Example #1: Two point charges

$$q_1 = +2 \text{ C}, q_2 = +5 \text{ C}, r = 10 \text{ m}$$

$$\vec{F}_{1 \text{ on } 2} = k_e \frac{|q_1||q_2|}{r^2} \hat{r}_{12}$$

$$= (8.99 \times 10^9 \text{ Nm}^2/\text{C}^2) \frac{(2\text{C})(5\text{C})}{(10 \text{ m})^2} \hat{r}_{12}$$



$$= (8.99 \times 10^8 \text{ N}) \hat{r}_{12}$$

$$= 8.99 \times 10^8 (\cos 30^\circ \hat{i} + \sin 30^\circ \hat{j}) \text{ N}$$

$$\vec{F}_{2 \text{ on } 1} = k_e \frac{|q_1||q_2|}{r^2} \hat{r}_{21}$$


$$= (8.99 \times 10^8 \text{ N}) \hat{r}_{21}$$

$$= 8.99 \times 10^8 (-\cos 30^\circ \hat{i} - \sin 30^\circ \hat{j}) \text{ N}$$



Force is fully defined.

Example #2 (last section's homework): Three point Charges



Total electric force on q_1 ?

$$|\vec{F}_{31}| = 8.99 \times 10^{-9} \text{ N}$$

$$|\vec{F}_{21}| = 2.25 \times 10^{-9} \text{ N}$$


$$\vec{F}_{net} = (-6.74 \times 10^{-9} \text{ N}) \hat{i}$$

Example #3: Building blocks of electric force

$$\vec{F} = \frac{KQq}{r^2} \hat{r}$$





$+Q$



● = positive charge q at the position indicated


Example #3: Building blocks of electric force

$$\vec{F} = \frac{-KQq}{r^2} \hat{r}$$



● = positive charge q at the position indicated


Example #4: Two point charges (Electric Dipole)



$$\left| \vec{F}_{net} \right| = 2 \left(\frac{KqQ}{\left(x^2 + (d/2)^2 \right)^{3/2}} \right) \sin \theta = \frac{KqQd}{\left(x^2 + (d/2)^2 \right)^{3/2}}$$


Superposition with Building Blocks

The vector represents the magnitude and direction of the electric force on the charge q **at that point**. It comes from superposition of the individual forces from $+Q$ and $-Q$.



Superposition with Building Blocks

The vector represents the magnitude and direction of the electric force on the charge q **at that point**. It comes from superposition of the individual forces from $+Q$ and $+Q$.



This section we talked about:

Chapter 21.1-3: Examples

See you on Wednesday

