

Monday Jan 23, 2017

Labatorials start this week

- Check your schedule
- Print write-up
- Prepare
- Take advantage of the small group (24 not 200 students in the room) and group work (3-4)
- Checkpoints test your understanding
- Come on time (late arrival = group working without you = mark deduction)

Last time


- The electric field: conceptually difficult but much more useful
- TopHat questions related to direction of net E-field
- Finishing up Coloumb's Law: Group Activity

This time

- More on electric fields: how to calculate them
- Example: electric field of a dipole.

Electric field of point charges


$$\vec{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \hat{r}$$



© 2013 Pearson Education, Inc.

+ charge: \vec{E} points away from Q

$$\vec{E} = \frac{-1}{4\pi\epsilon_0} \frac{Q}{r^2} \hat{r}$$



© 2013 Pearson Education, Inc.


– charge: \vec{E} points toward $-Q$

Electric Field Vectors

The vector represents the magnitude and direction of the electric field **at that point**.

But \vec{E} is not a spatial quantity that stretches from one end of the arrow to the other.

Instead, think of \vec{E} as a spatial quantity at every point with a direction at that point given by the arrow.




Copyright © 2008 Pearson Education, Inc., publishing as Pearson Addison-Wesley.

Electric Field building blocks

The electric field around a **point charge, q** , is given by

$$\vec{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{r}$$

\hat{r} is a **unit vector** that always points **away from q** .




We can use this with superposition to find the electric field of more complicated objects.

TopHat Questions


TopHat Question

What is the direction of the electric field at the point indicated?




TopHat Question

What is the direction of the electric field at the point indicated?




TopHat Question

What is the direction of the electric field at the point indicated?



TopHat Question

An electric dipole: if the electric field strength at point P is $E = 6068 \text{ N/C}$, what is the charge q ?



For single point charge:

$$E = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2}$$

$$\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m}^2$$

A. 0.36 nC


C. 0.22 nC

B. 0.27 nC

D. 0.13 nC

TopHat Question Feedback

An electric dipole: if the electric field strength at point P is $E = 6068 \text{ N/C}$, what is the charge q ?



For single point charge:

$$E = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2}$$


$$\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m}^2$$

$$E_+ = \frac{1}{4\pi\epsilon_0} \frac{q}{r_+^2} \quad E_- = -\frac{1}{4\pi\epsilon_0} \frac{q}{(2r_+)^2}$$

$$E_{net} = \frac{1}{4\pi\epsilon_0} q \left(\frac{1}{r_+^2} - \frac{1}{4r_+^2} \right) = \frac{1}{4\pi\epsilon_0} \frac{3q}{4r_+^2}$$


$$q = 4\pi\epsilon_0 \frac{4r_+^2 E_{net}}{3}$$

Cases we've already seen



TopHat Question

An infinitely long wire: if the electric field strength at point P is $E = 3670 \text{ N/C}$, how much charge is contained in a 0.500 m length of the wire?



$$E_{\text{wire}} = \frac{1}{4\pi\epsilon_0} \frac{2\lambda}{r}$$

$$\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m}^2$$

A. 18.37 nC


C. 61.23 nC

B. 30.62 nC

D. 9.19 nC

TopHat Question Feedback

An infinitely long wire: if the electric field strength at point P is $E = 3670 \text{ N/C}$, how much charge is contained in a 0.500 m length of the wire?



$$\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m}^2$$

$$E_{\text{wire}} = \frac{1}{4\pi\epsilon_0} \frac{2\lambda}{r}$$


$$E_{\text{wire}} = \frac{1}{2\pi\epsilon_0} \frac{\lambda}{r}$$

$$\lambda = 2\pi\epsilon_0 E_{\text{wire}} r$$

$$\lambda = \frac{\Delta Q}{\Delta L}$$

$$Q = \lambda L$$

Charged disk



$$E = \frac{\sigma}{2\epsilon_0} \left[1 - \frac{z}{\sqrt{z^2 + R^2}} \right]$$