

Last time

- Top hat questions
- Electric field pattern due to point charges
- Electric field pattern for a dipole
- Electric field due to a charged 90° arc of radius R at its center, constant charge density
- Electric field due to a charged semi-circle of radius R at its center using superposition principle
- Electric field due to a charged circle of radius R at its center using superposition principle

This time

- Electric field of a thin ring on a point along the axis of the ring
- Activity #3

Infinitely thin ring of charge

$$\lambda = \frac{|Q|}{2\pi R} \quad \text{Uniformly charged ring}$$


ds Infinitesimal arc length

dQ Infinitesimal charge on ds

$$dQ = \lambda ds$$

$$r^2 = (R^2 + h^2)$$

$$\cos \theta = \frac{h}{r} = \frac{h}{(R^2 + h^2)^{1/2}}$$




Copyright © 2014 John Wiley & Sons, Inc. All rights reserved.

halliday_10e_fig_22_11


$d\vec{E}$ is the electric field generated by dQ at point P.

Do this for a large number
pieces around the ring.

$d\vec{E}$ traces a cone.




Now consider the cone and project out $d\vec{E}$ in the xy-plane.



$$\lambda = \frac{Q}{2\pi R}, Q = 2\pi\lambda R$$

$$r^2 = (R^2 + h^2)$$

$$\cos \theta = \frac{h}{(R^2 + h^2)^{1/2}}$$



Copyright © 2014 John Wiley & Sons, Inc. All rights reserved.

halliday_10e_fig_22_11

$$dE_z = dE \cos \theta$$

$$dE_z = dE \cos \theta = k_e \frac{dQ}{r^2} \cos \theta = k_e \frac{dQ}{(R^2 + h^2)} \frac{h}{(R^2 + h^2)^{1/2}} = k_e \frac{h\lambda ds}{(R^2 + h^2)^{3/2}}$$

$$E_z = \int_{\text{Around the circle}} k_e \frac{h\lambda ds}{(R^2 + h^2)^{3/2}} = k_e \frac{h\lambda}{(R^2 + h^2)^{3/2}} \int_{\text{Around the circle}} ds$$

$$E_z = k_e \frac{h\lambda}{(R^2 + h^2)^{3/2}} 2\pi R = k_e \frac{hQ}{(R^2 + h^2)^{3/2}} \quad \vec{E} = k_e \frac{hQ}{(R^2 + h^2)^{3/2}} \hat{k}$$

$$\vec{E} = k_e \frac{hQ}{(R^2 + h^2)^{3/2}} \hat{k}$$

Move down to the center of the ring.


$$h \rightarrow 0? \Rightarrow E_z = 0$$

Move very far away from the ring.

$$h \gg R? \quad (R^2 + h^2) \approx h^2$$

$$E_z = k_e \frac{hQ}{(R^2 + h^2)^{3/2}} \approx k \frac{hQ}{h^3} = k \frac{Q}{h^2}$$


Appears as a point charge Q located at the center of the disk.



Copyright © 2014 John Wiley & Sons, Inc. All rights reserved.

halliday_10e_fig_22_11

What is the change in the area of a circle if its radius r is increased by an infinitesimal amount (dr)?



$$\Delta A = \pi(r + \Delta r)^2 - \pi r^2 = \pi r^2 + 2\pi r\Delta r + \Delta r^2 - \pi r^2$$

$$\Delta A = 2\pi r\Delta r + \Delta r^2 \approx 2\pi r\Delta r \quad \text{When can we make the } \approx \text{ sign } =?$$

$$\lim_{\Delta r \rightarrow 0} \Delta r = dr$$

$$\lim_{\Delta A \rightarrow 0} \Delta A = dA$$

$$dA = 2\pi r dr$$

