

Basic Definitions

Definition. (Inner Product) An *inner product* on V is a function that takes each ordered pair (x, y) of elements of V to a number $\langle x, y \rangle \in \mathbb{F}$ and, $\forall x, y, z \in V, \lambda \in \mathbb{F}$, has the following properties:

Positivity	$\langle y, y \rangle \geq 0$
Definiteness	$\langle y, y \rangle = 0 \iff y = 0$
Additivity	$\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$
Homogeneity	$\langle \lambda x, y \rangle = \lambda \langle x, y \rangle$
Conjugate Symmetry	$\langle x, y \rangle = \overline{\langle y, x \rangle} = \langle y, x \rangle^*$

Definition. (Norm) Given a vector space V over $\mathbb{F} = \mathbb{R}$ or \mathbb{C} , a *norm* is a mapping $\| \cdot \| : V \rightarrow [0, \infty)$ for which

- (i) $\|f\| \geq 0$ and $\|f\| = 0 \iff f = 0$
- (ii) $\|\lambda f\| = |\lambda| \|f\|$
- (iii) $\|f + g\| \leq \|f\| + \|g\|$

for all $f, g \in V, \lambda \in \mathbb{F}$. If V is an inner product space, there is a trivial norm:

$\|f\| = \sqrt{\langle f, f \rangle}$

Normed Spaces

Theorem. (Triangle Inequality)

$\|x + a\| \leq \|x\| + \|a\|$

Theorem. (Reverse Triangle Inequality)

$|\|x\| - \|a\|| \leq \|x - a\|$

Example. (Example Normed Spaces)

$V = \mathbb{R}^n, \quad \|x\|_2 = \sqrt{\sum_{j=1}^n x_j^2}$
 $V = \mathbb{R}^2, \quad \|x\|_1 = |x_1| + |x_2|$
 $V = \mathbb{R}^2, \quad \|x\|_\infty = \max\{|x_1|, |x_2|\}$
 $V = C([0, 1], \mathbb{R}), \quad \|f\|_p = \left(\int_0^1 |f|^p\right)^{\frac{1}{p}}$

Definition. (Open and Closed Balls) With $(X, \| \cdot \|)$ a normed linear space, with $x_0 \in X$, and $r > 0$:

- Open Ball** With centre x_0 and radius r as $\mathcal{B}(x_0, r) = \{x \in X : \|x - x_0\| < r\}$
- Closed Ball** With centre x_0 and radius r as $\mathcal{B}[x_0, r] = \{x \in X : \|x - x_0\| \leq r\}$
- Sphere** With centre x_0 and radius r as $\mathcal{S}(x_0, r) = \{x \in X : \|x - x_0\| = r\}$

Inner Product Spaces

Example. (Inner-Product Space Examples)

- ℓ^p space:

$\ell^2 = \ell^2(\mathbb{N}, \mathbb{C}) = \left\{ \{c_k\}_{k=1}^\infty : \sum_{k=1}^\infty |c_k|^2 < \infty \right\}$

And hence for ℓ^2 (the only ℓ^p space that is a Hilbert space)

$\langle x, y \rangle_{\ell^2} = \sum_n x_n \overline{y_n}$

- $L^2(\mathbb{R})$ space:

$\langle x, y \rangle_{L^2} = \int_{-\infty}^\infty f(x) \overline{g(x)} dx$

Theorem. (Cauchy-Schwarz Inequality) Given that V is an inner-product space over \mathbb{F} , then for all $f, g \in V$:

$|\langle f, g \rangle| \leq \|f\| \|g\|$

When $\{f, g\}$ is a linearly dependent set, we have equality.

Sequences and Convergence

Definition. (Cauchy Sequence) A sequence $\{a_n\}$ of real numbers is a *Cauchy Sequence* if

$\forall \varepsilon > 0, \exists N > 0 : m, n > N \implies \|a_m - a_n\| < \varepsilon$

Given that $n > N \implies \|x_n - x\| < \varepsilon$, then $\lim_{n \rightarrow \infty} x_n = x$.

Definition. (Geometric Series) A geometric series (aka “geometric progression”) has sequence elements of the form $a_n = ar^n$, where r is the “*common ratio*”. The partial sum is given by:

$s_k = a \left(\frac{1 - r^k}{1 - r} \right)$

and so, given that $|r| < 1$, the partial sum limits to:

$\lim_{n \rightarrow \infty} \sum_{k=1}^n ar^k = \frac{a}{1 - r}$

Operator Theory

Proposition. (Operator Continuity) Let X and Y be normed linear spaces and $T : X \rightarrow Y$ be a linear mapping. Then, the following are equivalent:

- (i) T is continuous at every point in X
- (ii) T is continuous at 0
- (iii) T is bounded

Hilbert Spaces

Theorem. (Riesz Representation Theorem) Given a Hilbert space \mathcal{H} over \mathbb{C} and a linear functional $F : \mathcal{H} \rightarrow \mathbb{C}$, there exists a unique $y = y_f$, for $y_f \in \mathcal{H}$ such that

$f(x) = \langle x, y_f \rangle, \quad \forall x \in \mathcal{H}$

Definition. (Adjoint Operator) Let $T : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ be a bounded linear mapping between Hilbert spaces. The mapping $T^* : \mathcal{H}_2 \rightarrow \mathcal{H}_1$ is called the *adjoint* of the operator T and, for all $x \in \mathcal{H}_1, y \in \mathcal{H}_2$, is defined:

$\langle Tx, y \rangle_2 = \langle x, T^*y \rangle_1$

T is said to be *self-adjoint* if $T = T^*$.

Definition. (Unitary Operator) Let \mathcal{H} be a Hilbert space and $U \in \mathcal{B}(\mathcal{H})$. We say that U is *unitary* if U is invertible and $U^{-1} = U^*$. Equivalently, $UU^* = U^*U = I$.

Definition. (Bounded Linear Operator) Let $T : X \rightarrow Y$ be a bounded linear operator between normed linear spaces. The *norm* of T is defined to be:

$\|T\| = \inf\{M > 0 : \forall x \in X, \|Tx\| \leq M\|x\|\}$

Note that $\|Tx\| \leq \|T\|\|x\|$ and $\|T\|$ is the smallest M such that this is valid.

Proposition. Let $T : X \rightarrow Y$ be a bounded linear operator between normed linear spaces. Then

$\|T\| = \sup\{\|Tx\| : \|x\| = 1\}$

Definition. (Normal Operator) $T \in \mathcal{L}(V)$ is *normal* if $TT^* = T^*T$. Furthermore, T is normal if and only if $\|Tv\| = \|T^*v\|$.

Example. (Orthogonal Sets)

$\mathcal{H} = \mathbb{C}^N$ with $\left\{ f_j(k) = \frac{1}{\sqrt{N}} e^{2\pi i j k / N} \right\}_{j=0}^{N-1}$
 $\mathcal{H} = L^2\left(\left[-\frac{1}{2}, \frac{1}{2}\right]\right)$ with $\left\{ f_j(x) = e^{2\pi i j x} \right\}_{j=-\infty}^\infty$
 $\mathcal{H} = L^2(\mathbb{R})$ with $\left\{ f_j(x) = \chi_{[j, j+1]}(x) \right\}_{j=-\infty}^\infty$
 $\mathcal{H} = \ell^2(\mathbb{N})$ with $\{e_j(k) = \delta_{ij}\}_{j=1}^\infty$

Proposition. For $\{x_n\}$ an orthonormal set in \mathcal{H} :

$\sum_{n=1}^\infty |\langle x, x_n \rangle|^2 \leq \|x\|^2$

Spectral Theory

Theorem. (Normal Operator Eigenvectors) Suppose $T \in \mathcal{L}(V)$ is normal. Then, eigenvectors of T corresponding to distinct eigenvalues are orthogonal.

Theorem. (Real Spectral Theorem) Suppose $\mathbb{F} = \mathbb{R}$ and $T \in \mathcal{L}(V)$. Then, the following are equivalent:

- (i) T is self-adjoint.
- (ii) V has an orthonormal basis consisting of eigenvectors of T .
- (iii) T has a diagonal matrix with respect to some orthonormal basis of V .

Theorem. (Complex Spectral Theorem) Suppose $\mathbb{F} = \mathbb{C}$ and $T \in \mathcal{L}(V)$. Then, the following are equivalent:

- (i) T is normal.
- (ii) V has an orthonormal basis consisting of eigenvectors of T .
- (iii) T has a diagonal matrix with respect to some orthonormal basis of V .

Theorem. (Diagonalisation) A square $n \times n$ matrix, A , with entries in a field \mathbb{F} is called diagonalisable or non-defective if there exists an $n \times n$ invertible matrix (i.e. an element of the general linear group $\text{GL}_n(\mathbb{F})$), P , such that $P^{-1}AP$ is a diagonal matrix.

Given that Λ is diagonal matrix with eigenvalue entries of A on the main diagonal, then $AP = P\Lambda$ implies that $P = V$, the matrix whose columns are eigenvectors to A , corresponding to the eigenvalue in the same column of Λ . In this case:

$A = V\Lambda V^{-1}$

Note that one of the *spectral theorems* may guarantee the existence of Λ .

Function Spaces

Definition. (Lebesgue Spaces) Given two measurable functions f and g , we say that $f \sim g$ if $f = g$ *almost everywhere*.

For $1 \leq p < \infty$:

$L^p(\mathbb{R}) = \left\{ f : \mathbb{R} \rightarrow \mathbb{C} : \|f\|_p = \left(\int |f|^p \right)^{\frac{1}{p}} < \infty \right\} / \sim$

For $p = \infty$:

$L^\infty(\mathbb{R}) = \{f : \mathbb{R} \rightarrow \mathbb{C} : \|f\|_\infty = \text{ess sup } |f(x)| < \infty\} / \sim$

where

$\text{ess sup } |f| = \inf\{\alpha \in \mathbb{R} : \mu\{x \in \mathbb{R} : |f(x)| > \alpha\} = 0\}$

Inequalities

Theorem. (Arithmetic-Geometric Mean Inequality) For positive x and y :

$4xy \leq (x + y)^2$

Notice that by expanding the RHS:

$2xy \leq x^2 + y^2$

Trigonometry

Definition.

$\sin z = \frac{e^{iz} - e^{-iz}}{2i}$
 $\cos z = \frac{e^{iz} + e^{-iz}}{2}$
 $\sinh z = \frac{e^z - e^{-z}}{2}$
 $\cosh z = \frac{e^z + e^{-z}}{2}$

Definition. (Double Angle Formulae)

$\sin 2\theta = 2 \sin \theta \cos \theta$
 $\cos 2\theta = \cos^2 \theta - \sin^2 \theta$
 $= 2 \cos^2 \theta - 1$
 $= 1 - 2 \sin^2 \theta$
 $\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$

Definition. (Half Angle Formulae)

$\sin^2 \theta = \frac{1 - \cos 2\theta}{2}$
 $\sin\left(\frac{\theta}{2}\right) = \pm \sqrt{\frac{1 - \cos \theta}{2}}$
 $\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$
 $\cos\left(\frac{\theta}{2}\right) = \pm \sqrt{\frac{1 + \cos \theta}{2}}$
 $\tan\left(\frac{\theta}{2}\right) = \pm \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}}$

Definition. (Ptolemy's Identities (Difference Formulas))

$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$
 $\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$

Definition. (Trigonometric Integrals)

$\int \frac{1}{\sin ax} dx = -\frac{1}{a} \ln|\csc ax + \cot ax| + C$
 $\int \frac{1}{\cos ax} dx = -\frac{1}{a} \ln\left|\tan\left(\frac{1}{2}ax + \frac{1}{4}\pi\right)\right| + C$
 $\int \tan ax dx = -\frac{1}{a} \ln|\cos ax| + C = \frac{1}{a} \ln|\sec ax| + C$
 $\int \tan^2 ax dx = \tan x - x + C$
 $\int \sec ax dx = \frac{1}{a} \ln|\sec ax + \tan ax| + C$
 $= \frac{1}{a} \ln\left|\tan\left(\frac{1}{2}ax + \frac{1}{4}\pi\right)\right| + C$
 $= \frac{1}{a} \arctan(\sin ax) + C$
 $\int \frac{1}{x^2 + a^2} dx = \frac{1}{a} \arctan\left(\frac{x}{a}\right) + C$

Fourier Series

Definition. (Fourier Series) For $f \in \mathcal{H}$ and let $\{e_n\}_{n=-\infty}^\infty$ be an orthonormal basis in \mathcal{H} , the **Fourier series** of f is

$$f(x) = \sum_{n=1}^\infty \langle x, e_n \rangle e_n$$

Theorem. (Parseval Equality) With $f \in \mathcal{H}$ and $\{e_n\}_{n=1}^\infty$ an orthonormal basis for \mathcal{H} , then for $\mathcal{F} : \mathcal{H} \rightarrow \ell^2(\mathbb{N})$:

$$(\mathcal{F}f)_n = \langle f, e_n \rangle$$

is unitary. Furthermore, we have

$$\langle f, g \rangle_{\mathcal{H}} = \langle \mathcal{F}f, \mathcal{F}g \rangle_{\ell^2}$$

Theorem. (Plancherel Equality) With $f \in \mathcal{H}$ and $\{e_n\}_{n=1}^\infty$ an orthonormal basis for \mathcal{H} :

$$\|f\|_{\mathcal{H}}^2 = \sum_{n=1}^\infty |\langle f, e_n \rangle|^2 = \|\mathcal{F}f\|_{\ell^2}^2$$

L1 Fourier Transform

Definition. (L^1 Fourier Transform) For $f \in L^1(\mathbb{R})$, the **Fourier Transform** \hat{f} of f is defined:

$$\mathcal{F}f(\xi) = \hat{f}(\xi) = \int_{-\infty}^\infty f(t)e^{-2\pi i \xi t} \, dt \quad (\xi \in \mathbb{R})$$

- Proposition.** (Properties)
- $\|\hat{f}\|_1 \leq \|f\|_1$
 - $\hat{f} \in \mathcal{C}(\mathbb{R})$
 - $\lim_{|\xi| \rightarrow \infty} |\hat{f}(\xi)| = 0$
 - $\hat{f} \equiv 0 \iff f = 0$ almost everywhere

Theorem. (In the Schwarz Space) For $f \in \mathcal{S}(\mathbb{R})$:

- $\mathcal{F} : \mathcal{S} \rightarrow \mathcal{S}$
- If $f \in \mathcal{S}$ then, for all $t \in \mathbb{R}$:

$$f(t) = \int_{-\infty}^\infty \hat{f}(\xi)e^{2\pi i \xi t} \, d\xi$$

- If $f, g \in \mathcal{S}$ then $\langle f, g \rangle = \langle \hat{f}, \hat{g} \rangle$

Remark. (Differentiation)

$$\mathcal{F}Df(\xi) = 2\pi i \xi \mathcal{F}f(\xi)$$
$$\mathcal{F}(-2\pi i t f) = D\mathcal{F}f$$

Example. With $B(t) = \chi_{[-\frac{1}{2}, \frac{1}{2}]}(t)$:

$$\hat{B}(\xi) = \text{sinc}(\xi) = \frac{\sin(\pi \xi)}{\pi \xi}$$

And for $k \in \mathbb{Z}$, $\hat{B}(k) = \delta_{k0}$.

L2 Fourier Transform

Proposition. (Properties) With \mathcal{F} and \mathcal{F}^{-1} defined on $L^2(\mathbb{R})$, then:

- \mathcal{F} and \mathcal{F}^{-1} are linear operators.
- \mathcal{F} and \mathcal{F}^{-1} are unitary operator on $L^2(\mathbb{R})$, and so for all $f, g \in L^2(\mathbb{R})$

$$\langle \hat{f}, \hat{g} \rangle = \langle f, g \rangle$$

Equivalently, $\mathcal{F}^* = \mathcal{F}^{-1}$.

Convolution

Definition. (Convolution) For $f, g \in L^1(\mathbb{R})$, we define the **convolution**:

$$f * g(t) = \int_{-\infty}^\infty f(s)g(t-s) \, ds$$

Theorem. If $f, g \in L^1(\mathbb{R})$ then

- $f * g \in L^1(\mathbb{R})$ and $\|f * g\|_1 \leq \|f\|_1 \|g\|_1$.
- $(f * g)(\xi) = \hat{f}(\xi)\hat{g}(\xi)$.

Example. Let $f = \chi_{[-\frac{1}{2}, \frac{1}{2}]}$. Then,

$$f * f(x) = g(x) = \max\{1 - |x|, 0\}$$
$$\hat{g}(\xi) = \hat{f}(\xi)^2 = \frac{\sin^2(\pi \xi)}{\pi^2 \xi^2}$$

Covariance

Definition. Given $t_0, \xi_0 \in \mathbb{R}$ we define the following operators on $L^2(\mathbb{R})$:

$$\tau_{t_0}f(t) = f(t - t_0)$$
$$M_{\xi_0}f(t) = e^{-2\pi i \xi_0 t}f(t)$$
$$D_a f(t) = \frac{1}{\sqrt{a}}f\left(\frac{t}{a}\right)$$

Proposition. (Properties)

- Each operator has an inverse, as follows:
- $$\tau_{t_0}^{-1} = \tau_{-t_0} \quad M_{\xi_0}^{-1} = M_{-\xi_0} \quad D_a^{-1} = D_{a^{-1}}$$
- Each operator is unitary.
 - Each operator has a relationship with the Fourier transform:

$$\mathcal{F}\tau_{t_0} = M_{t_0}\mathcal{F}$$
$$\mathcal{F}M_{\xi_0} = \tau_{-\xi_0}\mathcal{F}$$
$$\mathcal{F}D_a = D_{a^{-1}}\mathcal{F}$$

Sampling

Definition. (Bandlimit) Given $\Omega > 0$, let

$$B_\Omega = \left\{f \in L^2(\mathbb{R}) : \hat{f}(\xi) = 0 \text{ for } \xi > \frac{\Omega}{2}\right\}$$

Functions in B_Ω are said to be **bandlimited** with **bandwidth** Ω .

Theorem. (Shannon Sampling Theorem) If $f \in B_\Omega$, then

$$f(t) = \sum_{n=-\infty}^\infty f\left(\frac{n}{\Omega}\right) \text{sinc}(\Omega t - n)$$

where $\text{sinc}(x) = \frac{\sin(\pi x)}{\pi x}$.

Theorem. (Poisson Summation Formula) If $f, \hat{f} \in L^1(\mathbb{R})$ and

$$\sum_{n=-\infty}^\infty f(t+n) \quad \text{and} \quad \sum_{n=-\infty}^\infty \hat{f}(\xi+n)$$

are absolutely convergent for all $t, \xi \in \mathbb{R}$, then

$$\sum_{n=-\infty}^\infty f(t+n) = \sum_{m=-\infty}^\infty \hat{f}(m)e^{2\pi i m t}$$

Remark. This can be used to show that certain infinite sums have some particular value.

Uncertainty

Theorem. (Heisenberg Uncertainty Inequality)

$$\int_{-\infty}^\infty (t - t_0)^2 |f(t)|^2 \, dt \cdot \int_{-\infty}^\infty (\xi - \xi_0)^2 |\hat{f}(\xi)|^2 \, d\xi \geq \frac{\|f\|_2^4}{16\pi^2}$$

Time-Frequency Analysis

Definition. (Short-Time Fourier Transform) Let $\varphi \in L^2(\mathbb{R})$. Given $b, \xi \in \mathbb{R}$, let

$$\varphi_{b,\xi}(t) = M_{-\xi}\tau_b\varphi(t) = e^{2\pi i \xi t}\varphi(t - b)$$

Then the short-time Fourier transform $S_\varphi f$ of f with respect to φ is defined by

$$S_\varphi f(b, \xi) = \langle f, \varphi_{b,\xi} \rangle$$

Remark. The short-time Fourier transform is also known as the sliding window Fourier transform, the windowed Fourier transform and the Gabor transform (especially in the case where the window φ is a Gaussian). Note that

$$S_\varphi f(b, \xi) = \int_{-\infty}^\infty f(t)e^{-2\pi i \xi t}\overline{\varphi(t - b)} \, dt = \left(\mathcal{F}f\overline{\varphi(\cdot - b)}\right)(\xi)$$

$\varphi_{b,\xi}$ is obtained from φ by a translation in time by b and in frequency by ξ , i.e., a translation in “phase space” (the time-frequency plane) by (b, ξ) .

Also, if $\varphi \equiv 1$, $S_\varphi f(b, \xi) = \hat{f}(\xi)$, but in fact since φ is assumed to have decay properties, $S_\varphi f$ represents a windowed version of the Fourier transform around $t = b$. By the Plancherel formula we may write the integral as

$$S_\varphi f(b, \xi) = \int_{-\infty}^\infty \hat{f}(s)e^{2\pi i b(s - \xi)}\overline{\hat{\varphi}(s - \xi)} \, ds$$

$S_\varphi f(b, \xi)$ represents time-frequency localised information about f , i.e., the information about f near the time b and the frequency ξ .

Theorem. If $f \in L^2(\mathbb{R})$, then $S_\varphi : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}^2)$ is a *multiple of an isometry*

$$\int_{-\infty}^\infty \int_{-\infty}^\infty |S_\varphi f(b, \xi)|^2 \, db \, d\xi = \|\varphi\|_2^2 \int_{-\infty}^\infty |f(t)|^2 \, dt = \|\varphi\|_2^2 \|f\|_2^2$$