

ME140A: Numerical Analysis in Engineering

Lecture Notes

Alexander Meiburg

9/22/22 - 9/22/22

Course Organization

- ▶ Goal: **Numerical solution of integrals and differential equations**
- ▶ Homework will rely significantly on you programming these methods you learn. MATLAB recommended, alternatives welcome
- ▶ Collaborate on the homework! You learn more that way. Just make sure that you are, in fact, learning. 😊
- ▶ HW: 10% of grade. Exams: 30%/30%/30%.
- ▶ Submit homework via email
- ▶ Office hours by appointment or Zoom, but my schedule is very open!
- ▶ Full syllabus available [here](#)
- ▶ These notes will be continually updated [here](#)

Numerical Integration

Recall: **Differentiation** systematically lets you take a function $F(x)$ and find its derivative $f(x) = F'(x)$.

$$\frac{d}{dx} \left(e^{\sin(x+\log x)} \right) = e^{\sin(x+\log x)} \cos(x+\log x) (1 + 1/x)$$

Numerical Integration

Recall: **Differentiation** systematically lets you take a function $F(x)$ and find its derivative $f(x) = F'(x)$.

$$\frac{d}{dx} \left(e^{\sin(x+\log x)} \right) = e^{\sin(x+\log x)} \cos(x+\log x) (1 + 1/x)$$

Integration asks for the opposite. You have a handful of rules(!), but they can't cover every case. Often impossible, and we resort to defining new functions or using the computer

$$\int e^{-x^2} dx := \Phi(x)$$


Numerical Integration

Recall: **Differentiation** systematically lets you take a function $F(x)$ and find its derivative $f(x) = F'(x)$.

$$\frac{d}{dx} \left(e^{\sin(x+\log x)} \right) = e^{\sin(x+\log x)} \cos(x+\log x) (1 + 1/x)$$

Integration asks for the opposite. You have a handful of rules(!), but they can't cover every case. Often impossible, and we resort to defining new functions or using the computer

$$\int e^{-x^2} dx := \Phi(x)$$



Numerical Integration

Example: Computing the position of an object after some movement.

$y(t)$ = Position as a function of time

$v(t)$ = Velocity

$$v = \frac{dy(t)}{dt}, \quad y(t) = \int_0^t v(t) dt$$

Numerical Integration

Example: Computing the position of an object after some movement.

$y(t)$ = Position as a function of time

$v(t)$ = Velocity

$$v = \frac{dy(t)}{dt}, \quad y(t) = \int_0^t v(t) dt$$

What form are we given $v(t)$? Could be:

- Explicit function of t (from theory, specifications...)

Numerical Integration

Example: Computing the position of an object after some movement.

$y(t)$ = Position as a function of time

$v(t)$ = Velocity

$$v = \frac{dy(t)}{dt}, \quad y(t) = \int_0^t v(t) dt$$

What form are we given $v(t)$? Could be:

- ▶ Explicit function of t (from theory, specifications...)
- ▶ Data (samples at certain points)

Numerical Integration

Example: Computing the position of an object after some movement.

$y(t)$ = Position as a function of time

$v(t)$ = Velocity

$$v = \frac{dy(t)}{dt}, \quad y(t) = \int_0^t v(t) dt$$

What form are we given $v(t)$? Could be:

- ▶ Explicit function of t (from theory, specifications...)
- ▶ Data (samples at certain points)
- ▶ A function of t **and** of y , or something else

This last one forms a differential equation, and will need different methods. But many ideas will transfer!

Numerical Integration

Example: Computing the position of an object after some movement.

$y(t)$ = Position as a function of time

$v(t)$ = Velocity

$$v = \frac{dy(t)}{dt}, \quad y(t) = \int_0^t v(t) dt$$

What form are we given $v(t)$? Could be:

- ▶ Explicit function of t (from theory, specifications...)
- ▶ Data (samples at certain points)
- ▶ A function of t **and** of y , or something else

This last one forms a differential equation, and will need different methods. But many ideas will transfer!

May also have y as an integral over *several* variables, not just one. e.g. Dust accumulating on a surface varies with x , y , and t . Can do three integrals in a row (analytically), or one 3D integral (numerically).

Aside: Numerical Differentiation

Differentiation is easy if we have an exact formula, but what about for data points?

Aside: Numerical Differentiation

Differentiation is easy if we have an exact formula, but what about for data points?

$$v(t) \approx \frac{y(t + 1\text{s}) - y(t)}{1\text{s}}$$

Aside: Numerical Differentiation

Differentiation is easy if we have an exact formula, but what about for data points?

$$v(t) \approx \frac{y(t + 1s) - y(t)}{1s}$$

But consider:

t	y(t)
0	5
1	6.1
2	7.3
3	8.4
4	9.8
7	15.3
8	17.4
9	59.8
10	138.7
11	138.8

Issues such as irregular data, or gaps in time too large to understand what happened. Big question in its own right, Week 2!

Newton-Cotes

Problem: given $f(t)$, find $F(t) = \int_0^t f(t) dt$. If $f(t)$ is too complicated, let's find something simpler we can integrate. What's simple? Polynomials!

Newton-Cotes

Problem: given $f(t)$, find $F(t) = \int_0^t f(t) dt$. If $f(t)$ is too complicated, let's find something simpler we can integrate. What's simple? Polynomials!


Idea: **Sample** the function at several points, **estimate** the function in between with a simpler formula, **analytically integrate** the estimate.

Newton-Cotes

Problem: given $f(t)$, find $F(t) = \int_0^t f(t) dt$. If $f(t)$ is too complicated, let's find something simpler we can integrate. What's simple? Polynomials!


Idea: **Sample** the function at several points, **estimate** the function in between with a simpler formula, **analytically integrate** the estimate.

Simplest: Linear fit through two points. ("Trapezoidal rule")



Newton-Cotes

Fit quadratic ("Simpson's rule"):



Credit: Wikimedia

Newton-Cotes

In general, find

$$f_n(x) = a_0 + a_1x + a_2x^2 + \dots a_nx^n$$

and integrate

$$\int_a^b f_n(x) dx$$

Turns out: a_i depend linearly on the $f(x_i)$, so the result is some weighted sum of the $f(x_i)$.

Newton-Cotes

In general, find

$$f_n(x) = a_0 + a_1x + a_2x^2 + \dots a_nx^n$$

and integrate

$$\int_a^b f_n(x) dx$$

Turns out: a_i depend linearly on the $f(x_i)$, so the result is some weighted sum of the $f(x_i)$.

Trapezoidal:

$$\int_a^b f(x) dx \approx \frac{b-a}{2} (f(b) + f(a))$$

Simpson's:

$$\int_a^b f(x) dx \approx \frac{b-a}{6} \left(f(b) + 4f\left(\frac{a+b}{2}\right) + f(a) \right)$$

Newton-Cotes


If our function is too complicated over $[a, b]$, then subdivide and do each separately.

$$\int_{x=a}^b f(x) = \int_{x=a}^{(a+b)/2} f(x) + \int_{x=(a+b)/2}^b f(x)$$

Newton-Cotes

If our function is too complicated over $[a, b]$, then subdivide and do each separately.

$$\int_{x=a}^b f(x) = \int_{x=a}^{(a+b)/2} f(x) + \int_{x=(a+b)/2}^b f(x)$$



Error Analysis

Write

$$f(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2 + \dots$$

First and second terms accurate, third isn't. Fitting gives

$$E_{trap} \approx \frac{1}{12}|f''(\xi)|(b - a)^2$$

Error Analysis

Write

$$f(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2 + \dots$$

First and second terms accurate, third isn't. Fitting gives

$$E_{trap} \approx \frac{1}{12}|f''(\xi)|(b - a)^2$$

Refine this with subintervals