

Primary Care Office Visit Note

Patient Name: Michael Reynolds

DOB: 1967-03-14

Age: 58

Sex: Male

MRN: MR-472918

Date of Service: 2026-05-21

Provider: Sarah Whitman, MD

Location: ValleySteer Primary Care – Internal Medicine

Chief Complaint

Follow-up for high blood pressure.

History of Present Illness

Michael Reynolds is a 58-year-old male with a history of essential hypertension who presents today for blood pressure follow-up.

At his prior visit, his blood pressure was elevated, and lisinopril was increased from 20 mg daily to 30 mg daily. He reports taking lisinopril consistently and denies dizziness, cough, chest pain, shortness of breath, headache, vision changes, or leg swelling.

He has been checking blood pressure at home several times per week. Most home readings have been in the 130s systolic and 80s diastolic. He reports reducing sodium intake and walking several evenings per week.

He also has type 2 diabetes mellitus treated with metformin, but today's visit is primarily for hypertension follow-up.

Past Medical History

- Essential hypertension
 - Type 2 diabetes mellitus without known complications
 - Hyperlipidemia
-

Current Medications

- Lisinopril 30 mg tablet, take 1 tablet by mouth once daily
 - Metformin 1000 mg tablet, take 1 tablet by mouth twice daily with meals
 - Atorvastatin 20 mg tablet, take 1 tablet by mouth nightly
-

Vital Signs

Date/Time	BP	Pulse	Resp	Temp	SpO2	Weight
2026-05-21 08:42 AM	138/88 mmHg	74 bpm	16	98.0°F	99%	211 lb

BP Measurement Details:

Blood pressure was obtained in clinic while seated, left arm, adult cuff, after the patient rested for 5 minutes. Reading documented as 138/88 mmHg.

Physical Examination

General: Alert, pleasant male in no acute distress.

Cardiovascular: Regular rate and rhythm. No murmurs. No peripheral edema.

Respiratory: Clear to auscultation bilaterally.

Neurologic: Alert and oriented. No focal deficits.

Extremities: No edema.

Assessment

1. Essential hypertension

Improved control after lisinopril dose increase. Clinic BP today is 138/88 mmHg.
Patient is tolerating current medication without side effects.

2. Type 2 diabetes mellitus without complication

Chronic condition, treated with metformin. Not the primary focus of today's visit.

3. Hyperlipidemia

Stable on atorvastatin.

Plan

Hypertension

- Continue lisinopril 30 mg once daily.
- Continue home BP monitoring 2–3 times per week.
- Continue low-sodium diet and regular walking.
- No medication change today.

Type 2 Diabetes Mellitus

- Continue metformin 1000 mg twice daily.
- Routine diabetes follow-up as previously scheduled.

Hyperlipidemia

- Continue atorvastatin 20 mg nightly.
-

Follow-Up

Return in 3 months for hypertension and chronic disease follow-up.

Electronic Signature

Electronically signed by Sarah Whitman, MD

Date Signed: 2026-05-21 09:27 AM