

A dramatic space scene featuring a black hole on the right side, with a bright, glowing accretion disk. In the center of the frame, a small, dark sphere is visible against the dark background of space. The overall color palette is dominated by the dark blues and blacks of space, contrasted with the bright yellows and oranges of the accretion disk.

# How space travel is revolutionized with this one weird trick from chaos theory


Ingo Blechschmidt  
thanking Sven Prüfer and Matthias Hutzler

Institut für Mathematik  
Universität Augsburg  
December 30th, 2016

- 1 A crash course on orbital mechanics
  - Basic facts
  - Changing orbits
  - The tyranny of the rocket equation
  
- 2 One weird trick from chaos theory
  - Lagrangian points
  - Weak stability boundaries
  - The rescue of the Hiten
  - In nature

# Part I

## A crash course on orbital mechanics




## Basic facts

- Getting to space is easy.  
The hard part is staying there.
- Gravitational acceleration at the height of the ISS is still  $\approx 8.7 \text{ m/s}^2$ .



# Basic facts

- Getting to space is easy.  
The hard part is staying there.



$$F_{\text{centripetal}} = F_{\text{gravitation}} \leadsto v_1 = \sqrt{GM/r}$$
$$E_{\text{kinetic}} = E_{\text{gravitation}} \leadsto v_2 = \sqrt{2} v_1$$

## Basic facts

- Getting to space is easy.  
The hard part is staying there.

body	second escape velocity
Earth	$11.2 \text{ km/s} \approx 40\,000 \text{ km/h}$
Moon	$2.4 \text{ km/s}$
Sun	$618 \text{ km/s}$
Milky Way	$\approx 550 \text{ km/s}$

# Basic facts

- Getting to space is easy.  
The hard part is staying there.
- Velocity is very important.

# Basic facts

- Getting to space is easy.  
The hard part is staying there.
- Velocity is very important.
- In the **one-body problem**, there are only three kinds of orbits: elliptic, parabolic, and hyperbolic.



# Basic facts


- Getting to space is easy.  
The hard part is staying there.
- Velocity is very important.
- In the **one-body problem**, there are only three kinds of orbits: elliptic, parabolic, and hyperbolic.
- Have your models straight: Earth is ...
  - 1 a perfect ball?
  - 2 has atmosphere?
  - 3 rotating?

# Changing orbits

## “Live demo”

- Changing the phase
- Changing the eccentricity
- Changing the radius
- Changing inclination

# The tyranny of the rocket equation



Konstantin Tsiolkovsky (\* 1857, † 1935)


$$m_{\text{total}} = m_{\text{payload}} \cdot e^{\Delta v / v_{\text{eff. exhaust}}}$$

# Part II


## One weird trick from chaos theory




# Lagrangian points




# Lagrangian points



# Weak stability boundaries




# Weak stability boundaries





# The rescue of the Hiten



## In nature



Figure 10. Stars stream outward from the Tadpole Galaxy (Arp 188) along a tubelike channel that stretches for some 280,000 light-years. This conduit (the galactic equivalent of the tubes making up the interplanetary transport network) arose through gravitational interaction with a compact galaxy that can now be seen lurking behind one of the Tadpole's spiral arms. (Courtesy of ACS Science & Engineering Team and NASA.)