

MADS-MMS – Mathematics and Multivariate Statistics

Analytic Geometry

Prof. Dr. Stephan Doerfel



FACHHOCHSCHULE KIEL
University of Applied Sciences



Moodle (SoSe 2025)

Agenda

Motivation

Inner Products

Norms and Distances

Angles

Lines, Planes, Hyperplanes

Outline

Motivation

Inner Products

Norms and Distances


Angles

Lines, Planes, Hyperplanes

Motivation

- ▶ one heavily exploited feature in machine learning are geometric relationships between instances
- ▶ kNN exploits distances between instances
- ▶ SVMs measure distances between planes and instances
- ▶ k-means groups instances with short distances between each other
- ▶ text-mining uses the angle between long vectors as similarity measure

Example 1 – \mathbb{R}^2 – Geometric Interpretation



Example 1 – \mathbb{R}^2 – Geometric Interpretation

- ▶ the geometric interpretation works similarly in \mathbb{R}^n with $n > 2$
- ▶ it allows computing various geometric entities, like planes, volumes, distances, angles, ...

Chapter Goals

- ▶ mathematical foundations of geometry
- ▶ inner products (e.g. for distance functions, for SVMs)
- ▶ distance functions (e.g. for clustering)
- ▶ geometric figures (e.g. for SVMs)

Outline

Motivation

Inner Products

Norms and Distances

Angles

Lines, Planes, Hyperplanes

Inner Product

Definition 1 (Inner Product)

Let V be a real-valued vector space. Then a function $\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{R}$ is called an **inner product**, if for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$ and $r, s \in \mathbb{R}$ holds

1. $\langle r\mathbf{u} + s\mathbf{v}, \mathbf{w} \rangle = r\langle \mathbf{u}, \mathbf{w} \rangle + s\langle \mathbf{v}, \mathbf{w} \rangle$ and $\langle \mathbf{u}, r\mathbf{v} + s\mathbf{w} \rangle = r\langle \mathbf{u}, \mathbf{v} \rangle + s\langle \mathbf{u}, \mathbf{w} \rangle$ (**bilinear**)
2. $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$ (**symmetric**)
3. $\langle \mathbf{v}, \mathbf{v} \rangle > 0$ for $\mathbf{v} \neq 0$ and $\langle 0, 0 \rangle = 0$ (**positive definite**)

Example in \mathbb{R}^2 :

$$\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_1 - (u_1 v_2 + u_2 v_1) + 2u_2 v_2$$

The Dot Product

- In the vector space \mathbb{R}^n , the dot product or scalar product

$$\langle \mathbf{u}, \mathbf{v} \rangle := \sum_{i=1}^n u_i v_i$$

is an inner product.

- \mathbb{R}^n together with the dot product is called a **Euclidean vector space**.

Outline

Motivation

Inner Products

Norms and Distances

Angles

Lines, Planes, Hyperplanes

Vector Norm

Definition 2

A **norm** on a vector space is a function $\| \cdot \| : V \rightarrow \mathbb{R}$, such that for all $\mathbf{u}, \mathbf{v} \in V$ and $r \in \mathbb{R}$ holds

1. $\|r\mathbf{v}\| = |r| \cdot \|\mathbf{v}\|$
2. $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$ (**triangle inequality**)
3. $\|\mathbf{v}\| \geq 0$ and $\|\mathbf{v}\| = 0 \iff \mathbf{v} = \mathbf{0}$

The norm of a vector can be understood as its **length**.

Example: Manhattan Norm

$$\|\mathbf{v}\|_1 = \sum_{i=1}^n |v_i|$$

- ▶ sum the absolute values of a vector
- ▶ use case: travel distance in manhattan (taxicab norm)
- ▶ use case: regularization (in machine learning)
- ▶ What does a circle look like?

Example: Euclidean Norm

$$\|\mathbf{v}\|_2 := \sqrt{\sum_{i=1}^n v_i^2}$$

- ▶ square root of the sum of squares
- ▶ corresponds to the “intuitive” length of vector
- ▶ the length from Euclidean geometry
- ▶ What does a circle look like?

Observation:

$$\|\mathbf{v}\|_2 = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$$

Induced Norm

- ▶ If $\langle \cdot, \cdot \rangle$ is the inner product of a vector space, then it induces a norm on the vector space: $\|\mathbf{v}\| := \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$.
- ▶ The dot product induces the Euclidean norm.
- ▶ Not each norm is an induced norm (e.g. Manhattan norm).

Unit Vectors

Given a vector \mathbf{v} . How do we get a vector with the same orientation, but with length 1?

$$\frac{1}{\|\mathbf{v}\|} \mathbf{v}$$

Proof: $\left\| \frac{1}{\|\mathbf{v}\|} \mathbf{v} \right\| = \frac{1}{\|\mathbf{v}\|} \|\mathbf{v}\| = 1.$

Distance Function

Definition 3

Let O be a set of objects. A **distance function** is a function $d : O \times O \rightarrow \mathbb{R}_{\geq 0}$ such that

1. $d(o_1, o_2) = d(o_2, o_1)$
2. $d(o_1, o_2) = 0 \iff o_1 = o_2$

Definition 4

A distance function d is called a metric, if for all o_1, o_2, o_3 the triangle inequality holds:

$$d(o_1, o_3) \leq d(o_1, o_2) + d(o_2, o_3)$$



The general definition of distance allows for various inclusions and combinations of an objects features.

Induced Metric


- ▶ In a vector space V with inner product $\langle \cdot, \cdot \rangle$,

$$d : (\mathbf{u}, \mathbf{v}) \mapsto \sqrt{\langle \mathbf{u} - \mathbf{v}, \mathbf{u} - \mathbf{v} \rangle}$$

is a distance metric.

- ▶ In a vector space V , a norm induces a distance metric by

$$d : (\mathbf{u}, \mathbf{v}) \mapsto \|\mathbf{u} - \mathbf{v}\|.$$

 The different inner products and their distance function properties are the key ingredient in SVMs using the kernel trick.

Distance functions: Minkowski-Metrics

Definition 5 (Minkowski-Metric)

Let $\mathbf{u} = (u_1, \dots, u_n)$ and $\mathbf{v} = (v_1, \dots, v_n)$ be real-valued vectors and $p \in \mathbb{R}_{>0}$. For $p \geq 1$, the Minkowski-Metric (L_p -metric) is defined as:

$$d_p(\mathbf{u}, \mathbf{v}) := \sqrt[p]{\sum_{i=1}^n |u_i - v_i|^p}$$

Often used Minkowski-Metrics:


- ▶ Manhattan ($p = 1$): $d_1(\mathbf{u}, \mathbf{v}) = \sum_{i=1}^n |u_i - v_i|$
- ▶ Euclidean ($p = 2$): $d_2(\mathbf{u}, \mathbf{v}) = \sqrt{\sum_{i=1}^n (u_i - v_i)^2}$
- ▶ Maximum ($p = \infty$): $d_\infty(\mathbf{u}, \mathbf{v}) = \max_{i=1}^n |u_i - v_i|$

Weighted Minkowski-Metric

- ▶ Sometimes we want to emphasize particularly important dimensions.
- ▶ The Minkowski-Metrics can be extended using weights ω_k :

$$d_p^\omega(\mathbf{u}, \mathbf{v}) = \sqrt[p]{\sum_{i=1}^n \omega_i |u_i - v_i|^p}$$

Unit Circles in Different Minkowski-Metrics



Source: <https://commons.wikimedia.org/wiki/User:Waldir>

Other Distance Functions

- ▶ For categorical attributes, the Hamming-Distance:

$$d(\mathbf{u}, \mathbf{v}) = \sum_{i=1}^n \delta(u_i, v_i) \quad \text{with} \quad \delta(u_i, v_i) = \begin{cases} 0 & \text{if } u_i = v_i \\ 1 & \text{otherwise} \end{cases}$$

- ▶ For sets A and B , the Jaccard Distance:

$$J(A, B) = 1 - \frac{|A \cap B|}{|A \cup B|}$$

- ▶ For strings the Levenshtein distance and the Damerau–Levenshtein distance between two strings measure the number of edit operations for turning one string into the other.
- ▶ many more, see e.g. implementations of `sklearn.neighbors.DistanceMetric`

Similarity Function

- ▶ Distance functions measure the opposite of similarity. Higher means more distance, thus less similarity.
- ▶ When we talk about measuring similarity, we usually rather mean distance.

 Exercises 1–3

Outline

Motivation

Inner Products

Norms and Distances

Angles

Lines, Planes, Hyperplanes

The Cosine of Two Vectors


- ▶ Let V be a vector space with inner product $\langle \cdot, \cdot \rangle$, then for two vectors $\mathbf{u}, \mathbf{v} \neq 0$

$$\cos \alpha = \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{u}\| \cdot \|\mathbf{v}\|}$$

is the cosine of the angle between the two vectors

- ▶ Hereby the norm is induced by the inner product
- ▶ With the dot product (thus Euclidean norm), the angles correspond to our usual intuition of angles

Recap Cosine



- ▶ \cos is a trigonometric function
- ▶ $\cos(x + 2k\pi) = \cos(x)$ for $k \in \mathbb{Z}$ (periodic)
- ▶ $\cos(0) = 1$ (0°)
- ▶ $\cos(\frac{\pi}{2}) = 0$ (90°)
- ▶ $\cos(\pi) = -1$ (180°)

Orthogonal Vectors

Two vectors are orthogonal (angle 90°) if their inner product is zero.

Cosine Similarity

$$d(\mathbf{u}, \mathbf{v}) = \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{u}\| \cdot \|\mathbf{v}\|}$$

- ▶ an actual similarity function (higher means more similar)
- ▶ ignores the length of vectors, only orientation is relevant
- ▶ 1 means identical orientation
- ▶ -1 means inverse orientation
- ▶ $1 - d(\mathbf{u}, \mathbf{v})$ is often used as distance measure, however not a proper metric (triangle inequality)

Example Cosine Similarity for Texts

Representing text documents as vectors:

- ▶ Simple Version: dimensions are words, each entry is the frequency of the term in the document
- ▶ Tf-idf: dimensions are words, each entry is the product of its frequency in the document (tf) and the negative log of the share of documents that contain the term (idf).
 - ▶ Advantage: Reduce the influence of very common words, emphasize document-specific words.

In both cases: very long, sparse vectors.

- ▶ cosine similarity works well with texts of different length
- ▶ a text and the same text appended to itself have cosine distance 0 but would have high values with many other distance functions (e.g. Euclidean)

Outline

Motivation

Inner Products


Norms and Distances

Angles

Lines, Planes, Hyperplanes

A Line in \mathbb{R}^2

Let's consider \mathbb{R}^2 . How can we describe a line?



A Line in \mathbb{R}^2

- ▶ all vectors that are orthogonal to a given vector $\mathbf{w} \neq 0$ form a line through the origin in \mathbb{R}^2
- ▶ adding one vector \mathbf{u}_0 moves the line away from the origin
- ▶ thus, \mathbf{x} is on the line, if and only if $\mathbf{x} - \mathbf{u}_0$ is orthogonal to \mathbf{w} , meaning

$$\langle \mathbf{x} - \mathbf{u}_0, \mathbf{w} \rangle = 0$$

- ▶ equivalent:

$$\langle \mathbf{x}, \mathbf{w} \rangle - \langle \mathbf{u}_0, \mathbf{w} \rangle = 0$$

- ▶ In this model: \mathbf{w} and \mathbf{u}_0 are fix parameters that determine the line.

Alternative Line Descriptions in \mathbb{R}^2

$$\langle \mathbf{x} - \mathbf{u}_0, \mathbf{w} \rangle = 0 \quad \text{with } \mathbf{w}, \mathbf{u}_0 \in \mathbb{R}^2$$

$$\langle \mathbf{x}, \mathbf{w} \rangle + b = 0 \quad \text{with } \mathbf{w} \in \mathbb{R}^2, b \in \mathbb{R}$$

$$x_2 = mx_1 + n \quad \text{with } \mathbf{x} = (x_1, x_2), m, n \in \mathbb{R}$$

With $\langle \cdot, \cdot \rangle$ being the dot product, the first two are equivalent. The third is the linear function description of a line. It is however not fully equivalent to the other two.

 Exercises Homework: Prove $1 \iff 2$, Here: Prove $2 \iff 3$.

Hyperplane

Definition 6 (Hyperplane)

A **Hyperplane** in an n -dimensional vector space \mathbb{R}^n is an $(n - 1)$ -dimensional affine subspace.

- ▶ $n=1 \rightarrow$ point
- ▶ $n=2 \rightarrow$ line
- ▶ $n=3 \rightarrow$ (regular) plane

The construction of hyperplanes is exactly analogous to the hyperplanes in \mathbb{R}^2 (lines).

Distance between Point and Hyperplane

Given a hyperplane in \mathbb{R}^n as $\langle \mathbf{x}, \mathbf{w} \rangle + b = 0$ and a vector $\mathbf{v} \in \mathbb{R}^n$. Assuming $\mathbf{w} \neq 0$, what is the distance between \mathbf{v} and the hyperplane?

- ▶ to \mathbf{v} , add a vector \mathbf{z} that is orthogonal to the hyperplane, such that the sum is on the hyperplane:

$$\langle \mathbf{v} + \mathbf{z}, \mathbf{w} \rangle + b = 0 \text{ and } \mathbf{z} = r\mathbf{w} \text{ with } r \in \mathbb{R}$$

- ▶ Thus:

$$\langle \mathbf{v} + r\mathbf{w}, \mathbf{w} \rangle + b = 0 \implies |r| = \frac{-(b + \langle \mathbf{v}, \mathbf{w} \rangle)}{\langle \mathbf{w}, \mathbf{w} \rangle}$$

- ▶ The distance we are looking for is:

$$\|\mathbf{z}\| = \|r\mathbf{w}\| = |r| \|\mathbf{w}\| = \frac{-(b + \langle \mathbf{v}, \mathbf{w} \rangle)}{\|\mathbf{w}\|^2} \|\mathbf{w}\| = \frac{-(b + \langle \mathbf{v}, \mathbf{w} \rangle)}{\|\mathbf{w}\|}$$