

Chapter 4

Sets

Discrete Structures for Computing on January 4, 2023

Nguyen An Khuong, Tran Tuan Anh, Nguyen Tien Thinh, Mai
Xuan Toan, Tran Hong Tai
Faculty of Computer Science and Engineering
University of Technology - VNUHCM
trtanh@hcmut.edu.vn

Sets

Nguyen An Khuong,
Tran Tuan Anh, Nguyen
Tien Thinh, Mai Xuan
Toan, Tran Hong Tai



Contents

Sets

Set Operation

Contents

① Sets

② Set Operation

Sets

Nguyen An Khuong,
Tran Tuan Anh, Nguyen
Tien Thinh, Mai Xuan
Toan, Tran Hong Tai



Contents

Sets

Set Operation

Course outcomes

Sets

Nguyen An Khuong,
Tran Tuan Anh, Nguyen
Tien Thinh, Mai Xuan
Toan, Tran Hong Tai



Contents

Sets

Set Operation

Course learning outcomes

L.O.1	Understanding of logic and discrete structures
	L.O.1.1 – Describe definition of propositional and predicate logic
	L.O.1.2 – Define basic discrete structures: set, mapping, graphs
L.O.2	Represent and model practical problems with discrete structures
	L.O.2.1 – Logically describe some problems arising in Computing
	L.O.2.2 – Use proving methods: direct, contrapositive, induction
	L.O.2.3 – Explain problem modeling using discrete structures
L.O.3	Understanding of basic probability and random variables
	L.O.3.1 – Define basic probability theory
	L.O.3.2 – Explain discrete random variables
L.O.4	Compute quantities of discrete structures and probabilities
	L.O.4.1 – Operate (compute/ optimize) on discrete structures
	L.O.4.2 – Compute probabilities of various events, conditional ones, Bayes theorem

Set Definition

- Set is a **fundamental** discrete structure on which all discrete structures are built
- Sets are used to group objects, which often have the **same properties**

Example

- Set of all the students who are currently taking Discrete Mathematics 1 course.
- Set of all the subjects that K2011 students have to take in the first semester.
- Set of natural numbers \mathbb{N}

Definition

A **set** is an unordered collection of objects.

The objects in a set are called the **elements** (*phần tử*) of the set.

A set is said to **contain** (*chứa*) its elements.





Definition

- $a \in A$: a is an element of the set A
- $a \notin A$: a is **not** an element of the set A

Definition (Set Description)

- The set V of all vowels in English alphabet, $V = \{a, e, i, o, u\}$
- Set of all real numbers greater than 1???
 $\{x \mid x \in \mathbb{R}, x > 1\}$
 $\{x \mid x > 1\}$
 $\{x : x > 1\}$



Definition

Two sets are **equal** iff they have the same elements.


- $(A = B) \leftrightarrow \forall x(x \in A \leftrightarrow x \in B)$

Example

- $\{1, 3, 5\} = \{3, 5, 1\}$
- $\{1, 3, 5\} = \{1, 3, 3, 3, 5, 5, 5, 5\}$

Venn Diagram

- John Venn in 1881
- **Universal set** (*tập vũ trụ*) is represented by a rectangle
- **Circles** and other **geometrical figures** are used to represent sets
- **Points** are used to represent particular elements in set



Special Sets

- **Empty set** (*tập rỗng*) has no elements, denoted by \emptyset , or $\{\}$
- A set with one element is called a **singleton set**
- What is $\{\emptyset\}$?
- **Answer**: singleton






Definition

The set A is called a **subset** (*tập con*) of B iff every element of A is also an element of B , denoted by $A \subseteq B$.

If $A \neq B$, we write $A \subset B$ and say A is a **proper subset** (*tập con thực sự*) of B .

- $\forall x(x \in A \rightarrow x \in B)$
- For every set S ,
(i) $\emptyset \subseteq S$, (ii) $S \subseteq S$.



Cardinality

Definition

If S has exactly n distinct elements where n is non-negative integers, S is **finite set** (*tập hữu hạn*), and n is **cardinality** (*bản số*) of S , denoted by $|S|$.

Example

- A is the set of odd positive integers less than 10. $|A| = 5$.
- S is the letters in Vietnamese alphabet, $|S| = 29$.
- Null set $|\emptyset| = 0$.

Definition

A set that is **infinite** if it is not finite.

Example

- Set of positive integers is infinite





Definition

Given a set S , the **power set** (*tập lũy thừa*) of S is the set of all subsets of the set S , denoted by $P(S)$.

Example

What is the power set of $\{0, 1, 2\}$?

$$P(\{0, 1, 2\}) = \{\emptyset, \{0\}, \{1\}, \{2\}, \{0, 1\}, \{0, 2\}, \{1, 2\}, \{0, 1, 2\}\}$$

Example

- What is the power set of the empty set?
- What is the power set of the set $\{\emptyset\}$

Theorem

If a set has n elements, then its power set has 2^n elements.

Prove using induction!



Ordered n -tuples

Definition

The **ordered n -tuple** (*dãy sắp thứ tự*) (a_1, a_2, \dots, a_n) is the ordered collection that has a_1 as its first element, a_2 as its second element, \dots , and a_n as its n th element.

Definition

Two ordered n -tuples $(a_1, a_2, \dots, a_n) = (b_1, b_2, \dots, b_n)$ iff $a_i = b_i$, for $i = 1, 2, \dots, n$.

Example

2-tuples, or **ordered pairs** (*cặp*), (a, b) and (c, d) are equal iff $a = c$ and $b = d$



Cartesian Product

- René Descartes (1596–1650)

Definition

Let A and B be sets. The **Cartesian product** (*tích Đề-các*) of A and B , denoted by $A \times B$, is the set of ordered pairs (a, b) , where $a \in A$ and $b \in B$. Hence,

$$A \times B = \{(a, b) \mid a \in A \wedge b \in B\}$$

Example

Cartesian product of $A = \{1, 2\}$ and $B = \{a, b, c\}$. Then

$$A \times B = \{(1, a), (1, b), (1, c), (2, a), (2, b), (2, c)\}$$

Show that $A \times B \neq B \times A$



Cartesian Product

Definition

$$A_1 \times A_2 \times \cdots \times A_n = \{(a_1, a_2, \dots, a_n) \mid a_i \in A_i \text{ for } i = 1, 2, \dots, n\}$$

Example

$A = \{0, 1\}, B = \{1, 2\}, C = \{0, 1, 2\}$. What is $A \times B \times C$?

$$\begin{aligned} A \times B \times C = \{ & (0, 1, 0), (0, 1, 1), (0, 1, 2), (0, 2, 0), (0, 2, 1), \\ & (0, 2, 2), (1, 1, 0), (1, 1, 1), (1, 1, 2), (1, 2, 0), \\ & (1, 2, 1), (1, 2, 2) \} \end{aligned}$$




Union

Definition

The **union** (*hợp*) of A and B

$$A \cup B = \{x \mid x \in A \vee x \in B\}$$



- Example:
 - $\{1,2,3\} \cup \{2,4\} = \{1,2,3,4\}$
 - $\{1,2,3\} \cup \emptyset = \{1,2,3\}$




Intersection

Definition

The **intersection** (*giao*) of A and B

$$A \cap B = \{x \mid x \in A \wedge x \in B\}$$



Example:

- $\{1,2,3\} \cap \{2,4\} = \{2\}$
- $\{1,2,3\} \cap \mathbb{N} = \{1,2,3\}$



Union/Intersection

Sets

Nguyen An Khuong,
Tran Tuan Anh, Nguyen
Tien Thinh, Mai Xuan
Toan, Tran Hong Tai



Contents

Sets

Set Operation

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \dots \cup A_n = \{x \mid x \in A_1 \vee x \in A_2 \vee \dots \vee x \in A_n\}$$


$$\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap \dots \cap A_n = \{x \mid x \in A_1 \wedge x \in A_2 \wedge \dots \wedge x \in A_n\}$$

Difference

Definition

The **difference** (*hiệu*) of A and B

$$A - B = \{x \mid x \in A \wedge x \notin B\}$$



Example:

- $\{1,2,3\} - \{2,4\} = \{1,3\}$
- $\{1,2,3\} - \mathbb{N} = \emptyset$




Complement

Definition

The **complement** (*phần bù*) of A

$$\overline{A} = \{x \mid x \notin A\}$$



Example:

- $A = \{1,2,3\}$ then $\overline{A} = ???$
- **Note that** $A - B = A \cap \overline{B}$



Set Identities

$A \cup \emptyset$	$=$	A	Identity laws
$A \cap U$	$=$	A	Luật đồng nhất
$A \cup U$	$=$	U	Domination laws
$A \cap \emptyset$	$=$	\emptyset	Luật nuốt
$A \cup A$	$=$	A	Idempotent laws
$A \cap A$	$=$	A	Luật lũy đẳng
$\overline{(\overline{A})}$	$=$	A	Complementation law
			Luật bù



Set Identities

$A \cup B$	=	$B \cup A$	Commutative laws
$A \cap B$	=	$B \cap A$	Luật giao hoán
$A \cup (B \cup C)$	=	$(A \cup B) \cup C$	Associative laws
$A \cap (B \cap C)$	=	$(A \cap B) \cap C$	Luật kết hợp
$A \cup (B \cap C)$	=	$(A \cup B) \cap (A \cup C)$	Distributive laws
$A \cap (B \cup C)$	=	$(A \cap B) \cup (A \cap C)$	Luật phân phối
$\overline{A \cup B}$	=	$\overline{A} \cap \overline{B}$	De Morgan's laws
$\overline{A \cap B}$	=	$\overline{A} \cup \overline{B}$	Luật De Morgan



Method of Proofs of Set Equations

To prove $A = B$, we could use

- Venn diagrams
- Prove that $A \subseteq B$ and $B \subseteq A$
- Use **membership table**
- Use set builder notation and logical equivalences

Sets

Nguyen An Khuong,
Tran Tuan Anh, Nguyen
Tien Thinh, Mai Xuan
Toan, Tran Hong Tai




Contents

Sets

Set Operation

Example (1)



Example

Verify the distributive rule $P \cup (Q \cap R) = (P \cup Q) \cap (P \cup R)$



Example (2)

Example

Prove: $\overline{A \cap B} = \overline{A} \cup \overline{B}$

(1) Show that $\overline{A \cap B} \subseteq \overline{A} \cup \overline{B}$

Suppose that $x \in \overline{A \cap B}$

By the definition of complement, $x \notin A \cap B$

So, $x \notin A$ or $x \notin B$

Hence, $x \in \overline{A}$ or $x \in \overline{B}$

We conclude, $x \in \overline{A} \cup \overline{B}$

Or, $\overline{A \cap B} \subseteq \overline{A} \cup \overline{B}$

(2) Show that $\overline{A} \cup \overline{B} \subseteq \overline{A \cap B}$



Example (3)

Prove: $\overline{A \cap B} = \overline{A} \cup \overline{B}$

A	B	$A \cap B$	$\overline{A \cap B}$	$\overline{A} \cup \overline{B}$
1	1	1	0	0
1	0	0	1	1
0	1	0	1	1
0	0	0	1	1



Example (4)

Prove: $\overline{A \cap B} = \overline{A} \cup \overline{B}$

$$\begin{aligned}\overline{A \cap B} &= \{x | x \notin A \cap B\} \\ &= \{x | \neg(x \in A \cap B)\} \\ &= \{x | \neg(x \in A \wedge x \in B)\} \\ &= \{x | \neg(x \in A) \vee \neg(x \in B)\} \\ &= \{x | x \notin A \vee x \notin B\} \\ &= \{x | x \in \overline{A} \vee x \in \overline{B}\} \\ &= \{x | x \in \overline{A} \cup \overline{B}\}\end{aligned}$$

