| PRODUCT                        | QUANTITY                                | DOSAGE                                | TIMING OF<br>DOSE   | APPLICATION                         | INDICATION                                 |
|--------------------------------|---|---------------------------------------|---|-------------------------------------|--|
| 2-Methoxyestradiol             |   |                                       |   |                                     |  |
| Capsules                       | 3mL                                     | 100mcg daily                          | Before bed  | Suppository                         | Cancer                                     |
| 3-Desoxy DHEA                  |   |                                       |   |                                     |  |
| Capsules                       | 100mg (30<br>capsules)                  | 1 capsule<br>daily                    | Patient<br>preference   | Oral                                | Aromatase inhibitor                        |
| 5-amino-1MQ                    |   |                                       |   |                                     |  |
| Injection                      | 10ml at<br>300mg/ml                     | 150mcg (.5ml)<br>daily for 20<br>days | 30 minutes<br>after oral<br>administratio<br>n of NAD+<br>precursor | Subcutaneous injection              | Weight Loss<br>and<br>Performance          |
| Amlexanox                      |   | ,                                     |   | ,                                   |  |
| Capsules                       | 40mg (90<br>capsules)                   | 1 capsule TID                         | Patient<br>preference   | Oral                                | Insulin<br>sensitivity,<br>weight loss     |
| Ammonium<br>Tetrathiomolybdate |   |                                       |   |                                     |  |
| Capsules                       | 40mg (90<br>capsules)                   | 1 capsule TID                         | Patient<br>preference   | Oral                                | Cancer                                     |
| Aniracetam                     |   |                                       |   |                                     |  |
| Capsules                       | 375mg (60<br>capsules)                  | 2 capsules<br>daily                   | Patient<br>preference   | Oral                                | AMPA receptor<br>modulator<br>(Neurogenic) |
| AOD-9604                       |   | ,                                     |   | ,                                   |  |
| Cream                          | 30mL at<br>600mcg/mL                    | 4 clicks (1mL)<br>daily               | Patient<br>preference   | Rubbed<br>between inner<br>forearms | Repair                                     |
| Injection                      | 5mL at<br>1200mcg/mL                    | 0.25mL daily                          | Patient<br>preference   | Subcutaneous<br>Injection           | Repair                                     |
| Injection with HA              | AOD<br>1000mcg/mL<br>+ HA10mg/mL        | 0.5mL-0.75mL<br>weekly for 4<br>weeks | Done by<br>physician  | Intra-articular                     | Repair hyaline<br>cartilage                |
| Argireline/Leuphasyl           |   |                                       |   |                                     |  |
| Cream                          | 0.5%/0.5%<br>cream in 30ml<br>TopiClick | 1ml applied<br>daily                  | Patient<br>preference   | Topical                             | Reduce wrinkle<br>depth and fine<br>lines  |
| Argireline/GHK-Cu/Leuphasyl    |   |                                       |   |                                     |  |

| Cream               | 0.5%/0.2%/<br>0.5% cream in<br>30ml TopiClick | 1ml applied<br>daily              | Patient<br>preference                            | Topical  | Reduce wrinkle<br>depth and fine<br>lines |
|---------------------|---|-----------------------------------|--|--|---|
| BPC-157             |   |                                   |  |  |   |
| Injection           | 3mL at<br>2000mcg/mL                          | 0.15mL daily                      | Patient<br>preference                            | Subcutaneous<br>injection into<br>site of injury | Repair tendon<br>+ ligaments              |
| Injection           | 5mL at<br>2000mcg/mL                          | 0.15mL daily                      | Patient<br>preference                            | Subcutaneous injection into site of injury       | Repair tendon<br>+ ligaments              |
| Capsules            | 500mcg (30<br>capsules)                       | 1 capsule<br>daily                | Patient<br>preference                            | Oral   | Repair<br>bowel/gut                       |
| Cerebrolysin        |   |                                   |  |  |   |
| Injection           | 4x10 mL at<br>215mg/mL                        | 1mL daily                         | Patient<br>preference                            | Subcutaneous injection                           | Neurogenic                                |
| CJC 1295            |   |                                   |  |  |   |
| Injection           | 2mL at<br>2000mcg/mL                          | 0.10mL<br>5 nights a<br>week      | Nightly, on<br>an empty<br>stomach<br>before bed | Subcutaneous injection                           | GHRH                                      |
| Injection           | 5mL at<br>2000mcg/mL                          | 0.10mL<br>5 nights a<br>week      | Nightly, on<br>an empty<br>stomach<br>before bed | Subcutaneous injection                           | GHRH                                      |
| CJC 1295/Ipamorelin |   |                                   |  |  |   |
| Injection           | 2mL (or 5ml)<br>at 1000mcg/<br>1000mcg/mL     | 0.10mL<br>5 nights a<br>week      | Nightly, on<br>an empty<br>stomach<br>before bed | Subcutaneous injection                           | GHRH, weight loss                         |
| Injection           | 2mL (or 5ml)<br>at 2000mcg/<br>1000mcg/mL     | 0.10mL<br>5 nights a<br>week      | Nightly, on<br>an empty<br>stomach<br>before bed | Subcutaneous injection                           | GHRH, weight loss                         |
| Injection           | 2mL (or 5ml)<br>at 2000mcg/<br>2000mcg/mL     | 0.10mL<br>5 nights a<br>week      | Nightly, on<br>an empty<br>stomach<br>before bed | Subcutaneous injection                           | GHRH, weight loss                         |
| DHH-B               |   |                                   |  |  |   |
| Capsules            | 30 7.5mg<br>capsules                          | Take 1-2<br>capsules as<br>needed | Patient<br>Preference                            | Oral   | Anti-Anxiety                              |
| DIHEXA              |   |                                   |  |  |   |
| Capsules            | 5mg (30                                       | 1-2 capsules                      | Patient  | Oral   | Neurological                              |

|   | capsules)  | daily                               | preference                                   |   | function/repair                                    |
|---|--|-------------------------------------|--|---|--|
| Cream   | 30mL at<br>20mg/mL   | 4 clicks (1mL)<br>daily             | Patient<br>preference                        | Transdermal                             | Neurological<br>function/repair                    |
| DSIP  |  |                                     |  |   |  |
| Injection   | 3mL at<br>1000mcg/mL   | 0.1mL daily,<br>2-3 times a<br>week | Patient<br>preference                        | Subcutaneous injection                  | Endocrine<br>regulation,<br>sleep                  |
| Enclomiphene  |  |                                     |  |   |  |
| Capsules  | 25mg (30<br>capsules)  | 1 capsule<br>daily                  | Patient<br>preference                        | Oral                                    | Aromatase<br>inhibitor<br>(Reproductive<br>system) |
| Epicatechin   |  |                                     |  |   |  |
| Injection   | 100mg/ml   | 100mg (1ml)<br>daily                | Patient<br>preference                        | Subcutaneous injection into the abdomen | Sarcopenia,<br>Muscle gain                         |
| Epitalon  |  |                                     |  |   |  |
| Injection   | 5mL at<br>3000mcg/mL   | 0.1mL daily                         | In the<br>morning                            | Subcutaneous injection into the abdomen | Hormone<br>regulation,<br>telomere<br>extension    |
| Fat Loss Cream<br>(Aminophylline/Glycrrhetinic<br>Acid) |  |                                     |  |   |  |
| Cream   | 60mL at 0.5%<br>aminophylline/<br>glycrrhetinic<br>acid 2.5% | 2 pumps<br>(1mL) twice<br>daily     | Patient<br>preference                        | Transdermal                             | Fat reduction                                      |
| FGL(I)  |  |                                     |  |   |  |
| Injection   | 5mL at<br>10mg/mL  | 0.25mL daily                        | Patient<br>preference                        | Subcutaneous injection into the abdomen | Neurogenic/<br>neurorestorativ<br>e                |
| FOXO4-DRI   |  |                                     |  |   |  |
| Injection   | 10ml vial at<br>10mg/ml                                      | 300mcg/kg                           | 3 doses<br>every other<br>day over 5<br>days | Intravenous                             | Clear senescent<br>cells                           |
| GHK-Cu  |  |                                     |  |   |  |
| Injection   | 5mL at<br>10mg/mL  | 0.2mL daily                         | Patient preference                           | Subcutaneous injection                  | Skin elasticity                                    |
| Foam  | 50mL at<br>5mg/mL  | 2 pumps<br>(1mL) daily              | Patient<br>preference                        | Foam (for scalp application)            | Hair loss  |

| IGF-1         |                                    |  |  |   |   |
|---------------|------------------------------------|--|--|---|---|
| Injection     | Two 6.2mL<br>vials at<br>620mcg/mL | 0.4mL-0.8ml<br>daily                       | Patient<br>preference                      | Subcutaneous injection into the abdomen       | Muscle<br>building,<br>weight loss  |
| IGF-1 LR3     |                                    |  |  |   |   |
| Injection     | 6.2mL at<br>100mcg/mL              | 0.4-0.8mL<br>daily                         | Post workout<br>if applicable              | Subcutaneous<br>injection into<br>the abdomen | Weight loss,<br>muscle<br>building,<br>autocrine +<br>paracrine<br>hormone<br>signaling |
| Ipamorelin    |                                    |  |  |   |   |
| Injection     | 5mL at<br>2000mcg/mL               | 0.1mL daily, 5<br>days a week              | Patient<br>preference                      | Subcutaneous injection                        | GHRH, weight loss   |
| iRGD          |                                    |  |  |   |   |
| Injection     | 10mL at<br>2.5mg/mL                | 40mcg/kg<br>daily                          | Dose<br>alongside<br>concurrent<br>therapy | Subcutaneous injection                        | Cancer  |
| Kisspeptin-10 |                                    |  |  |   |   |
| Injection     | 5mL at<br>100mcg/mL                | 0.1mL daily                                | Patient<br>preference                      | Subcutaneous injection into abdomen           | LH increase   |
| KPV           |                                    |  |  |   |   |
| Topical       | 30ml TopiClick<br>at 15mg/ml       | 0.5ml applied twice daily                  | Patient<br>preference                      | Transdermal<br>Cream                          | Psoriasis   |
| Liraglutide   |                                    |  |  |   | _   |
| Injection     | 5mL at<br>3600mcg/mL               | Varies                                     | Varies                                     | Subcutaneous injection                        | Insulin<br>resistance,<br>weight loss   |
| LL-37         |                                    |  |  |   |   |
| Injection     | 5mL at<br>2000mcg/mL               | Varies with indication and patient         | Physician<br>preference                    | Subcutaneous injection                        | Antimicrobial   |
| Melanotan I   |                                    |  |  |   |   |
| Injection     | 5mL at<br>2000mcg/mL               | 0.15mL daily<br>with UVB light<br>exposure | Patient<br>preference                      | Subcutaneous injection into the abdomen       | Vitiligo  |
| Melanotan II  |                                    |  |  |   |   |
| Injection     | 5mL at<br>2000mcg/mL               | 0.15mL daily<br>for 1-2 weeks              | Patient<br>preference                      | Subcutaneous injection into                   | Libido, tanning,<br>weight loss   |

|                            |                             | I   | T   | I   | 1                                   |
|----------------------------|-----------------------------|---|---|---|-------------------------------------|
|                            |                             | then 0.25mL<br>weekly for<br>maintenance  |   | the abdomen                               |                                     |
| Met-Enkephalin             |                             | ,   | ,   |   |                                     |
| Injection                  | 10mg<br>Lyophilized<br>vial | 10mg IV once<br>weekly                    | N/A   | Intravenous                               | Cancer                              |
| MK-677                     |                             |   |   |   |                                     |
| Capsules                   | 25 mg (30<br>capsules)      | 1 capsule<br>daily                        | Patient<br>preference<br>on empty<br>stomach  | Oral                                      | Growth<br>hormone IGF-1<br>increase |
| MOTS-c                     |                             | ,   |   |   |                                     |
| Injection                  | 4mL at<br>10mg/mL           | 1mL once<br>weekly                        | Patient<br>preference                         | Subcutaneous injection into the abdomen   | Energy, weight loss                 |
| Myristyl                   |                             |   |   |   |                                     |
| Injection                  | 5ml vial at<br>6mg/ml       | 100mcg/kg<br>once weekly<br>intravenously | N/A   | Intravenous                               | Cholesterol<br>Reduction            |
| NMN                        |                             |   |   |   |                                     |
| Injection                  | 10ml vial at<br>200mg/ml    | 200mg (1ml)<br>BID<br>subcutaneous<br>-ly | Patient<br>preference                         | Oral                                      | Anti-aging/<br>Weight loss          |
| Capsule                    | 30 capsules at 500mg        | 1 capsule<br>daily                        | Patient<br>preference                         | Oral                                      | Anti-aging/<br>Weight loss          |
| PEG-MGF                    |                             |   |   |   |                                     |
| Injection                  | 5mL at<br>2000mcg/mL        | 0.1mL daily, 5<br>days a week             | Patient<br>preference                         | Subcutaneous injection into the abdomen   | Building muscle                     |
| Pentosan Polysulfate (PPS) |                             |   |   |   |                                     |
| Injection                  | 10mL at<br>250mg/mL         | 250mg                                     | N/A   | Intramuscular                             | Osteoarthritis                      |
| PNC-27                     |                             |   |   |   |                                     |
| Injection                  | 5mL at<br>1000mcg/mL        | 2.5mg-5mg<br>daily                        | Patient<br>preference                         | Subcutaneous injection                    | Cancer                              |
| PT-141 (Bremelanotide)     |                             |   |   |   |                                     |
| Injection                  | 2mL at<br>10mg/mL           | 0.2mL<br>(female) as<br>needed            | 1-5 hours<br>before sexual<br>activity (first | Subcutaneous<br>injection into<br>abdomen | Erectile<br>dysfunction,<br>libido  |

|                          |                        |  | dose will<br>establish<br>timing)  |   |                                     |
|--------------------------|------------------------|--|--|---|-------------------------------------|
| Injection                | 2mL at<br>10mg/mL      | 0.1mL (male)<br>as needed                            | 1-5 hours<br>before sexual<br>activity (first<br>dose will<br>establish<br>timing) | Subcutaneous<br>injection into<br>abdomen | Erectile<br>dysfunction,<br>libido  |
| PTD-DBM                  |                        |  |  |   |                                     |
| Cream                    | 5mL at 0.5%<br>spray   | Spray a<br>sufficient<br>amount to<br>cover area     | Applied after<br>microneedli-<br>ng  | Transdermal                               | Hair loss                           |
| RG3                      |                        |  |  |   |                                     |
| Nasal                    | 15mL                   | One spray<br>into each<br>nostril 2-4<br>times daily | Patient<br>preference  | Nasal                                     | Neurogenic/<br>neurorestorativ<br>e |
| Nasal                    | 30mL                   | One spray<br>into each<br>nostril 2-4<br>times daily | Patient<br>preference  | Nasal                                     | Neurogenic/<br>neurorestorativ<br>e |
| RG3/Methylcobalamin/NAD+ |                        |  |  |   |                                     |
| Nasal                    | 15mL                   | One spray<br>into each<br>nostril 2-4<br>times daily | Patient<br>preference  | Nasal                                     | Neurogenic/<br>neurorestorativ<br>e |
| Nasal                    | 30mL                   | One spray<br>into each<br>nostril 2-4<br>times daily | Patient<br>preference  | Nasal                                     | Neurogenic/<br>neurorestorativ<br>e |
| SARMS LGD-4033           |                        |  |  |   |                                     |
| Capsules                 | 0.5mg (30<br>capsules) | 1 month on, 1<br>month off<br>rotation               | Patient<br>preference  | Oral                                      | Muscle Building                     |
| Selank                   |                        |  |  |   |                                     |
| Injection                | 5mL at<br>1000mcg/1mL  | 0.1mL daily  | Patient<br>preference  | Subcutaneous injection into the abdomen   | Neurorestorati-<br>ve               |
| Nasal                    | 3mL at<br>7500mcg/mL   | One spray<br>into each<br>nostril once<br>daily      | Patient<br>preference  | Nasal                                     | Neurogenic                          |
| Semax                    |                        |  |  |   |                                     |

| Injection        | 5mL at                                   | 0.1mL daily                                     | Patient  | Subcutaneous                                  | Neurorestorati-                       |
|------------------|--|---|--|---|---------------------------------------|
|                  | 1000mcg/1mL                              |   | preference   | injection into the abdomen                    | ve                                    |
| Nasal            | 3mL at<br>7500mcg/mL                     | One spray<br>into each<br>nostril once<br>daily | Patient<br>preference                                  | Nasal   | Neurogenic                            |
| Tesamorelin      |  |   |  |   |                                       |
| Injection        | 24 vials<br>(lyophilized) at<br>1mg/vial | 1 (1mg) vial 6<br>days a week                   | Before bed,<br>90 minutes<br>after last<br>food intake | Subcutaneous<br>injection into<br>the abdomen | Muscle<br>building,<br>weight loss    |
| Tesofensine      |  |   |  |   |                                       |
| Capsules         | 500mcg (30<br>capsules)                  | 1 capsule<br>daily                              | In the morning   | Patient<br>preference                         | Weight loss                           |
| Thymosin Alpha-1 |  |   |  |   |                                       |
| Injection        | 5mL at<br>3000mcg/mL                     | 0.15mL daily<br>or 1.5mg<br>twice a week        | Patient<br>preference                                  | Subcutaneous injection into the abdomen       | Immune<br>modulation                  |
| Thymosin Beta-4  |  |   |  |   |                                       |
| Injection        | 5mL at<br>3000mcg/mL                     | 0.25mL daily<br>for 20 days                     | Patient<br>preference                                  | Subcutaneous injection into the abdomen       | Repair soft<br>tissue                 |
| ТМВ              |  |   |  |   |                                       |
| Capsules         | 40mg (30<br>capsules)                    | 40mg twice daily                                | Patient<br>preference                                  | Oral  | Cancer                                |
| TTA/Amlexanox    |  |   |  |   |                                       |
| Capsules         | 200mg/40mg<br>(30 capsules)              | 1 capsule 3 times daily                         | Breakfast,<br>lunch, dinner                            | Oral  | Insulin<br>resistance,<br>weight loss |
| VIP              |  |   |  |   |                                       |
| Nasal            | 12mL at<br>50mcg/mL                      | One spray into each nostril up to 4 times daily | Patient<br>preference                                  | Nasal   | Mold toxicity                         |
| Zinc Thymulin    |  |   |  |   |                                       |
| Foam             | 30mL at 50<br>mcg/mL                     | 2 pumps<br>(1mL) daily                          | Applied right before bed                               | Foam (for scalp application)                  | Hair loss                             |