To: From:

RE Patient: Date:

Phone Number: Sender's Phone Number:

Fax Number: Sender's Fax Number:

Pages Including Cover: 2

| PLEASE COMPLE | Patient Inform TE AND FAX B | | |
|---------------------------------|--------------------------------|------|--|
| Date of Last Visit: | | | |
| Reason for Visit: | | | |
| Allergies: | | | |
| Current Medications: | | | |
| | | | |
| Affected Area: | | | |
| e remember to circle quantity a | | | |

The documents accompanying this fax transmission are confidential. Information contained in this fax transmission belongs to the company sending the data and is legally privileged. The information accompanying this fax transmission is intended only for the use of the individual (or company) identified as "recipient." The recipient of this information is prohibited from disclosing, copying, distributing, or using this information except as permitted by current law governing privacy of information issues. Such information must be destroyed after its stated need has been fulfilled, unless otherwise prohibited by law if you have received this fax transmission in error, please notify the "sender" immediately for return instructions.

Fax back to:

| | Patient: | | | |
|--|--|--|--|--|
| | | | | |
| LOCAL ANESTHETIC | | | | |
| Medication: Lidocaine 5% Ointment Dispense: 300 Grams Sig: Apply 2.3 gms to affected area(s) 3.4 times a day (1 gram = 1) | | | | |
| Sig: Apply 2-3 gms to affected area(s) 3-4 times a day. (1 gram = 1 dime size). | | | | |
| INFLAMMATION MANAGEMENT | | | | |
| Medication: Clobetasol 0.05% Ointment Dispense: 360 Grams | | | | |
| Sig: Apply 2-3 gms to affected area(s) 3-4 times daily (1 gm =1 dime size). | | | | |
| ORAL NSAID | | | | |
| Medication: Naproxen Oral Suspenion (125MG/5ML Dispense: 946 ML Sig: Take 10-20ml by mouth twice daily as needed for pain. | | | | |
| ANTI-FUNGAL | • | | | |
| Medication: Naftifin Dispense: 240 grams Sig: Apply 2-3 gram | | | | |
| Signature: | Date: | | | |
| Doctor: | | | | |
| | | | | |
| NPI: | | | | |
| Phone: | Fax: | | | |
| Dispense as Written: | | | | |
| | Medication: Lidocain Dispense: 300 Grams Sig: Apply 2-3 gms to dime size). INFLAMMATION MA Medication: Clobetas Dispense: 360 Grams Sig: Apply 2-3 gms to size). ORAL NSAID Medication: Naproxo Dispense: 946 ML Sig: Take 10-20ml b ANTI-FUNGAL Medication: Naftifin Dispense: 240 grams Sig: Apply 2-3 gram Sig: Apply 2-3 gram Sig: Apply 2-3 gram Sig: Apply 2-3 gram | | | |

The documents accompanying this telecopy transmission contain confidential information belonging to the sender that is legally privileged. This information is intended for the use of the individual or entity referred above. The authorized recipient of the information is prohibited from disclosing the information to any other party and is required to destroy the information after its stated need has been fulfilled, unless otherwise required by state law.

If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action in reference on the contents of these documents is strictly prohibited. If you have received this telecopy in error, please notify the sender immediately to arrange for return of these documents.