

ClinVar Report

James Diao

February 1, 2017

Contents

1	Collect and Merge ClinVar Data	2
1.1	Import ClinVar VCF	2
1.2	Merge ClinVar with 1000 Genomes and ExAC	2
2	Summary Statistics	3
2.1	Fraction of Individuals with Pathogenic Non-Reference Sites	3
3	Penetrance Estimates	4
3.1	Max/Min Penetrance as a Function of $P(D)$ and $P(V D)$	4
3.2	Penetrance Estimates by Ancestry	4

Sourcing ClinVar input from: clinvar_2013-01-14.vcf

Sending output to: Report_2013-01-14.pdf

1 Collect and Merge ClinVar Data

1.1 Import ClinVar VCF

1.2 Merge ClinVar with 1000 Genomes and ExAC

2 Summary Statistics

2.1 Fraction of Individuals with Pathogenic Non-Reference Sites

3 Penetrance Estimates

3.1 Max/Min Penetrance as a Function of $P(D)$ and $P(V|D)$

The left end of the boxplot indicates $P(V|D) = 0.01$,
the bold line in the middle indicates $P(V|D) = \text{point value}$,
the right end of the boxplot indicates $P(V|D) = 1$.

Note: Some diseases have mean theoretical penetrance = 1 because the assumed allelic heterogeneity is greater than is possible, given the observed prevalence and allele frequencies.

3.2 Penetrance Estimates by Ancestry