

POLI 30 D: Political Inquiry
Professor Umberto Mignozzetti
(Based on DSS Materials)

**Lecture 06 | Measuring Population
Characteristics I**

Before we start

Announcements:

- ▶ Quizzes and Participation: On Canvas.
- ▶ Github page:
<https://github.com/umbertomig/POLI30Dpublic>
- ▶ Piazza forum: <https://piazza.com/ucsd/winter2023/17221>

Before we start

Recap:

- ▶ We learned the definitions of Theory, Scientific Theory, and Hypotheses.
- ▶ Data, datasets, variables, and how to compute means.
- ▶ Causal effect, treatments, outcomes, randomization, and ATE.

Great job!

- ▶ Do you have any questions about these contents?

Plan for Today

- Sample vs. Population
- Representative samples
- Random Sampling
- Random Treatment Assignment vs. Random Sampling
- Exploring One Variable At a Time
 - Table of frequencies
 - Table of proportions
 - Histogram
 - Descriptive Statistics: mean, median, standard deviation, and variance

Why Do We Analyze Data?

1. MEASURE: **To infer population characteristics via survey research**
 - what proportion of constituents support a particular policy?
2. PREDICT: **To make predictions**
 - who is the most likely candidate to win an upcoming election?
3. EXPLAIN: **To estimate the causal effect of a treatment on an outcome**
 - what is the effect of small classrooms on student performance?

Why Do We Analyze Data?


1. MEASURE: To infer population characteristics via survey research
 - what proportion of constituents support a particular policy?
2. PREDICT: To make predictions
 - who is the most likely candidate to win an upcoming election?
3. EXPLAIN: To estimate the causal effect of a treatment on an outcome
 - what is the effect of small classrooms on student performance?

Sample vs. Population

- ▶ We often want to know the characteristics of a large **population** such as the residents of a country
- ▶ Yet collecting data from every individual in the population is either prohibitively expensive or infeasible.
- ▶ In the US, we try to collect data from each individual every ten years
 - ▶ The 2020 census cost \$14.2 billion, approximately (the population at that time was around 331 mi)
 - ▶ This is not feasible for research purposes!
- ▶ We use surveys to collect data from a small subset of observations in order to understand the population

Sample vs. Population

- The subset of individuals chosen for study is called a **sample**



- Researchers typically survey about 1,000 people to infer the characteristics of more than 200 million US citizens
- $n=1,000$, $N=200$ million

Representative Samples

- ▶ In survey research, the sample needs to be representative of the population of interest
- ▶ **Representative sample:** Accurately reflects the characteristics of the population from which it is drawn
- ▶ If the sample is not representative, our inferences regarding the population characteristics will be wrong

Representative Samples

- ▶ Are you a representative sample of US residents?
- ▶ Are you a representative sample of UCSD students?
- ▶ Are you a representative sample of UCSD Poli majors?
- ▶ Are you a representative sample of POLI 30 D students?
- ▶ What would be the best way to draw a representative sample of UCSD students?

Representative Samples

- You should be careful about how representative your sample is:



Random Sampling

- ▶ The best way to draw a representative sample is to select individuals at *random* from the population.

Random sampling makes the **sample** and the **target population** *on average* identical.

- ▶ Random sampling: enables us to infer *valid* population characteristics from the sample,

Random Treatment Assignment vs. Random Sampling

- ▶ They both use a random process but are different concepts.
- ▶ **Random treatment assignment** means that treatment is assigned at random:
 - ▶ makes treatment and control groups comparable.
- ▶ **Random sampling** means that individuals are selected at random from the population into the sample:
 - ▶ makes the sample representative of the population.
- ▶ For this class, we assume we are always studying a representative sample.

Exploring One Variable At a Time

- ▶ Suppose we have collected data from a sample. Now what?
- ▶ To understand the content and distribution of each variable, we can:
 - ▶ Create a table of frequencies
 - ▶ Create a table of proportions
 - ▶ Create a histogram
 - ▶ Compute descriptive statistics
- ▶ Let us return to the voting experiment
 - ▶ data collected from a sample of registered voters in the state of Michigan

The *voting* dataset

Unit of observation: registered voters

Description of variables:

variable	description
<i>birth</i>	year of birth of registered voter
<i>message</i>	whether registered voter received message: "yes", "no"
<i>voted</i>	whether registered voter voted: 1=voted, 0=didn't vote

Table of Frequencies

- ▶ The **frequency table** shows the values the variable takes and the number of times each value appears in the variable
- ▶ R function: `table()`

```
table(voting$voted)
##
##      0      1
## 158276 71168
```

- ▶ Interpretation?

Table of Proportions

- ▶ The **table of proportions** shows the proportion of observations that take each value in the variable
- ▶ The proportions in the table should add up to 1
- ▶ R function: `prop.table(table())`

```
prop.table(table(voting$voted))
```

```
##
```

```
##           0           1
```

```
## 0.6898241 0.3101759
```


- ▶ Interpretation?

Histogram

- ▶ The **histogram** is the visual representation of a variables distribution through bins of different heights
- ▶ The position of the bins along the x-axis indicates the interval of values
- ▶ The height of the bins indicates the frequency (or count) of the interval of values
- ▶ R functions: `hist()` or `ggplot() + geom_histogram()`
- ▶ Great for quantitative variables (the numeric R data types)

Histogram

```
ggplot(voting, aes(x = birth)) +  
  geom_histogram(binwidth = 5) + theme_light()
```




► Interpretation?

Barplots

- ▶ The **barplot** is similar to a histogram, but discretizes the variation
- ▶ The position of the bins along the x-axis indicates a value
- ▶ The height of the bins indicates the frequency (or count) of the values
- ▶ R functions: `barplot(table())` or `ggplot() + geom_bar()`
- ▶ Great for qualitative variables (numeric binary or character)

Barplots

```
ggplot(voting, aes(x = voted)) +  
  geom_bar(aes(x = as.character(voted))) + theme_light()
```



► Interpretation?

Descriptive Statistics

- ▶ The **descriptive statistics** of a variable numerically summarizes the main characteristics of its distribution
- ▶ Measures of centrality
(center of the distribution):
 - ▶ mean
 - ▶ median
- ▶ Measures of spread
(amount of variation from the center):
 - ▶ standard deviation
 - ▶ variance

Mean

- ▶ The **mean** of a variable equals the sum of the values across all observations divided by the total number of observations
- ▶ What is the function in R?
- ▶ Example:

```
mean(voting$birth)
## [1] 1956.18
mean(voting$voted)
## [1] 0.3101759
```

- ▶ Interpretations?

Median

- ▶ The **median** of a variable is the value at the midpoint of the distribution that divides the data into two equal-size groups
- ▶ When the variable contains an odd number of observations, the median is the middle value of the distribution
- ▶ When the variable contains an even number of observations, the median is the average of the two middle values

Median

- ▶ Example, if $X = \{10, 4, 6, 8, 22\}$, what is the median of X ?
 - ▶ First, we need to sort the values of X in ascending order (as they would be in the distribution):
 $\{4, 6, 8, 10, 22\}$
 - ▶ The value in the middle of the distribution is 8 so the median is 8.
- ▶ R function: `median()`

```
median(voting$birth)
## [1] 1956
```

- ▶ Interpretations?

Standard Deviation

- The **standard deviation** of a variable is a measure of the spread of its distribution

$$sd(X) = \sqrt{\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n}}$$

- $sd(X)$ stands for the standard deviation of X
- X_i is a particular observation of X
- \bar{X} stands for the mean of X
- n is the total number of observations in the variable
- $\sum_{i=1}^n (X_i - \bar{X})^2$ means the sum of all $(X_i - \bar{X})^2$ from $i = 1$ to $i = n$

Standard Deviation

The **standard deviation** of variable measures the average distance of the observations to the mean.

- ▶ The larger the standard deviation, the flatter the distribution
- ▶ It gives us a sense of the range of the data, especially when dealing with bell-shaped distributions
- ▶ In bell-shaped (normal) distributions, 95% of the observations fall within two standard deviations from the mean

Standard Deviation

- ▶ R function: `sd()`

```
sd(voting$birth)
## [1] 14.46019
```

- ▶ If *birth* were normally distributed, about 95% of the registered voters would have been born between 1927 and 1985:
 - ▶ $\bar{X} - 2 \times \text{sd}(X) = 1956 - 2 \times 14.5 = 1927$
 - ▶ $\bar{X} + 2 \times \text{sd}(X) = 1956 + 2 \times 14.5 = 1985$

Variance

- ▶ Another measure of the spread of the distribution
- ▶ The **variance** of a variable is simply the square of the standard deviation

$$\text{var}(X) = [\text{sd}(X)]^2$$

- $\text{var}(X)$ stands for the variance of X
- $\text{sd}(X)$ stands for the standard deviation of X

Variance

- ▶ R function: `var()`

```
var(voting$birth)
## [1] 209.0971
```

- ▶ Alternatively: `sd()^2`

```
sd(voting$birth)^2
## [1] 209.0971
```

- ▶ We are usually better off using standard deviations as our measure of spread:
- ▶ Same unit of measurement as the variable

Summary

- ▶ **Today's Class:**
 - ▶ Sample vs. Population
 - ▶ Representative Samples and Random Sampling
 - ▶ Exploring a single variable
- ▶ **Next class:**
 - ▶ Correlations
 - ▶ Scatter-plots

Questions?

See you in the next class!