

BACHELOR OF COMPUTER APPLICATIONS (BCA_NEW)

Course Code: BCS-012

Course Title: BASIC MATHEMATICS

Assignment Number: BCA (I)/012/Assignment/2024-25

Last Dates for Submission: 31st October, 2024 (For July Session)

: 30th April, 2025 (For January Session)

Maximum Marks: 100

Weightage : 25%

Please **NOTE THAT** these solutions/answers of **BCS-012 (July 2024 and January 2025 session)** are just the general answers to show the format of the answers of this assignment. So solve or check this assignment by yourself for submitting, It will help you to understand the subject **more better** and it will also **help** you in your exams. You can take the **references** from this PDF **AS THIS SOLUTION SET IS JUST LIKE “TEMPLATE”** for the BCS-012 assignment's solution.

This will take the 15-20 or more Double side ruled page. This assignment is made with the help of resources available on the internet as well as the textbook of BCS-012 (IGNOU book).

NOTE: THIS ASSIGNMENT IS AVAILABLE FOR STUDENTS **FREE**, IF ANYONE RE-UPLOAD THIS ASSIGNMENT, GIVE PROPER CREDITS TO “**VIDEO OF EDUCATION VOE**” YOUTUBE CHANNEL.

GENERAL INSTRUCTION:

- QUESTIONS ARE WRITTEN AS **EXAMPLE QUESTION?** (IN **BLACK BALL PEN**)
- ANSWERS ARE WRITTEN AS **EXAMPLE ANSWER.** (IN **BLUE BALL PEN**)
- **DO NOT COPY** THE EXACT ANSWER FROM THIS SOLVED ASSIGNMENT.
- THERE IS THE **TIMESTAMP** FOR EACH ANSWER IN THE DESCRIPTION AND COMMENT BOX FOR SAVING YOUR PRECIOUS TIME.
- THIS SOLVED ASSIGNMENT IS SOLELY PROVIDED TO **HELP** THE IGNUO BCA STUDENTS.

- THIS PDF LINK IS GIVEN IN THE DESCRIPTION BOX FOR DOWNLOAD. IF YOU WANT A **PDF** FILE WITHOUT ANY WATERMARK AND WITH GRAPHS YOU CAN **BUY** IT AND HELP US TO CREATE MORE ASSIGNMENTS IN LESS TIME BY SUPPORTING US OR YOU CAN USE THE ANSWERS FROM THIS VIDEO TO SOLVE YOUR ASSIGNMENT, DON'T FORGET TO SHARE THIS VIDEO IN YOUR BCA GROUPS AND HELP YOUR FRIENDS.
- FOR MAKING THIS PDF WE TAKE THE QUESTIONS FROM **INTERNET** AND PREVIOUS **SOLVED ASSIGNMENT** FROM THE INTERNET AND WE DID **NOT** SAY THAT THE ANSWERS OF THESE QUESTIONS ARE 100% CORRECT WE FIND THE ANSWERS FROM INTERNET SOURCES, **SO PLEASE READ THE ANSWERS CAREFULLY BEFORE YOU USE.**
- IF YOU DID NOT UNDERSTAND THE ANSWER, THE PHOTO IS BLURED/NOT CLEAR OR YOU FIND ANYTHING IS WRONG OR YOU WANT TO BUY THIS ASSIGNMENT PDF THEN **DM ME ON INSTAGRAM** (ID: **@the_vor_**) OR LINK IN **CHANNEL INFO**.
- OUR PREVIOUS VIDEO IS AVAILABLE FREE ON YOUTUBE, LINK IS [**BEVAE-181: ENVIRONMENTAL STUDIES- SOLVED ASSIGNMENT- BCA_NEW**](#) PLAYLIST OF 1ST SEM BCA_NEW SOLVED ASSIGNMENTS [**IS 1ST SEM SOLVED ASSIGNMENT**](#) AND THE 2ND SEM BCA SOLVED ASSIGNMENT PLAYLIST IS [**2ND SEM BCA SOLVED ASSIGNMENT**](#)
- [**1ST SEMESTER SOLVED ASSIGNMENT PLAYLIST IS PLAYLIST**](#)
- [**2ND SEMESTER SOLVED ASSIGNMENT PLAYLIST IS PLAYLIST**](#)
- OUR **CHANNEL** LINK IS [**VIDEO OF EDUCATION VOE**](#)
- PLEASE **SUBSCRIBE** OUR CHANNEL FOR MORE INFORMATIVE VIDEOS, **NEXT VIDEO WILL BE UPLOAD WHEN WE HAVE 92 SUBSCRIBERS ON THE CHANNEL.**
- NEXT VIDEO IS **SOLVED ASSIGNMENT** FOR JUNE 2024 SESSION, SO **STAY TUNE** WITH US.
- The sources of this PDF are **Wikipedia pages**, BCS-012 **textbook** and some other **online sources**.
- If something is **wrong** or **mistype** just tell us so we can correct that mistake.
- **IF YOU WANT TO BUY BCS-012 WATERMARK FREE AND UNIQUE SOLVED ASSIGNMENT (IMAGE format), DM ME ON INSTAGRAM.**

Enjoy your learning.



BCS-013-Basic Mathematics

Q-1: For what value of 'K' the points $(-K+1, 3K)$, $(K, 3-3K)$ and $(-4-K, 6-3K)$ are collinear.

Ans: The given points are collinear if

$$\begin{vmatrix} -K+1 & 3K & 1 \\ K & 3-3K & 1 \\ -4-K & 6-3K & 1 \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} K & 3-3K & 1 \\ -3K+1 & 4K-3 & 0 \\ -4-3K & 4 & 0 \end{vmatrix} = 0 \quad \left[\begin{matrix} R_2 \rightarrow R_2 - R_1 \\ R_3 \rightarrow R_3 - R_1 \end{matrix} \right]$$

$$\Rightarrow 4(-3K+1) - (-4-3K)(4K-3) = 0$$

$$\Rightarrow (1-3K)(4-8-4K) = 0$$

$$\Rightarrow (1-3K)(-4-4K) = 0$$

$$\Rightarrow (1-3K)(K+1) = 0$$

$$\Rightarrow \boxed{K = \frac{1}{3} \text{ or } K = -1}$$

$\Rightarrow -1$ and $\frac{1}{3}$ are the value of 'K' at which points are collinear.

Q-2 Solve the following system of equations by using matrix Inverse method.

$$3x + 4y + 7z = 14$$

$$2x - y + 3z = 4$$

$$3x + 3y - 3z = 0$$

Ans: These equations can be written in a matrix as:

$$\begin{bmatrix} 3 & 4 & 7 \\ 2 & -1 & 3 \\ 3 & 3 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 14 \\ 4 \\ 0 \end{bmatrix}$$

OR

$$AX = B$$

$$A = \begin{bmatrix} 3 & 4 & 7 \\ 2 & -1 & 3 \\ 3 & 3 & -3 \end{bmatrix}, \quad X = \begin{bmatrix} X \\ Y \\ Z \end{bmatrix} \text{ and } B = \begin{bmatrix} 14 \\ 4 \\ 0 \end{bmatrix}$$

$$\begin{aligned} |A| &= 3 \begin{vmatrix} -1 & 3 & -4 \\ 3 & -3 & 3 \\ 3 & 3 & -1 \end{vmatrix} = 3(3(-6) - 4(-6 - 6)) + 7(4 + 3) \\ &= 3(-3) - 4(-13) + 7(6) \\ &= -9 + 48 + 42 \\ |A| &= 81 \end{aligned}$$

So for the Unique ... solution we used

$$X = A^{-1}B$$

$$A^{-1} = \frac{\text{adj}(A)}{|A|}$$

Cofactors of A are :-

$C_{11} = (-1)^{1+1} \cdot$	$3-6 = -3 = -3$
$C_{31} = (-1)^{3+1} \cdot$	$-13-14 = -(-36) = 26$
$C_{31} = (-1)^{3+1} \cdot$	$13+7 = 19 = 19$
$C_{13} = (-1)^{1+2} \cdot$	$-6-6 = -(-12) = 12$
$C_{23} = (-1)^{2+3} \cdot$	$-9-14 = +(-23) = 23$
$C_{32} = (-1)^{3+2} \cdot$	$9-14 = -(-5) = 5$
$C_{13} = (-1)^{1+3} \cdot$	$4+3 = 6 = 6$
$C_{23} = (-1)^{2+3} \cdot$	$6-8 = -(-2) = 2$
$C_{33} = (-1)^{3+3} \cdot$	$-3-8 = -11 = -11$

$$\text{adj } A = \begin{bmatrix} -3 & 13 & 6 \\ 26 & -23 & 2 \\ 19 & 5 & -11 \end{bmatrix}^T = \begin{bmatrix} -3 & 26 & 19 \\ 13 & -23 & 5 \\ 6 & 2 & -11 \end{bmatrix}$$

$$\text{Now, } X = A^{-1}B = \frac{1}{81} \begin{bmatrix} -3 & 36 & 19 \\ 13 & -23 & 5 \\ 6 & 3 & -11 \end{bmatrix} \begin{bmatrix} 14 \\ 4 \\ 0 \end{bmatrix}$$

$$X = \frac{1}{81} \begin{bmatrix} -3(14) + 36(4) + 19(0) \\ 13(14) - (-23)(4) + 5(0) \\ 6(14) + 3(4) + (-11)(0) \end{bmatrix}$$

$$X = \frac{1}{81} \begin{bmatrix} 62 \\ 76 \\ 92 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 62/81 \\ 76/81 \\ 92/81 \end{bmatrix}$$

So, $x = \frac{62}{81}$, $y = \frac{76}{81}$, and $z = \frac{92}{81}$ are the solution

Q-3 Use principle of mathematical induction to prove that
 $\frac{1}{1 \times 3} + \frac{1}{2 \times 3} + \dots + \frac{1}{n(n+1)} = \frac{n}{n+1}$

Sol:-

Let

$$P(n) = \frac{1}{1 \times 3} + \frac{1}{2 \times 3} + \dots + \frac{1}{n(n+1)} = \frac{n}{n+1}$$

$$\text{For } n = 1, \text{ LHS} = \frac{1}{1 \times 3} = \frac{1}{3}; \text{ RHS} = \frac{1}{1+1} = \frac{1}{2}$$

$$\therefore \text{LHS} = \text{RHS}$$

$\therefore P(1)$ is True

Let us assume $P(K)$ is true for some $K \in N$

$$\text{i.e., } \frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \dots + \frac{1}{K(K+1)} = \frac{K}{K+1}$$

Adding $(K+1)^{\text{th}}$ term $= \frac{1}{(K+1)(K+2)}$ on both sides,

We get,

$$\frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \dots + \frac{1}{K(K+1)} + \frac{1}{(K+1)(K+2)}$$

$$= \frac{K}{K+1} + \frac{1}{(K+1)(K+2)}$$

$$= \frac{K(K+2) + 1}{(K+1)(K+2)} \Rightarrow \frac{K^2 + 2K + 1}{(K+1)(K+2)}$$

$$= \frac{(K+1)^2}{(K+1)(K+2)} = \frac{K+1}{K+2} \Rightarrow \frac{K+1}{(K+1)+1}$$

which is $P(K+1)$

Thus, $P(K) \Rightarrow P(K+1)$

Hence, by mathematical induction $P(n)$ is true for all $n \in N$.

Q-4: How many terms of G.P $\sqrt{3}, 3, 3\sqrt{3}, \dots$

Add upto $39+13$

$$\text{Sol: } a = \sqrt{3}, r = \frac{3}{\sqrt{3}} = \sqrt{3}$$

$$S_n = \frac{a(r^n - 1)}{r - 1}$$

$$= 39+13 = \frac{\sqrt{3}(\sqrt{3}^n - 1)}{\sqrt{3} - 1}$$

$$53 = \frac{\sqrt{3}(\sqrt{3}^n - 1)}{\sqrt{3} - 1}$$

$$\begin{aligned}
 53(\sqrt[7]{3}-1) &= \sqrt[7]{3}(\sqrt[7]{3^n}-1) \\
 53\sqrt[7]{3}-53 &= \sqrt[7]{3}(\sqrt[7]{3^n}-1) \\
 53\sqrt[7]{3}-53+\sqrt[7]{3} &= \sqrt[7]{3^{n+1}}-\sqrt[7]{3} \\
 53\sqrt[7]{3}-53 &= \sqrt[7]{3^{n+1}}
 \end{aligned}$$

Now, $53\sqrt[7]{3}-53 \approx (\sqrt[7]{3})^6$
 Let's put $(\sqrt[7]{3})^6$ in our eq

$$(\sqrt[7]{3})^6 = \sqrt[7]{3^{n+1}}$$

Comparing the power

$$\begin{array}{|c|l}
 \hline
 6 & = n+1 \\
 \hline
 n & = 5 \\
 \hline
 \end{array}$$

So, 5 terms needed.

Q-5: If $y = ae^{mx} + be^{-mx}$, prove that $\frac{d^2y}{dx^2} = m^2y$
Solt: let $y = ae^{mx} + be^{-mx}$ — eq(1)
 Differentiate with respect to x ,

$$\frac{dy}{dx} = ame^{mx} - b m e^{-mx} \quad \left[\because \frac{d}{dx} e^x = e^x \right]$$

Differentiate again with respect to x .

$$\frac{d^2y}{dx^2} = am^2e^{mx} + bm^2e^{-mx}$$

$$\Rightarrow \frac{d^2y}{dx^2} = m^2(ae^{mx} + be^{-mx})$$

$$\Rightarrow \frac{d^2y}{dx^2} = m^2y \quad [\text{from 1}]$$

Hence proved.

Q-6: Integrate function $f(x) = \frac{x}{[(x+1)(3x-1)]}$ w.r.t x .

Sol

We will use partial fraction decomposition

$$= \frac{x}{(x+1)(3x-1)} = \frac{A}{x+1} + \frac{B}{3x-1} \quad \text{--- eq (1)}$$

Taking RHS and by adding both fractions, we get

$$\frac{A}{x+1} + \frac{B}{3x-1} = \frac{A(3x-1) + B(x+1)}{(x+1)(3x-1)}$$

from eq (1)

$$\frac{x}{(x+1)(3x-1)} = \frac{\cancel{A}(3x-1) + \cancel{B}(x+1)}{(x+1)(3x-1)}$$

$$\Rightarrow x = A(3x-1) + B(x+1)$$

$$\Rightarrow x = (3A)x - A + Bx + B$$

$$\Rightarrow x = (3A+B)x + (-A+B)$$

Now, match the coefficients of same terms
For the x coefficient

$$1 = 3A + B \quad \text{--- eq (2)}$$

For the constant term :

$$0 = -A + B \quad \text{--- eq (3)}$$

From eq (3)

$$B = A$$

Substitute $B = A$ into the eq (2) :

$$1 = 3A + A \Rightarrow 1 = 3A = A = \frac{1}{3}$$

Since $B = A$:

$$B = \frac{1}{3}$$

Now put the value of A and B in the eq(1) 7

$$\frac{x}{(x+1)(3x-1)} = \frac{1/3}{x+1} + \frac{1/3}{3x-1}$$

Taking RHS

$$\frac{1}{3} \left(\frac{1}{x+1} \right) + \frac{1}{3} \left(\frac{1}{3x-1} \right)$$

To integrate;

$$\begin{aligned} & \frac{1}{3} \int \frac{1}{x+1} dx + \frac{1}{3} \int \frac{1}{3x-1} dx \\ &= \frac{1}{3} \left[\ln|x+1| + (1/3)dx + C \right] \quad [\text{by substitution method}] \end{aligned}$$

$$V = 3x-1$$

$$dU = 3dx$$

Hence

$$dx = \frac{du}{3}$$

So,

$$\int \frac{1}{3x-1} dx = \int \frac{1 \cdot du}{3 \cdot 3} = \frac{1}{3} \ln|u| = \frac{1}{3} \ln|3x-1| + C$$

$$\Rightarrow \frac{1}{3} (\ln|x+1| + \frac{1}{3} \ln|3x-1|) + C$$

$$\Rightarrow \boxed{\frac{1}{3} \ln|x+1| + \frac{1}{6} \ln|3x-1| + C}$$

Q-7: If $1, \omega, \omega^2$ are Cube Roots of unity show that $(1+\omega)^2 - (1+\omega)^3 + \omega^2 = 0$.

Sol =

$$(1+\omega)^2 = 1^2 + 2 \cdot 1 \cdot \omega + \omega^2 = 1 + 2\omega + \omega^2$$

$$(1+\omega)^3 = (1+\omega)(1+\omega^2) = (1+\omega)(1+2\omega+\omega^2)$$

$$\begin{aligned} &= (1+\omega)(1+2\omega+\omega^2) = 1 + 2\omega + \omega^2 + \omega(1+2\omega+\omega^2) \\ &= 1 + 2\omega + \omega^2 + \omega + 2\omega^2 + \omega^3 \end{aligned}$$

$$= 1 + 3w + 3w^2 + w^3$$

$$\Rightarrow 1 + 3w + 3w^2 + 1$$

[Property of cube root]
 $w^3 = 1$

$$= 3w^2 + 3w + 2$$

Substitute the expanded forms into the original eq

$$(1+w)^2 - (1+w)^3 + w^2 = (1+3w+w^2) - (2+3w+3w^2) + w^2$$

Taking RHS

$$\begin{aligned} &= (1+3w+w^2) - (2+3w+3w^2) + w^2 \\ &= 1+3w+w^2 - 2 - 3w - 3w^2 - w^2 \\ &= (1-2) + (3w-3w) + (w^2-3w^2-w^2) \\ &= -1 - w - w^2 \end{aligned}$$

Use the property $1+w+w^2 = 0$

$$-(1+w+w^2) = 0$$

our RHS is 0 so the expression is

$$(1+w)^2 - (1+w)^3 + w^2 = 0$$

Q-8: If α, β are roots of equation $3x^2 - 3x - 5 = 0$, then find a quadratic equation whose roots are α^2, β^2 .

Sol: α and β are two roots of the given quadratic equation $3x^2 - 3x - 5 = 0$

$$\Rightarrow 3x^2 - 3x - 5 = 0$$

$$= 3x^2 - 5x + 3x - 5 = 0$$

$$= x(3x-5) + 1(3x-5) = 0$$

$$= (3x-5)(x+1) = 0$$

$$= 3x-5 = 0 \quad \text{and} \quad x+1 = 0$$

$$\therefore x = \frac{5}{3} \text{ and } x = -1$$

$$\therefore \alpha = \frac{5}{3} \text{ and } \beta = -1$$

$$\alpha^2 = \left(\frac{5}{3}\right)^2 = \frac{25}{9} \quad \text{and} \quad \beta^2 = (-1)^2 = 1$$

\therefore New roots are $\frac{5}{4}$ and 1

New quadratic equation,

$$(x - \frac{5}{4})(x - 1) = 0$$

$$x^2 - x - \frac{35x}{4} + \frac{5}{4} = 0$$

$$4x^2 - 4x - 35x + 5 = 0$$

$$\Rightarrow 4x^2 - 39x + 5 = 0$$

Q-9 : Solve the inequality $\frac{3}{5}(x-3) \leq \frac{5}{3}(3-x)$ and set.

Sol:

$$\frac{3(x-3)}{5} \leq \frac{5(3-x)}{3}$$

$$3 \times 3(x-3) \leq 5 \times 5(3-x)$$

$$9(x-3) \leq 25(3-x)$$

$$9x - 27 \leq 75 - 25x$$

$$9x - 27 + 25x \leq 75 - 25x + 18 + 25x$$


$$\therefore 34x \leq 68$$

$$x \leq \frac{68}{34}$$

$$x \leq 2$$

Hence, x is a real number which less than or equal to 2.
Hence, $x \in (-\infty, 2]$ is the solution.

Graph of the solution set



Q-10: If positive number exceeds its positive square root by 13, then find the number.

Sol - Let the required number be x .

Given: Number is greater than its positive square root]

$$= x - 13 = \sqrt{x}$$

Squaring on both sides, we get

$$(x-13)^2 = x$$

$$x^2 + 13^2 - 2x(13) = x$$

$$= x^2 - 25x + 144 = 0$$

$$x(x-16) - 9(x-16) = 0$$

$$(x-16)(x-9) = 0$$

$$\therefore x = 16, x = 9$$

As $9 < 13$, it was given that the number will be greater than 13.


So, $x = 16$, is the required number.

Q-11: Find the area bounded by the curves $y = x^2$ and $y = x$

11

Sol

By solving equation $y = x^2$ and $y = x$



$$\therefore x^2 = x$$

$$\therefore x^2 - x = 0$$

$$x(x-1) = 0$$

$$\therefore x = 0 \text{ or } x = 1$$

Points of intersection of both curves are $(0, 0)$ and $(1, 1)$

Required area A

$$= \int_a^b [f_1(x) - f_2(x)] dx = \int_0^1 [x - x^2] dx$$

$$= \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = \frac{1}{2} - \frac{1}{3} = \frac{1}{6}$$

$$A = |I| = \frac{1}{6} \text{ sq unit.}$$

Q-12: find the inverse of the matrix $A =$

$$\begin{bmatrix} 1 & 6 & 4 \\ 3 & 4 & -1 \\ -1 & 3 & 5 \end{bmatrix}, \text{ if it exists.}$$

Sol- Let's first find the determinant to check if inverse exists or not.

$$|A| = \begin{vmatrix} 1 & 6 & 4 \\ 3 & 4 & -1 \\ -1 & 3 & 5 \end{vmatrix}$$

$$|A| = 1 \begin{vmatrix} 4 & -1 \\ 3 & 5 \end{vmatrix} - 6 \begin{vmatrix} 3 & -1 \\ -1 & 5 \end{vmatrix} + 4 \begin{vmatrix} 3 & 4 \\ -1 & 3 \end{vmatrix}$$

$$\begin{aligned} & |(20+30) - 6(10-1) + 4(4+4)| \\ &= 22 - 54 + 32 \\ &\quad 54 - 54 \end{aligned}$$

$$|A| = 0$$

So,

Determinant is zero, therefore inverse matrix doesn't exist.

Q-13: If m times the n^{th} term of an A.P. is n times its n^{th} term, show that $(m+n)^{\text{th}}$ term of the A.P. is zero.

Sol: Given,

$$n^{\text{th}} \text{ term of AP} = t_n = a + (n-1)d$$

$$m^{\text{th}} \text{ term of AP} = t_m = a + (m-1)d$$

$$\Rightarrow mt_m = nt_m$$

$$m[a + (m-1)d] = n[a + (n-1)d]$$

$$m[a + (m-1)d] - n[a + (n-1)d] = 0$$

$$a(m-n) + d[m(m+n)(m-n) - (m-n)] = 0$$

$$(m-n)[a + d((m+n) - 1)] = 0$$

$$a + [(m+n)-1]d = 0$$

$$\text{But } t_{m+n} = a + [(m+n)-1]d$$

$$\therefore t_{m+n} = 0$$

Q-14: Show that

i) $\lim_{x \rightarrow 0} \frac{|x|}{x}$ does not exist

Sol

$$\text{LHL} = \lim_{h \rightarrow 0} \frac{|0-h|}{0-h} = \lim_{h \rightarrow 0} \frac{h}{-h} = -1$$

RHL

$$\lim_{h \rightarrow 0} \frac{|0+h|}{0+h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1.$$

LHL \neq RHL∴ Limit does not exist at $x=0$.(ii) $f(x) = |x|$ is continuous at $x=0$
solto check the continuity of $f(x)$ at $x=a$,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} x = 0$$

and

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (-x) = 0 = f(0)$$

$$\text{Now, } Rf(0) = Lf(0) = f(0)$$

So, the function $f(x)$ is continuous at $x=0$.

Q-15: Suriti wants to invest at most 13000 in Saving certificates and National Saving Bonds. She has to invest at least 3000 in Saving certificates and at least 4000 in National Saving Bonds. If Rate of Interest on Saving certificates is 8% per annum and ROI on National Saving bond is 10% annum. How much money should she invest to earn maximum yearly income? Find also the maximum yearly income.

Ans:

Let Suriti invests Rs x in Saving certificates and
Rs y in National bonds.

Therefore,

$$x, y \geq 0$$

Suriti wants to invest at most Rs 12000 in Saving Certificates and National Saving bonds.

$$x+y \leq 12000$$

According to rules, she has to invest at least Rs 3000 in Saving Certificates and at least Rs 4000 in National Saving Bonds.

$$x \geq 3000$$

$$y \geq 4000$$

If the rate of interest on Saving certificate is 8% per annum and the rate of interest on National Saving Bond is 10% per annum.

Total earning from investment =

$$Z = \frac{8x}{100} + \frac{10y}{100} \text{ which is to be maximized.}$$

Thus, the mathematical formulation of the given linear programming problem is

$$\max Z = \frac{8x}{100} + \frac{10y}{100}$$


Subject to

$$\begin{aligned} x+y &\leq 19000 \\ x &\geq 3000 \\ y &\geq 4000 \end{aligned}$$

$$x, y \geq 0$$

First we will convert inequations in equations
 $x+y = 19000, x=3000, y=4000, x=0, y=0$

Region represented by $x+y \leq 19000$:



The line $x+y=13000$ meets the coordinate axes at $A(13000, 0)$ and $B(0, 13000)$ respectively. By joining these points we obtain the line $x+y=13000$. Clearly $(0,0)$ satisfies the inequation $x+y \leq 13000$. So, the region which contains the origin represents the solution set of the inequation $x+y \leq 13000$.

Region represented by $x \geq 3000$:

The line $x=3000$ is the line that passes through $(3000, 0)$ and is parallel to y axis. The region to the right of the line $x=3000$ will satisfy the inequation

Region represented by $y \geq 4000$:

The line $y=4000$ is the line that passes through $(0, 4000)$ and is parallel to x axis. The region above the line $y=4000$ will satisfy the inequation $y \geq 4000$.

Region represented by $x \geq 0$ and $y \geq 0$:

Since, every point in the first quadrant satisfies these inequations. So, the first quadrant is the region represented by the inequations $x \geq 0, y \geq 0$.

The feasible region determined by the system of constraints is

The corner points are $E(3000, 10000)$, $D(8000, 4000)$, $C(3000, 4000)$

The values of Z at these points are as follows

corner points	$Z = (8x/100) + (10y/100)$
E	1160
D	1040
C	560

The maximum value of Z is 1160 which is attained at $E(3000, 10000)$.

Thus, the maximum earning is Rs 1160 obtained when Rs 3000 were invested in Saving's certificate and Rs 10000 were invested in National Saving Bond.

Q-16: A spherical balloon is being inflated at the rate of $900 \text{ cm}^3/\text{Sec}$. How fast Radius of the balloon Increasing when the Radius is 15 cm.

Sol: Let r be the radius of the balloon and V be its volume at any time t .

$$\Rightarrow V = \frac{4}{3} \pi r^3$$

Differentiate both sides w.r.t t , we get,

$$\frac{dV}{dt} = \frac{4}{3} \pi \cdot 3r^2 \frac{dr}{dt}$$

$$\Rightarrow \frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

It is given that $\frac{dV}{dt} = 900 \text{ cm}^2/\text{sec}$

$$\Rightarrow 900 = 4\pi r^2 \frac{dr}{dt}$$

$$= \frac{dr}{dt} > \frac{995}{\pi r^2}$$

When $r = 15 \text{ cm}$

$$\Rightarrow \frac{dr}{dt} = \frac{225}{\pi \times (15)^2}$$

$$\Rightarrow \frac{dr}{dt} = \frac{1}{\pi} \text{ cm/s}$$

Radius of the balloon is increasing at the rate of $\frac{1}{\pi}$ cm/s or 0.3183 cm/s

When the radius is 15 cm.