References

J. Aeles, F. Horst, S. Lapuschkin, L. Lacourpaille, and F. Hug, 'Revealing the unique features of each individual's muscle activation signatures', <i>The Royal Society Publishing Interface</i> , vol. 18, no. 174, p. 20200770, Jan. 2021, doi: 10.1098/rsif.2020.0770.
[2] M. J. Antony <i>et al.</i> , 'Classification of EEG Using Adaptive SVM Classifier with CSP and Online Recursive Independent Component Analysis', <i>Sensors (Basel)</i> , vol. 22, no. 19, p. 7596, Oct. 2022, doi: 10.3390/s22197596.
A. Biasiucci <i>et al.</i> , 'Brain-actuated functional electrical stimulation elicits lasting arm motor recovery after stroke', <i>Nature Communications</i> , vol. 9, no. 2421, June 2018, doi: 10.1038/s41467-018-04673-z.
[4] A. Buerkle, W. Eaton, N. Lohse, T. Bamber, and P. Ferreira, 'EEG based arm movement intention recognition towards enhanced safety in symbiotic Human-Robot Collaboration', <i>Robotics and Computer-Integrated Manufacturing</i> , vol. 70, p. 102137, Aug. 2021, doi: 10.1016/j.rcim.2021.102137.
[5] R. Byrne, 'Development of a Low Cost, Open-source, Electroencephalograph-Based Brain-Computer Interface'. Dublin Institute of Technology, May 28, 2018. Accessed: Jan. 10, 2025. [Online]. Available: https://github.com/RonanB96/Low-Cost-EEG-Based-BCI/blob/master/Thesis/Development%20of%20a%20Low%20Cost%2C%20Open-source%2C%20Electroencephalograph-Based%20Brain-Computer%20Interface.pdf
[6] M. Conti <i>et al.</i> , 'Brain Functional Connectivity in de novo Parkinson's Disease Patients Based on Clinical EEG', <i>Frontiers in Neurology</i> , vol. 13, p. 844745, Mar. 2022, doi: 10.3389/fneur.2022.844745.
[7] A. Cuesta-Gómez, F. Molina-Rueda, M. Carratala-Tejada, E. Imatz-Ojanguren, D. Torricelli, and J. C. Miangolarra-Page, 'The Use of Functional Electrical Stimulation on the Upper Limb and Interscapular Muscles of Patients with Stroke for the Improvement of Reaching Movements: A Feasibility Study', <i>Frontiers in Neurology</i> , vol. 8, May 2017, doi: 10.3389/fneur.2017.00186.
[8] K. Desai, 'Parkinson's Disease Detection via Resting-State Electroencephalography Using Signal Processing and Machine Learning Techniques', <i>arxiv</i> , Mar. 2023, doi: 10.48550/arXiv.2304.01214.
[9] B. M. Doucet, A. Lam, and L. Griffin, 'Neuromuscular Electrical Stimulation for Skeletal Muscle Function', <i>Yale Journal of Biology and Medicine</i> , vol. 85, no. 2, pp. 201–215, June 2012.
[10] S. Farashi, A. Sarihi, M. Ramezani, S. Shahidi, and M. Mazdeh, 'Parkinson's disease tremor prediction using EEG data analysis-A preliminary and feasibility study', <i>BMC Neurology</i> , vol. 23, p. 420, Nov. 2023, doi: 10.1186/s12883-023-03468-0.
(11)

[1]

F. Haddad, M. Sawalha, Y. Khawaja, A. Najjar, and R. Karaman, 'Dopamine and Levodopa Prodrugs for the Treatment of Parkinson's Disease', *Molecules*, vol. 23, no. 1, p. 40, Dec. 2017, doi: 10.3390/molecules23010040.

[12]

F. Karimi, J. Niu, K. Gouweleeuw, Q. Almeida, and N. Jiang, 'Movement-related EEG signatures associated with freezing of gait in Parkinson's disease: an integrative analysis', *Brain Communications*, vol. 3, no. 4, p. fcab277, Nov. 2021, doi: 10.1093/braincomms/fcab277.

[13]

M. B. Kebaetse, A. E. Turner, and S. A. Binder-Macleod, 'Effects of stimulation frequencies and patterns on performance of repetitive, nonisometric tasks', *Journal of Applied Physiology*, vol. 92, no. 1, pp. 109–116, Jan. 2002, doi: 10.1152/jappl.2002.92.1.109.

[14]

M. A. Khan *et al.*, 'A systematic review on functional electrical stimulation based rehabilitation systems for upper limb post-stroke recovery', *Frontiers in Neurology*, vol. 14, no. 1272992, Dec. 2023, doi: 10.3389/fneur.2023.1272992.

[15]

N. Kueper *et al.*, 'EEG and EMG dataset for the detection of errors introduced by an active orthosis device', *Frontiers in Human Neuroscience*, vol. 18, p. 1304311, Jan. 2024, doi: 10.3389/fnhum.2024.1304311.

[16]

D. Liao, F. Li, D. Lu, and P. Zhong, 'Activation of Piezo1 Mechanosensitive Ion Channel In HEK293T Cells by 30 MHz Vertically Deployed Surface Acoustic Waves', *Biochemical and Biophysical Research Communications*, vol. 518, no. 3, pp. 541–547, Oct. 2019, doi: 10.1016/j.bbrc.2019.08.078.

[17]

M. Mahmoodi, Bahador Makkiabadi, M. Mahmoudi, and S. Sanei, 'A new method for accurate detection of movement intention from single channel EEG for online BCI', *Computer Methods and Programs in Biomedicine Update*, vol. 1, p. 100027, Aug. 2021, doi: 10.1016/j.cmpbup.2021.100027.

[18]

A. Miladinović *et al.*, 'EEG changes and motor deficits in Parkinson's disease patients: Correlation of motor scales and EEG power bands', *Procedia Computer Science*, vol. 192, pp. 2616–2623, 2021, doi: 10.1016/j.procs.2021.09.031.

[19]

C. R. Oehrn *et al.*, 'Chronic adaptive deep brain stimulation versus conventional stimulation in Parkinson's disease: a blinded randomized feasibility trial', *Nature Medicine*, vol. 30, pp. 3345–3356, Aug. 2019, doi: doi:oio.org/10.1038/s41591-024-03196-z.

[20]

M. Popovic, T. Thrasher, M. Adams, V. Takes, V. Zivanovic, and M. Tonack, 'Functional electrical therapy: retraining grasping in spinal cord injury', *Spinal Cord*, vol. 44, pp. 143–151, Mar. 2006, doi: 10.1038/sj.sc.3101822.

[21]

A. Saikia, A. Saikia, Amit Ranjan Barua, and S. Paul, 'EEG-EMG Correlation for Parkinson's disease', *International Journal of Engineering and Advanced Technology (IJEAT)*, vol. 8, no. 6, Aug. 2019, doi: 10.35940/ijeat.F8360.088619.

[22]

V. del C. Silva-Acosta, I. Román-Godínez, S. Torres-Ramos, and R. A. Salido-Ruiz, 'Automatic estimation of continuous elbow flexion-extension movement based on

electromyographic and electroencephalographic signals', *Biomedical Signal Processing and Control*, vol. 70, p. 102950, Sept. 2021, doi: 10.1016/j.bspc.2021.102950.

[23]

P. Solla, F. Marrosu, and M. G. Marrosu, 'Therapeutic interventions and adjustments in the management of Parkinson disease: Role of combined carbidopa/levodopa/entacapone (Stalevo)', *Neuropsychiatric Disease and Treatment*, vol. 6, no. 1, pp. 483–490, July 2010.

[24]

A. A. P. Suárez, S. B. Batista, I. P. Ibáñez, E. C. Fernández, M. F. Campos, and L. M. Chacón, 'EEG-Derived Functional Connectivity Patterns Associated with Mild Cognitive Impairment in Parkinson's Disease', *Behavioral Sciences (Basel)*, vol. 11, no. 3, p. 11030040, Mar. 2021, doi: 10.3390/bs11030040.

[25]

M. Sun *et al.*, 'FES-UPP: A Flexible Functional Electrical Stimulation System to Support Upper Limb Functional Activity Practice', *Frontiers in Neuroscience*, vol. 12, no. 449, July 2018, doi: 10.3389/fnins.2018.00449.

[26]

H.-C. Wang, A. J. Lees, and P. Brown, 'Impairment of EEG desynchronisation before and during movement and its relation to bradykinesia in Parkinson's disease', *Journal of Neurology, Neurosurgery & Psychiatry*, vol. 66, no. 4, pp. 442–446, Apr. 1999, doi: 10.1136/jnnp.66.4.442.

[27]

R. A. Zanini, E. L. Colombin, and M. C. F. de Castro, 'Parkinson's disease EMG signal prediction using Neural Networks', 2019 IEEE International Conference on Systems, Man and Cybernetics (SMC), pp. 2446–2453, Sept. 2019.

[28]

D. Zhang, C. Hansen, F. D. Fréne, S. P. Kærgaard, and W. Qian, 'Automated labeling and online evaluation for self-paced movement detection BCI', *Knowledge-Based Systems*, vol. 265, p. 110383, Feb. 2023, doi: 10.1016/j.knosys.2023.110383.

[29]

L. Zhang *et al.*, 'Activation of Piezo1 by ultrasonic stimulation and its effect on the permeability of human umbilical vein endothelial cells', *Biomedicine & Pharmacotherapy*, vol. 131, p. 110796, Nov. 2020, doi: 10.1016/j.biopha.2020.110796.