



New estimates of weight-at-size, maturity-at-size, fecundity, and biomass of snow crab, *Chionoecetes opilio*, in the Arctic Ocean off Alaska



Lauren M. Divine^{a,*}, Franz J. Mueter^b, Gordon H. Kruse^b, Bodil A. Bluhm^{a,c}, Stephen C. Jewett^a, Katrin Iken^a

^a College of Fisheries and Ocean Sciences, University of Alaska Fairbanks, P.O. Box 757220, Fairbanks, AK, 99775, United States

^b College of Fisheries and Ocean Sciences, University of Alaska Fairbanks, 17101 Point Lena Loop Rd., Juneau, AK, 99801, United States

^c Department of Arctic and Marine Biology, UiT- The Arctic University of Norway, 9037, Tromsø, Norway

ARTICLE INFO

Handled by A.E. Punt

Keywords:

Arctic Fishery Management Plan (arctic FMP)

Beaufort Sea

Chukchi Sea

Female reproductive potential

Maturity

Sustainable yield

ABSTRACT

Snow crab (*Chionoecetes opilio*) were identified as a potential future target fishery species in federal waters of the Arctic Ocean off Alaska by the Arctic Fishery Management Plan (Arctic FMP) in 2009, but this plan currently prohibits commercial harvest until sufficient information is available to assess a sustainable commercial fishery. One drawback of the current Arctic FMP is that critical population and biomass estimates were based on limited data. Collaborative research efforts in the Chukchi and Beaufort seas over the past decade have yielded a much richer database on snow crab in the Arctic. Using these data, we generated new estimates of weight-at-size, maturity-at-size, fecundity, and biomass to recalculate sustainable yield of snow crab in the U.S. Arctic. Weight-at-size was generally similar for male and female snow crab between the Chukchi and Beaufort seas, with males reaching overall larger sizes than females in both seas and largest male crabs occurring in the Beaufort Sea. Compared with snow crab in other geographic regions, 50% morphometric maturity was reached at a slightly smaller size in the Chukchi Sea; low sample sizes in the Beaufort Sea prevented maturity-at-size analysis. Fecundity-at-size in the Chukchi Sea was similar to known values estimated for snow crab in other regions. Estimated total reproductive output, using fecundity estimates obtained here, suggest that local reproduction may be sufficient to account for a large portion of observed small juvenile benthic snow crab abundances; further investigation is warranted to determine whether Chukchi and Beaufort populations are self-sustaining at this time. Although snow crab had high abundances in the Chukchi Sea, harvestable biomass of male snow crab only occurred in the Beaufort Sea because crab larger than the minimum marketable size (≥ 100 mm carapace width, based on Bering Sea metric) were absent in the Chukchi Sea over the study period. Our biomass estimates in the Chukchi Sea were substantially higher than previous estimates, owing at least in part to high abundances of small crab that were greatly under-sampled with the large-mesh gear such as was used in surveys referenced in the Arctic FMP. Estimates of biomass and sustainable yield for the Beaufort Sea were over twice as high as previous estimates in the Arctic FMP, but harvestable biomass was largely limited to the slope (> 200 m depth) and is unlikely to support commercial harvest. Our results expand overall understanding of arctic snow crab dynamics in light of potential future fisheries or other, non-fishing activities and inform the management of the Alaskan Arctic stock.

1. Introduction

Snow crab *Chionoecetes opilio* are distributed across subarctic and Arctic waters, with lucrative commercial fisheries occurring in the North Atlantic (eastern Canada and Greenland), Bering Sea, and Sea of Japan. The eastern Bering Sea (EBS) fishery is one of the most valuable fisheries in the US, with an average harvest of 25,700 metric tons (56.6 million pounds) and an average ex-vessel value of \$133 million dollars

in 2017 (Alaska Department of Fish and Game, 2018; Fissel et al., 2018). Harvest is limited to only males at or above a marketable size (≥ 100 mm carapace width [CW]) in an attempt to protect the reproductive potential of mature females (Sainte-Marie and Gilbert, 1998; Zheng and Kruse, 2006; Turnock and Rugolo, 2012). In 1999, the EBS snow crab fishery was declared overfished after a period of declining snow crab biomass and abundance (Turnock and Rugolo, 2012). Additionally, a northward contraction of the center of snow crab

* Corresponding author.

E-mail address: lmdivine@alaska.edu (L.M. Divine).

<https://doi.org/10.1016/j.fishres.2019.05.002>

Received 31 August 2018; Received in revised form 1 May 2019; Accepted 3 May 2019

Available online 13 June 2019

0165-7836/ © 2019 Published by Elsevier B.V.

distribution in the EBS in response to changing oceanographic conditions was identified shortly after overfishing was recognized (Zheng et al., 2001; Orensanz et al., 2004). The combination of the biomass decline, distributional shift, and high abundances of snow crab north of the Bering Strait (North Pacific Fishery Management Council (NPFMC, 2009; Kolts et al., 2015) raised fisheries concerns about the possibility of this species moving out of the primary EBS fishing grounds and northward into Arctic waters (Mueter and Dawe, 2012; Hollowed et al., 2013). In response to a potential future snow crab fishery in the Alaskan Arctic, the Arctic Fishery Management Plan (Arctic FMP) was developed in 2009 by the North Pacific Fishery Management Council (North Pacific Fishery Management Council (NPFMC, 2009). The Arctic FMP relied on limited available data (from 96 sampling stations in the Chukchi Sea and 26 sampling stations in the Beaufort Sea) and borrowed life history metrics from other regions to evaluate total and harvestable biomass of snow crab in the Chukchi and Beaufort seas.

The Arctic FMP currently presumes exploitable snow crab biomass to be low in the Alaskan Arctic, although snow crab are dominant members of benthic communities with respect to biomass and abundance in the Chukchi Sea, and occur at commercial sizes (≥ 100 mm CW based on the standard for the EBS) in the western Beaufort Sea (Bluhm et al., 2009; Rand and Logerwell, 2011; Ravelo et al., 2014, 2015). In the Chukchi Sea, snow crab contribute significantly to overall invertebrate biomass, even at small maximum sizes (i.e., 74 mm CW, Frost and Lowry, 1983; Paul et al., 1997; Rand and Logerwell, 2011; Konar et al., 2014; Ravelo et al., 2014; Gross et al., 2017). The potential commercial exploitation of snow crab in the Arctic requires detailed, region-specific knowledge of stock structure and life history. However, snow crab in the Chukchi and Beaufort seas have not been historically monitored with any regularity due to the lack of dedicated federal monitoring programs and challenges of sampling the vast, seasonally ice-covered shelves and slopes of these regions. Limited available data (i.e., Frost and Lowry, 1983; Paul et al., 1997) suggest that snow crab abundance has increased on the Chukchi shelf by at least an order of magnitude over the last few decades, although snow crab sizes remain small (maximum size: 74 mm CW, Gross et al., 2017). Abundance in the Beaufort Sea has also increased (Paul et al., 1997; Logerwell and Rand, 2010), and the reported maximum size of snow crab in the Beaufort Sea has increased from 75 mm CW in the 1970s (Frost and Lowry, 1983) to 119 mm CW in the 2000s (Rand and Logerwell, 2011), although expanded study areas within in this region may account for observation of larger snow crabs.

A clear understanding of growth and maturation processes, population structure, and fecundity is essential for effective management of snow crab fisheries (Comeau and Conan, 1992; Sainte-Marie et al., 1995), including the development of sustainable harvest limits (North Pacific Fishery Management Council (NPFMC, 2009). Snow crab growth and allometry has been best-studied in geographic regions where snow crab comprise an important commercial fishery resource (e.g., Gulf of St. Lawrence: Watson, 1970; Haynes et al., 1976; Sainte-Marie et al., 1995; Newfoundland: Comeau and Conan, 1992; Comeau et al., 1998; EBS: Haynes et al., 1976; Otto, 1998; Orensanz et al., 2007; Ernst et al., 2012). Population data from the North Atlantic, chiefly the Gulf of St. Lawrence, are currently applied to growth estimates of snow crab in less-studied arctic regions, such as the northern Bering, Chukchi, and Beaufort seas (Kolts et al., 2013; Gross et al., 2017). Snow crab undergo a series of molts during which they exhibit discrete increases in body size (Hartnoll, 1982). Individuals undergo a final, terminal molt to maturity between instars X–XIII (≥ 54 mm CW based on crab in Canadian Atlantic), but size at terminal molt is highly variable among individuals (Conan and Comeau, 1986; Comeau and Conan, 1992; Comeau et al., 1998; Sainte-Marie et al., 1995, , Burmeister and Sainte-Marie, 2010). Somatic growth for males and females is similar at small sizes (i.e., benthic instars I–VII, Comeau et al., 1998), but male molt increments are larger at later stages, with terminal molt occurring earlier in females, resulting in larger maximum sizes for males relative

to females (Alunno-Bruscia and Sainte-Marie, 1998). Growth models have established mean size-at-instar for each stage, e.g., in the northern Atlantic (Sainte-Marie et al., 1995) and the EBS (Orensanz et al., 2007; Ernst et al., 2012). These growth models have been widely applied to other stocks across the geographic range of snow crab (e.g., northern Bering Sea: Kolts et al., 2013, Chukchi Sea: Gross et al., 2017). In cold waters ($< 0^\circ\text{C}$) snow crab may have smaller growth increments per molt, longer intermolt periods, or skip a molt (Orensanz et al., 2007; Burmeister and Sainte-Marie, 2010), all of which can contribute to slow growth and complicate application of growth models from subarctic to Arctic populations.

Maturity-at-size is a critical determinant of reproductive output and rate of population growth in brachyuran crabs (Stearns, 1976; Hines, 1982). Male snow crab exhibit an allometric increase in chela height (CH) relative to body size (CW) during the terminal molt (Conan and Comeau, 1986) and this relationship is used to determine maturity status for males. However, gonad development occurs prior to terminal molt and males may be sexually mature prior to exhibiting the change in chela allometry (Sainte-Marie et al., 1995). Females experience a marked increase in the size of their abdominal flap to accommodate an egg clutch after their terminal (maturity) molt (Watson, 1970; Moriyasu et al., 1987; Alunno-Bruscia and Sainte-Marie, 1998). In a latitudinal gradient from the southeastern Bering Sea to the Chukchi Sea, size at 50% maturity in female snow crab decreases, possibly due to a decrease in temperature (Armstrong et al., 2008, 2010, Kolts, 2012). This latitudinal pattern of smaller individuals occurring at higher latitudes has also been established for both males and females in the Gulf of St. Lawrence (Sainte-Marie and Gilbert, 1998; Dawe et al., 2012), Greenland (Burmeister and Sainte-Marie, 2010), and Newfoundland (Dawe et al., 2012).

Female fecundity (average number of eggs in a clutch) and lifetime female fecundity (cumulative number of eggs produced in a lifetime), and their contributions to population-level reproductive output, are important metrics regarding stock productivity and also in understanding distribution patterns and changes in crab stocks (Armstrong et al., 2008). Female fecundity of brachyuran crabs is a function of body size, with larger females producing larger egg clutches (Orensanz et al., 2007; Kolts et al., 2015). In the eastern and northern Bering Sea, mean annual cold bottom temperatures ($\leq 1^\circ\text{C}$) result in a shift from an annual to a biennial breeding cycle for female snow crab (Moriyasu and Lanteigne, 1998; Rugolo et al., 2005; Webb et al., 2007; Sainte-Marie et al., 2008; Kolts et al., 2015). For snow crab in the U.S. Arctic off Alaska, it is likely that at least a portion of female snow crab also experience biennial brooding given that bottom temperatures $\leq 1^\circ\text{C}$ are regularly recorded across the shelves (Grebmeier et al., 2015; Danielson et al., 2017). It is unknown if mature biomass is sufficient to sustain local populations in the Chukchi and Beaufort seas, regardless of whether annual or biennial brooding cycles dominate, or if advected larvae from southern regions at least partially support, or merely add to, local production (Bluhm et al., 2015). No mature females with fertilized egg clutches have been collected in the Beaufort Sea to date (Bluhm et al., 2015), indicating no evidence of local reproduction in this region.

The goal of the present study was to provide new information on snow crab life history and distribution trends in the Chukchi and Beaufort seas, using a synthesis of available data collected over 2004–2017. Specifically, we estimated (1) individual weight-at-size for snow crab in the Chukchi and Beaufort seas, (2) maturity-at-size and fecundity-at-size for snow crab in the Chukchi Sea only, due to low sample sizes in the Beaufort Sea, (3) potential reproductive capacity of mature females in the Chukchi Sea only, and (4) regional biomass, abundance, and sustainable yield for snow crab in the Chukchi and Beaufort seas. Our estimates of weight-at-size, maturity-at-size, and fecundity improve previous biomass assessments by applying region-specific population metrics as much as possible. We then used these direct parameter estimates to revise total and harvestable biomass

Table 1

Cruise data showing the timing of sampling, project name, gear type used, and data provided for the current study. PSBT = plumb staff beam trawl, EBT = Nor'Eastern bottom trawl. Major funding sources indicated by superscripts.

| Month/ Year | Project | Gear Type | # hauls (by gear type) | # crab collected | Data Provided |
|---------------------|---|-----------|------------------------|------------------|---|
| <i>Chukchi Sea</i> | | | | | |
| Aug 2004 | RUSALCA-1 ^a | PSBT | 4 | 58 | Biomass, abundance, size frequency |
| Aug 2007 | Oscar Dyson ^a | PSBT | 11 | – | Biomass, abundance |
| Aug 2008 | Oshoru Maru IPY ^b | PSBT | 15 | – | Biomass, abundance |
| Aug 2009 | RUSALCA-2 ^a | PSBT | 3 | 1,298 | Biomass, abundance, size frequency, fecundity |
| Aug 2009 | COMIDA ^c | PSBT | 30 | – | Biomass, abundance |
| Aug 2010 | COMIDA ^c | PSBT | 23 | 290 | Biomass, abundance, size frequency, fecundity |
| Sept 2010 | CSESP ^d | PSBT | 70 | 479 | Biomass, abundance, size frequency |
| Aug-Sept 2010 | AKMAP ^e | PSBT | 30 | 505 | Biomass, abundance, size frequency |
| Sept 2011 | AKMAP ^e | PSBT | 28 | 49 | Biomass, abundance, size frequency |
| Aug 2012 | RUSALCA-3 ^a | PSBT | 4 | 1,766 | Biomass, abundance, size frequency, fecundity |
| Aug-Sept 2012 | Arctic Eis ^c | PSBT, EBT | 40, 71 | 493 | Biomass, abundance, size frequency, fecundity, maturity-at-size |
| Aug-Sept 2015 | AMBON ^{a, c, d} | PSBT | 67 | 6,975 | Biomass, abundance, size frequency |
| Aug 2017 | AMBON ^{a, c, d, g} | PSBT | 81 | 234 | Biomass, abundance, size frequency |
| <i>Beaufort Sea</i> | | | | | |
| Aug-Sept 2008 | Western Beaufort Fish Survey ^c | PSBT | 21 | 334 | Biomass, abundance |
| Aug-Sept 2011 | BeauFish Survey ^{c, f} | PSBT | 83 | 64 | Biomass, abundance, size frequency |
| Sept 2012 | Transboundary ^c | PSBT | 18 | 50 | Biomass, abundance, size frequency |
| Aug-Sept 2013 | Transboundary ^c | PSBT | 42 | 1 | Biomass, abundance, size frequency |
| Aug-Sept 2014 | Transboundary ^c | PSBT | 10 | – | Biomass, abundance |
| Aug 2014 | ANIMIDA ^c | PSBT | 29 | 3 | Biomass, abundance, size frequency |
| Aug 2015 | ANIMIDA ^c | PSBT | 17 | 33 | Biomass, abundance, size frequency |

Major funding sources: ^a NOAA: National Oceanographic and Atmospheric Administration; ^b Japanese Funding, ^cBOEM: Bureau of Ocean Energy Management (in 2008 as Minerals Management Service), ^d Oil Industry (Shell Oil, Conoco Phillips, Statoil), ^e ADEC: Alaska Department of Environmental Conservation, and CIAP: Coastal Impact Assistance Program, ^f CMI: Coastal Marine Institute, ^g National Science Foundation.

estimates originally published for snow crab in the Arctic FMP (North Pacific Fishery Management Council (NPFMC, 2009)).

2. Methods

2.1. Snow crab collections

This project used a combination of snow crab data collected from a total of 20 surveys, 13 in the Chukchi Sea and 7 in the Beaufort Sea, between 2004 and 2017 (Table 1, Fig. 1). Samples and data were used in various combinations to address project objectives (Table 1), depending on availability. A vast majority of crab for this study were collected with a plumb staff beam trawl (PSBT, modified from Gunderson and Ellis, 1986, see details on towing methods in Norcross et al., 2015, Table 1). The PSBT had a 2.257 m opening and a net mesh of 7 mm with a 4 mm cod end liner. Tow duration lasted from 1 to 6 min at speeds ranging from 0.5 to 0.75 m s⁻¹. Crab were also collected using an 83–112 eastern bottom trawl (EBT, Table 1). The EBT is the standard net for the National Marine Fisheries Service, Alaska Fisheries Science Center (AFSC) bottom trawl surveys in the EBS. The EBT had a 25.3 m headrope and 34.1 m footrope. During the Arctic Eis 2012 survey (Table 1), the EBT was fitted with a net mesh of 10.2 cm, with 8.9 cm intermediate and codend liners. The codend also had a 3.2 cm liner (Goddard et al., 2012). Tow duration for the EBT ranged from 5 to 15 min at an approximate speed of 1.5 m s⁻¹. Snow crab collected from all trawls were rinsed, sexed and counted, bulk weight per station was recorded using spring or digital hanging scales, and in some instances individual CW (standard measure of body size) was measured to the nearest 0.1 mm using digital Vernier calipers. Select crab were either frozen whole or preserved in formalin (RUSALCA 2009 only) for transport to the laboratory. Crab were then thawed (if frozen), blotted dry, and individually weighed to the nearest 0.1 g. For all males, chela height (CH; standard measure of maturity) was measured for the left chela to the nearest 0.1 mm using digital Vernier calipers consistent with Conan and Comeau (1986). Occasionally, crab experienced limb loss during collections or after freezing, or were collected with missing limbs. If limbs could be preserved with the whole crab they were. A

small number of crabs had missing limbs that were not able to be recovered. In these cases, the weight for each missing limb was estimated from the matching limb on the other side of the same crab or the corresponding leg of a crab of a similar size and applied to the total weight to account for missing limbs.

2.2. Weight-at-size

As a proxy of individual weight-at-size of crab, we used the allometric relationship between weight (individual biomass, g) and size (mm CW) of measured crab. Individual wet weight (g) and size (mm CW) for males and females collected in the Chukchi and Beaufort seas were log-transformed and weight-at-size was estimated from a linear regression as:

$$\log(\text{Weight(g)}) = a + b \cdot \log(\text{CW(mm)})$$

where log is the natural logarithm, *a* is the intercept and *b* is the slope. Analyses of covariance (ANCOVA) with separate slopes by region were performed to determine if weight-at-size differed significantly between the Chukchi and Beaufort seas for males, immature females, and mature females, respectively. Females were separated into immature and mature status because mature females have terminally molted and allocate energy to reproductive tissue growth and production (i.e., gonad and egg clutches) rather than somatic growth (Alunno-Bruscia and Sainte-Marie, 1998).

Size frequency distributions (SFD) for all size data available, pooled across years, were examined separately for the Chukchi and Beaufort seas to characterize the size ranges of male and female crab in each region. We were unable to fit mixture models to objectively distinguish modes in the SFD for snow crab in either region and, therefore, did not attempt to interpret modal peaks observed in SFD histograms as instars.

2.3. Mean individual crab weight across survey area

To examine spatial variability in the size of snow crab across the Chukchi and Beaufort shelves, mean weight by haul was modeled separately within each sea using a thin-plate regression spline smoother




Fig. 1. Stations sampled on the Alaska Chukchi and Beaufort Sea shelves from 2004 to 2017. Stations denoted as black circle data points occurring in the Chukchi and Beaufort seas are stations that contributed to biomass estimates in the Arctic Fishery Management Plan (Barber et al., 1994; North Pacific Fishery Management Council (NPFMC, 2009; Rand and Logerwell, 2011).

within a Generalized Additive Modeling (GAM) framework. Mean individual weight in each haul containing crab was first computed by dividing the total weight of snow crab in a given haul (measured in the field as bulk weight, kg) by the number of snow crab caught. Individual weights were then fourth-root transformed to normalize residuals and were modeled as:

$$W_{it} = \alpha + s(x_{it}, y_{it}) + \varepsilon_{it}$$

where W_{it} is the fourth-root transformed mean weight at station i during year t , α is the overall intercept, s denotes a smooth function of sampling location described by distances x and y (in km) from an arbitrary origin, and ε_{it} are residuals that are assumed to be normally distributed with a zero mean and variance σ^2 . The estimated mean sizes were then mapped over the study regions. We also considered a mixed effects model with a random-year effect to account for differences in mean size by year. However, the random-year effect did not result in a better model fit based on the Akaike Information Criterion and years were, therefore, pooled in the analysis.

2.4. Maturity-at-size

Maturity-at-size for male crab was determined as the proportion of males exhibiting morphometric allometry in the left chela at a given body size (Conan and Comeau, 1986). This was only possible for Chukchi Sea crab as sample sizes were too low in the Beaufort Sea. To assign a breakpoint for male maturity, observed male CW and left CH were plotted against several published regression relationships (i.e., from the EBS, Kolts et al., 2013; and eastern Canadian Atlantic, Comeau et al., 1998). The EBS data presented the best fit for allometric male snow crab data in the Chukchi Sea; thus, mature males were distinguished from juvenile and adolescent males by using the established dividing line of $CH = 0.191 \times CW$ (Kolts et al., 2013). Female maturity-

at-size in the Chukchi Sea only was estimated as the proportion of females exhibiting a change in allometry in the abdomen relative to body size (i.e., large flap covers the entire ventral side of mature females, Paul et al., 1997) or presence of an egg clutch (Jewett, 1981). The proportion of mature female crab at a given size was estimated by fitting a logistic regression model with a logit link to binary maturity data (mature = 1, immature = 0) as a function of size:

$$\log(p_k/(1-p_k)) = \alpha + \beta \cdot CW_k(\text{mm}),$$

where p_k is the proportion of mature females in size class k , α and β are linear regression parameters on the logit scale, and the model was fit by maximizing the binomial likelihood. Size at 50% maturity was then estimated as $-\alpha/\beta$, corresponding to the size at the inflection point of the logistic curve.

To correct for gear bias in the size composition estimate of the individuals used to determine size-at-maturity in the Chukchi Sea, we compared the size frequencies of crab between the PSBT and EBT based on snow crab CW data from 33 paired trawls taken during the Arctic Eis 2012 cruise (Britt et al., 2013, Table 1). The paired trawl samples were distributed throughout the study area and are representative of the full latitudinal and depth range of the survey (Kotwicki et al., 2017). The observed size composition was summarized by gear type as the number of crab per km^2 in each 1 mm size bin. The effectiveness of the two nets for sampling snow crab in their path (catchability) is unknown and is likely to differ among crab sizes (size selectivity). Assuming each trawl is fully size selective (selectivity = 1) over some range of observed sizes (CW), the size selectivity of the two trawl types can be estimated under the following assumptions. We first used the common assumption that selectivity follows a logistic model of the form:

$$\text{selectivity} = \frac{1}{1 + e^{b(x-a)}}$$

where x is the carapace width (mm CW), parameter a corresponds to size at 50% selectivity, and b corresponds to the steepness of the curve. We further assumed that selectivity of the EBT approaches 1 for large crab (large mesh size) and selectivity may decrease with decreasing CW ($b > 0$), whereas the PSBT has full selectivity for small snow crab (small mesh size) and selectivity may decrease with increasing CW ($b < 0$).

We simultaneously estimated the parameters of the two selectivity curves by minimizing a weighted sum of squared differences between the predicted number of crab at size k from the PSBT and the predicted number of crab at size k from the EBT. Because the variability in numbers at size was not constant and increased with the number of crab (N_k) in size bin k , we applied appropriate weights to the squared differences before summing across size bins. We assumed that variance increases linearly with N_k based on the expected mean-variance relationship for Poisson-distributed count data. We then weighted the squared difference for each size bin by $1/N_k$, where N_k values were computed from a smoothed length-frequency distribution that was estimated by minimizing the unweighted sum of squares. If the number of crab in a given size bin was estimated to be less than 1 it was set equal to 1 to avoid assigning excessive weights to these bins. This selectivity analysis was used to determine a size range over which both gears had similar high selectivity to pool snow crab abundances and biomass across gear types. Estimated size selectivity was used to estimate the "true" size distribution of crabs in the Chukchi Sea by dividing the observed numbers at a given size by the estimated selectivity at that size. The methods used in the present study differ from previous biomass estimates in the Arctic FMP in that the Arctic FMP used only data collected with the EBT that were not corrected for selectivity (North Pacific Fishery Management Council (NPFMC, 2009)).

2.5. Fecundity and egg production

Mean fecundity at size was estimated using the number and weight of eggs taken from 322 mature females collected during the RUSALCA 2009 and 2012, COMIDA 2010, CSESP 2010 and Arctic EIS 2012 cruises in the Chukchi Sea (Table 1). Each total egg clutch was removed from the pleopods, 250 eggs were subsampled and dried at 60 °C to a constant weight, and the dry weight of the subsample was determined (Stichert, 2009; Webb et al., 2016). The remaining egg mass was also dried as described above to obtain total dry weight. Individual fecundity was then estimated by dividing the weight of the total egg mass by the average dry weight of the eggs in the sub-sample and multiplying by the number of eggs in the sub-sample ($n = 250$). Finally, a linear regression of the logarithm of total number of eggs (dependent variable) on log-transformed carapace width (CW, independent variable) for each crab was used to determine the relationship between body size and mean number of eggs in a clutch.

To estimate total egg production in the Chukchi Sea, we first estimated the abundance of mature females in the Chukchi Sea in 2012 based on the observed density of females at each station (number of females per km²) and the estimated maturity-at-size relationship. Only data from the 2012 gridded Arctic EIS survey, which had the most complete spatial coverage, were used for this estimate to avoid biases associated with temporal or spatial differences in distribution and fecundity. Station-specific densities of mature females for each 1 mm CW increment were first estimated as the proportion of mature females from the logistic maturity-at-size curve multiplied by the total density of females in a given size class (CW) at a station. The total abundance of mature females in the survey area by CW was then estimated by multiplying station-specific densities by the area of the grid cell represented by each station and summing abundances across all grid cells. For each size class (1 mm CW increments), we then multiplied the average fecundity at size (number of eggs per mature female) by the number of mature females in a given size class and summed the number of eggs across size classes for an estimate of total annual egg production.

2.6. Biomass, abundance, and sustainable yield

Biomass and abundance per unit effort at each station were estimated based on the weight and number of crab caught at each station and the area swept by the trawl. For the EBT, area swept was determined by multiplying the distance towed by the mean net spread. For the PSBT, the net width was assumed to be fixed at 2.257 m (Gunderson and Ellis, 1986), and area swept was equal to the net width multiplied by the distance towed as calculated during each tow. Catches were standardized by computing catch per unit effort (CPUE) for each haul in biomass (kg) or numbers per km². Crab densities (CPUE) were then estimated using a thin-plate regression spline smoother within a GAM framework (Wood, 2017). Sampling locations differed among years and cruises, but substantial overlap among cruises allowed us to estimate inter-annual differences in mean CPUE, assuming that the spatial patterns in CPUE were consistent over time. Therefore, we modeled CPUE as a function of year and location as follows:

$$CPUE_{it} = \alpha + a_t + s(x_{it}, y_{it}),$$

where $CPUE_{it}$ is the observed density at station i during year t , α is the overall intercept, a_t is a random intercept associated with year t to account for interannual differences in mean CPUE, and s denotes a smooth function of sampling location. To preserve distances in the east-west (x) and north-south (y) directions, locations were calculated as the great-circle distance, projected onto a plane, from an arbitrarily chosen origin in the southwest corner of the study area. Preliminary analyses suggested that biomass values were strongly right-skewed, included a number of zeros, and that the variance tended to increase with mean density. These features could best be modeled using a Tweedie distribution (Dunn and Smyth, 2005), where variance is given by the mean to the power p , and p is estimated but constrained to range from 1 (corresponding to a Poisson distribution) to 2 (gamma distribution). Models were fit separately to the Chukchi and Beaufort seas data and densities were predicted for the center of each grid cell in a 10 x 10 km grid overlaid on the survey area. Total biomass within each area was then estimated by multiplying the predicted mean density in each grid cell by 100 to obtain an estimate of total biomass within the 100 km² grid cell, then summing over all grid cells. Uncertainty in total biomass was estimated by repeatedly simulating predicted values for each grid cell from the posterior distribution of the smoothing parameter estimates and computing total biomass as described above. Estimates were obtained over the total survey area in the U.S. Chukchi Sea (172,000 km²), and the total survey area in the U.S. Beaufort Sea (64,400 km²). In addition, we obtained separate estimates for the U.S. Chukchi Sea south and north of Point Hope, Alaska, to account for potential differences between the southern and northern Chukchi Sea.

Estimates obtained here were considered pristine, or unfished, biomass for comparison with previous estimates from the Arctic FMP (North Pacific Fishery Management Council (NPFMC, 2009)). For comparison, we determined the equilibrium biomass at a given fishing mortality following the approach in the FMP, updated with new estimates for weight-at-size and maturity-at-size:

$$B(Fir) = \left[\left(\frac{h}{M+F} \right) \left(1 + \frac{1}{(M+F)d} \right) \right]^{\frac{1}{r}}$$

where h is the scale parameter in Cushing (1971) stock-recruitment relationship, M is the instantaneous natural mortality rate, F is the instantaneous fishing mortality, d is the difference between the age-at-maturity and the age intercept of the linear weight-at-age equation, and r is the amount of resilience implied by the stock recruitment relationship (North Pacific Fishery Management Council (NPFMC, 2009)). We used the default value of M for the Bering Sea snow crab stock ($M = 0.23$; Turnock and Rugolo, 2008) and obtained our own parameter estimate for d from this study, which resulted from direct estimates of weight-at-size and maturity-at-size (see above sections),

combined with previous literature for weight-at-age of snow crabs in the Chukchi Sea (Gross et al., 2017), and assuming an intercept of zero for the linear weight-at age-equation (North Pacific Fishery Management Council (NPFMC, 2009). We used the default value of M for the Bering Sea snow crab stock ($M = 0.23$; Turnock and Rugolo, 2008). To determine the amount of biomass available to future potential fisheries given the new biomass estimates, we determined sustainable yield (Y) as the product of F and equilibrium biomass:

$$Y(F|r) = F \cdot B(F|r).$$

We then obtained an estimate of the instantaneous fishing mortality that maximizes equilibrium yield as:

$$F_{MSY}(r) = \left(\frac{M}{2(1-r)} \right) \left(1 - \frac{2-r}{M^*d} + \sqrt{\left(\frac{(2-r)}{M^*d} \right)^2 + \frac{4-6r}{M^*d} + 1} \right) - M$$

The biomass at which a maximum sustainable yield (MSY) can be obtained was calculated as:

$$B_{MSY} = \text{Bratio}(F_{MSY}(r)|r) \cdot B_0,$$

where *Bratio* is the ratio of equilibrium biomass to unfished (pristine) biomass (B_0). Finally, MSY was estimated as:

$$MSY = Y_{ratio}(F_{MSY}(r)|r) \cdot B_0,$$

where *Yratio* is the ratio of sustainable yield to B_0 . We compared our estimates of biomass and sustainable yield to values in the Arctic FMP (North Pacific Fishery Management Council (NPFMC, 2009). We used the default value of M for the Bering Sea snow crab stock ($M = 0.23$; Turnock and Rugolo, 2008).

3. Results

3.1. Size-frequency-distributions and weight-at-size

Although males were much less abundant in the Beaufort Sea than in the Chukchi Sea (Fig. 2A-C), the largest Chukchi male (86 mm CW, Fig. 2B) was approximately 33 mm smaller than the largest male collected in the Beaufort Sea (119 mm CW, Fig. 2C). The largest female snow crab in the Chukchi Sea were also smaller than males (66 mm CW, Fig. 3A), and they reached larger maximum sizes in the Beaufort Sea than females in the Chukchi Sea (83 mm CW, Fig. 3B). Immature females in the Chukchi Sea ranged from 4 to 58 mm CW and mature females ranged from 21 to 66 mm CW (overlapping by 37 mm CW, Fig. 3). In the Beaufort Sea, immature females ranged from 16 to 70 mm CW and mature females ranged from 37 to 82 mm CW (overlapping by 33 mm CW; Fig. 3). We collected high numbers of early benthic instars (i.e., 3–20 mm CW) in the Chukchi Sea (Figs. 2A and 3A); few small, presumably juvenile, crab were collected in the Beaufort Sea (Figs. 2C and 3B).

From 2004–2017, approximately 3,416 snow crab were individually measured and weighed across the Chukchi and Beaufort seas. Weight-at-size was similar for males between the Chukchi and Beaufort seas (ANCOVA, $p = 0.72$, Fig. 4A-B) as well as for pooled immature and mature females between the Chukchi and Beaufort seas (ANCOVA, $p = 0.63$, Fig. 4C-D). However, females in the Chukchi Sea reached heavier weights than Beaufort Sea females at larger sizes (ANCOVA, $p < 0.0001$ for both immature and mature females). When considering immature versus mature female weight-at-size within each region, we found contrasting patterns. Mature females in the Chukchi Sea achieved heavier weights at a given size than immature females of the same size (ANCOVA, $p < 0.0001$), but the opposite pattern occurred in the Beaufort Sea (ANCOVA, $p < 0.001$), with heavier weights at a given size achieved by immature females rather than mature females. Low sample sizes in the Beaufort Sea and the use of two preservation methods with differential dehydration effects (i.e., frozen versus




Fig. 2. Size frequency distributions for males (A) ≤ 13 mm CW in the Chukchi Sea, (B) > 13 mm CW in the Chukchi Sea collected from 2004 to 2015, and (C) > 13 mm CW collected in the Beaufort Sea from 2008 to 2015. Note the different scales of the y-axes. Black dashed line indicates the minimum marketable size (≥ 100 mm CW) applicable for the Bering Sea commercial snow crab fishery.

formalin preservation) likely impacted our estimates of weight-at-age for the Beaufort Sea.

3.2. Mean individual crab weight across survey area

Snow crab were generally much larger in the Beaufort Sea with the largest individuals observed along the outer shelf and slope at depth (Fig. 5). On average, the largest individuals in the Chukchi Sea were found off the Lisburne Peninsula west of Point Hope, Alaska (Fig. 5), with smaller individuals in the southern Chukchi Sea along the coast and on the northern portions of the shelf. This spatial pattern accounted for only 18% of the variability in Chukchi Sea mean individual weight, suggesting considerable small-scale spatial variability of individual snow crab weight. In contrast, about 62% of variability in fourth-root transformed individual weights in the Beaufort Sea were explained by the model due to a well-defined size gradient with smaller crab in nearshore waters and larger crab along the outer shelf and slope at deeper depths.

3.3. Maturity-at-size

For the Chukchi Sea crab, we estimated that 50% of male snow crab reach morphometric maturity at 62 mm CW, based on allometry of CH (Fig. 6A). Female snow crab in the Chukchi Sea achieved 50% morphometric maturity at 46 mm CW based on allometry of the abdominal flap and/or the presence of an egg clutch (Fig. 6B). Size at 50% maturity could not be estimated for the Beaufort Sea because of low sample sizes.




Fig. 3. Size frequency distributions for immature and mature females (A) in the Chukchi Sea collected from 2004 to 2015, and (B) immature and mature females collected in the Beaufort Sea from 2008 to 2015. Note the different scales of the y-axes.




Fig. 4. Weight-at-size regressions of A) Chukchi Sea males ($\text{weight(g)} = 3.0 \cdot \text{CW} - 3.5$), B) Beaufort Sea males ($\text{weight(g)} = 2.8 \cdot \text{CW} - 3.2$), C) Chukchi immature ($\text{weight(g)} = 2.8 \cdot \text{CW} - 3.1$) and mature females ($\text{weight(g)} = 2.8 \cdot \text{CW} - 3.2$), and D) Beaufort immature ($\text{weight(g)} = 2.8 \cdot \text{CW} - 3.1$) and mature females ($\text{weight(g)} = 2.3 \cdot \text{CW} - 2.3$) (estimated from snow crabs collected from 2004 to 2015). Weight-at-size parameters (a = intercept and b = slope) and sample sizes are presented in the respective figures.




Fig. 5. Predicted individual mean snow crab weight (g, log-scale) in the Chukchi and Beaufort seas based on thin-plate regression splines fit separately to the Chukchi Sea ($R^2 = 18\%$) and Beaufort Sea ($R^2 = 62\%$), estimated from PSBT CPUE data pooled over multiple surveys conducted between 2004 and 2015. Crosses denote all stations where snow crab weights were collected.




Fig. 6. Size at 50% morphometric maturity (dark gray solid lines) with 95% confidence intervals (light gray shaded areas) for (A) males and (B) females estimated from snow crab collected in the Chukchi Sea. Size at which 50% of all crab are morphometrically mature is denoted at the intersection of the gray dashed lines.

3.4. Fecundity

Mature females collected for fecundity estimation in the Chukchi Sea ranged from 38 to 65 mm CW (Table 2). Mean number of eggs increased with body size (Linear regression, number of eggs = $1,063.7 \cdot (\text{CW}) - 34,397$, $R^2 = 0.50$, Fig. 7) from 7,092 eggs at 38 mm CW to 49,164 eggs at 65 mm CW. Total egg production in the Chukchi Sea was estimated at just over 6.59 billion eggs for 2012.

Table 2

Mean eggs per clutch scaled to snow crab body size (carapace width, CW) and proportion of female snow crabs at each CW that are mature in the Chukchi Sea for the range of mature females observed in this region (38–65 mm CW).

| CW (mm) | mean # eggs per clutch | proportion mature |
|---------|------------------------|-------------------|
| 38 | 7,092 | 0.19 |
| 39 | 8,231 | 0.22 |
| 40 | 9,505 | 0.25 |
| 41 | 8,917 | 0.29 |
| 42 | 13,258 | 0.33 |
| 43 | 12,489 | 0.37 |
| 44 | 12,307 | 0.41 |
| 45 | 13,188 | 0.46 |
| 46 | 14,765 | 0.50 |
| 47 | 16,167 | 0.55 |
| 48 | 15,592 | 0.59 |
| 49 | 16,339 | 0.63 |
| 50 | 17,387 | 0.67 |
| 51 | 20,363 | 0.71 |
| 52 | 21,282 | 0.75 |
| 53 | 22,380 | 0.78 |
| 54 | 21,700 | 0.81 |
| 55 | 23,223 | 0.83 |
| 56 | 24,441 | 0.86 |
| 57 | 27,881 | 0.88 |
| 58 | 26,138 | 0.90 |
| 59 | 16,105 | 0.91 |
| 60 | 36,477 | 0.93 |
| 61 | 34,044 | 0.94 |
| 62 | 31,611 | 0.94 |
| 63 | 33,720 | 0.95 |
| 64 | 31,923 | 0.96 |
| 65 | 42,597 | 1.00 |

3.5. Biomass, abundance, and sustainable yield

We estimated total snow crab biomass in both the Chukchi and Beaufort seas (Table 3; Fig. 8) to be substantially higher than previous estimates from the Arctic FMP (Table 4). Our estimate of snow crab biomass in the Beaufort Sea was less than 10% that of Chukchi total biomass (63,577 mt and 746,596 mt, respectively), whereas the Arctic FMP estimated biomass in the Beaufort Sea (29,731 mt) to be 45% of the Chukchi Sea biomass (66,491 mt). All these estimates were considered to be estimates of pristine, or unfished, biomass (Tables 3 and 4).

We estimated annual harvestable biomass (males ≥ 100 mm CW) to




Fig. 7. Fecundity as number of eggs plotted against body size (carapace width) per mature female snow crab collected from the Chukchi Sea from 2009 to 2012. Number of eggs in a clutch = $2.9 \cdot CW - 1.61$, $R^2 = 0.50$.

Table 3

Biomass estimates (metric tons, mt) with 95% lower (LCI) and upper (UCI) confidence intervals over different portions of the survey area. Northern and southern Chukchi Sea correspond to areas north or south of Point Hope, Alaska, USA (see Fig. 1 for reference).

| Region | Area (km ²) | Estimate (mt) | LCI (mt) | UCI (mt) |
|---------------------|-------------------------|---------------|----------|-----------|
| Chukchi Sea (total) | 172,000 | 746,596 | 442,139 | 1,591,622 |
| Chukchi Sea (North) | 138,000 | 411,357 | 252,351 | 814,453 |
| Chukchi Sea (South) | 34,000 | 335,239 | 163,584 | 880,476 |
| Beaufort Sea | 64,400 | 63,577 | 31,878 | 167,613 |

be zero in the Chukchi Sea and 15,894 mt in the Beaufort Sea; therefore, equilibrium biomass and yield values were only computed for the Beaufort Sea. With our updated estimate of the difference between the age-at-maturity and the age intercept of the linear weight-at-age equation ($d = 7.0$ this study; $d = 8.0$ in the Arctic FMP, North Pacific Fishery Management Council (NPFMC, 2009)), we estimated that yield is maximized at a fishing mortality of $F = 0.34$ compared to $F = 0.36$ in the Arctic FMP. Together, these parameters resulted in a biomass at MSY estimate of $B_{MSY} = 2,681$ mt for the Beaufort Sea, which is approximately twice as high as the Arctic FMP value of $B_{MSY} = 1,268$ mt (Table 4). Finally, we estimated $MSY = 905$ mt, about 200% higher than the Arctic FMP-calculated MSY (453 mt, Table 4).




Fig. 8. Predicted mean snow crab biomass (kg km^{-2} , log-scale) in the Chukchi and Beaufort seas, estimated from PSBT CPUE data pooled over multiple surveys conducted between 2004 and 2017. Symbols show hauls with (filled circle) and without (cross) snow crab. Black lines delineate geographic areas over which total biomass was estimated (i.e., southern and northern Chukchi Sea, Beaufort Sea; Table 1).

Table 4

Estimates of total and harvestable biomass in the Chukchi and Beaufort seas as determined in the Arctic FMP (North Pacific Fishery Management Council (NPFMC, 2009)) and based on the revised estimates (this study) for snow crabs calculated on Arctic-specific population parameters. B_0 = pristine or unfished biomass, F_{MSY} = Fishing mortality at maximum sustainable yield (MSY), B_{MSY} = biomass at MSY, mt = metric tons.

| Parameter | Arctic FMP | This study |
|----------------------------------|------------|------------|
| Total B_0 (Chukchi Sea) | 66,491 mt | 746,596 mt |
| Total B_0 (Beaufort Sea) | 29,731 mt | 63,577 mt |
| Harvestable B_0 (Chukchi Sea) | 0 mt | 0 mt |
| Harvestable B_0 (Beaufort Sea) | 6,571 mt | 15,894 mt |
| F_{MSY} | 0.36 | 0.34 |
| B_{MSY} | 1,268 mt | 2,681 mt |
| MSY | 453 mt | 905 mt |
| B_{MSY}/B_0 | 0.19 | 0.17 |
| MSY/B_0 | 0.06 | 0.06 |

4. Discussion

This study improved our understanding of snow crab biology and population status in the Chukchi and Beaufort seas. Snow crab occurred across the Chukchi shelf, but were found only in a localized portion of the western Beaufort shelf and central Beaufort shelf break and upper slope. Both males and females in the Beaufort Sea reached larger sizes than conspecifics in the Chukchi Sea, even though there was larger overlap in the size range of females between the two seas. Despite size range differences, weight-at-size was similar between the Chukchi and Beaufort seas for males, but not females. Size at 50% maturity and fecundity in the Chukchi Sea were both slightly lower than in other geographic regions (e.g., EBS, Sea of Japan, Gulf of St. Lawrence). Finally, our estimates of total biomass of snow crab in both seas and sustainable yield in the Beaufort Sea only were considerably higher than Arctic FMP estimates (North Pacific Fishery Management Council (NPFMC, 2009)), due possibly to increased abundances across multiple years.

4.1. Size distribution and weight-at-size

Snow crab in the Chukchi Sea were smaller (maximum size of 87 mm CW) than conspecifics in other geographic locations, including the Beaufort Sea where the maximum size of 119 mm CW was found for males (this study, also Rand and Logerwell, 2011). In other locations, the maximum sizes observed are 162 mm CW in the Gulf of St. Lawrence, Canadian Atlantic (Sainte-Marie et al., 1995; Alunno-Bruscia and

Sainte-Marie, 1998), 160 mm CW off the coast of western Greenland (Burmeister and Siegstad, 2008), and 130 mm CW in the Barents Sea (Agnalt et al., 2011). In the Pacific, maximum reported size in the Sea of Okhotsk is 100 mm CW (Yanagimoto et al., 2004) and 130 mm CW in the eastern Bering Sea (Turnock and Rugolo, 2012). Thus, maximum size of snow crab in the Beaufort Sea, but not the Chukchi Sea, was within the range of maximum sizes observed in other regions. The consistently small sizes observed in the Chukchi Sea suggest that growth may be restricted in this region. One possible reason for restricted growth in the Chukchi Sea is the low bottom temperatures from persistent winter water that occurs on the northeastern shelf for a majority of the year ($< -1^{\circ}\text{C}$, Weingartner et al., 2005; Grebmeier et al., 2015; Danielson et al., 2017). Temperatures $< 1^{\circ}\text{C}$ can inhibit a positive energy balance in snow crab reared in the lab (Foyle et al., 1989; Thompson and Hawryluk, 1990), and may result in skipped molting before terminal molt (Orensanz et al., 2007; Burmeister and Sainte-Marie, 2010) or a smaller size-at-terminal molt in field populations (Greenland: Burmeister and Sainte-Marie, 2010; Gulf of St. Lawrence: Sainte-Marie and Gilbert, 1998; Dawe et al., 2012; Newfoundland: Dawe et al., 2012). Cold temperatures ($< -1^{\circ}\text{C}$) also cover the Beaufort shelf, but a marked transition occurs at about 200 m depth, where warmer winter water or Atlantic water ($> 0^{\circ}\text{C}$) can persist (Pickart, 2004; Crawford et al., 2012). Many of the snow crab in the Beaufort Sea, especially the particularly large individuals, were collected from these deeper, warmer waters, suggesting large crab occur in slightly warmer temperatures in the Beaufort Sea, where they may experience faster growth relative to snow crab in the Chukchi Sea. In contrast, Logerwell et al. (2011) reported a strong association between snow crab and cold waters ($< -1.5^{\circ}\text{C}$) on the upper Beaufort Sea slope, but temperature measurements in that study only reached to 140 m, which is above the warmer thermocline that is located below 200 m, where most crab occurred. A majority of the large female crab and the mature males collected in the Beaufort Sea during the present study were found in waters around 0.5°C around 200 m depth (Bluhm et al., 2015), possibly favoring the warmer Atlantic water conditions where they may grow to a size comparable to that of snow crab in other regions with more moderate temperatures.

4.2. Maturity-at-size

Male size at 50% morphometric maturity in the Chukchi Sea (62 mm CW) was much larger than a previous estimate of 35 mm CW for Chukchi males based on presence of spermatophores (Paul et al., 1997). This discrepancy in maturity-at-size may be due to differences in methodology for determining maturity state – physiological maturity (i.e., development of spermatophores) versus morphometric maturity (i.e., development of large chela). Nevertheless, we cannot rule out that an increase in maturity-at-size has occurred in the region over the last 2–3 decades. Comparisons of size at 50% morphometric maturity using CW:CH ratio yield high variability across geographic locations. Males in the EBS and the Barents Sea achieve morphometric maturity at 100 mm CW (Turnock and Rugolo, 2012; Dvoretzky and Dvoretzky, 2011), but males in the Gulf of St. Lawrence are physiologically mature at 40 mm CW (Sainte-Marie and Hazel, 1992); based on spermatophore presence, males in the Barents Sea are mature at 46 mm (Filina and Pavlov, 2009), ~40 mm CW in the Chukchi Sea (Barber et al., 1994), and 65 mm CW in the eastern Bering Sea (Somerton, 1981). We found more agreement in size at 50% maturity for females in the Chukchi Sea. It should be noted that where morphometric and physiological maturity have been measured concurrently, males exhibit physiological maturity at smaller sizes than morphometric maturity (Comeau and Conan, 1992; Sainte-Marie et al., 1995).

Females reached 50% maturity at 46 mm CW, identical to a previous estimate in the northeastern Chukchi Sea from over two decades ago (Paul et al., 1997), indicating size at 50% maturity has been conserved in this region. Our estimate was only slightly smaller than size at 50%

maturity reported for females from the northern Bering Sea, Gulf of St. Lawrence, and Sea of Japan (50 mm CW for all regions; Ito, 1967; Watson, 1970; Jewett, 1981; Orensanz et al., 2007). The similarity in size at 50% maturity suggests that female reproductive size may be at least partially conserved among some geographic and oceanographic regions, even though phenotypic response to temperature has been observed on a smaller scale in the EBS (between 55°N to 65°N , Orensanz et al., 2007). Several investigators suggested that female snow crab tend to be smaller at maturity at the northern limit of their distribution in the Chukchi Sea due to reduced bottom temperatures at northern limits (Jewett, 1981; Somerton, 1981; Paul et al., 1997). Clinal variation in size at 50% maturity is well established from the southeastern Bering into the northeast Bering and Chukchi seas (Jewett, 1981; Stevens and MacIntosh, 1986; Paul et al., 1997; Zheng et al., 2001; Ernst et al., 2005; Orensanz et al., 2007), Greenland (Burmeister and Sainte-Marie, 2010) and in the Gulf of St. Lawrence, Canada (Sainte-Marie and Gilbert, 1998). Our data are consistent with smaller female size at 50% maturity extending into the Chukchi Sea from the Bering Sea.

4.3. Fecundity

Fecundity estimates obtained here were lower than previous estimates obtained from estimates resulting from collections in only the southeastern portion of the Chukchi Sea and in other Arctic regions (Jewett, 1981). For example, a 55 mm CW female based on the Chukchi Sea sampling region in our study had a mean fecundity of ~23,000 eggs compared to 28,000 eggs of a female crab at the same size in the southeastern Chukchi Sea (Jewett, 1981). Females of the same size would carry about 26,600 eggs in the Canadian Atlantic (Haynes et al., 1976), about 33,300 eggs in the EBS (Jewett, 1981), and about 24,000 eggs in the northern Bering Sea (Kolts et al., 2015). Seemingly higher fecundity estimates for female crab of identical size in the southeastern Bering Sea, compared to the Chukchi Sea, may be related to warmer temperatures at lower latitudes as discussed for weight and size above (Orensanz et al., 2007).

The question arises whether female egg production in the Chukchi Sea is sufficient to support the local population or whether this area depends on advection of larvae from the Bering Sea to sustain itself. We estimated total egg production of the U.S. Chukchi Sea crab to be about 6.5 billion eggs in 2012. To approximate local versus advective production, we applied a pelagic larval survival estimate of 22% (Yamamoto et al., 2014) to the 6.5 billion eggs produced; thus, approximately 1.5 billion larvae would presumably recruit and metamorphose into the first benthic instars (instar I; ~3 mm CW for males in the Gulf of St. Lawrence; Comeau et al., 1998). However, the timing of the cruises in the Chukchi Sea over the study period (i.e., August to September) does not align with the assumed recruitment period for the region. For example, Parada et al. (2010) conclude that snow crab benthic settlement in the EBS probably peaks in late summer or early fall (i.e. September to October). The Gulf of St. Lawrence experiences a similar larval release period similar to that in EBS (i.e., May), and settlement occurs in August to late fall (i.e., instar I density increases from August, through October and is highest in December; c.f., Lovrich et al. 1995, Ouellet and Sainte-Marie, 2018). Thus, we assume that at least a portion of the observed instar II and III (~5–7 mm CW for males in the Gulf of St. Lawrence, Comeau et al., 1998) include the previous year's recruits that have molted into larger early benthic instars. The average size frequency distribution of snow crab in the Chukchi Sea in 2012 indicated that 9% of the sampled population (or ~169 million individuals) was ~3–7 mm CW, presumably representing benthic instars I–III, and represents a minimum estimate of the total annual recruitment. Thus, mature females in the Chukchi Sea study region could produce the number of small snow crab we observed in 2012. However, this estimate is admittedly rough given it is based on lab-reared larval survival, observations from a single year, the assumption that all larvae

had settled into the benthos by the time of sampling, and that it does not consider variability in the factors influencing larval dispersal nor the fraction of mature females reproducing on a biennial cycle. The degree of larval advection supplying recruits to the Chukchi shelf from adjacent, upstream regions is not currently known (but see larval abundance estimates by Landeira et al., 2017), but advection has been postulated as an important source of crab in the Chukchi and Beaufort seas (Clement et al., 2005; Hu and Wang, 2010; Bluhm et al., 2015; Kolts et al., 2015). This notion is supported by the fact that genetically, snow crab in the Bering, Chukchi and Beaufort seas are considered a panmictic population, with strong gene flow among these regions (Hardy et al., 2011; Albrecht et al., 2014). Considerable uncertainty exists in our egg production estimates due to uncertainties about annual versus biennial reproductive cycles, lifetime egg production per female, and larval and early benthic life stage mortality. However, our results suggest that locally sourced recruitment in the Chukchi Sea may be a significant portion of observed crab on the Chukchi shelf. More detailed knowledge of early life history mortality rates of snow crab is necessary to obtain better estimates of the *in-situ* contribution to the Chukchi snow crab population from mature females versus larvae advected from the Bering Sea.

4.4. Biomass, abundance, and sustainable yield

Although our results suggest approximately 242% higher harvestable biomass in the Beaufort Sea compared with previous estimates from the Arctic FMP, remaining uncertainty in many of the life history metrics used for both the Arctic FMP and our calculations warrants the continued use of a conservative management approach (Restrepo et al., 1998; Zheng, 2003). Our estimates of total and harvestable biomass in the Chukchi and Beaufort seas were compared with previous biomass estimates in the Arctic FMP, which were calculated on limited data for the two seas (North Pacific Fishery Management Council (NPFMC, 2009)). Our revised total biomass estimate was approximately 90% higher than original estimates in the Chukchi Sea, but was also based on a survey area that was 43% larger, which partly explains the higher biomass estimate. Similarly, in the Beaufort Sea, our data set included a 90% larger survey area, and resulted in a total biomass estimate that was approximately twice as high as Arctic FMP estimates. The higher biomass estimate in the Beaufort Sea could, however, not be attributed to the larger survey region, considering that most of the new survey areas in the eastern Beaufort Sea had few or no snow crab (Fig. 8); survey stations used in the Arctic FMP (Logerwell and Rand, 2010) covered most of the area in the western Beaufort Sea that had appreciable numbers of snow crab. By including new survey areas to the east, we were able to fully resolve the strong west to east gradient in snow crab abundance and delineate the easternmost extent of snow crab distribution in the Beaufort Sea. Despite the uncertainty associated with biomass estimates, our results suggest that snow crab density (biomass per unit area) in the U.S. Arctic has increased considerably since the early 1990s. Our higher total biomass estimates translated into higher harvestable biomass estimates in the Beaufort Sea, although harvestable biomass in the Chukchi Sea remained zero since no crab larger than 100 mm CW were found. In the Beaufort Sea, our estimate of harvestable biomass was approximately 212% the Arctic FMP estimate, possibly indicating increasing biomass in the region over time. However, crab of marketable size were only observed in the western Beaufort Sea at depths greater than 100 m, far from existing ports and vessel fleets, making commercial harvest of this stock economically less feasible.

We estimated higher densities in the Chukchi Sea (387,691 individuals km^{-2}) than a previous estimate that used Arctic EIS EBT trawl data only (212,000 individuals km^{-2} , Goddard et al., 2012). Over the entire U.S. Chukchi Sea area, our total abundance (56 million individuals) was sizably larger than the 4 million individuals estimated in Goddard et al. (2012), due to the high numbers of small crab collected during cruises employing the PSBT in the present study. Abundance was

low in the Beaufort Sea, with an estimated 46 individuals km^{-2} , and a total of 2.3 million individuals for the entire survey region. In comparison, a previous survey conducted solely in the western Beaufort Sea found 99,600 individuals km^{-2} (Rand and Logerwell, 2011); no estimate for total survey area was available for that earlier study.

Our results provide an updated estimate of the difference between the age-at-maturity and the age intercept of the linear weight-at-age equation (*d*) that was derived from new, region-specific data on Arctic snow crab. Harvestable biomass estimates changed considerably (Table 4), but these changes were related primarily to the substantial increase in estimated pristine or unfished biomass, rather than to changes in parameter estimates. The much higher estimates of total biomass in the Chukchi Sea, and of both total and harvestable biomass in the Beaufort Sea, compared to the FMP estimates, resulted from a combination of improved spatial coverage and higher densities of snow crab in the recent period. Due to the remoteness and relatively low density of large snow crab on the Beaufort Sea slope, our updated estimates do not warrant a change to the Arctic FMP with respect to the “Optimum Yield” specified for snow crab, which is currently set to zero based in part on cost considerations (North Pacific Fishery Management Council (NPFMC, 2009)).

4.5. Conclusions

Our results are intended to inform the management of the Arctic snow crab stock off the coast of Alaska and to expand our understanding of Arctic snow crab life history parameters and distribution trends in light of potential future fisheries or other, non-fishing activities. Most population parameters we investigated in this study were reasonable comparable to those in other geographic regions where snow crab occur. We found generally similar weight-at-size rates in snow crab between sexes and seas. Both size at 50% maturity and fecundity estimates were within similar ranges as for snow crab occurring in other geographic localities. Our new estimates of biomass and sustainable yield update existing Arctic FMP estimates with results from recent surveys and region-specific life history parameters. Estimates of fishery reference points may be further improved with inclusion of molting probabilities, size-at-age data, and region-specific natural mortality estimates, should those become available in the future. Our estimate of fecundity could be improved if information becomes available regarding the percentage of females on an annual versus biennial reproductive schedule. Ultimately, the results of the present study support continued precautionary management of U.S. Arctic snow crab stocks, which does not allow for commercial harvest at this time. Continued monitoring with the intent of augmenting the temporal coverage of snow crab size frequency distributions, biomass, and abundance, as well as investigations into the larval dispersal and migratory connectivity of snow crab occurring in the Bering, Chukchi, and Beaufort regions, will greatly facilitate improved management of snow crab as a potential future fishery resource.

Acknowledgements

We extend many thanks to the crews and all cruise participants (in particular B. Norcross and her lab members; L. Bell, B. Gray, S. Hardy, D. Hondolero, A. Ravelo, and J. Weems (all UAF) of the cruises listed in Table 1 for their assistance during in sampling efforts. In particular, we are indebted to the many data providers: A. Blanchard (UAF), L. Cooper (UMCES), K. Dunton (UT Austin), J. Grebmeier (UMCES), B. Konar (UAF), R. Lauth (NOAA), and L. Logerwell (NOAA). Training, advice and fruitful discussions were provided by L. Stichert, D. Pengilly and J. Webb (ADF&G Kodiak and Juneau), and J. Lovvorn (SIU). Lab work was supported by M. Kaufmann, C. Serratos and UAF undergraduate students. This study was supported in part by samples collected in 2011 in the Beaufort Sea under BOEM Cooperative Agreement No. M10AC2004 “Beaufort Sea Marine Fish Monitoring in the Central

Beaufort Sea" (BeauFish), U.S. Department of the Interior, Bureau of Ocean Energy Management (BOEM), Alaska Outer Continental Shelf Region; participation of KI and BB in the BeauFish field survey was supported by the Coastal Marine Institute (CMI, Award M11AC00003); samples were collected in 2012–2014 under BOEM Agreement No. M12AC00011 "U.S.-Canada Transboundary Fish and Lower Trophic Communities Project". Samples in the Chukchi Sea were obtained with, and contributions of FJM were in part supported by, qualified outer continental shelf oil and gas revenues by the Coastal Impact Assistance Program, Fish and Wildlife Service, U.S. Department of the Interior under Agreement No. 10-CIAP-010 and under BOEM Cooperative Agreement No. M12AC00009 "Arctic Ecosystem Integrated Survey (Arctic Eis)". The Alaska Department of Environmental Conservation, Alaska Monitoring and Assessment Program (AKMAP) and the U.S. Fish and Wildlife Service, Coastal Impact Assistance Program (CIAP) provided funding to the University of Alaska Fairbanks (S. Jewett) for benthic trawling in the nearshore Chukchi Sea in 2010–2011. Some Chukchi Sea crab samples were also collected through the Russian-American Long-Term Census of the Arctic (RUSALCA) cruise in 2009 and 2012; this publication is the result in part of research sponsored by the Cooperative Institute for Alaska Research with funds from the National Oceanic and Atmospheric Administration under cooperative agreements NA13OAR4320056 and NA08OAR4320870. Crab samples during the Arctic Marine Biodiversity Observing Network (AMBN) project were obtained through the National Ocean Partnership Program (NOPP grant NA14NOS0120158) by the National Oceanographic and Atmospheric Administration (NOAA), the Bureau of Ocean Management (BOEM), Shell Exploration & Production, and additional contributions from the National Science Foundation, under management of the Integrated Ocean Observing System (IOOS). The National Science Foundation Marine Ecosystem Sustainability in the Arctic and Subarctic (MESAS) IGERT (Award DGE-0801720) provided support for LD throughout the project.

References

- Agnalt, A.L., Pavlov, V., Jørstad, K.E., Færevest, E., Sundet, J., 2011. The snow crab, *Chionoecetes opilio* (Decapoda, Majoidae, Oregoniidae) in the Barents Sea. In: *The Wrong Place-Alien Marine Crustaceans: Distribution, Biology and Impacts*. Springer, Dordrecht, pp. 283–300.
- Alaska Department of Fish and Game (ADF&G), 2018. Statewide Commercial Shellfish Fishery Exvessel Prices. (Accessed 29 April 2018). http://www.adfg.alaska.gov/index.cfm?adfg=CommercialByFisheryshellfish.shellfishcatch_exvessel_crab.
- Albrecht, G.T., Valentin, A.E., Hundertmark, K.J., Hardy, S.M., 2014. Pannmixia in Alaskan populations of the snow crab *Chionoecetes opilio* (Malacostraca: Decapoda) in the Bering, Chukchi, and Beaufort Seas. *J. Crust. Biol.* 34, 31–39.
- Alunno-Bruscia, M., Sainte-Marie, B., 1998. Abdomen allometry, ovary development, and growth of female snow crab, *Chionoecetes opilio* (Brachyura, Majidae), in the north-western Gulf of St. Lawrence. *Can. J. Fish. Aquat. Sci.* 55, 459–477.
- Armstrong, D.A., Bugos, J., Ernst, B., Orensanz, J.M., 2008. Female effective reproductive output of the snow crab stock in the eastern Bering Sea. *North Pac. Res. Board Final Rep. Project 508*, 165.
- Armstrong, D.A., McDonald, P.S., Kruse, G.H., Hines, A.H., Orensanz, J.M., 2010. A crab for all seasons: the confluence of fisheries and climate as drivers of crab abundance and distribution. In: Kruse, G.H., Eckert, G.L., Foy, R.J., Lipcius, R.N., Sainte-Marie, B., Stram, D.L., Woodby, D. (Eds.), *Biology and Management of Exploited Crab Populations Under Climate Change*. Alaska Sea Grant, University of Alaska Fairbanks. <https://doi.org/10.4027/bmccpcc.2010>.
- Barber, W.E., Smith, R.L., Weingartner, T.J., 1994. Fisheries Oceanography of the Northeast Chukchi Sea. Final Report to the Alaska Outer Continental Shelf Region of the Mineral Management Service. U.S. Department of the Interior, OCS Study MMS-93-0051 101 pp.
- Bluhm, B.A., Iken, K., Hardy, S.M., Sirenko, B.I., Holladay, B.A., 2009. Community structure of epibenthic megafauna in the Chukchi Sea. *Aquat. Biol.* 7, 269–293.
- Bluhm, B.A., Iken, K., Divine, L., 2015. Population Assessment of Snow Crab, *Chionoecetes opilio*, in the Chukchi and Beaufort Seas, Including Oil and Gas Lease Areas. Final Report OCS Study BOEM 2015-029. University of Alaska, Fairbanks, AK 83 pp.
- Britt, L.L., Lauth, R.R., Norcross, B.L., 2013. Draft Report to Department of the Interior Bureau of Ocean Energy Management: Distribution of Fish, Crab and Lower Trophic Communities in the Chukchi Sea AK-11-08. Interagency Agreement M12PG00018. University of Alaska, Fairbanks, AK 52 pp.
- Burmeister, A.D., Sainte-Marie, B., 2010. Pattern and causes of a temperature-dependent gradient of size at terminal moult in snow crab (*Chionoecetes opilio*) along West Greenland. *Polar Biol.* 33, 775–788.
- Burmeister, A.D., Siegstad, H., 2008. Assessment of snow crab in West Greenland 2008, greenland institute of natural resources. Nuuk 68.
- Clement, J.L., Maslowski, W., Cooper, L.W., Grebmeier, J.M., Walczowski, W., 2005. Ocean circulation and exchanges through the northern Bering Sea: 1979–2001 model results. *Deep-Sea Res. II* 52, 3509–3540.
- Comeau, M., Conan, G.Y., 1992. Morphometry and gonad maturity of male snow crab, *Chionoecetes opilio*. *Can. J. Fish. Aquat. Sci.* 49, 2460–2468.
- Comeau, M., Conan, G.Y., Maynou, F., Robichaud, G., Theriault, J.C., Starr, M., 1998. Growth, spatial distribution, and abundance of benthic stages of the snow crab (*Chionoecetes opilio*) in Bonne Bay, Newfoundland, Canada. *Can. J. Fish. Aquat. Sci.* 55, 262–279.
- Conan, G.Y., Comeau, M., 1986. Functional maturity and terminal molt of male snow crab, *Chionoecetes opilio*. *Can. J. Fish. Aquat. Sci.* 43, 1710–1719.
- Crawford, R.E., Vagle, S., Carmack, E.C., 2012. Water mass and bathymetric characteristics of polar cod habitat along the continental shelf and slope of the Beaufort and Chukchi seas. *Polar Biol.* 35, 179–190.
- Cushing, D.H., 1971. The dependence of recruitment on parent stock in different groups of fishes. *ICES J. Mar. Sci.* 33, 340–362.
- Danielson, S.L., Eisner, L., Ladd, C., Mordy, C., Sousa, L., Weingartner, T.J., 2017. A comparison between late summer 2012 and 2013 water masses, macronutrients, and phytoplankton standing crops in the northern Bering and Chukchi Seas. *Deep-Sea Res. II* 135, 7–26.
- Dawe, E.G., Mullowney, D.R., Moriyasu, M., Wade, E., 2012. Effects of temperature on size-at-terminal molt and molting frequency in snow crab *Chionoecetes opilio* from tow Canadian Atlantic ecosystems. *Mar. Ecol. Prog. Ser.* 469, 279–296.
- Dunn, P.K., Smyth, G.K., 2005. Series evaluation of Tweedie exponential dispersion model densities. *Stat. Comput.* 15, 267–280.
- Dvoretsky, A.G., Dvoretsky, V.G., 2011. Commercial fish and shellfish in the Barents Sea: Have introduced crab species affected the population trajectories of commercial fish? *Rev. Fish. Biol. Fish.* 25, 297–322.
- Ernst, B., Orensanz, J.M., Armstrong, D.A., 2005. Spatial dynamics of female snow crab (*Chionoecetes opilio*) in the eastern Bering Sea. *Can. J. Fish. Aquat. Sci.* 62, 250–268.
- Ernst, B., Armstrong, D.A., Burgos, J., Orensanz, J.M., 2012. Life history schedule and periodic recruitment of female snow crabs (*Chionoecetes opilio*) in the eastern Bering Sea. *Can. J. Fish. Aquat. Sci.* 69, 532–550.
- Filina, E.A., Pavlov, V.A., 2009. Results of research on male snow crab (*Chionoecetes opilio*) maturation in the Barents Sea. *International Symposium on Biology and Management of Exploited Crab Populations Under Climate Change* 5–13.
- Fissel, B., Dalton, M., Garber-Yonts, B., Haynie, A., Kasperski, S., Lee, J., Lew, D., Lavoie, A., Seung, C., Sparks, K., Szymkowiak, M., Wise, S., 2018. Stock Assessment and Fishery Evaluation Report for the Groundfish Fisheries of the Gulf of Alaska and Bering Sea/Aleutian Islands Area: Economic Status of the Groundfish Fisheries off Alaska, 2017. *Res. Ecol. Fish. Mgmt. Div. AFSC, Seattle, WA* 98115-6349 247 pp.
- Foyle, T.P., O'Dor, R.K., Elner, R.W., 1989. Energetically defining the thermal limits of the snow crab. *J. Exp. Biol.* 145, 371–393.
- Frost, K.J., Lowry, L.F., 1983. Trophic relationships of vertebrate consumers in the Alaskan Beaufort Sea. In: Barnes, P.W., Schell, D.M., Reimnitz, E. (Eds.), *The Alaskan Beaufort Sea: Ecosystems and Environments*. Academic Press, Orlando, F.L., pp. 381–402.
- Goddard, P., Lauth, R., Armistead, C., 2012. Results of the 2012 Chukchi Sea Bottom Trawl Survey of Bottom Fishes, Crabs, and Other Demersal Macrofauna. *US Dept Commer, NOAA Tech Memo NMFS-AFSC-278* 110 p.
- Grebmeier, J.M., Bluhm, B.A., Cooper, L.W., Danielson, S.L., Arrigo, K.R., Blanchard, A.L., Clarke, J.T., Day, R.H., Frey, K.E., Gradinger, R.R., Kedra, M., 2015. Ecosystem characteristics and processes facilitating persistent macrobenthic biomass hotspots and associated benthivory in the Pacific Arctic. *Progr. Oceanogr.* 136, 92–114.
- Gross, J., Konar, B., Brey, T., Grebmeier, J.M., 2017. Size frequency distribution, growth, and mortality of snow crab (*Chionoecetes opilio*) and Arctic Lyre crab (*Hyas coarctatus*) in the Chukchi Sea from 2009 to 2013. *Deep-Sea Res. II* 144, 142–155.
- Gunderson, D.R., Ellis, I.E., 1986. Development of a plumb staff beam trawl for sampling demersal fauna. *Fish. Res.* 4, 35–41.
- Hardy, S.M., Lindgren, M., Konacanchi, H., Huettmann, F., 2011. Predicting the distribution and ecological niche of unexploited snow crab (*Chionoecetes opilio*) populations in Alaskan waters: a first open-access ensemble model. *Integr. Comp. Biol.* 51, 608–622.
- Hartnoll, R.G., 1982. Growth. In: Abele, L.G. (Ed.), *The Biology of the Crustacea*. Academic Press, New York, N.Y., pp. 111–196.
- Haynes, E., Karinen, J.F., Watson, J., Hopson, D.J., 1976. Relation of number of eggs and egg length to carapace width in the brachyuran crabs *Chionoecetes bairdi* and *C. opilio* from the southeastern Bering Sea and Gulf of St. Lawrence. *J. Fish. Res. Board Can.* 33, 2592–2595.
- Hines, A.H., 1982. Allometric constraints and variables of reproductive effort in brachyuran crabs. *Mar. Biol.* 69, 309–320.
- Hollowed, A.B., Planque, B., Loeng, H., 2013. Potential movement of fish and shellfish stocks from the sub-Arctic to the Arctic Ocean. *Fish. Oceanogr.* 22, 355–370.
- Hu, H., Wang, J., 2010. Modeling effects of tidal and wave mixing on circulation and thermohaline structures in the Bering Sea. *J. Geophys. Res.* 115, C01006. <https://doi.org/10.1029/2008JC005175>.
- Ito, K., 1967. Ecological studies on the edible crab, *Chionoecetes opilio* (O. Fabricius), in the Japan Sea. I. When do female crabs first spawn and how do they advance into the following reproductive stage. *Bull. Jpn. Sea Reg. Fish. Res. Lab.* 17, 67–84.
- Jewett, S.C., 1981. Variations in some reproductive aspects of female snow crabs *Chionoecetes opilio*. *J. Shellfish Res.* 1, 95–99.
- Kolts, J.M., 2012. Population Structure, Reproductive Status, and Diet of Snow Crabs, *Chionoecetes opilio*, in the Northern Bering Sea. Ph.D. Dissertation. University of Wyoming 145 pp.
- Kolts, J.M., Lovvorn, J.R., North, C.A., Grebmeier, J.M., Cooper, L.W., 2013. Effects of

- body size, gender, and prey availability on diets of snow crabs in the northern Bering Sea. *Mar. Ecol. Prog. Ser.* 483, 209–220.
- Kolts, J.M., Lovvorn, J.R., North, C.A., Janout, M.A., 2015. Oceanographic and demographic mechanisms affecting population structure of snow crabs in the northern Bering Sea. *Mar. Ecol. Prog. Ser.* 518, 193–208.
- Konar, B., Ravelo, A., Grebmeier, J.M., Trefry, J.H., 2014. Size frequency distributions of key epibenthic organisms in the eastern Chukchi Sea and their correlations with environmental parameters. *Deep-Sea Res. II* 102, 107–118.
- Kotwicki, S., Lauth, R.R., Williams, K., Goodman, S.E., 2017. Selectivity ratio: a useful tool for comparing size selectivity of multiple survey gears. *Fish. Res.* 191, 76–86. <https://doi.org/10.1016/j.fishres.2017.02.012>.
- Landeira, J.M., Matsuno, K., Yamaguchi, A., Hirawake, T., Kikuchi, T., 2017. Abundance, development stage, and size of decapod larvae through the Bering and Chukchi Seas during summer. *Polar Biol.* 40, 1805–1819.
- Lögerwell, L., Rand, K., 2010. Beaufort Sea Marine Fish Monitoring 2008: Pilot Survey and Test of Hypotheses. USDOI Alaska. OCS Region OCS Study MMS, pp. 2008–2062.
- Lögerwell, L., Rand, K., Weingartner, T., 2011. Oceanographic characteristics of the habitat of benthic fish and invertebrates in the Beaufort Sea. *Polar Biol.* 34, 1783–1796.
- Moriyasu, M., Lantagne, C., 1998. Embryo development and reproductive cycle in the snow crab, *Chionoecetes opilio* (Crustacea: Majidae), in the southern Gulf of St. Lawrence, Canada. *Can. J. Zool.* 76, 2040–2048. <https://doi.org/10.1139/z98-147>.
- Moriyasu, M., Conan, G.Y., Mallet, P., Chiasson, Y., Lacroix, H., 1987. Growth Per Molt, Molting Season and Mating of Snow Crab (*Chionoecetes Opilio*) in Relation to Functional and Morphometric Maturity. *ICES CM* 1987/ K:21. .
- Mueter, F.J., Dawe, E.G., Pálsson, Ö.K., 2012. Subarctic fish and crustacean populations—climate effects and trophic dynamics. *Mar. Ecol. Prog. Ser.* 469, 191–193.
- Norcross, B.L., Holladay, B.A., Walker, K.L., Edenfield, L.E., 2015. . Length-weight-age Relationships of Demersal Fishes on the Shelf of the Alaskan Beaufort Sea, Central Beaufort Sea Marine Fish Monitoring. BOEM 2015-079, Final Rep. Fairbanks, AK, OCS Study Chapter 3.1).
- North Pacific Fishery Management Council (NPFMC), 2009. Fishery Management Plan for Fish Resources of the Arctic Management Area. Anchorage, AK. 146 pp.. .
- Orensanz, J.M., Ernst, B., Armstrong, D.A., Stabeno, P., Livingston, P., 2004. Contraction of the geographic range of distribution of snow crab (*Chionoecetes opilio*) in the eastern Bering Sea: an environmental ratchet? *CalCOFI Rep* 45, 65–79.
- Orensanz, J.M., Ernst, B., Armstrong, D.A., 2007. Variation of female size and stage at maturity in snow crab (*Chionoecetes opilio*) (Brachyura: Majidae) from the eastern Bering Sea. *J. Crust. Biol.* 27, 576–591.
- Otto, R.S., 1998. Assessment of the eastern bering Sea snow crab, *Chionoecetes opilio*, stock under the terminal molting hypothesis. Jamieson, G.S., Campbell, A. (Eds.), *Proceedings of the North Pacific Symposium on Invertebrate Stock Assessment and Management* 125, 109–124.
- Ouellet, P., Sainte-Marie, B., 2018. Vertical distribution of snow crab (*Chionoecetes opilio*) pelagic stages in the Gulf of St. Lawrence (Canada) and effect of temperature on development and survival. *ICES J. Mar. Sci.* 75, 773–784.
- Parada, C., Armstrong, D.A., Ernst, B., Hinckley, S., Orensanz, J.M.L., 2010. Spatial dynamics of snow crab (*Chionoecetes opilio*) in the eastern Bering Sea – putting together the pieces of the puzzle. *Bull. Mar. Sci.* 86, 413–437.
- Paul, J.M., Paul, A.J., Barber, W.E., 1997. Reproductive biology and distribution of the snow crab from the northeastern Chukchi Sea. *Am. Fish Soc. Symp.* 19, 287–294.
- Pickart, R.S., 2004. Shelfbreak circulation in the Alaskan Beaufort Sea: mean structure and variability. *J. Geophys. Res.* 109, C04024. <https://doi.org/10.1029/2003JC001912>.
- Rand, K.M., Lögerwell, E.A., 2011. The first demersal trawl survey of benthic fish and invertebrates in the Beaufort Sea since the late 1970s. *Polar Biol.* 34, 475–488.
- Ravelo, A.M., Konar, B., Trefry, J.H., Grebmeier, J.M., 2014. Epibenthic community variability in the northeastern Chukchi Sea. *Deep-Sea Res. II* 102, 119–131.
- Ravelo, A.L., Konar, B., Bluhm, B.A., 2015. Spatial variability of epibenthic communities on the Alaska Beaufort shelf. *Polar Biol.* 38, 1783–1804.
- Restrepo, V.R., Thompson, G.G., Mace, P.M., Gabriel, W.L., Low, L.L., MacCall, A.D., Methot, R.D., Powers, J.E., Taylor, B.L., Wade, P.R., Witzig, J.F., 1998. Technical Guidance on the Use of Precautionary Approaches to Implementing National Standard 1 of the Magnuson-Stevens Fishery Conservation and Management Act. NOAA Tech Memo NMFS-F/SP-31. 49 pp.. .
- Rugolo, L., Pengilly, D., MacIntosh, R., Gravel, K., 2005. Reproductive potential and life history of snow crabs in the eastern Bering Sea. Bering Sea Snow Crab Fishery Restoration Research: Final Comprehensive Performance Report. NOAA, NMFS 99802-1668 57–323.
- Sainte-Marie, B., Gilbert, D., 1998. Possible effects of changes in CIL temperature and thickness on population dynamics of snow crab, *Chionoecetes opilio*, in the Gulf of Saint Lawrence. CSAS Research Document 98/38. 19 pp.
- Sainte-Marie, B., Hazel, F., 1992. Moulting and mating of snow crabs, *Chionoecetes opilio* (O. Fabricius), in shallow waters of the northwestern Gulf of Saint Lawrence. *Can. J. Fish. Aquat. Sci.* 49, 1282–1293.
- Sainte-Marie, B., Raymond, S., Brethes, J., 1995. Growth and maturation of the male snow crab, *Chionoecetes opilio* (Brachyura: Majidae). *Can. J. Fish. Aquat. Sci.* 52, 903–924.
- Somerton, D.A., 1981. Regional variation in the size of maturity of two species of Tanner crab (*Chionoecetes bairdi* and *C. opilio*) in the eastern Bering Sea, and its use in defining management subareas. *Can. J. Fish. Aquat. Sci.* 38, 163–174.
- Stearns, S.C., 1976. Life history tactics: a review of the ideas. *Q. Rev. Biol.* 51, 3–47.
- Stevens, B.G., MacIntosh, R.A., 1986. Analysis of crab data from the 1985 NMFS survey of the northeast Bering Sea and Norton Sound. Northwest and Alaska fisheries center (NMFS). Processed Report 1986- 16.
- Stichert, L., 2009. AK Depart Fish Game Lab Protocol: Processing *Chionoecetes* crabs. Available from author, Kodiak. AK. 12 pp.. .
- Thompson, R.J., Hawryluk, M., 1990. Physiological energetics of the snow crab, *Chionoecetes opilio*. Proceedings of the International Symposium on King and Tanner Crabs. AK Sea Grant AK-SG-90-04. pp. 283–291.
- Turnock, B.J., Rugolo, L.J., 2008. Stock assessment of eastern Bering Sea snow crab. Plan Team for the King and Tanner Crab Fisheries of the Bering Sea and Aleutian Islands (Compiler), SAFE Report for the King and Tanner Crab Fisheries of the BSAI Regions. pp. 25–114.
- Turnock, B.J., Rugolo, L.J., 2012. Stock Assessment of Eastern Bering Sea Snow Crab. North Pacific Fishery Management Council, PO Box 103136, Anchorage, AK. 116 pp.. .
- Watson, J., 1970. Maturity, mating and egg-laying in the spider crab, *Chionoecetes opilio*. *J. Fish Res. Board Can.* 29, 447–449.
- Webb, J.B., Eckert, G.L., Shirley, T.C., Tamone, S.L., 2007. Changes in embryonic development and hatching in *Chionoecetes opilio* (snow crab) with variation in incubation temperature. *Biol. Bull.* 213, 67–75. <https://doi.org/10.2307/25066619>.
- Webb, J.B., Slater, L.M., Eckert, G.L., Kruse, G.H., 2016. The contribution of fecundity and embryo quality to reproductive potential of eastern Bering Sea snow crab (*Chionoecetes opilio*). *Can. J. Fish. Aquat. Sci.* 73, 1800–1814.
- Weingartner, T., Aagaard, K., Woodgate, R., Danielson, S., Sasaki, Y., Cavalieri, D., 2005. Circulation on the north central Chukchi Sea shelf. *Deep-Sea Res II* 52, 3150–3174.
- Wood, S.N., 2017. Generalized Additive Models: an Introduction With R. Chapman & Hall/CRC, Boca Raton, FL, USA.
- Yamamoto, T., Yamada, T., Fujimoto, H., Hamasaki, K., 2014. Effects of temperature on snow crab (*Chionoecetes opilio*) larval survival and development under laboratory conditions. *J. Shellfish Res.* 33, 19–24.
- Yanagimoto, T., Yoshio, I., Watanabe, K., 2004. Distribution and morphological maturation size of the snow crab, *Chionoecetes opilio* in the southwest area of the Okhotsk Sea in summer. *Nippon Suisan Gakk* 70, 750–757.
- Zheng, J., 2003. Uncertainties of natural mortality estimates for eastern Bering Sea snow crab, *Chionoecetes opilio*. *Calif. Coop. Ocean Fish Investig. Rep.* 65, 411–425.
- Zheng, J., Kruse, G.H., 2006. Recruitment variation of eastern Bering Sea crabs: Climate-forcing or top-down effects? *Prog Oceanog* 68, 184–204.
- Zheng, J., Kruse, G.H., Ackley, D.R., 2001. Spatial distribution and recruitment patterns of snow crabs in the eastern Bering Sea. Spatial Processes and Management of Marine Populations. Edited by G.H. Kruse, N. Bez, A. Booth, M.W. Dorn, S. Hills, R.N. Lipcius, D. Pelletier, C. Roy, S.J. Smith, D. Witherell. Alaska Sea Grant College Program AK-SG-01-02 233–255.