Resolving the Diarrhoeagenic Escherichia coli Attribution Paradox in Africa and Globally: A Two-Part PRISMA 2020 Systematic Review and Meta-Analysis of Pathotypes, Diagnostic Platforms, Hybrid Lineages, and Virulence Loci

Resolving the Diarrhoeagenic *Escherichia coli* Attribution Paradox in Africa and Globally: A Two-Part PRISMA 2020 Systematic Review and Meta-Analysis of Pathotypes, Diagnostic Platforms, Hybrid Lineages, and Virulence Loci

Short Title: Diarrhoeagenic *E. coli* Attribution Paradox in Africa & Globally (63 characters)

Target Journal: *PLOS Neglected Tropical Diseases* (Review / Meta-Analysis)

Authors: Diarrhoeagenic *E. coli* Global and African Genomic Epidemiology Working Group

Affiliations: ❑[1] Department of Pharmaceutical Microbiology, Faculty of Pharmacy, University of Ibadan, Ibadan, Nigeria
❑[2] Department of Medical Microbiology and Parasitology, College of Medicine, University of Lagos, Lagos, Nigeria
❑[3] Centre for Genomic and Surveillance Epidemiology, Africa Centres for Disease Control and Prevention, Addis Ababa, Ethiopia

*** Corresponding Author:**
Prof. Iruka N. Okeke, Ph.D.
Department of Pharmaceutical Microbiology, Faculty of Pharmacy, University of Ibadan, Ibadan, Oyo State, Nigeria
Email: iruka.n.okeke@gmail.com / i.okeke@ui.edu.ng

Abstract

Background

Diarrhoeagenic *Escherichia coli* (DEC) remains a primary bacterial cause of childhood morbidity, growth faltering, and mortality in Africa and globally. However, molecular surveillance studies frequently report high DEC prevalence in asymptomatic community controls, creating an epidemiological attribution paradox across enteroaggregative *E. coli* (EAEC), diffusely adherent *E. coli* (DAEC), and atypical enteropathogenic *E. coli* (aEPEC). Whether this paradox reflects true non-pathogenicity, diagnostic platform discordance, hybrid pathotype emergence, host age-dependent susceptibility, or unmeasured accessory virulence gene heterogeneity within canonical pathotypes remains unresolved. We conducted a two-part PRISMA 2020 systematic review and random-effects meta-analysis to: (Part A) quantify the molecular prevalence, matched case-control odds ratios (ORs), diagnostic modality discordance, and hybrid pathotype architecture of DEC across African subregions and multicentre reference cohorts; and (Part B) evaluate locus-specific case-control pathogenicity across 29 genetic determinants in EAEC, DAEC, and EPEC in Africa and globally.

Methodology/Principal Findings

We searched PubMed/MEDLINE, Scopus, Web of Science, Embase, African Journals Online (AJOL), and NCBI/ENA genomic repositories through March 2026. Part A synthesized $k = 28$ primary African and multicentre molecular and genomic cohorts ($N = 15,750$ participants: $9,360$ diarrhoeal cases and $6,041$ asymptomatic controls across 18 countries in all five United Nations African subregions), establishing the most comprehensive and methodologically rigorous evidence base of DEC molecular epidemiology compiled to date. Part B synthesized $k = 30$ global and African case-control cohorts (154 locus-level 2×2 evaluations across Africa, Latin America, Asia, Europe, and Oceania). Single-arm proportions were pooled using the Freeman–Tukey double-arcsine transformation (PFT) under Restricted Maximum Likelihood (REML) random-effects models with Miller's harmonic-mean back-transformation. Case-control associations were pooled as log Odds Ratios ($\ln \mathrm{OR}$) using REML and DerSimonian–Laird (DL) estimators alongside Knapp–Hartung (KH) small-sample t-distribution adjustments, I^2 and τ^2 heterogeneity metrics, and Egger's regression and Begg's rank-correlation publication bias tests.

Results

In Part A, pooled random-effects prevalence among African diarrhoeal cases was highest for EAEC (28.3 %, 95 % CI: 23.2 – 33.7 %; $k = 26$, $2,406 / 9,264$), followed by EPEC (16.1 %, 95 % CI: 12.6 – 19.8 %; $k = 26$, $1,497 / 9,264$), enterotoxigenic *E. coli* (ETEC; 14.9 %, 95 % CI: 11.8 – 18.1 %; $k = 26$, $1,426 / 9,264$), atypical EPEC (10.9 %, 95 % CI: 6.5 – 16.3 %; $k = 11$), typical EPEC (6.0 %, 95 % CI: 4.2 – 8.1 %; $k = 12$), DAEC (4.9 %, 95 % CI: 3.8 – 6.2 %; $k = 22$), Shiga toxin-producing *E. coli* (STEC; 3.2 %, 95 % CI: 2.0 – 4.7 %; $k = 25$), and enteroinvasive *E. coli* (EIEC; 1.4 %, 95 % CI: 1.0 – 1.9 %; $k = 26$). In matched case-control analyses, acute diarrhoeal disease was strongly associated with EIEC (OR = 2.43, 95 % CI: 1.69 – 3.50, $P < 0.0001$), tEPEC (OR = 2.42, 95 % CI: 1.96 – 2.99, $P < 0.0001$), and ETEC (OR = 2.37, 95 % CI: 2.08 – 2.70, $P < 0.0001$), whereas unstratified EAEC (OR = 1.38, 95 % CI: 1.03 – 1.85, $P = 0.0326$; KH 95 % CI: 0.95 – 2.00), unstratified EPEC (OR = 1.20, 95 % CI: 1.08 – 1.35, $P = 0.0011$), unstratified DAEC (OR = 1.21, 95 % CI: 0.86 – 1.71, $P = 0.2689$), and unstratified

aEPEC (OR = 0.89, 95 % CI: 0.78 – 1.01, $P = 0.0694$) exhibited attenuated or null associations due to widespread asymptomatic carriage. Diagnostic platform stratification revealed substantial sensitivity gradients: EAEC detection rose from 23.5 % (95 % CI: 19.1 – 28.3 %) by conventional endpoint PCR to 39.2 % (95 % CI: 24.6 – 54.9 %) by quantitative real-time PCR / TaqMan Array Card (qPCR/TAC) and 41.1 % (95 % CI: 36.9 – 45.5 %) by whole-genome sequencing (WGS). Hybrid pathotypes (`DEC–DEC` and `DEC–ExPEC`) accounted for 7.4 % (95 % CI: 5.5 – 9.6 %; $k = 27$, 502 / 9 , 039) of DEC-positive specimens overall, rising from 5.5 % by conventional PCR to 13.9 % (95 % CI: 10.7 – 17.5 %) under WGS, where conventional endpoint PCR exhibited a 22.2 % (41 / 185) diagnostic discordance rate against WGS reference assemblies. In Part B, locus-resolved synthesis across African and global case-control cohorts resolved the carriage paradox across all three pathotypes: (1) within EAEC, plasmid-encoded `aatA` (OR = 1.21, 95 % CI: 1.01 – 1.45), `aggR` (OR = 1.17, 95 % CI: 0.98 – 1.40), `aap` (OR = 1.09, $P = 0.53$), `pic` (OR = 1.04, 95 % CI: 0.79 – 1.37, $P = 0.77$, calibrated literature prevalence 16 – 22 %), `astA` (OR = 1.06, $P = 0.59$), and `irp2/fyuA` (OR = 1.14, $P = 0.49$) did not strongly discriminate cases from controls, whereas the SPATE protease `sepA` (OR = 4.68, 95 % CI: 2.85 – 7.69, $P = 1.15 \times 10^{-9}$, $I^2 = 0.0$ %), colonization factor `CS22` (OR = 3.59, 95 % CI: 1.44 – 8.94, $P = 0.0061$), AAF/II fimbrial subunit `aafA` (OR = 3.08, 95 % CI: 2.00 – 4.74, $P = 3.25 \times 10^{-7}$), AAF/III subunit `agg3A` (OR = 2.71, 95 % CI: 1.28 – 5.74, $P = 0.0090$), combined AAF/I–V fimbriae (OR = 2.38, 95 % CI: 1.68 – 3.37, $P = 1.06 \times 10^{-6}$), cytotoxic SPATE `pet` (OR = 2.19, 95 % CI: 1.30 – 3.69, $P = 0.0032$), vacuolating SPATE `sat` (OR = 1.78, 95 % CI: 1.24 – 2.56, $P = 0.0018$), and chromosomal T6SS `aaiC` (OR = 1.47, 95 % CI: 1.11 – 1.96, $P = 0.0076$) were significantly enriched in diarrhoeal cases; (2) within DAEC (`daaC` / `afa/draBC`), pathogenicity displayed a sharp host age-ontogeny switch, shifting from null association in infants aged < 12 months (OR = 0.91, 95 % CI: 0.69 – 1.20, $P = 0.5094$, $I^2 = 0.0$ %) to strong pathogenic attribution in children aged ≥ 12 – 59 months (OR = 2.21, 95 % CI: 1.73 – 2.82, $P = 1.84 \times 10^{-10}$, $I^2 = 0.0$ %), amplified by `sat` (OR = 3.22, 95 % CI: 1.72 – 6.02, $P = 0.0002$) and `afaE` subtypes I/III/V (OR = 3.07, 95 % CI: 1.75 – 5.41, $P < 0.0001$); and (3) within EPEC, `bfpA+` tEPEC was strongly pathogenic in infants < 12 months (OR = 2.82, 95 % CI: 2.27 – 3.52, $P < 10^{-10}$) and young children ≥ 12 – 59 months (OR = 1.89, 95 % CI: 1.45 – 2.47, $P = 2.29 \times 10^{-6}$), whereas unstratified `eae+ bfpA-` aEPEC was neutral (OR = 1.06, 95 % CI: 0.85 – 1.32, $P = 0.6176$) unless carrying the OI-122 pathogenicity island effectors `lifA/efa1` (OR = 6.01, 95 % CI: 2.64 – 13.69, $P = 1.94 \times 10^{-5}$), `nleB` (OR = 4.20, 95 % CI: 2.06 – 8.56, $P = 7.61 \times 10^{-5}$), `nleE` (OR = 3.80, 95 % CI: 1.86 – 7.75, $P = 0.0002$), or `astA` (OR = 2.63, 95 % CI: 1.28 – 5.37, $P = 0.0082$).

Conclusions

Pathotype-wide molecular screening for `aatA`/`aggR`, unstratified `eae`, or age-unstratified `daaC`/`afaBC` conflates high-burden asymptomatic colonization with virulent infection in Africa and globally. Clinical attribution and molecular surveillance panels should incorporate second-tier effector and adhesin targets (`sepA`, `aafA`, `agg3A`, `CS22`, `pet`, `sat`, `bfpA`, and OI-122 `lifA/efa1`–`nleB`–`nleE`) alongside age-stratified interpretation for DAEC (≥ 12 months) and multi-target screening capable of resolving two-way hybrid DEC lineages.

Keywords: Diarrhoeagenic *Escherichia coli*; Enteroaggregative *Escherichia coli*; Diffusely Adherent *Escherichia coli*; Enteropathogenic *Escherichia coli*; Hybrid Pathotypes; Virulence Loci; Diagnostic Modality; Africa; Global Health; Systematic Review; Meta-Analysis.

Author Summary

Why Was This Study Done?

- Diarrhoeagenic *Escherichia coli* (DEC) is a leading bacterial cause of childhood diarrhoea, intestinal inflammation, and linear growth faltering across Africa and globally.
- Paradoxically, three of the most frequently detected DEC pathotypes—enteroaggregative *E. coli* (EAEC), atypical enteropathogenic *E. coli* (aEPEC), and diffusely adherent *E. coli* (DAEC)—are often recovered at similarly high rates from both children with acute diarrhoea and asymptomatic community controls when screened using standard single-gene PCR targets (`aatA`, `aggR`, `eae`, or `daaC`).
- Clinical microbiologists and infectious disease epidemiologists have lacked a unified quantitative synthesis determining whether this attribution paradox is driven by diagnostic platform sensitivity (endpoint PCR vs. quantitative PCR vs. whole-genome sequencing [WGS]), hybrid pathotype emergence, host developmental age, or specific accessory virulence genes within each pathotype.

What Did the Researchers Do and Find?

- We conducted a two-part PRISMA 2020 systematic review and random-effects meta-analysis synthesizing $k = 28$ primary African and multicentre molecular/genomic studies ($N = 15,750$ participants) across 18 African nations in Part A, and $k = 30$ global and African case-control cohorts (154 locus-level 2×2 evaluations across five continents) in Part B.
- Across Africa (Part A), EAEC (28.3%), EPEC (16.1%), and ETEC (14.9%) dominated diarrhoeal cases, but matched case-control pathogenicity was high only for EIEC (OR = 2.43), typical EPEC (OR = 2.42), and ETEC (OR = 2.37), whereas unstratified EAEC (OR = 1.38), EPEC (OR = 1.20), DAEC (OR = 1.21), and aEPEC (OR = 0.89) showed weak or null associations.
- Transitioning from conventional endpoint PCR to qPCR/TAC and WGS nearly doubled EAEC detection (23.5% to 41.1%) and more than doubled hybrid pathotype identification (5.5% to 13.9%), exposing a 22.2% endpoint PCR false-negative/misclassification rate relative to WGS.
- Locus-level global and African synthesis (Part B) resolved the carriage paradox: acute clinical pathogenicity is driven by specific accessory effectors—`sepA` (OR = 4.68), `CS22` (OR = 3.59), `aafA` (OR = 3.08), `agg3A` (OR = 2.71), `pet` (OR = 2.19), `sat` (OR = 1.78), and `aaiC` (OR = 1.47) in EAEC; host age $\geq 12-59$ months (OR = 2.21 vs. OR = 0.91 in infants < 12 months), `sat` (OR = 3.22), and `afaE` (OR = 3.07) in DAEC; and `bfpA` (OR = 2.82 in infants) or the OI-122 island effectors `lifA/efa1` (OR = 6.01), `nleB` (OR = 4.20), and `nleE` (OR = 3.80) in aEPEC.

What Do These Findings Mean?

- Routine diagnostic panels that rely solely on canonical backbone markers (`aatA`/`aggR` for EAEC, unstratified `eae` for aEPEC, or age-unstratified `daaC`/`afaBC` for DAEC) cannot distinguish widespread endemic colonization from virulent infection.

- Clinical microbiology laboratories, enteric surveillance networks, and vaccine developers in Africa and globally should incorporate second-tier adhesin and toxin targets (`sepA`, `aafA`, `agg3A`, `CS22`, `pet`, `sat`, `bfpA`, and OI-122 `lifA/efa1`–`nleB`–`nleE`), apply age-stratified interpretation for DAEC in children aged ≥ 12 months, and adopt multi-locus molecular or genomic workflows capable of detecting two-way hybrid DEC lineages.

Introduction

Global and African Burden of Diarrhoeagenic *Escherichia coli* (DEC)

Diarrhoeal disease remains a leading cause of preventable morbidity, linear growth faltering, and under-five mortality worldwide, with the heaviest clinical burden concentrated in sub-Saharan Africa and South Asia (Kotloff et al., 2013; Platts-Mills et al., 2015; Troeger et al., 2018). Among bacterial enteropathogens, diarrhoeagenic *Escherichia coli* (DEC) represents the most common and genetically diverse group colonizing and infecting infants and young children in low- and middle-income countries (LMICs) (Nataro & Kaper, 1998; Croxen et al., 2013; Kaper et al., 2004). Since the classic synthesis by Nataro and Kaper (1998), DEC isolates have been classified into six canonical pathotypes based on conserved virulence plasmids, chromosomal pathogenicity islands, and in vitro tissue-culture adherence phenotypes: 1. **Enteropathogenic *E. coli* (EPEC):** Characterized by the chromosomal Locus of Enterocyte Effacement (LEE) encoding the outer-membrane adhesin intimin (`eae`) that induces attaching-and-effacing (A/E) lesions, further subdivided by the presence (`tEPEC`) or absence (`aEPEC`) of the EPEC adherence factor (EAF) plasmid encoding the bundle-forming pilus (`bfpA`) (Kaper et al., 2004; Trabulsi et al., 2002). 2. **Enterotoxigenic *E. coli* (ETEC):** Defined by plasmid-encoded heat-labile (`elt` / LT) and/or heat-stable (`estA` / STp, STh) enterotoxins paired with over 25 antigenically distinct colonization factors (CFs) that mediate small-intestinal attachment and secretory diarrhoea (Qadri et al., 2005; Kotloff et al., 2013). 3. **Enteroaggregative *E. coli* (EAEC):** Phenotypically defined by stacked-brick aggregative adherence to HEp-2 cells and molecularly identified by the pAA virulence plasmid encoding the AraC/XylS-family master transcriptional regulator `aggR`, the anti-aggregation protein transporter `aatA` (CVD432 probe), dispersin (`aap`), aggregative adherence fimbriae (`AAF/I`–`V`), and chromosomal type VI secretion systems (`aaiC`) (Harrington et al., 2006; Estrada-Garcia & Navarro-Garcia, 2012; Hebbelstrup Jensen et al., 2014). 4. **Shiga toxin-producing / Enterohaemorrhagic *E. coli* (STEC/EHEC):** Defined by lambdoid bacteriophage-encoded Shiga toxins (`stx1`, `stx2`), with the EHEC subset additionally harbouring the LEE island (`eae`) and enterohaemolysin (`ehxA`) associated with haemorrhagic colitis and haemolytic uraemic syndrome (Kaper et al., 2004; Lupindu, 2018). 5. **Enteroinvasive *E. coli* (EIEC):** Closely related to *Shigella* spp., harbouring the large invasion plasmid pINV (`ipaH`, `ial`, `virF`) that directs colonic epithelial invasion and dysentery (Croxen et al., 2013). 6. **Diffusely Adherent *E. coli* (DAEC):** Characterized by diffuse adherence across the entire HEp-2 cell surface mediated by Afa/Dr diffusely adhering fimbrial and afimbrial adhesins (`afa/draBC`, `daaC`, `daaE`, `afaE` subtypes) that bind human decay-accelerating factor (DAF/CD55) and carcinoembryonic antigen-related cell adhesion molecules (CEACAMs) (Servin, 2005;Servin, 2014).

The Epidemiological Attribution Paradox in Africa and Globally

Over the past two decades, landmark multicentre studies—including the seven-site Global Enteric Multicenter Study (GEMS) across four African countries (The Gambia, Mali, Mozambique, and Kenya) and three Asian sites (Kotloff et al., 2013; Liu et al., 2016), the eight-country Malnutrition and Enteric Disease Study (MAL-ED) (Platts-Mills et al., 2015), and the Vaccine Impact on Diarrhea in Africa (VIDA) study in The Gambia, Mali, and Kenya (Kasumba et al., 2023; Ochieng et al., 2023)—alongside single-country investigations in Nigeria (Okeke et al., 2000; Onanuga et al., 2014; Saka et al., 2019; Akinlabi et al., 2023a), Mozambique (Mandomando et al., 2007; Garrine et al., 2020; Manhique-Coutinho et al., 2022), Kenya (Sang et al., 2012; Shah et al., 2016), South Africa (Tau et al., 2012; Tau et al., 2012), Ghana (Opintan et al., 2010), Tanzania (Moyo et al., 2007; Vargas et al., 2004), and Latin America (Girón et al., 1991; Levine et al., 1993; Scaletsky et al., 2002; Ochoa et al., 2009; Lima et al., 2013) have documented a striking epidemiological paradox.

When single-arm case prevalence is measured in children presenting with diarrhoea, **EAEC** ($20-45\%$), **aEPEC** ($8-20\%$), and **DAEC** ($4-15\%$) consistently rank among the most frequently detected bacterial enteropathogens in Africa and globally (Kalule et al., 2024; Lima et al., 2013). Yet when matched asymptomatic community controls are simultaneously enrolled, the unstratified case-control odds ratios for EAEC (`aatA`/`aggR`), aEPEC (`eae+ bfpA-`), and overall DAEC (`daaC`/`afaBC`) frequently hover near unity ($OR \approx 0.85-1.38$), contrasting sharply with the robust case-control pathogenicity of ETEC ($OR > 2.0$), tEPEC ($OR > 2.0$), and EIEC ($OR > 2.0$) (Kotloff et al., 2013; Platts-Mills et al., 2015; Akinlabi et al., 2023a). Even in the absence of overt acute watery diarrhoea, subclinical colonization with EAEC and aEPEC is strongly associated with intestinal inflammation (elevated faecal myeloperoxidase, calprotectin, and neopterin), environmental enteric dysfunction (EED), and linear growth stunting across African and Latin American birth cohorts (Lima et al., 2013; Rogawski et al., 2017; Havt et al., 2017).

Four Competing Hypotheses Underlying the Attribution Paradox

Four non-mutually exclusive hypotheses have been proposed in the global literature to explain the divergence between high single-arm prevalence and attenuated case-control attribution: 1. **Diagnostic Modality Sensitivity and Primer Discordance:** Conventional endpoint multiplex PCR assays targeting conserved plasmid markers suffer from primer-binding mismatches in diverse African lineages and fail to quantify pathogen load, whereas quantitative real-time PCR (qPCR / TaqMan Array Card [TAC]) and whole-genome sequencing (WGS) reveal substantially higher pathogen prevalence, co-infection rates, and allelic diversity (Liu et al., 2016; Platts-Mills et al., 2015; Akinlabi et al., 2023b). 2. **Emergence of Two-Way Hybrid Pathotypes (`DEC–DEC` and `DEC–ExPEC`):** Horizontal transfer of conjugative virulence plasmids and bacteriophages across *E. coli* phylogroups generates hybrid pathotypes—such as `EAEC/ETEC`, `EAEC/EPEC`, `EAEC/STEC`, `EPEC/ETEC`, and `EAEC/UPEC`—that combine the mucosal biofilm persistence of EAEC or ExPEC with acute enterotoxins or cytotoxins, confounding rigid single-marker diagnostic schemes (Karch et al., 2012; Santos et al., 2020; Akinlabi et al., 2025; Medugu et al., 2025). 3. **Host Developmental Age-Ontogeny:** Mucosal receptor expression (such as brush-border DAF/CD55 and CEACAMs bound by Afa/Dr DAEC adhesins) and passive maternal secretory IgA immunity against plasmid-encoded fimbriae evolve rapidly across the first two

years of life, potentially rendering a pathotype non-pathogenic in young infants (< 12 months) while driving acute diarrhoea in weaned toddlers (≥ 12 – 59 months) (Levine et al., 1993; Scaletsky et al., 2002; Ochoa et al., 2009; Garrine et al., 2020). 4. **Accessory Virulence Locus Heterogeneity Within Canonical Pathotypes:** Canonical pathotype-defining markers (`aatA`, `aggR`, `eae`, `daaC`) capture broad genetic backbones that include both avirulent commensal-like strains and hypervirulent clinical clones harbouring specific Serine Protease Autotransporters of *Enterobacteriaceae* (SPATEs; `sepA`, `pet`, `sat`), fimbrial adhesin variants (`aafA` [AAF/II], `agg3A` [AAF/III], `CS22`, `afaE-I/III/V`), or non-LEE-encoded type III secretion system effectors on genomic island OI-122 (`lifA/efa1`, `nleB`, `nleE`) (Boisen et al., 2012; Boisen et al., 2020; Afset et al., 2006; Hazen et al., 2016; Ochieng et al., 2023).

Rationale and Objectives of the Two-Part Meta-Analysis

While earlier regional reviews (e.g., Kalule et al., 2024) provided valuable single-arm prevalence compilations across African hospital series through 2022, critical epidemiological, diagnostic, and genetic enigmas have remained unaddressed. Specifically, no quantitative meta-analysis has simultaneously: (i) updated the African evidence base through March 2026 to incorporate recent large-scale WGS and TAC cohorts ($k = 28$ primary studies, $N = 15,750$); (ii) directly contrasted single-arm Freeman–Tukey random-effects case prevalence against matched case-control odds ratios across all DEC pathotypes; (iii) quantified diagnostic discordance and two-way hybrid pathotype detection across conventional PCR, qPCR/TAC, and WGS platforms; and (iv) synthesized the global and African case-control literature at the individual virulence gene level ($k = 30$ cohorts, 154 locus-level 2×2 evaluations across five continents) to identify which specific genetic determinants and host age strata resolve the EAEC, DAEC, and EPEC carriage paradox.

Here, we report a comprehensive, two-part PRISMA 2020 systematic review and random-effects meta-analysis addressing DEC pathotypes, diagnostic modalities, hybrid lineages, and virulence loci in Africa and globally.

Methods

Ethics Statement

This study is a secondary systematic review and meta-analysis of previously published peer-reviewed epidemiological cohorts and publicly accessible genomic repositories. No primary human participants were recruited, and no individualized patient health identifiers were accessed. Ethical approvals for primary study enrolment, informed consent, and biological specimen collection were obtained by the principal investigators of the original included studies under protocols approved by their respective institutional review boards, as detailed in the original publications and Supplementary Table S3.

Study Design and Reporting Framework

This investigation was designed and executed as a two-part quantitative systematic review and meta-analysis in strict accordance with the Preferred Reporting Items for Systematic Reviews

and Meta-Analyses (PRISMA 2020) statement (Page et al., 2021) (**S1 Table**) and the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) guidelines. The dual-track workflow is illustrated in **Fig 1**: - **Part A (African & Multicentre DEC Pathotype Epidemiology, Diagnostic Platforms, and Hybrid Lineages):** Synthesizes primary molecular and genomic studies conducted across all five United Nations African subregions (Western, Eastern, Southern, Central, and Northern Africa), alongside multicentre reference cohorts with substantial African enrolment (GEMS, MAL-ED, and VIDA), published between January 1, 2000, and March 25, 2026, establishing an updated, high-resolution continental evidence base across 18 countries. - **Part B (Global & African Virulence Locus Case-Control Meta-Analysis):** Synthesizes matched and unmatched case-control studies from Africa, Latin America, Asia, Europe, and Oceania (January 1, 1990 – March 25, 2026) reporting extractable 2 × 2 contingency counts (cases positive/total vs. asymptomatic controls positive/total) for individual virulence loci within EAEC (`sepA`, `CS22`, `aafA`, `agg3A`, `agg4A`, `aggA`, `AAF_any`, `pet`, `sat`, `aaiC`, `aatA`, `aggR`, `irp2`, `fyuA`, `aap`, `pic`, `astA`), DAEC (`daaC`/`afaBC` overall and stratified by `<12` vs. `>=12–59` months, `afaE`, and `sat`), and EPEC (`bfpA` stratified by `<12` vs. `>=12–59` months, unstratified `eae`, and OI-122 effectors `lifA/efa1`, `nleB`, `nleE`, and `astA`).

Information Sources and Search Strategy

Systematic literature searches were executed across six bibliographic databases and genomic repositories: PubMed/MEDLINE, Scopus, Web of Science Core Collection (WoS), Embase, EBSCOhost, and African Journals Online (AJOL), supplemented by targeted screening of NCBI Sequence Read Archive (SRA) / European Nucleotide Archive (ENA) BioProjects and reference lists of prior reviews (Hebbelstrup Jensen et al., 2014; Lupindu, 2018; Kalule et al., 2024).

For **Part A**, the Boolean search string combined pathotype terms, diagnostic terms, and geographic filters for all 54 African sovereign states: > `("diarrhoeagenic Escherichia coli" OR "diarrheagenic Escherichia coli" OR "DEC" OR "enteroaggregative" OR "EAEC" OR "enteropathogenic" OR "EPEC" OR "enterotoxigenic" OR "ETEC" OR "enteroinvasive" OR "EIEC" OR "Shiga toxin-producing" OR "STEC" OR "enterohaemorrhagic" OR "EHEC" OR "diffusely adherent" OR "DAEC" OR "hybrid Escherichia coli" OR "heteropathogenic") AND ("PCR" OR "multiplex PCR" OR "qPCR" OR "TaqMan Array Card" OR "whole genome sequencing" OR "WGS" OR "virulence genes") AND ("Africa" OR "sub-Saharan Africa" OR "West Africa" OR "East Africa" OR "Southern Africa" OR "Central Africa" OR "North Africa" OR "Algeria" OR "Angola" OR "Benin" OR "Botswana" OR "Burkina Faso" OR "Burundi" OR "Cabo Verde" OR "Cameroon" OR "Central African Republic" OR "Chad" OR "Comoros" OR "Congo" OR "Cote d'Ivoire" OR "Djibouti" OR "Egypt" OR "Equatorial Guinea" OR "Eritrea" OR "Eswatini" OR "Ethiopia" OR "Gabon" OR "Gambia" OR "Ghana" OR "Guinea" OR "Guinea-Bissau" OR "Kenya" OR "Lesotho" OR "Liberia" OR "Libya" OR "Madagascar" OR "Malawi" OR "Mali" OR "Mauritania" OR "Mauritius" OR "Morocco" OR "Mozambique" OR "Namibia" OR "Niger" OR "Nigeria" OR "Rwanda" OR "Sao Tome" OR "Senegal" OR "Seychelles" OR "Sierra Leone" OR "Somalia" OR "South Africa" OR "South Sudan" OR "Sudan" OR "Tanzania" OR "Togo" OR "Tunisia" OR "Uganda" OR "Zambia" OR "Zimbabwe")`.

For **Part B**, the global search removed geographic restrictions and targeted locus-specific case-control attribution: > `("enteroaggregative Escherichia coli" OR "EAEC" OR "diffusely adherent Escherichia coli" OR "DAEC" OR "enteropathogenic Escherichia coli" OR "EPEC" OR "atypical EPEC" OR "aEPEC") AND ("case-control" OR "healthy controls" OR "asymptomatic controls" OR "diarrhea" OR "diarrhoea") AND ("aggR" OR "aatA" OR "CVD432" OR "aap" OR "aaiC" OR "sepA" OR "pet" OR "sat" OR "pic" OR "astA" OR "aafA" OR "aggA" OR "agg3A" OR "agg4A" OR "CS22" OR "irp2" OR "fyuA" OR "daaC" OR "afaBC" OR "draBC" OR "afaE" OR "bfpA" OR "eae" OR "lifA" OR "efa1" OR "nleB" OR "nleE" OR "OI-122")`.

Eligibility Criteria (PICOS Framework)

1. **Population:** Human patients presenting with acute or persistent diarrhoea (cases) and, where applicable, asymptomatic community or hospital controls without diarrhoea for at least 7–14 days prior to sampling. Studies restricted exclusively to animal reservoirs, food matrices, or environmental water without human clinical isolates were excluded.
2. **Intervention / Exposure (Diagnostic Ascertainment):** Molecular confirmation of DEC pathotypes or individual virulence genes via endpoint PCR, multiplex PCR, quantitative real-time PCR (qPCR / TAC), DNA hybridization probes (`CVD432`, `daaC`), or whole-genome sequencing (WGS). Studies relying solely on O:H serotyping or biochemical profiling without molecular/probe confirmation were excluded (Okeke et al., 2000; Kalule et al., 2024).
3. **Comparator:** For case-control odds ratio models in Part A and Part B, an explicitly defined asymptomatic human control group from the same geographic source population tested using the identical diagnostic platform.
4. **Outcomes:** Extractable numerator and denominator counts (r / n) for pathotype-level case prevalence (Part A), matched case-control pathotype counts (a, b, c, d; Part A), hybrid pathotype co-detections (Part A), and locus-specific 2×2 case-control counts across EAEC, DAEC, and EPEC (Part B).
5. **Handling of Unassayed Pathotypes and Overlapping Cohorts:** When a primary study assayed only a subset of DEC pathotypes (e.g., EAEC-specific WGS cohorts or EPEC/ETEC-focused surveys), unassayed pathotypes were coded as missing (`NA` / `-1`) rather than zero so that pathotype-specific random-effects denominators included only cohorts that actively tested for that pathotype. Where multiple publications analysed overlapping subsets of the same parent cohort (e.g., GEMS primary TAC re-analysis vs. GEMS EAEC sub-studies), the primary non-overlapping denominator was used for Part A pathotype prevalence while locus-specific sub-studies were assigned to their corresponding gene models in Part B.

Data Extraction and Quality Assessment

From each eligible publication, we extracted first author, publication year, study country, UN subregion, study design (matched case-control vs. cross-sectional/genomic surveillance), participant age group, clinical setting, diagnostic modality (conventional/multiplex PCR, qPCR/TAC, or WGS), total sample size (N), diarrhoeal case denominator (n_{cases}), asymptomatic control denominator (n_{controls}), pathotype-specific positive counts in cases and controls (EAEC,

ETEC, EPEC, tEPEC, aEPEC, DAEC, STEC, EIEC), hybrid pathotype counts and gene combinations, and locus-specific 2×2 contingency tables (a = cases positive, b = cases negative, c = controls positive, d = controls negative). Methodological quality and risk of bias were evaluated using the Joanna Briggs Institute (JBI) Critical Appraisal Checklist for Prevalence Studies (9 items; Part A) (Munn et al., 2015) and the Newcastle–Ottawa Scale (NOS; 9-star system across Selection, Comparability, and Exposure ascertainment; Part A and Part B case-control cohorts) (Wells et al., 2014) (**S4 Table**; visual dashboard in **S8 Fig**).

Statistical Meta-Analysis Models

2.5.1 Single-Arm Pooled Prevalence (PFT Transformation with REML and DL)

Because pathotype and hybrid proportions frequently approach boundary values (0 % to 5 % for EIEC, STEC, and hybrids), raw proportions violate Gaussian normality assumptions. We applied the Freeman–Tukey double-arcsine variance-stabilizing transformation (PFT) (Freeman & Tukey, 1950) to each study's positive count r_i and sample size n_i:

$$\theta_i = \arcsin\left(\sqrt{\frac{r_i}{n_i+1}}\right) + \arcsin\left(\sqrt{\frac{r_i+1}{n_i+1}}\right), \quad v_i = \mathrm{Var}(\theta_i) = \frac{1}{n_i+0.5}$$

Between-study variance (τ^2) was estimated primarily via Restricted Maximum Likelihood (REML) optimizing the restricted log-likelihood:

$$L_{\mathrm{REML}}(\tau^2) = -\frac{1}{2}\left[\sum_{i=1}^{k} \ln(v_i+\tau^2) + \ln\left(\sum_{i=1}^{k} \frac{1}{v_i+\tau^2}\right) + \sum_{i=1}^{k} \frac{(\theta_i - \hat{\theta}_{\mathrm{REML}}(\tau^2))^2}{v_i+\tau^2}\right]$$

and cross-checked using the non-iterative DerSimonian–Laird (DL) method of moments estimator (DerSimonian & Laird, 1986). Pooled transformed estimates $\hat{\theta}$ and their 95 % confidence bounds were back-transformed to natural proportions ($\hat{p} \in [0,1]$) using Miller's exact inverse transformation incorporating the harmonic mean of study sample sizes $\tilde{n}_h = k / \sum_{i=1}^{k} n_i^{-1}$ (Miller, 1978; Barendregt et al., 2013).

2.5.2 Case-Control Odds Ratio Pooling (REML, DL, and Knapp–Hartung Adjustments)

For each 2×2 case-control comparison (a_i, b_i, c_i, d_i), the natural log Odds Ratio ($y_i = \ln \mathrm{OR}_i$) and its asymptotic within-study variance (v_i) were calculated as:

$$y_i = \ln\left(\frac{a_i d_i}{b_i c_i}\right), \quad v_i = \frac{1}{a_i} + \frac{1}{b_i} + \frac{1}{c_i} + \frac{1}{d_i}$$

In studies containing any zero cell ($a_i = 0, b_i = 0, c_i = 0$, or $d_i = 0$), the Haldane–Anscombe continuity correction (+0.5 added to all four cells of that study) was applied prior to logarithmic transformation (Haldane, 1956; Sweeting et al., 2004). Random-effects inverse-variance weights $w_i^* = (v_i + \hat{\tau}^2_{\mathrm{REML}})^{-1}$ yielded the pooled log-odds ratio $\hat{\mu}_{\mathrm{REML}} = \sum w_i^* y_i / \sum w_i^*$ and standard Wald z-test 95 % confidence intervals $\exp(\hat{\mu}_{\mathrm{REML}} \pm 1.96\,\mathrm{SE}(\hat{\mu}_{\mathrm{REML}}))$. To account for uncertainty in estimating $\hat{\tau}^2$ when the number of studies k is moderate or small ($k = 3-13$), we additionally

computed the Hartung–Knapp–Sidik–Jonkman (KH) small-sample variance scaling factor $q^{*} = max\left(1, \frac{1}{k-1}\sum w_i^{*}\left(y_i - \hat{\mu}_{\mathrm{REML}}\right)^2\right)$ and evaluated 95 % KH confidence intervals and two-sided P-values against a Student's t-distribution with $df = k - 1$ degrees of freedom (IntHout et al., 2014;Jackson et al., 2017).

2.5.3 Heterogeneity, Subgroup Stratification, and Publication Bias Diagnostics

Between-study heterogeneity was quantified using Cochran's Q statistic ($Q = \sum w_i\left(y_i - \hat{\mu}_{\mathrm{FE}}\right)^2$, χ^2_{k-1}), the Higgins–Thompson I^2 inconsistency index ($I^2 = max\left(0, \left(Q - df\right)/Q\right) \times 100\,\%$), and $\hat{\tau}^2_{\mathrm{REML}}$ (Higgins et al., 2003). Pre-specified subgroup analyses stratified studies by UN African subregion, diagnostic platform (conventional PCR vs. qPCR/TAC vs. WGS), and host developmental age stratum (< 12 months vs. ≥ 12 – 59 months). Small-study effects and publication bias were evaluated via funnel plots of $\ln \mathrm{OR}_i$ against $\mathrm{SE}\left(\ln \mathrm{OR}_i\right)$, Egger's weighted linear regression test of the standardized normal deviate ($\mathrm{SND}_i = y_i / \sqrt{v_i}$) on precision ($1 / \sqrt{v_i}$) using the exact standard error of the regression intercept (Egger et al., 1997), and Begg and Mazumdar's rank-correlation test (Begg & Mazumdar, 1994). All statistical computations and high-resolution figures were executed programmatically in Python (`scipy.stats`, `scipy.optimize`, `numpy`, `pandas`, `matplotlib`, and `seaborn`).

2.5.4 Multivariate Meta-Regression and Heterogeneity Variance Partitioning

To formally partition between-study variance and evaluate the independent explanatory power of prespecified methodological covariates, we constructed fixed- and mixed-effects meta-regression models (Borenstein et al., 2009; Higgins & Thompson, 2004):

$$y_i = x_i \beta + u_i + \epsilon_i, u_i \sim N\left(0, \tau^2_{\mathrm{res}}\right), \epsilon_i \sim N\left(0, v_i\right)$$

where y_i represents the transformed effect size (Freeman–Tukey double-arcsine proportion θ_i or logit prevalence), v_i is the study-specific sampling variance, x_i is the vector of study-level moderators, β is the vector of meta-regression slope coefficients, and τ^2_{res} is the residual between-study variance. Three primary moderators were specified: 1. **Diagnostic Modality:** Whole-genome sequencing (WGS) and culture-based methods (colony hybridization) versus conventional multiplex PCR (reference). 2. **Study Era:** Pre-2015 historical cohorts versus contemporary cohorts (2015–2024; reference). 3. **Age Restriction:** Pediatric-restricted cohorts (children < 59 months; reference) versus all-ages / mixed-population cohorts.

Parameters were estimated via restricted maximum likelihood (REML) utilizing numerical optimization algorithms (`scipy.optimize`). The proportion of true between-study variance explained by covariates was quantified through the pseudo-R^2 (R^2_{analog}) index:

$$R^2 = max\left(0, \frac{\tau^2_0 - \tau^2_{\mathrm{res}}}{\tau^2_0}\right) \times 100\,\%$$

where τ^2_0 is the total between-study variance from the unadjusted intercept-only random-effects model. The omnibus hypothesis that all moderator coefficients simultaneously equal zero (

$H_0: \beta_{1:p} = 0$) was tested using the Wald χ^2 statistic (Q_M) and Knapp–Hartung adjusted F-statistic ($F(p-1, k-p)$). Residual heterogeneity was evaluated via Cochran's Q_E test and residual inconsistency $I^2_{\text{res}} = max(0, (Q_E - df_E)/Q_E) \times 100\%$.

2.5.5 Evidence Certainty Grading (GRADE Framework)

To transition from quantitative effect pooling to structured clinical and public health guidance, we implemented the Grading of Recommendations Assessment, Development and Evaluation (GRADE) framework for prognostic, etiological, and observational epidemiological evidence (Guyatt et al., 2008; Schünemann et al., 2013; Iorio et al., 2015). For each DEC pathotype and high-attribution virulence locus, evidence certainty was systematically appraised across five downgrading domains: 1. **Risk of Bias:** Methodological quality and confounding control assessed via the Newcastle–Ottawa Scale (case-control comparability, community control selection) and JBI checklist. 2. **Inconsistency:** Unexplained statistical heterogeneity across studies ($I^2 \geq 50\%$, $\tau^2 > 0.05$, or non-overlapping confidence intervals). 3. **Indirectness:** Generalizability and applicability of study populations, diagnostic exposure definitions, and clinical diarrhoeal endpoints relative to endemic African and global pediatric settings. 4. **Imprecision:** Width of the 95% confidence interval (e.g., crossing null association thresholds or clinical equivalence bounds) and cumulative event burden. 5. **Publication Bias:** Systematic suppression or selective reporting detected through funnel plot asymmetry and Egger's regression tests.

Upgrading criteria included large magnitude of effect (mOR > 2.0 for moderate upgrade; mOR > 5.0 for strong upgrade) in the absence of plausible confounding, and consistent dose-response or developmental age-stratified gradients. Overall certainty of evidence for each pathotype and locus outcome was categorized into four standardized tiers: **High (⊕⊕⊕⊕)**, **Moderate (⊕⊕⊕○)**, **Low (⊕⊕○○)**, or **Very Low (⊕○○○)**, and summarized in a formal Summary of Findings (SoF) matrix.

Results

Systematic Evidence Architecture, Workflow Mindmap, and Dual-Track Study Inclusion

The comprehensive methodological workflow, systematic evidence architecture, and dual-track inclusion framework are presented in **Fig 1**. The synthesis integrates 55 years of molecular surveillance (1970–2026) across continental Africa and globally, organized into four strategic methodological clusters: (1) Continental African surveillance, pathotype diversity, and antimicrobial resistance mobilome ($k = 28$ primary cohorts, $N = 15,750$); (2) Global case-control virulome meta-analysis ($k = 47$ cohorts, 29 loci across 5 continents); (3) Biostatistical modeling featuring Freeman–Tukey REML random-effects pooling, continuity corrections, and mixed-effects meta-regression; and (4) GRADE multi-domain certainty profiling linked to Africa CDC PGI genomic architecture and WHO AFRO IDSR clinical guideline revisions.

In **Part A** (African and multicentre DEC molecular epidemiology), database searches across PubMed/MEDLINE, Scopus, Web of Science, Embase, EBSCOhost, and African Journals Online (AJOL) identified $2,640$ initial records published between January 2000 and March 2026. Following removal of 984 duplicates, $1,656$ unique titles and abstracts were screened against PICOS criteria, excluding $1,432$ records (non-human/environmental studies, non-African cohorts without African sub-data, or studies lacking molecular typing). Full-text eligibility assessment was performed on 224 articles, of which 196 were excluded (86 lacked pathotype-specific molecular confirmation, 76 had missing denominators or non-extractable counts, and 34 represented overlapping sub-analyses of identical cohorts). Twenty-eight primary African and multicentre cohorts ($k = 28$, comprising $N = 15,750$ participants: $9,360$ diarrhoeal cases and $6,041$ asymptomatic controls across 18 countries in all five UN African subregions) were included in the primary Part A synthesis, establishing the primary continental evidence base for Part A.

In **Part B** (global and African virulence locus case-control meta-analysis), $3,910$ candidate records were identified across bibliographic databases and NCBI/ENA BioProjects. After removing $1,645$ duplicates and screening $2,265$ unique abstracts, 197 full-text articles were evaluated, excluding 172 reports (119 lacked extractable gene-level 2×2 case-control counts, 31 enrolled controls with recent diarrhoea or antibiotic exposure, and 22 were duplicate reports of identical isolate collections). Thirty global and African case-control study entries ($k = 30$ distinct age- or pathotype-stratified cohorts from 25 unique study populations, contributing 154 locus-level 2×2 evaluations across five continents: Africa, Latin America, Asia, Europe, and Oceania) satisfied all inclusion criteria.




Fig. 1 PRISMA 2020 flow diagram depicting systematic study identification, screening, eligibility, and dual-track evidence synthesis.

Fig 1A: PRISMA 2020 Flow Diagram of Included Studies




Fig. 1 PRISMA 2020 flow diagram depicting systematic study identification, screening, eligibility, and dual-track evidence synthesis.

Fig 1B: Minimalist Methodological Evidence Architecture Workflow

Fig 1. Systematic evidence architecture, PRISMA 2020 flow diagrams, and evidence synthesis workflow for the African and global diarrhoeagenic *Escherichia coli* (DEC) systematic review.** (**A**) Dual-track PRISMA 2020 systematic review workflow detailing bibliographic database identification across PubMed/MEDLINE, Scopus, Web of Science Core Collection,

Embase, EBSCOhost, and African Journals Online (AJOL), supplemented by genomic repository mining (NCBI Sequence Read Archive [SRA] and European Nucleotide Archive [ENA] BioProjects). **Track A (Continental African & Multicentre DEC Molecular Epidemiology):** $2,640$ initial records identified, 984 duplicates removed, $1,656$ unique records screened, 224 full-text articles assessed against PICOS eligibility criteria, yielding $k = 28$ primary African and multicentre molecular/genomic cohorts ($N = 15,750$ subjects: $9,360$ acute diarrhoeal cases and $6,041$ matched asymptomatic community controls across 18 African nations) evaluating canonical pathotype prevalence, matched case-control odds ratios, diagnostic platform sensitivities, and two-way hybrid lineages. **Track B (Global & African Virulence Locus Case-Control Attribution):** $3,910$ records identified, $1,645$ duplicates removed, $2,265$ records screened, 197 full-text articles assessed, yielding $k = 30$ unique case-control cohort entries contributing 154 locus-level 2×2 contingency evaluations across five continents (Africa, Latin America, Asia, Europe, and Oceania) evaluating 29 discrete virulence loci within EAEC, DAEC, and EPEC. (**B**) High-level conceptual mindmap outlining the four strategic methodological pillars, biostatistical pooling engines, and genomic-to-epidemiological translation pathways.

Characteristics and Methodological Quality of Included African Cohorts (Part A)

The study-level demographic, clinical, diagnostic, and pathotype characteristics of the $k = 28$ primary Part A cohorts are detailed in **Table 1** and compiled with granular metadata across all 68 surveillance cohorts in **S3 Table**. The chronological deployment of all 12 diagnostic modalities across these 68 African cohorts is presented in **S18 Fig**. Geographically, the included cohorts spanned Western Africa ($k = 12$: Nigeria [$k = 8$: Okeke et al., 2000; Onanuga et al., 2014; Ifeanyi et al., 2015; Akinlabi et al., 2023a, 2023b, 2025; Olayinka et al., 2024; Medugu et al., 2025], Ghana [$k = 2$: Opintan et al., 2010; Dela et al., 2022], The Gambia [$k = 1$: Foster-Nyarko et al., 2021], and Burkina Faso [$k = 1$: Bonkoungou et al., 2021], alongside sites in Mali and The Gambia within GEMS, VIDA, and MAL-ED), Eastern Africa ($k = 8$: Kenya [$k = 2$: Sang et al., 2012; Kiiru et al., 2024], Mozambique [$k = 2$: Mandomando et al., 2007; Garrine et al., 2020], Ethiopia [$k = 2$: Zelelie et al., 2023; Wolde et al., 2024], Tanzania [$k = 1$: Moyo et al., 2007], and Zambia [$k = 1$: Mwape et al., 2023]), Southern Africa ($k = 2$: South Africa [Obi et al., 2004; Mkuhlu et al., 2020]), Central Africa ($k = 2$: Gabon [Mabika Mabika et al., 2021] and Cameroon [Mahamat et al., 2024]), Northern Africa ($k = 1$: Egypt [Ali et al., 2014]), and three landmark multicentre African/global networks ($k = 3$: GEMS [Kotloff et al., 2013], MAL-ED [Platts-Mills et al., 2015], and VIDA [Ochieng et al., 2023]).

Fourteen cohorts ($k = 14$) incorporated matched asymptomatic control arms ($5,449$ diarrhoeal cases and $6,041$ asymptomatic controls), while 14 cohorts ($k = 14$) were cross-sectional hospital/community surveillance or whole-genome sequencing investigations ($3,911$ diarrhoeal cases or characterized clinical isolates). Methodological quality was high across the evidence base (median Newcastle–Ottawa Scale score $8/9$, range $7–9$; median JBI score $8/9$, range $7–9$; **Table 1**, **S4 Table**, and **S8 Fig**).

Table 1. Characteristics, diagnostic modalities, pathotype distributions, hybrid lineages, and quality appraisal of primary African and multicentre studies included in Part A ($k = 28$ primary studies, $N = 15,750$).

Study (Author, Year)	Country (UN Subregion)	Study Design	Age Group	Diagnostic Modality	Sample Size N (Cases / Controls)	EAEC Cases n (%)	ETEC Cases n (%)	EPEC Cases n (%)	DAEC Cases n (%)	STEC Cases n (%)	EIEC Cases n (%)	Hybrid DEC n (%) & Dominant Combinations	Quality Score
Kotloff et al. (2013) [GEMS Africa Sites]	The Gambia, Mali, Mozambique, Kenya (Western/Eastern/Southern Africa)	Matched Case-Control	0–59m	Multiplex PCR & Quantitative TAC	4,110 (1,890 / 2,220)	397/1890 (21.0%)	284/1890 (15.0%)	246/1890 (13.0%)	NA	19/1890 (1.0%)	38/1890 (2.0%)	64/1890 (3.4%)	NOS 9/9
Ochieng et al. (2023) [VIDA Consortium]	The Gambia, Mali, Kenya (Western/Eastern Africa)	Matched Case-Control	0–59m	Quantitative TAC (qPCR)	942 (480 / 462)	280/480 (58.3%)	112/480 (23.3%)	168/480 (35.0%)	NA	24/480 (5.0%)	14/480 (2.9%)	28/480 (5.8%)	NOS 9/9
Platts-Mills et al. (2015) [MAL-ED Africa]	South Africa, Tanzania (Eastern/Southern Africa)	Matched Case-Control / Birth Cohort	0–24m	Quantitative TAC (qPCR)	1,420 (680 / 740)	272/680 (40.0%)	156/680 (22.9%)	163/680 (24.0%)	NA	14/680 (2.1%)	21/680 (3.1%)	34/680 (5.0%)	NOS 9/9
Akinlabi et al. (2023a)	Nigeria (Western Africa)	Matched Case-Control	0–59m	WGS & Multiplex PCR Platform Evaluation	360 (120 / 240)	46/120 (38.3%)	11/120 (9.2%)	15/120 (12.5%)	6/120 (5.0%)	1/120 (0.8%)	1/120 (0.8%)	12/120 (10.0%)	NOS 9/9
Akinlabi et al. (2023b)	Nigeria (Western Africa)	Matched Case-Control	0–59m	Multiplex PCR & WGS	497 (120 / 377)	44/120 (36.7%)	10/120 (8.3%)	16/120 (13.3%)	7/120 (5.8%)	1/120 (0.8%)	1/120 (0.8%)	11/120 (9.2%)	NOS 9/9
Akinlabi et al. (2025)	Nigeria (Western Africa)	Case-Control Genomic Survey	0–59m	WGS (Illumina)	96 (96 / 0)	NA	NA	NA	NA	NA	NA	15/96 (15.6%)	JBI 8/9
Olayinka et al. (2024)	Nigeria (Western Africa)	Matched Case-Control	0–59m	Multiplex PCR	501 (167 / 334)	52/167 (31.1%)	24/167 (14.4%)	21/167 (12.6%)	9/167 (5.4%)	3/167 (1.8%)	2/167 (1.2%)	9/167 (5.4%)	NOS 9/9
Zelelie et al. (2023)	Ethiopia (Eastern Africa)	Matched Case-Control	0–59m	Multiplex PCR	476 (391 / 85)	66/391 (16.9%)	35/391 (9.0%)	23/391 (5.9%)	1/391 (0.3%)	3/391 (0.8%)	20/391 (5.1%)	13/183 (7.1%)	NOS 9/9
Wolde et al. (2024)	Ethiopia (Eastern Africa)	Cross-Sectional WGS	Mixed	WGS (Illumina)	260 (79 / 0)	34/79 (43.0%)	18/79 (22.8%)	21/79 (26.6%)	6/79 (7.6%)	7/79 (8.9%)	2/79 (2.5%)	14/79 (17.7%)	JBI 8/9
Mwape et al. (2023)	Zambia (Eastern/Southern Africa)	Matched Case-Control	0–36m	Quantitative Real-Time PCR Array	385 (314 / 71)	124/314 (39.5%)	93/314 (29.6%)	143/314 (45.5%)	18/314 (5.7%)	14/314 (4.5%)	6/314 (1.9%)	62/314 (19.7%)	NOS 9/9
Kiiru et al. (2024, 2025)	Kenya (Eastern Africa)	Cross-Sectional WGS	0–59m	WGS (Illumina)	219 (75 / 0)	38/75 (50.7%)	19/75 (25.3%)	24/75 (32.0%)	5/75 (6.7%)	4/75 (5.3%)	1/75 (1.3%)	14/75 (18.7%)	JBI 8/9
Mkuhlu et al. (2020)	South Africa (Southern Africa)	Cross-Sectional Hospital	Mixed	Multiplex PCR	265 (265 / 0)	48/265 (18.1%)	106/265 (40.0%)	31/265 (11.7%)	37/265 (14.0%)	NA	0/265 (0.0%)	14/265 (5.3%)	JBI 8/9

Study (Author, Year)	Country (UN Subregion)	Study Design	Age Group	Diagnostic Modality	Sample Size N (Cases / Controls)	EAEC Cases n (%)	ETEC Cases n (%)	EPEC Cases n (%)	DAEC Cases n (%)	STEC Cases n (%)	EIEC Cases n (%)	Hybrid DEC n (%) & Dominant Combinations	Quality Score
Garrine et al. (2020)	Mozambique (Southern Africa)	Matched Case-Control	0–59m	Multiplex PCR	1,179 (588 / 591)	48/588 (8.2%)	42/588 (7.1%)	52/588 (8.8%)	24/588 (4.1%)	2/588 (0.3%)	1/588 (0.2%)	14/588 (2.4%)	NOS 9/9
Onanuga et al. (2014)	Nigeria (Western Africa)	Matched Case-Control	0–24m	Multiplex PCR	730 (201 / 529)	16/201 (8.0%)	8/201 (4.0%)	6/201 (3.0%)	NA	19/201 (9.5%)	1/201 (0.5%)	NA	NOS 9/9
Mahamat et al. (2024)	Cameroon (Central Africa)	Cross-Sectional Hospital	0–59m	Multiplex PCR	184 (184 / 0)	68/184 (37.0%)	31/184 (16.8%)	42/184 (22.8%)	14/184 (7.6%)	6/184 (3.3%)	3/184 (1.6%)	28/184 (15.2%)	JBI 8/9
Medugu et al. (2025)	Nigeria (Western Africa)	Cross-Sectional WGS	Mixed	WGS (Illumina)	107 (107 / 0)	44/107 (41.1%)	12/107 (11.2%)	19/107 (17.8%)	8/107 (7.5%)	2/107 (1.9%)	1/107 (0.9%)	20/107 (18.7%)	JBI 8/9
Foster-Nyarko et al. (2021)	The Gambia (Western Africa)	Community Genomic Cohort	36–60m	WGS (Illumina)	88 (0 / 88)	NA	NA	NA	NA	NA	NA	8/88 (9.1%)	NOS 9/9
Dela et al. (2022)	Ghana (Western Africa)	Cross-Sectional Hospital	0–59m	Multiplex PCR	122 (98 / 0)	18/98 (18.4%)	8/98 (8.2%)	9/98 (9.2%)	2/98 (2.0%)	1/98 (1.0%)	0/98 (0.0%)	3/98 (3.1%)	JBI 8/9
Mabika Mabika et al. (2021)	Gabon (Central Africa)	Cross-Sectional Hospital	0–59m	Multiplex PCR	120 (120 / 0)	31/120 (25.8%)	34/120 (28.3%)	26/120 (21.7%)	8/120 (6.7%)	34/120 (28.3%)	2/120 (1.7%)	21/120 (17.5%)	JBI 8/9
Bonkoungou et al. (2021)	Burkina Faso (Western Africa)	Cross-Sectional Clinical	0–59m	16-Plex PCR	285 (285 / 0)	61/285 (21.4%)	26/285 (9.1%)	27/285 (9.5%)	9/285 (3.2%)	11/285 (3.9%)	4/285 (1.4%)	18/285 (6.3%)	JBI 8/9
Ali et al. (2014)	Egypt (Northern Africa)	Matched Case-Control	0–59m	Multiplex PCR	105 (62 / 43)	19/62 (30.6%)	8/62 (12.9%)	11/62 (17.7%)	2/62 (3.2%)	3/62 (4.8%)	1/62 (1.6%)	4/62 (6.5%)	NOS 9/9
Mandomando et al. (2007)	Mozambique (Southern Africa)	Matched Case-Control	0–59m	Multiplex PCR	529 (352 / 177)	51/352 (14.5%)	34/352 (9.7%)	38/352 (10.8%)	11/352 (3.1%)	4/352 (1.1%)	2/352 (0.6%)	8/352 (2.3%)	NOS 9/9
Moyo et al. (2007)	Tanzania (Eastern Africa)	Cross-Sectional Hospital	0–59m	Multiplex PCR	647 (647 / 0)	95/647 (14.7%)	42/647 (6.5%)	39/647 (6.0%)	18/647 (2.8%)	7/647 (1.1%)	3/647 (0.5%)	14/647 (2.2%)	JBI 8/9
Okeke et al. (2000)	Nigeria (Western Africa)	Matched Case-Control	0–59m	Colony Hybridization & PCR	168 (84 / 84)	42/84 (50.0%)	12/84 (14.3%)	14/84 (16.7%)	5/84 (6.0%)	1/84 (1.2%)	1/84 (1.2%)	6/84 (7.1%)	NOS 9/9
Sang et al. (2012)	Kenya (Eastern Africa)	Cross-Sectional Hospital	0–59m	Multiplex PCR	855 (855 / 0)	248/855 (29.0%)	162/855 (18.9%)	179/855 (20.9%)	42/855 (4.9%)	26/855 (3.0%)	11/855 (1.3%)	29/855 (3.4%)	JBI 8/9
Obi et al. (2004)	South Africa (Southern Africa)	Cross-Sectional Survey	0–59m	Colony Hybridization & PCR	350 (350 / 0)	84/350 (24.0%)	49/350 (14.0%)	56/350 (16.0%)	21/350 (6.0%)	14/350 (4.0%)	7/350 (2.0%)	12/350 (3.4%)	JBI 8/9
Ifeanyi et al. (2015)	Nigeria (Western Africa)	Cross-Sectional Hospital	0–59m	Multiplex PCR	300 (300 / 0)	72/300 (24.0%)	36/300 (12.0%)	45/300 (15.0%)	12/300 (4.0%)	9/300 (3.0%)	3/300 (1.0%)	11/300 (3.7%)	JBI 8/9

Study (Author, Year)	Country (UN Subregion)	Study Design	Age Group	Diagnostic Modality	Sample Size N (Cases / Controls)	EAEC Cases n (%)	ETEC Cases n (%)	EPEC Cases n (%)	DAEC Cases n (%)	STEC Cases n (%)	EIEC Cases n (%)	Hybrid DEC n (%) & Dominant Combinations	Quality Score
Opintan et al. (2010)	Ghana (Western Africa)	Cross-Sectional Hospital	0–59m	Multiplex PCR	450 (450 / 0)	108/450 (24.0%)	54/450 (12.0%)	63/450 (14.0%)	18/450 (4.0%)	14/450 (3.1%)	5/450 (1.1%)	16/450 (3.6%)	JBI 8/9

Note: NA indicates that the specific pathotype was not assayed in that study and was therefore excluded from the denominator for that pathotype's random-effects pooling. Quality scores reflect the Newcastle–Ottawa Scale (NOS, /9) for case-control designs and the Joanna Briggs Institute (JBI, /9) checklist for cross-sectional prevalence designs.

Subregional Prevalence, Diagnostic Modality Gradients, and Hybrid Pathotype Detection (Part A)

Across all $k = 28$ primary studies in Part A (**Table 2**, **Table 3**, and **Fig 2**), **EAEC** was the most prevalent DEC pathotype in African diarrhoeal cases, yielding a pooled Freeman–Tukey random-effects (PFT REML) prevalence of 28.3% (95% CI: 23.2 – 33.7%; $k = 26$, $2,406 / 9,264$; $I^2 = 96.6\%$, $\tau^2 = 0.086$; individual study forest plots and subregional raincloud distributions in **S1 Fig**; age-stratified prevalence distributions across 9 DEC categories in **S20 Fig**), followed by **total EPEC** at 16.1% (95% CI: 12.6 – 19.8%; $k = 26$, $1,497 / 9,264$; $I^2 = 95.2\%$) and **ETEC** at 14.9% (95% CI: 11.8 – 18.1%; $k = 26$, $1,426 / 9,264$; $I^2 = 93.4\%$; **S2 Fig**). When EPEC was stratified by the EAF plasmid bundle-forming pilus gene (`bfpA`), **atypical EPEC (`eae+ bfpA-`)** accounted for nearly twice the case burden (10.9%, 95% CI: 6.5 – 16.3%; $k = 11$, $581 / 5,128$) of **typical EPEC (`eae+ bfpA+`)** (6.0%, 95% CI: 4.2 – 8.1%; $k = 12$, $319 / 5,329$; paired cohort comparison in **S11 Fig**). Lower pooled case prevalences were observed for **DAEC** (4.9%, 95% CI: 3.8 – 6.2%; $k = 22$, $283 / 6,013$; $I^2 = 77.2\%$), **STEC/EHEC** (3.2%, 95% CI: 2.0 – 4.7%; $k = 25$, $243 / 8,999$; $I^2 = 87.7\%$), and **EIEC** (1.4%, 95% CI: 1.0 – 1.9%; $k = 26$, $151 / 9,264$; $I^2 = 65.2\%$; study-by-study forest plots presented in **S15 Fig**).

Subregional meta-analysis (**Table 3**, **Fig 2A**, and **S9 Fig**) confirmed that EAEC was the leading DEC pathotype across all five UN African subregions: Eastern Africa (33.7%, 95% CI: 23.9 – 44.4%), Central Africa (31.6%, 95% CI: 21.4 – 42.9%), Northern Africa (30.9%, 95% CI: 20.1 – 42.8%), Western Africa (30.1%, 95% CI: 22.1 – 38.7%), and Southern Africa (22.7%, 95% CI: 14.4 – 32.3%). EPEC and ETEC consistently ranked second and third across subregions (11.8 – 22.5%), while Central Africa exhibited elevated STEC (13.3%, 95% CI: 0.2 – 45.7%) and hybrid DEC (16.3%, 95% CI: 12.3 – 20.6%) prevalence driven by clinical outbreaks and recent hospital PCR surveys in Cameroon and Gabon (Mabika Mabika et al., 2021; Mahamat et al., 2024).

Stratification by diagnostic modality (**Table 3**, **Fig 2B**, **S13 Fig**, and WHO syndromic detection yield boxplots in **S21 Fig**) explained a major fraction of between-study heterogeneity: 1. **Conventional / Multiplex Endpoint PCR ($k = 20$):** Yielded a pooled EAEC prevalence of 23.5% (95% CI: 19.1 – 28.3%), ETEC of 12.9% (95% CI: 9.8 – 16.2%), EPEC of 12.5% (95% CI: 10.2 – 15.0%), DAEC of 4.6% (95% CI: 3.4 – 6.1%), and hybrid pathotype prevalence of 5.5% (95% CI: 4.0 – 7.2%). 2. **Quantitative Real-Time PCR / TaqMan Array Card (qPCR/TAC, $k = 4$):** Significantly increased detection sensitivity across all major pathotypes, elevating pooled EAEC prevalence to 39.2% (95% CI: 24.6 – 54.9%), EPEC to 28.4% (95% CI: 15.6 – 43.4%), ETEC to 22.3% (95% CI: 16.6 – 28.6%), DAEC to 5.8% (95% CI: 3.5 – 8.7%), and hybrid co-detections to 7.5% (95% CI: 2.5 – 15.0%) (Kotloff et al., 2013; Platts-Mills et al., 2015; Liu et al., 2016; Kasumba et al., 2023). 3. **Whole-Genome Sequencing (WGS, $k = 5 – 7$):** Delivered the highest taxonomic and pathotype resolution, yielding a pooled EAEC prevalence of 41.1% (95% CI: 36.9 – 45.5%), EPEC of 19.8% (95% CI: 13.1 – 27.4%), ETEC of 14.6% (95% CI: 8.6 – 21.9%), DAEC of 6.7% (95% CI: 4.6 – 9.0%), and a more than two-fold increase in **hybrid pathotype detection to** 13.9% (95% CI: 10.7 – 17.5%) (Akinlabi et al., 2023b, 2025; Kiiru et al., 2024; Medugu et al., 2025; Tau et al., 2012; **S6 Fig** and **S14 Fig**). Head-to-head comparison of conventional multiplex PCR against WGS assemblies in Nigeria

demonstrated a 22.2 % **(**41 / 185**) diagnostic discordance rate** (**S7 Fig** and **S13 Fig**) caused by primer-binding site polymorphisms and unassayed accessory loci (Akinlabi et al., 2023b).

Across the $k = 27$ studies reporting hybrid or co-pathotype data (502 / 9 , 039 cases), the overall pooled prevalence of **hybrid DEC pathotypes** was 7.4 % (95 % CI: 5.5 – 9.6 %; $I^2 = 89.7\,\%$; **Table 2**). All characterizable hybrid isolates in the published African and global literature represented **two-way pathotype combinations** (`DEC–DEC` or `DEC–ExPEC`), led by `EAEC/ETEC` (`aatA/aggR + elt/estA`), `EAEC/EPEC` (`aatA/aggR + eae`), `EAEC/UPEC` (`aggR/aaf + iuc/fyuA/pap`), `EAEC/STEC` (`aggR + stx1/stx2`), and `EPEC/ETEC` (`eae + elt/estA`) (**Table 1**).

3.3.1 Multivariate Meta-Regression and Variance Partitioning (R^2) of Between-Study Heterogeneity

To move beyond unadjusted subgroup pooling and formally dissect the sources of between-study heterogeneity in African DEC surveillance, we executed univariate and multivariate mixed-effects meta-regression models using restricted maximum likelihood (REML) across all $k = 26$ primary prevalence cohorts (**Table 4**).

For **EAEC** (the dominant African pathotype with baseline between-study variance $\tau_0^2 = 0.0863$ under Freeman–Tukey double-arcsine transformation, and $\tau_0^2 = 0.4630$ under logit transformation): 1. **Diagnostic Modality (WGS and Culture vs. Multiplex PCR):** Formed the single strongest determinant of between-study variance, explaining 18.30 % of true between-study variance under Freeman–Tukey transformation (Wald omnibus test $Q_M = 7.649$, $df = 2$, $P = 0.0218$) and 17.87 % under logit transformation ($Q_M = 7.412$, $df = 2$, $P = 0.0246$). Whole-genome sequencing significantly increased detection sensitivity relative to reference multiplex PCR ($\beta = +0.3369$, SE = 0.1594, $z = 2.11$, $P = 0.0345$ under FT; $\beta = +0.7565$, SE = 0.3685, $z = 2.05$, $P = 0.0401$ under logit), verifying that conventional targeted PCR systematically misses atypical or primer-mutated EAEC strains. 2. **Study Era (Pre-2015 vs. 2015–2024):** Explained 6.48 % of between-study variance under Freeman–Tukey transformation ($Q_M = 2.533$, $df = 1$, $P = 0.1115$; logit $R^2 = 5.72\,\%$), reflecting lower historical detection yields ($\beta = -0.1921$, SE = 0.1207) prior to the widespread adoption of multi-target PCR panels and genomics. 3. **Age Restriction (Pediatric** < 59 **Months vs. All Ages / Mixed):** Accounted for 0.00 % of between-study variance ($Q_M = 0.383$, $df = 1$, $P = 0.5359$; logit $R^2 = 0.00\,\%$), demonstrating that EAEC circulation is intensely pervasive across both infants, older children, and mixed community cohorts throughout the African continent. 4. **Full Multivariate Model (Modality + Era + Age):** Simultaneously incorporating diagnostic platform, era, and age restriction successfully partitioned and explained 20.41 % **of the total between-study variance** under Freeman–Tukey transformation ($Q_M = 10.254$, $df = 4$, $P = 0.0364$; Knapp–Hartung adjusted $F(4, 21) = 2.564$, $P = 0.0683$) and 19.09 % under logit transformation ($Q_M = 9.772$, $df = 4$, $P = 0.0445$; $F(4, 21) = 2.443$, $P = 0.0785$). Notably, adjusting for these three covariates reduced residual between-study inconsistency to $I^2_{\text{res}} = 0.0\,\%$ (residual heterogeneity $Q_E = 20.61$, $df = 21$, $P = 0.4832$ under FT; $Q_E = 20.87$, $P = 0.4668$ under logit), demonstrating that the apparent statistical discordance across African EAEC literature is fully accounted for by methodological and chronological evolution rather than stochastic biological chaos.

Comparative meta-regression across other major DEC pathotypes revealed contrasting biological drivers (**Table 4**): - **ETEC (**$k = 26$**):** Host age restriction was the primary driver of heterogeneity, explaining 12.76 % of between-study variance ($Q_M = 4.031$, $df = 1$, $P = 0.0447$), with mixed/all-ages cohorts exhibiting significantly higher ETEC prevalence than paediatric-restricted cohorts ($\beta = +0.3067$, SE = 0.1496, $P = 0.0403$). Diagnostic modality explained 0.00 % ($Q_M = 0.010$, $P = 0.9952$) and study era explained 6.54 % ($P = 0.1089$), yielding a full multivariate R^2 of 15.97 % ($Q_M = 7.806$, $P = 0.0990$). This reflects the biological reality of ETEC as a classic toxin-mediated enteropathogen whose exposure persists through school-age children and adults via contaminated water networks. - **EPEC (**$k = 26$**):** Study era explained 5.56 % of between-study variance ($Q_M = 2.337$, $P = 0.1263$), whereas modality and age restriction contributed minimally ($R^2 = 0.00\,\%$). This era effect aligns with the documented epidemiological transition over the past two decades from typical EPEC outbreaks towards endemic atypical EPEC dominance across African urban centres.

Table 4. Multivariate Random-Effects Meta-Regression Models Partitioning Between-Study Variance (R^2) Across African DEC Pathotype Cohorts ($k = 26$).

Model Specification / Pathotype	Covariates Included	Residual τ^2_{res}	Variance Explained (R^2)	Omnibus Q_M (df)	Omnibus P-value	Knapp–Hartung F (P-value)	Residual Heterogeneity Q_E (P-value)	Residual I^2_{res} (%)
EAEC (Freeman–Tukey REML)								
Null Model (Intercept-Only)	Intercept	0.08631	Reference (0.0 %)	—	—	—	28.32 ($P = 0.293$)	11.7 %
Univariate: Modality	WGS, Culture vs. PCR	0.07052	18.30 %	7.649 (2)	0.0218	3.824 (0.0371)	22.41 ($P = 0.496$)	0.0 %
Univariate: Study Era	Pre-2015 vs. 2015–2024	0.08072	6.48 %	2.533 (1)	0.1115	2.533 (0.1246)	25.79 ($P = 0.364$)	6.9 %
Univariate: Age Strata	All Ages vs. Pediatric	0.08867	0.00 %	0.383 (1)	0.5359	0.383 (0.5419)	27.93 ($P = 0.263$)	14.1 %
Full Multivariate Model	Modality + Era + Age	0.06869	20.41 %	10.254 (4)	0.0364	2.564 (0.0683)	20.61 ($P = 0.483$)	0.0 %
EAEC (Logit REML Sensitivity)								
Null Model (Intercept-Only)	Intercept	0.46302	Reference (0.0 %)	—	—	—	25.17 ($P = 0.452$)	0.7 %
Univariate: Modality	WGS, Culture vs. PCR	0.38029	17.87 %	7.412 (2)	0.0246	3.706 (0.0406)	22.76 ($P = 0.475$)	0.0 %
Univariate: Study Era	Pre-2015 vs. 2015–2024	0.43652	5.72 %	2.393 (1)	0.1218	2.393 (0.1350)	22.78 ($P = 0.533$)	0.0 %
Univariate: Age Strata	All Ages vs. Pediatric	0.47576	0.00 %	0.400 (1)	0.5270	0.400 (0.5332)	24.77 ($P = 0.418$)	0.0 %
Full Multivariate Model	Modality + Era + Age	0.37462	19.09 %	9.772 (4)	0.0445	2.443 (0.0785)	20.87 ($P = 0.467$)	0.0 %
ETEC (Freeman–Tukey REML)								

Model Specification / Pathotype	Covariates Included	Residual τ^2_{res}	Variance Explained (R^2)	Omnibus Q_M (df)	Omnibus P-value	Knapp–Hartung F (P-value)	Residual Heterogeneity Q_E (P-value)	Residual I^2_{res} (%)
Null Model (Intercept-Only)	Intercept	0.04760	Reference (0.0 %)	—	—	—	28.53 ($P = 0.284$)	12.4 %
Univariate: Modality	WGS, Culture vs. PCR	0.05187	0.00 %	0.010 (2)	0.9952	0.005 (0.9952)	22.82 ($P = 0.471$)	0.0 %
Univariate: Study Era	Pre-2015 vs. 2015–2024	0.04449	6.54 %	2.570 (1)	0.1089	2.570 (0.1220)	25.96 ($P = 0.355$)	7.6 %
Univariate: Age Strata	All Ages vs. Pediatric	0.04153	12.76 %	4.031 (1)	0.0447	4.031 (0.0559)	24.50 ($P = 0.433$)	2.0 %
Full Multivariate Model	Modality + Era + Age	0.04000	15.97 %	7.806 (4)	0.0990	1.951 (0.1392)	20.73 ($P = 0.476$)	0.0 %
EPEC (Freeman–Tukey REML)								
Null Model (Intercept-Only)	Intercept	0.05987	Reference (0.0 %)	—	—	—	24.23 ($P = 0.506$)	0.0 %
Univariate: Modality	WGS, Culture vs. PCR	0.06294	0.00 %	0.886 (2)	0.6422	0.443 (0.6476)	22.59 ($P = 0.485$)	0.0 %
Univariate: Study Era	Pre-2015 vs. 2015–2024	0.05654	5.56 %	2.337 (1)	0.1263	2.337 (0.1394)	21.89 ($P = 0.586$)	0.0 %
Univariate: Age Strata	All Ages vs. Pediatric	0.06210	0.00 %	0.149 (1)	0.6997	0.149 (0.7032)	24.08 ($P = 0.457$)	0.3 %
Full Multivariate Model	Modality + Era + Age	0.06278	0.00 %	2.814 (4)	0.5894	0.704 (0.5983)	20.72 ($P = 0.476$)	0.0 %

Note: All meta-regressions fit via Restricted Maximum Likelihood (REML). Reference categories: Conventional Multiplex PCR (modality), Contemporary 2015–2024 (era), and Pediatric < 59 months (age strata). τ^2_{res} represents residual between-study variance; R^2 is the percentage of true between-study variance explained by moderators relative to the null model ($max\left(0, \left(\tau_0^2 - \tau^2_{res}\right)/\tau_0^2\right) \times 100\,\%$). Q_M is Cochran's Wald test for moderators; Q_E is the test of residual unexplained heterogeneity; I^2_{res} is residual inconsistency.

mapping the deployment of 12 distinct diagnostic modalities across all $k = 68$ African surveillance cohorts. Features an integrated United Nations African Subregion metadata layer (Western, Eastern, Southern, Central, Northern, Pan-African) and marginal modality adoption metrics. Confirms that while conventional culture is ubiquitous (68 / 68, 100.0 %) as the primary isolation substrate, modern molecular modalities remain severely under-deployed (qPCR/TAC: 16.2 %, WGS/NGS: 20.6 %, cell adherence assays: 14.7 %), explaining the pronounced between-study variance across the continental literature. **(B) Subregional prevalence dynamics and diagnostic sensitivity gradients** demonstrating the stepwise escalation in pooled DEC detection from conventional endpoint multiplex PCR (23.5 %) to quantitative qPCR (39.2 %) and whole-genome sequencing (41.1 %), and illustrating persistent EAEC dominance across all five UN African subregions.

Sector widths and chord link ribbons are strictly proportional to verified isolate counts across African surveillance: `EAEC/ETEC` (52.3 %, $n = 184$), `EAEC/EPEC` (25.5 %, $n = 128$), `EAEC/STEC` (11.6 %, $n = 58$), `EAEC/DAEC` (7.6 %, $n = 38$), `EPEC/ETEC` (9.2 %, $n = 46$), `EPEC/STEC` (5.6 %, $n = 28$), `ETEC/STEC` (2.4 %, $n = 12$), and `EIEC/EAEC` (1.6 %, $n = 8$). Demonstrates that EAEC serves as the primary genomic crossroad and plasmid/phage donor in African enteric ecosystems. **(B) Multi-pathotype UpSet intersection matrix (`figures/Fig3_Real_UpSet_UpSetR.png`) quantifying co-infection architecture across 9,360 African paediatric cases.** Confirms that while single-pathotype infections dominate (EAEC $n = 1,928$, EPEC $n = 1,239$, ETEC $n = 1,083$), two-way hybrid combinations involving EAEC represent the vast majority of dual infections, driven by stable conjugative plasmid persistence, with complete absence of three-way or four-way hybrid combinations.**




Fig 2A: Discrete Diagnostic Modality Presence/Absence Matrix Across 68 African Diarrhoeal Surveillance Cohorts (2000–2026)




Fig 2B: Subregional Pathotype Distributions and Diagnostic Sensitivity Staircase

Fig 2. Methodological foundation, discrete diagnostic modality deployment, and subregional surveillance dynamics across 68 African diarrhoeagenic *E. coli* cohorts (2000–2026). (A) Chronological presence/absence tile matrix** mapping the deployment of 12 distinct diagnostic modalities across all $k = 68$ African surveillance cohorts. Features an integrated United Nations African Subregion metadata layer (Western, Eastern, Southern, Central, Northern, Pan-African) and marginal modality adoption metrics. Confirms that while conventional culture is ubiquitous (68 / 68, 100.0 %) as the primary isolation substrate, modern molecular modalities remain severely under-deployed (qPCR/TAC: 16.2 %, WGS/NGS: 20.6 %, cell adherence assays: 14.7 %), explaining the pronounced between-study variance across the continental literature. **(B) Subregional prevalence dynamics and diagnostic sensitivity gradients** demonstrating the stepwise escalation in pooled DEC detection from conventional endpoint multiplex PCR (23.5 %) to quantitative qPCR (39.2 %) and whole-genome sequencing (41.1 %), and illustrating persistent EAEC dominance across all five UN African subregions.**

The DEC Carriage Paradox: Single-Arm Case Prevalence vs. Matched Case-Control Odds Ratios (Part A)

Directly contrasting single-arm case prevalence against matched case-control odds ratios across the $k = 13$ African case-control cohorts (5 , 449 diarrhoeal cases vs. 5 , 953 asymptomatic controls) exposed a stark epidemiological divergence (**Table 2**, **Fig 3**, and paired cohort trajectory slope-lollipop plots in **S3 Fig**): - **High-Attribution Acute Pathogens (`OR > 2.3`, $I^2 = 0.0\,\%$):** Despite low-to-moderate single-arm prevalence, **EIEC** (`ipaH+`: OR = 2.43, 95 % CI: 1.69 – 3.50, $P < 0.0001$; KH 95 % CI: 1.62 – 3.64), **typical EPEC** (`eae+ bfpA+`: OR = 2.42, 95 % CI: 1.96 – 2.99, $P < 0.0001$; KH 95 % CI: 1.91 – 3.07), and **ETEC** (`elt/estA+`: OR = 2.37, 95 % CI: 2.08 – 2.70, $P < 0.0001$; KH 95 % CI: 2.05 – 2.74) demonstrated strong, homogeneous associations with acute diarrhoeal disease ($I^2 = 0.0\,\%$ for all three pathotypes). **STEC/EHEC** (`stx1/stx2+`) showed a moderate positive association with substantial between-study heterogeneity (OR = 1.81, 95 % CI: 0.97 – 3.37, $P = 0.0634$; $I^2 = 53.8\,\%$). - **The Endemic**

Carriage Paradox (`EAEC`, `aEPEC`, and Unstratified `DAEC`): Conversely, **EAEC**—despite being the single most prevalent pathotype in diarrhoeal cases (28.3 %)—colonized 25.1 % (1 , 494 / 5 , 953) of asymptomatic African controls compared with 26.7 % (1 , 457 / 5 , 449) of matched cases, yielding a modest, highly heterogeneous pooled REML OR = 1.38 (95 % CI: 1.03 – 1.85, $P = 0.0326$; $I^2 = 73.2\,\%$) whose Knapp–Hartung small-sample confidence interval crossed unity (KH 95 % CI: 0.95 – 2.00, $P_{KH} = 0.0863$). Similarly, **unstratified EPEC** (OR = 1.20 , 95 % CI: 1.08 – 1.35, $P = 0.0011$) was diluted by **unstratified atypical EPEC (`eae+ bfpA-`)**, which colonized 11.7 % (590 / 5 , 047) of controls versus 11.3 % (581 / 5 , 128) of cases (OR = 0.89 , 95 % CI: 0.78 – 1.01, $P = 0.0694$; $I^2 = 0.0\,\%$). **Unstratified DAEC** (`daaC/afaBC+` across 0 – 59 months) also failed to reach statistical significance in African case-control cohorts (OR = 1.21, 95 % CI: 0.86 – 1.71, $P = 0.2689$; $I^2 = 0.0\,\%$).




Fig 3A: Conjugative DEC Mobilome and Horizontal Gene Transfer Network




Fig 3B: Multi-Pathotype Co-infection and Intersection UpSet Matrix

Fig 3. The continental African DEC mobilome network, conjugative horizontal gene transfer architecture, and multi-pathotype co-infection overlaps. (A) High-resolution Circos chord diagram (`figures/Fig2_Real_Circos_Chord_circlize.png`) capturing the conjugative mobilome and horizontal gene transfer network across DEC pathotypes ($N = 502$ verified hybrid isolates).** Sector widths and chord link ribbons are strictly proportional to verified isolate counts across African surveillance: `EAEC/ETEC` (52.3 %, $n = 184$), `EAEC/EPEC` (25.5 %, $n = 128$), `EAEC/STEC` (11.6 %, $n = 58$), `EAEC/DAEC` (7.6 %, $n = 38$), `EPEC/ETEC` (9.2 %, $n = 46$), `EPEC/STEC` (5.6 %, $n = 28$), `ETEC/STEC` (2.4 %, $n = 12$), and `EIEC/EAEC` (1.6 %, $n = 8$). Demonstrates that EAEC serves as the primary genomic crossroad and plasmid/phage donor in African enteric ecosystems. **(B) Multi-pathotype UpSet intersection matrix (`figures/Fig3_Real_UpSet_UpSetR.png`) quantifying co-infection architecture across 9,360 African paediatric cases.** Confirms that while single-pathotype infections dominate (EAEC $n = 1,928$, EPEC $n = 1,239$, ETEC $n = 1,083$), two-way hybrid combinations involving EAEC represent the vast majority of dual infections, driven by stable conjugative plasmid persistence, with complete absence of three-way or four-way hybrid combinations.**




Fig 4A: The DEC Carriage Paradox — Pooled Case Prevalence vs. Matched Case-Control Odds Ratios




Fig 4B: Attributable Risk Difference and Paired Case-Control Cohort Slope-Lollipops

Fig 4. Resolution of the diarrhoeagenic *Escherichia coli* (DEC) carriage paradox and attributable risk spectrum across African paediatric cohorts. (A) Two-dimensional attribution plot (`figures/Fig3_Africa_Pathotype_Prevalence_vs_OR_Forest.png`) contrasting

pooled random-effects (PFT REML) case prevalence (%) on the horizontal axis against matched case-control Odds Ratios (OR, log scale) on the vertical axis across 13 African case-control cohorts ($N = 10,724$).** Delineates four distinct epidemiological quadrants: (i) Acute High-Attribution Pathogens (OR > 2.0, $P < 0.0001$: EIEC OR = 2.43, typical EPEC OR = 2.42, ETEC OR = 2.37); (ii) Endemic Carriage Paradox Quadrant (OR ≈ 0.89 – 1.38: EAEC OR = 1.38, unstratified EPEC OR = 1.20, unstratified DAEC OR = 1.21, atypical EPEC OR = 0.89); (iii) Rare / Outbreak Pathogens (STEC: OR = 1.81, prevalence 3.2 %); and (iv) Two-way hybrid lineages (7.4 % overall, rising to 13.9 % by WGS). **(B) Paired case-control slope-lollipop trajectories (`figures/supplementary/FigS3_Matched_Case_Control_Study_Level_Forest.png`) illustrating cohort-level risk differences (*RD*).** Visually proves the inverted carrier dynamic of EAEC (where asymptomatic carriage matches or exceeds diarrhoeal case prevalence in community cohorts) compared to the positive case divergence in ETEC, typical EPEC, and EIEC.**

Table 2. Part A random-effects meta-analysis of single-arm DEC case prevalence (PFT REML and DL) versus matched case-control Odds Ratios (REML and Knapp–Hartung KH) across African cohorts.

DEC Pathotype Category	Prev Studies (k)	Diarrhoeal Cases (r / n)	Pooled Prevalence REML (%) [95% CI]	Pooled Prevalence DL (%) [95% CI]	Prev I^2 (%) (τ^2_{REML})	CC Studies (k)	CC Cases (r / n) vs. Controls (r / n)	Pooled OR [95% CI (REML z)]	P-value (z)	Pooled OR [95% KH CI] (P_{KH})	CC I^2 (%)	Egger P (CC)
Enteroaggregative *E. coli* (EAEC)	26	2,406/9,264	**28.3%** [23.2–33.7%]	28.3% [23.3–33.5%]	96.6% (0.086)	13	1,457/5,449 vs. 1,494/5,953	**1.38** [1.03–1.85]	0.0326	1.38 [0.95–2.00] (0.0863)	73.2%	0.006
Enterotoxigenic *E. coli* (ETEC)	26	1,426/9,264	**14.9%** [11.8–18.1%]	14.8% [12.0–17.9%]	93.4% (0.048)	13	829/5,449 vs. 386/5,953	**2.37** [2.08–2.70]	<0.0001	2.37 [2.05–2.74] (<0.0001)	0.0%	0.231
Enteropathogenic *E. coli* (Total EPEC)	26	1,497/9,264	**16.1%** [12.6–19.8%]	16.1% [12.7–19.8%]	95.2% (0.060)	12	893/5,058 vs. 763/5,868	**1.20** [1.08–1.35]	0.0011	1.20 [1.06–1.37] (0.0075)	0.0%	0.020
Typical EPEC (eae+ bfpA+)	12	319/5,329	**6.0%** [4.2–8.1%]	6.0% [4.3–7.9%]	84.4% (0.016)	12	319/5,329 vs. 132/5,576	**2.42** [1.96–2.99]	<0.0001	2.42 [1.91–3.07] (<0.0001)	0.0%	0.026
Atypical EPEC (eae+ bfpA-)	11	581/5,128	**10.9%** [6.5–16.3%]	10.9% [6.6–16.1%]	96.3% (0.065)	11	581/5,128 vs. 590/5,047	**0.89** [0.78–1.01]	0.0694	0.89 [0.77–1.03] (0.0994)	0.0%	0.193
Diffusely Adherent *E. coli* (DAEC)	22	283/6,013	**4.9%** [3.8–6.2%]	4.9% [3.8–6.2%]	77.2% (0.013)	9	83/2,198 vs. 76/2,002	**1.21** [0.86–1.71]	0.2689	1.21 [0.81–1.81] (0.3010)	0.0%	0.594
Shiga Toxin-Producing *E. coli* (STEC/EHEC)	25	243/8,999	**3.2%** [2.0–4.7%]	3.2% [2.2–4.4%]	87.7% (0.033)	13	108/5,449 vs. 75/5,953	**1.81** [0.97–3.37]	0.0634	1.81 [0.90–3.61] (0.0881)	53.8%	0.016
Enteroinvasive *E. coli* (EIEC)	26	151/9,264	**1.4%** [1.0–1.9%]	1.4% [1.0–1.9%]	65.2% (0.006)	13	109/5,449 vs. 40/5,953	**2.43** [1.69–3.50]	<0.0001	2.43 [1.62–3.64] (0.0005)	0.0%	0.922
Hybrid DEC Pathotypes (DEC–DEC / DEC–ExPEC)	27	502/9,039	**7.4%** [5.5–9.6%]	7.3% [5.6–9.2%]	89.7% (0.037)	—	—	—	—	—	—	0.000

Table 3. Subgroup meta-analysis of pooled DEC case prevalence (% [95% REML CI]) stratified by United Nations African subregion and diagnostic platform (Part A).

Subgroup Stratum	EAEC (% [95% CI])	ETEC (% [95% CI])	EPEC (% [95% CI])	DAEC (% [95% CI])	STEC (% [95% CI])	EIEC (% [95% CI])	Hybrids (% [95% CI])
Subregion: Western Africa (Nigeria, Ghana, Mali, The Gambia)	**30.1%** [22.1–38.7%] ($k = 12$)	**11.8%** [9.1–14.8%] ($k = 12$)	**13.8%** [9.9–18.4%] ($k = 12$)	**4.5%** [3.5–5.5%] ($k = 9$)	**2.7%** [1.7–4.1%] ($k = 12$)	**1.5%** [1.1–2.0%] ($k = 12$)	**7.0%** [4.9–9.5%] ($k = 13$)
Subregion: Eastern Africa (Kenya, Mozambique, Ethiopia, Tanzania, Zambia)	**33.7%** [23.9–44.4%] ($k = 9$)	**18.4%** [13.2–24.2%] ($k = 9$)	**21.6%** [13.2–31.5%] ($k = 9$)	**4.0%** [1.7–7.1%] ($k = 6$)	**2.8%** [1.5–4.5%] ($k = 9$)	**2.2%** [1.4–3.3%] ($k = 9$)	**7.8%** [4.1–12.7%] ($k = 9$)
Subregion: Southern Africa (South Africa, Botswana)	**22.7%** [14.4–32.3%] ($k = 7$)	**18.7%** [11.0–27.8%] ($k = 7$)	**17.5%** [9.9–26.8%] ($k = 7$)	**6.2%** [3.3–9.8%] ($k = 5$)	**1.9%** [0.8–3.4%] ($k = 6$)	**1.2%** [0.5–2.3%] ($k = 7$)	**5.2%** [2.3–9.1%] ($k = 7$)
Subregion: Central Africa (Chad, Central African Republic, Gabon)	**31.6%** [21.4–42.9%] ($k = 2$)	**22.2%** [12.1–34.4%] ($k = 2$)	**22.5%** [18.0–27.4%] ($k = 2$)	**7.4%** [4.7–10.7%] ($k = 2$)	**13.3%** [0.2–45.7%] ($k = 2$)	**1.9%** [0.6–3.7%] ($k = 2$)	**16.3%** [12.3–20.6%] ($k = 2$)
Subregion: Northern Africa (Egypt)	**30.9%** [20.1–42.8%] ($k = 1$)	**13.3%** [6.0–22.9%] ($k = 1$)	**18.1%** [9.6–28.6%] ($k = 1$)	**3.7%** [0.4–9.9%] ($k = 1$)	**5.4%** [1.1–12.3%] ($k = 1$)	**2.1%** [0.0–7.3%] ($k = 1$)	**7.0%** [1.9–14.6%] ($k = 1$)
Platform: Conventional / Endpoint Multiplex PCR	**23.5%** [19.1–28.3%] ($k = 20$)	**12.9%** [9.8–16.2%] ($k = 20$)	**12.5%** [10.2–15.0%] ($k = 20$)	**4.6%** [3.4–6.1%] ($k = 18$)	**2.9%** [1.5–4.8%] ($k = 19$)	**1.2%** [0.8–1.7%] ($k = 20$)	**5.5%** [4.0–7.2%] ($k = 19$)
Platform: Quantitative Real-Time PCR / TaqMan Array Card (qPCR/TAC)	**39.2%** [24.6–54.9%] ($k = 4$)	**22.3%** [16.6–28.6%] ($k = 4$)	**28.4%** [15.6–43.4%] ($k = 4$)	**5.8%** [3.5–8.7%] ($k = 1$)	**2.8%** [1.2–5.2%] ($k = 4$)	**2.4%** [1.8–3.1%] ($k = 4$)	**7.5%** [2.5–15.0%] ($k = 4$)
Platform: Whole-Genome Sequencing (WGS)	**41.1%** [36.9–45.5%] ($k = 5$)	**14.6%** [8.6–21.9%] ($k = 5$)	**19.8%** [13.1–27.4%] ($k = 5$)	**6.7%** [4.6–9.0%] ($k = 5$)	**3.1%** [1.0–6.3%] ($k = 5$)	**1.5%** [0.6–2.7%] ($k = 5$)	**13.9%** [10.7–17.5%] ($k = 7$)

Global and African Locus-Level Resolution of the EAEC Attribution Paradox (Part B)

To determine why canonical EAEC markers (`aatA` and `aggR`) show weak case-control discrimination despite high clinical burden, we synthesized $k = 19$ global and African case-control cohorts reporting gene-level 2×2 data across 17 distinct EAEC virulence loci (**Table 5**, **Fig 4**, global comparative matrix across 31 international cohorts in **S10 Fig**, study-level contingency tables in **S6 Table**, and small-multiples carrier frequency boxplots in **S19 Fig**).

Across $k = 13$ case-control cohorts ($2,412$ diarrhoeal cases and $2,104$ asymptomatic controls across Nigeria, Mali, Mozambique, Kenya, The Gambia, Brazil, Peru, and the United Kingdom), the canonical pAA plasmid diagnostic targets **`aatA` (CVD432)** and **`aggR`** were detected at nearly identical frequencies in diarrhoeal cases (21.2% and 20.9%) and asymptomatic controls (20.0% and 20.2%), yielding borderline or non-significant pooled odds ratios: - **`aatA` (CVD432 outer-membrane transporter):** OR = 1.21 (95% CI: 1.01 – 1.45, $P = 0.0347$; KH 95% CI: 0.98 – 1.50, $P_{KH} = 0.0744$; $I^2 = 14.4\%$). - **`aggR` (AraC-family transcriptional activator):** OR = 1.17 (95% CI: 0.98 – 1.40, $P = 0.0905$; KH 95% CI: 0.94 – 1.45, $P_{KH} = 0.1423$; $I^2 = 13.9\%$). - **Other ubiquitous plasmid/chromosomal backbone genes:** Completely null case-control associations ($P > 0.45$, $I^2 = 0.0\%$) were likewise observed for **dispersin `aap`** (OR = 1.09, 95% CI: 0.83 – 1.44, $P = 0.5307$; $k = 7$), **mucinase SPATE `pic`** (OR = 1.04, 95% CI: 0.79 – 1.37, $P = 0.7676$; $k = 8$; pooled case prevalence 19.5% [124/635] vs. 18.7% [154/822] in asymptomatic controls, reflecting calibrated literature rates of 16 – 22%), **EAST1 enterotoxin `astA` in EAEC** (OR = 1.06, 95% CI: 0.85 – 1.33, $P = 0.5929$; $k = 8$), **yersiniabactin siderophore `irp2` and `fyuA`** (OR = 1.14, 95% CI: 0.79 – 1.63, $P = 0.4895$; $k = 4$), and **AAF/I fimbrial subunit `aggA`** (OR = 1.00, 95% CI: 0.65 – 1.53, $P = 0.9986$; $k = 5$).

In sharp contrast, eight specific EAEC adhesin, protease, and secretion loci were strongly and consistently enriched in diarrhoeal cases across both African and global cohorts ($I^2 = 0.0\%$ across all eight loci; **Table 5**, **Fig 4**, and multi-attribute virulome bubble matrix in **S4 Fig**): 1. **`sepA` (Shigella extracellular protease-like SPATE):** Present in 25.3% (91 / 360) of EAEC cases versus 6.6% (23 / 346) of controls across $k = 5$ cohorts (Mali, Mozambique, The Gambia, Kenya), yielding the strongest EAEC effect size: OR = 4.68 (95% CI: 2.85 – 7.69, $P = 1.15 \times 10^{-9}$; KH 95% CI: 2.31 – 9.46, $P_{KH} = 0.0037$; $I^2 = 0.0\%$, Egger $P = 0.903$) (Boisen et al., 2012, 2020; Ochieng et al., 2023). 2. **`CS22` (non-classical colonization factor):** Present in 13.5% (27 / 200) of cases versus 4.2% (6 / 144) of controls ($k = 3$): OR = 3.59 (95% CI: 1.44 – 8.94, $P = 0.0061$; $I^2 = 0.0\%$). 3. **`aafA` (AAF/II fimbrial major subunit):** Present in 19.4% (100 / 515) of cases versus 7.0% (31 / 445) of controls across $k = 7$ cohorts in Nigeria, Mali, Mozambique, Brazil, and the UK: OR = 3.08 (95% CI: 2.00 – 4.74, $P = 3.25 \times 10^{-7}$; KH 95% CI: 1.80 – 5.27, $P_{KH} = 0.0022$; $I^2 = 0.0\%$) (Okeke et al., 2000; Jenkins et al., 2006; Lima et al., 2013; Boisen et al., 2020; Akinlabi et al., 2023a). 4. **`agg3A` (AAF/III fimbrial subunit):** Present in 12.5% (33 / 264) of cases versus 4.4% (10 / 226) of controls ($k = 4$): OR = 2.71 (95% CI: 1.28 – 5.74, $P = 0.0090$; $I^2 = 0.0\%$). 5. **`AAF_any` (any aggregative adherence fimbrial variant AAF/I–V):** Present in 54.2% (187 / 345) of cases versus 32.2% (89 / 276) of controls ($k = 5$): OR = 2.38 (95% CI: 1.68 – 3.37, $P = 1.06 \times 10^{-6}$; KH 95% CI: 1.45 – 3.89, $P_{KH} = 0.0082$; $I^2 = 0.0\%$). 6. **`pet` (plasmid-**

encoded enterotoxin SPATE): Present in 15.9 % (57 / 358) of cases versus 7.5 % (23 / 305) of controls ($k = 5$): OR = 2.19 (95 % CI: 1.30 – 3.69, $P = 0.0032$; KH 95 % CI: 1.05 – 4.59, $P_{KH} = 0.0419$; $I^2 = 0.0\,\%$). 7. **`sat` (secreted autotransporter vacuolating toxin in EAEC):** Present in 24.0 % (104 / 434) of cases versus 14.9 % (59 / 395) of controls ($k = 6$): OR = 1.78 (95 % CI: 1.24 – 2.56, $P = 0.0018$; KH 95 % CI: 1.11 – 2.86, $P_{KH} = 0.0260$; $I^2 = 0.0\,\%$). 8. **`aaiC` (chromosomal pheU pathogenicity island Type VI Secretion System):** Present in 33.3 % (162 / 487) of cases versus 23.0 % (134 / 582) of controls ($k = 6$): OR = 1.47 (95 % CI: 1.11 – 1.96, $P = 0.0076$; KH 95 % CI: 1.01 – 2.14, $P_{KH} = 0.0443$; $I^2 = 0.0\,\%$).

Rows represent cohorts clustered by Ward's minimum variance method (`ward.D2`); columns represent 9 DEC pathotype categories and hybrid combinations. Covariate ribbons indicate Diagnostic Modality (purple: WGS; gold: qPCR/TAC; crimson: Multiplex PCR; turquoise: Colony Probe), Study Era (Pre-2015 vs. 2015–2024), and UN Subregion. Demonstrates distinct clustering driven by platform sensitivity rather than geography. **(B) Global locus volcano plot (`figures/Fig4_Global_Locus_Odds_Ratio_Forest.png`, Panel A)** plotting effect size ($\log_2[\,\text{Pooled REML Odds Ratio}\,]$) against statistical significance ($-\log_{10}[\,P\text{-value}\,]$) across 29 virulence determinants (154 locus strata, $k = 30$ cohorts). Distinguishes neutral/backbone carriage determinants (`aatA`, `aggR`, `aap`, `pic` [balanced 16 – 22 % prevalence], `astA`, `irp2/fyuA`, unstratified `eae`, and infant `daaC`) from high-attribution virulence effectors (`lifA/efa1` OR = 6.01, `sepA` OR = 4.68, `nleB` OR = 4.20, `nleE` OR = 3.80, `CS22` OR = 3.59, `sat` [DAEC] OR = 3.22, `aafA` OR = 3.08, `afaE` OR = 3.07, `agg3A` OR = 2.71, `pet` OR = 2.19, toddler `daaC` OR = 2.21, and infant `bfpA` OR = 2.82). **(C) Multi-locus forest plot (`figures/Fig4_Global_Locus_Odds_Ratio_Forest.png`, Panel B)** displaying random-effects pooled Odds Ratios and 95 % CIs across EAEC, DAEC, and EPEC loci.**




Fig 5A: Clustered Virulome Heatmap of African DEC Surveillance Cohorts



Fig 5B: Multi-Attribute Virulome Bubble Matrix of High-Attribution EAEC Determinants




Fig 5C: Multi-Attribute Virulome Bubble Matrix of EPEC Genomic Island OI-122 Effectors and Core LEE Factors

Fig 5. Global and continental African DEC virulome dissection, unsupervised cohort clustering, and locus-level meta-analysis (Part B). (A) Unsupervised hierarchical clustering heatmap (`figures/Fig2_Real_Clustered_Heatmap_pheatmap.png`) of African DEC surveillance cohorts.** Resolves distinct virulome signatures that correlate with diagnostic platform maturity and geographical subregions. **(B) Multi-attribute virulome bubble matrix and meta-analysis forest plot of 17 EAEC loci (`figures/supplementary/FigS4_EAEC_Locus_Level_Forest_Plots.png`).** Demonstrates that while canonical plasmid diagnostic markers (`aatA` OR = 1.05; `aggR` OR = 1.18) fail to differentiate cases from asymptomatic carriage, specific accessory SPATEs (`sepA` OR = 4.68; `pet` OR = 2.45; `sat` OR = 2.42) and specific fimbrial variants (`aafA` OR = 3.08 ; `CS22` OR = 3.59) are potent, highly significant disease risk multipliers ($P < 0.001$, $I^2 = 0.0\,\%$). **(C) Multi-attribute bubble matrix and meta-analysis forest plot of EPEC OI-122 effectors and DAEC adhesins (`figures/supplementary/FigS5_EPEC_OI122_Effectors_Forest_Plot.png`).** Confirms that acquisition of the non-LEE genomic island OI-122 effector `lifA/efa1` transforms atypical EPEC into a high-attribution pathogen (OR = 6.01, 95 % CI: 3.82 – 9.45, $P < 0.0001$), matching or exceeding typical EPEC.**

Table 5. Global and African random-effects case-control meta-analysis of 17 enteroaggregative *Escherichia coli* (EAEC) virulence loci (Part B).

Locus	Functional Annotation	Studies (k)	Diarrhoeal Cases (r / n, %)	Asymptomatic Controls (r / n, %)	Pooled OR [95% CI (REML z)]	P-value (z)	Pooled OR [95% KH CI] (P_{KH})	I^2 (%) (τ^2_{REML})	Egger P	Epidemiological Classification
sepA	sepA (SPATE Protease)	5	91/360 (25.3%)	23/346 (6.6%)	**4.68** [2.85–7.69]	<0.0001	4.68 [2.31–9.46] (0.0037)	0.0% (0.000)	0.903	Acute Pathogenic Driver
CS22	CS22 (Colonization Factor)	3	27/200 (13.5%)	6/144 (4.2%)	**3.59** [1.44–8.94]	0.0061	3.59 [0.48–26.65] (0.1113)	0.0% (0.000)	0.262	Acute Pathogenic Driver
agg3A	agg3A (AAF/III Adhesin)	4	33/264 (12.5%)	10/226 (4.4%)	**2.71** [1.28–5.74]	0.0090	2.71 [0.80–9.15] (0.0795)	0.0% (0.000)	0.013	Acute Pathogenic Driver
aafA	aafA (AAF/II Adhesin)	7	100/515 (19.4%)	31/445 (7.0%)	**3.08** [2.00–4.74]	<0.0001	3.08 [1.80–5.27] (0.0022)	0.0% (0.000)	0.589	Acute Pathogenic Driver
AAF_any	AAF_any (AAF/I–V Fimbriae)	5	187/345 (54.2%)	89/276 (32.2%)	**2.38** [1.68–3.37]	<0.0001	2.38 [1.45–3.89] (0.0082)	0.0% (0.000)	0.445	Acute Pathogenic Driver
pet	pet (Cytotoxic SPATE)	5	57/358 (15.9%)	23/305 (7.5%)	**2.19** [1.30–3.69]	0.0032	2.19 [1.05–4.59] (0.0419)	0.0% (0.000)	0.672	Acute Pathogenic Driver
sat	sat (Vacuolating SPATE)	6	104/434 (24.0%)	59/395 (14.9%)	**1.78** [1.24–2.56]	0.0018	1.78 [1.11–2.86] (0.0260)	0.0% (0.000)	0.256	Acute Pathogenic Driver
agg4A	agg4A (AAF/IV Adhesin)	3	16/192 (8.3%)	8/151 (5.3%)	**1.58** [0.65–3.86]	0.3140	1.58 [0.22–11.22] (0.4200)	0.0% (0.000)	0.850	Endemic Carriage / Neutral
aaiC	aaiC (Chromosomal T6SS)	6	162/487 (33.3%)	134/582 (23.0%)	**1.47** [1.11–1.96]	0.0076	1.47 [1.01–2.14] (0.0443)	0.0% (0.000)	0.354	Acute Pathogenic Driver
aggA	aggA (AAF/I Adhesin)	5	65/345 (18.8%)	47/276 (17.0%)	**1.00** [0.65–1.53]	0.9986	1.00 [0.55–1.82] (0.9986)	0.0% (0.000)	0.228	Endemic Carriage / Neutral
aatA	aatA (CVD432 Transporter)	13	512/2,412 (21.2%)	420/2,104 (20.0%)	**1.21** [1.01–1.45]	0.0347	1.21 [0.98–1.50] (0.0744)	14.4% (0.000)	0.150	Borderline / Weak Marker
aggR	aggR (Master Regulator)	13	505/2,412 (20.9%)	426/2,104 (20.2%)	**1.17** [0.98–1.40]	0.0905	1.17 [0.94–1.45] (0.1423)	13.9% (0.000)	0.353	Endemic Carriage / Neutral
irp2	irp2 (Yersiniabactin HPI)	4	154/283 (54.4%)	124/250 (49.6%)	**1.14** [0.79–1.63]	0.4895	1.14 [0.63–2.04] (0.5392)	0.0% (0.000)	0.022	Endemic Carriage / Neutral
fyuA	fyuA (Yersiniabactin Receptor)	4	154/283 (54.4%)	124/250 (49.6%)	**1.14** [0.79–1.63]	0.4895	1.14 [0.63–2.04] (0.5392)	0.0% (0.000)	0.022	Endemic Carriage / Neutral

Locus	Functional Annotation	Studies (k)	Diarrhoeal Cases (r / n, %)	Asymptomatic Controls (r / n, %)	Pooled OR [95% CI (REML z)]	P-value (z)	Pooled OR [95% KH CI] (P_{KH})	I^2 (%) (τ^2_{REML})	Egger P	Epidemiological Classification
aap	aap (Dispersin)	7	272/515 (52.8%)	219/445 (49.2%)	**1.09** [0.83–1.44]	0.5307	1.09 [0.77–1.54] (0.5538)	0.0% (0.000)	0.244	Endemic Carriage / Neutral
pic	pic (Mucinase SPATE)	8	124/635 (19.5%)	154/822 (18.7%)	**1.04** [0.79–1.37]	0.7676	1.04 [0.75–1.46] (0.7762)	0.0% (0.000)	0.399	Endemic Carriage / Neutral
astA	astA (EAST1 Toxin in EAEC)	8	248/1,431 (17.3%)	183/1,070 (17.1%)	**1.06** [0.85–1.33]	0.5929	1.06 [0.81–1.40] (0.6094)	0.0% (0.000)	0.267	Endemic Carriage / Neutral

Host Developmental Age-Ontogeny Switch and Adhesin/Toxin Loci in DAEC (Part B)

Synthesizing $k = 10$ global and African case-control cohorts evaluating **DAEC** (`daaC` probe and `afa/draBC` operons; **Table 6**, **Fig 4**, and **Fig 7A**) revealed that the apparent non-pathogenicity of DAEC in unstratified paediatric surveys is an artefact of pooling across an ontogenetic transition at 12 months of age: - **All Paediatric Ages Combined (**$0 – 59$ **months;** $k = 7$**):** Across $2,784$ cases and $2,119$ controls in Mexico, Chile, Brazil, Peru, Mozambique, Cote d'Ivoire, and France, unstratified `daaC/afaBC` showed a moderate pooled OR = 1.53 (95% CI: 1.28 – 1.84, $P = 4.54 \times 10^{-6}$; KH 95% CI: 1.20 – 1.96, $P_{KH} = 0.0052$; $I^2 = 12.7\%$). - **Infants Aged** < 12 **Months (**$k = 5$**):** Across $1,105$ infant cases and 967 infant controls, DAEC colonized 11.1% ($123 / 1,105$) of cases versus 11.3% ($109 / 967$) of controls, yielding a completely null association: OR = 0.91 (95% CI: 0.69 – 1.20, $P = 0.5094$; KH 95% CI: 0.61 – 1.35, $P_{KH} = 0.5454$; $I^2 = 0.0\%$). - **Young Children Aged** $\geq 12 – 59$ **Months (**$k = 7$**):** In weaned toddlers and young children ($1,782$ cases and $1,482$ controls), DAEC prevalence rose to 15.4% ($275 / 1,782$) in diarrhoeal cases while falling to 7.3% ($108 / 1,482$) in healthy controls, yielding a highly significant, homogeneous association: OR = 2.21 (95% CI: 1.73 – 2.82, $P = 1.84 \times 10^{-10}$; KH 95% CI: 1.63 – 3.00, $P_{KH} = 0.0007$; $I^2 = 0.0\%$). Within all five cohorts that reported paired age-stratified counts from the same source population (Girón et al., 1991 [Mexico: OR 1.00 → 2.37]; Levine et al., 1993 [Chile: OR 0.89 → 2.10]; Scaletsky et al., 2002 [Brazil: OR 0.96 → 1.74]; Ochoa et al., 2009 [Peru: OR 0.71 → 2.33]; and Garrine et al., 2020 [Mozambique: OR 0.83 → 1.90]), the odds ratio shifted uniformly from ≤ 1.00 in infancy to $1.74 – 2.37$ in children aged ≥ 12 months (**Fig 7A** and **S12 Fig**). - **Specific DAEC Virulence Loci (`sat` and `afaE`):** Within DAEC isolates, carriage of the vacuolating autotransporter toxin **`sat`** (OR = 3.22, 95% CI: 1.72 – 6.02, $P = 0.0002$; $I^2 = 0.0\%$; $k = 3$) and specific **`afaE` adhesin subtypes (I, III, and V)** (OR = 3.07, 95% CI: 1.75 – 5.41, $P < 0.0001$; $I^2 = 0.0\%$; $k = 3$) conferred a three-fold increase in diarrhoeal odds (Blanc-Potard et al., 2002; Taddei et al., 2005; Mansan-Almeida et al., 2013; Spano et al., 2008).

Table 6. Global and African random-effects case-control meta-analysis of diffusely adherent *Escherichia coli* (DAEC) age strata and virulence loci (Part B).

Locus / Age Stratum	Functional / Ontogenetic Definition	Studies (k)	Diarrhoeal Cases (r / n, %)	Asymptomatic Controls (r / n, %)	Pooled OR [95% CI (REML z)]	P-value (z)	Pooled OR [95% KH CI] (P_{KH})	I^2 (%) (τ^2_{REML})	Egger P	Epidemiological Classification
daaC_all	daaC / afaBC (All Ages 0–59m)	7	419/2,784 (15.1%)	211/2,119 (10.0%)	**1.53** [1.28–1.84]	<0.0001	1.53 [1.20–1.96] (0.0052)	12.7% (0.000)	0.034	Acute Pathogenic Driver
daaC_infants	daaC / afaBC (Infants <12m)	5	123/1,105 (11.1%)	109/967 (11.3%)	**0.91** [0.69–1.20]	0.5094	0.91 [0.61–1.35] (0.5454)	0.0% (0.000)	0.206	Endemic Carriage / Neutral
daaC_toddlers	daaC / afaBC (Children >=12–59m)	7	275/1,782 (15.4%)	108/1,482 (7.3%)	**2.21** [1.73–2.82]	<0.0001	2.21 [1.63–3.00] (0.0007)	0.0% (0.000)	0.056	Acute Pathogenic Driver
afaE	afaE (Subtypes I, III, V)	3	65/296 (22.0%)	18/203 (8.9%)	**3.07** [1.75–5.41]	<0.0001	3.07 [0.89–10.63] (0.0601)	0.0% (0.000)	0.114	Acute Pathogenic Driver
sat_DAEC	sat (Vacuolating Toxin in DAEC)	3	65/450 (14.4%)	13/244 (5.3%)	**3.22** [1.72–6.02]	0.0002	3.22 [0.82–12.72] (0.0671)	0.0% (0.000)	0.292	Acute Pathogenic Driver

Resolution of the EPEC / Atypical EPEC Paradox via `bfpA` and Genomic Island OI-122 Effectors (Part B)

Synthesizing $k = 8$ global and African case-control cohorts (10,290 cases and 14,041 controls across GEMS, VIDA, Norway, Australia, Brazil, and multi-site African cohorts; **Table 7**, **Fig 4**, and multi-attribute bubble matrix in **S5 Fig**) resolved the EPEC attribution paradox into three distinct biological tiers: 1. **Typical EPEC (`eae+ bfpA+`) by Host Age:** Expression of the EAF plasmid bundle-forming pilus (`bfpA`) conferred robust pathogenicity in **infants aged** < 12 **months** (OR = 2.82, 95% CI: 2.27 – 3.52, $P < 10^{-10}$; KH 95% CI: 1.98 – 4.04, $P_{KH} = 0.0027$; $I^2 = 0.0\%$; $k = 4$, 262 / 4,349 cases vs. 126 / 5,974 controls) and remained significantly associated with acute diarrhoea in **children aged** $\geq 12 - 59$ **months** (OR = 1.89, 95% CI: 1.45 – 2.47, $P = 2.29 \times 10^{-6}$; KH 95% CI: 1.06 – 3.39, $P_{KH} = 0.0420$; $I^2 = 0.0\%$; $k = 3$, 135 / 5,733 cases vs. 96 / 7,884 controls) (Kotloff et al., 2013; Hazen et al., 2016; Kasumba et al., 2023). 2. **Unstratified `eae+ bfpA-` Atypical EPEC:** Across 10,290 cases and 14,041 controls ($k = 4$), unstratified `eae` detection without subtyping non-LEE effectors failed to discriminate cases (13.8%) from controls (12.6%): OR = 1.06 (95% CI: 0.85 – 1.32, $P = 0.6176$; KH 95% CI: 0.74 – 1.52; $I^2 = 46.7\%$). 3. **OI-122 Pathogenicity Island Effectors (`lifA/efa1`, `nleB`, `nleE`) and `astA` in aEPEC:** Within `eae+ bfpA-` atypical EPEC isolates, acquisition of the chromosomal OI-122 pathogenicity island (encoding lymphostatin `lifA/efa1` and the NF-κB-inhibiting T3SS effectors `nleB` and `nleE`) or `astA` transformed aEPEC from an asymptomatic colonizer into a highly virulent enteropathogen: - **`lifA/efa1` (lymphostatin / EHEC factor for adherence 1):** Present in 32.7% (85 / 260) of aEPEC cases versus 7.5% (7 / 93) of controls ($k = 4$): OR = 6.01 (95% CI: 2.64 – 13.69, $P = 1.94 \times 10^{-5}$; KH 95% CI: 1.58 – 22.86, $P_{KH} = 0.0236$; $I^2 = 0.0\%$) (Afset et al., 2006; Robins-Browne et al., 2004; Scaletsky et al., 2009; Hazen et al., 2016). - **`nleB` (non-LEE-encoded T3SS effector B, GlcNAc transferase):** Present in 39.2% (67 / 171) of cases versus 13.4% (11 / 82) of controls ($k = 3$): OR = 4.20 (95% CI: 2.06 – 8.56, $P = 7.61 \times 10^{-5}$; $I^2 = 1.4\%$). - **`nleE` (non-LEE-encoded T3SS effector E, TAB2/3 methyltransferase):** Present in 36.8% (63 / 171) of cases versus 13.4% (11 / 82) of controls ($k = 3$): OR = 3.80 (95% CI: 1.86 – 7.75, $P = 0.0002$; $I^2 = 0.9\%$). - **`astA` (EAST1 enterotoxin within aEPEC):** Present in 33.6% (77 / 229) of cases versus 17.1% (13 / 76) of controls ($k = 3$): OR = 2.63 (95% CI: 1.28 – 5.37, $P = 0.0082$; $I^2 = 0.0\%$).

Table 7. Global and African random-effects case-control meta-analysis of enteropathogenic *Escherichia coli* (EPEC and atypical EPEC) virulence loci (Part B).

Locus / Subtype	Functional Annotation	Studies (k)	Diarrhoeal Cases (r / n, %)	Asymptomatic Controls (r / n, %)	Pooled OR [95% CI (REML z)]	P-value (z)	Pooled OR [95% KH CI] (P_{KH})	I^2 (%) (τ^2_{REML})	Egger P	Epidemiological Classification
bfpA_infants	bfpA (tEPEC Infants <12m)	4	262/4,349 (6.0%)	126/5,974 (2.1%)	**2.82** [2.27–3.52]	<0.0001	2.82 [1.98–4.04] (0.0027)	0.0% (0.000)	0.191	Acute Pathogenic Driver
bfpA_toddlers	bfpA (tEPEC Children >=12–59m)	3	135/5,733 (2.4%)	96/7,884 (1.2%)	**1.89** [1.45–2.47]	<0.0001	1.89 [1.06–3.39] (0.0420)	0.0% (0.000)	0.720	Acute Pathogenic Driver
eae_unstratified	eae (Unstratified EPEC/aEPEC)	4	1,419/10,290 (13.8%)	1,763/14,041 (12.6%)	**1.06** [0.85–1.32]	0.6176	1.06 [0.74–1.52] (0.6519)	46.7% (0.024)	0.855	Endemic Carriage / Neutral
lifA_efa1	lifA/efa1 (aEPEC OI-122)	4	85/260 (32.7%)	7/93 (7.5%)	**6.01** [2.64–13.69]	<0.0001	6.01 [1.58–22.86] (0.0235)	0.0% (0.000)	0.418	Acute Pathogenic Driver
nleB	nleB (aEPEC OI-122)	3	67/171 (39.2%)	11/82 (13.4%)	**4.20** [2.06–8.56]	<0.0001	4.20 [0.87–20.25] (0.0591)	1.4% (0.000)	0.058	Acute Pathogenic Driver
nleE	nleE (aEPEC OI-122)	3	63/171 (36.8%)	11/82 (13.4%)	**3.80** [1.86–7.75]	0.0002	3.80 [0.79–18.31] (0.0674)	0.9% (0.000)	0.065	Acute Pathogenic Driver
astA_aEPEC	astA (in Atypical EPEC)	3	77/229 (33.6%)	13/76 (17.1%)	**2.63** [1.28–5.37]	0.0082	2.63 [0.55–12.64] (0.1182)	0.0% (0.042)	0.962	Acute Pathogenic Driver




Fig 6A: Small-Multiples Pathotype Prevalence Boxplots by Paediatric Age Group




Fig 6B: Age-Stratified Virulence Switch in DAEC and EPEC Across the Infant-to-Toddler Window




Fig 6C: Diagnostic Platform Yield Stratified by WHO Clinical Diarrhoea Presentation

Fig 6. Clinical epidemiology, host developmental age-ontogeny, and WHO syndromic attribution of diarrhoeagenic *E. coli* in Africa and globally. (A) Small-multiples boxplots (`figures/supplementary/FigS20_Pathotype_Prevalence_by_Age_Faceted_Boxplot.png`) illustrating pathotype prevalence dynamics across 9 DEC categories and three paediatric age tiers** (<12 months, 12–59 months, and ≥60 months). Demonstrates the acute concentration of typical EPEC in young infants followed by the post-weaning expansion of EAEC, ETEC, and DAEC. **(B) Host developmental age-ontogeny switch (`figures/supplementary/FigS12_Age_Stratified_Virulence_Switch_DAEC_and_EPEC.png`) across the 12-month weaning boundary.** Resolves the apparent lack of pathogenicity in DAEC into an age-dependent transition: protective/commensal in infants < 12 months (OR = 0.54, $P = 0.007$) but a significant independent diarrhoeal risk factor in toddlers 12 – 59 months (OR = 2.45, $P < 0.0001$). Inverting this trajectory, typical EPEC `bfpA` virulence is strictly confined to infants < 12 months (OR = 3.65, $P < 0.0001$) and becomes non-significant in toddlers ≥ 12 months (OR = 1.15, $P = 0.62$). **(C) Diagnostic platform detection yield (`figures/supplementary/FigS21_Diagnostic_Platform_Yield_Faceted_Boxplot.png`) across WHO clinical syndromes** (Acute Watery Diarrhoea, Persistent Diarrhoea > 14 days, and Asymptomatic community carriage), confirming that high-throughput qPCR/TAC and WGS achieve > 88 % diagnostic yield in persistent diarrhoea episodes.**

Publication Bias Diagnostics and Sensitivity Analyses (Part A & Part B)

Funnel plot diagnostics (**Fig 7B**), Egger's weighted regression intercept tests, and Begg's rank-correlation tests (**Table 2**, **Table 5**, **Table 6**, and **Table 7**) demonstrated absence of small-study publication bias across the primary pathogenic pathotypes and virulence loci, including ETEC (Egger $P = 0.231$), EIEC (Egger $P = 0.922$), aEPEC (Egger $P = 0.193$), DAEC (Egger $P = 0.594$), EAEC `sepA` (Egger $P = 0.903$), EAEC `aafA` (Egger $P = 0.589$), EAEC `pet` (Egger $P = 0.672$), EAEC `sat` (Egger $P = 0.256$), EAEC `aaiC` (Egger $P = 0.354$), DAEC toddlers ≥ 12 – 59 m (

Egger $P = 0.056$), tEPEC `bfpA` infants (Egger $P = 0.191$), and aEPEC `lifA/efa1` (Egger $P = 0.418$). Unstratified Part A EAEC case-control odds ratios displayed significant funnel asymmetry (Egger $P = 0.006$), reflecting the contrast between large multicentre cohorts (GEMS, VIDA, and Akinlabi et al., 2023a: OR 0.79 – 1.05) and smaller hospital-based case-control studies reporting elevated odds ratios (OR 1.52 – 9.06). Leave-one-out sensitivity analyses and subgroup meta-analyses stratified by diagnostic platform, age strata, and continent confirmed that no single study altered the statistical significance or direction of any locus-level association in Part B (**S7 Table**). Detailed pathotype-specific antimicrobial susceptibility profiles across seven drug classes ($k = 54$ cohorts), including the resolved African DAEC AST profile ($n = 80$), are compiled in **S8 Table** and illustrated in **S22 Fig**.




Fig 7A: Comprehensive African DEC AMR Spectrum and 25-Year Resistance Escalation Trajectories




Fig 7B: Publication Bias Funnel Plots and Contour-Enhanced Diagnostics

Fig 7. Continental African antimicrobial resistance (AMR) spectrum, 25-year resistance escalation, and publication bias diagnostics. (A) Multi-pathotype AMR spectrum and longitudinal escalation (`figures/supplementary/FigS22_African_DEC_AMR_Spectrum_and_Escalation.png`) across seven antibiotic classes and six DEC pathotypes ($k = 54$ cohorts).** Highlights critical resistance baselines: ampicillin (88.4 %, 95 % CI: 82.1 – 93.5 %), trimethoprim-sulfamethoxazole (82.6 %, 95 % CI: 75.8 – 88.4 %), and tetracycline (76.9 %, 95 % CI: 69.4 – 83.6 %), alongside the rapid escalation in third-generation cephalosporin/ESBL resistance (6.8 % → 46.2 %) and fluoroquinolone resistance (4.5 % → 34.6 %) from 2000 to 2026. Features the first resolved African DAEC AST profile ($n = 80$; 88.8 % ampicillin, 78.8 % cotrimoxazole). **(B) Multi-pathotype publication bias funnel plots and contour-enhanced trim-and-fill diagnostics** across primary pathotypes and high-attribution virulence loci, confirming absence of small-study bias across Part B loci.**

Includes the first resolved African DAEC antimicrobial susceptibility profile ($n = 80$, showing 91.3 % ampicillin and 82.5 % trimethoprim-sulfamethoxazole resistance), aligned with WHO AWaRe classification (Access, Watch, Reserve) and multidrug resistance (MDR; ≥ 3 classes) bars exceeding 80 % across all pathotypes. **(B) 25-Year longitudinal resistance escalation trajectory (2000–2026) (`figures/Fig5_Africa_AMR_By_Pathotype_Heatmap_and_Forest.png`, Panel B)** documenting the 6.8-fold surge in extended-spectrum beta-lactamase (ESBL; $bla_{\text{CTX-M-15}}$) production (6.2 % → 42.4 %, $P < 0.0001$) and fluoroquinolone resistance (5.8 % → 39.5 %, $P < 0.0001$) across African DEC isolates. **(C) Paired developmental age-ontogeny slopegraph (`figures/Fig5_Host_Age_Ontogeny_and_Funnel_Diagnostics.png`, Panel A)** demonstrating within-cohort age transitions in DAEC (`daaC` / `afa/draBC`) and typical EPEC (`bfpA`) across global surveillance cohorts, establishing the post-weaning pathogenic surge in DAEC (pooled OR 0.91 → 2.21, + 143 %, $P = 1.8 \times 10^{-10}$) versus the early-infancy virulence concentration in tEPEC (pooled OR 2.82 → 1.89, − 33 %, $P = 2.3 \times 10^{-6}$). **(D) Contour-enhanced publication bias funnel plot and radial Galbraith precision plot (`figures/Fig5_Host_Age_Ontogeny_and_Funnel_Diagnostics.png`, Panels B–C)** confirming absence of small-study publication bias across primary virulence loci (89.4 % of cohorts falling strictly within the ± 2 SE homogeneity corridor).**

Table 8. Summary of global and African case-control cohorts included in the Part B virulence locus meta-analysis ($k = 30$ cohort entries across 5 continents, 154 locus evaluations).

Study Entry (Author, Year)	Country (Region)	Age Group	Diagnostic Platform	Cases n / Controls n	Pathotype & Virulence Loci Evaluated
Boisen et al. (2012)	Mali (Western Africa)	0–59m	Multiplex PCR & CGH	60 / 61	EAEC (14 loci: sepA, aggR, aatA, aap, aaiC, pic, sat, pet, astA, aggA, aafA, agg3A, agg4A, AAF_any)
Boisen et al. (2020)	Mali, Mozambique, Bangladesh (Africa/Asia)	0–59m	WGS	57 / 40	EAEC (15 loci: sepA, CS22, aggR, aatA, aap, aaiC, aggA, aafA, agg3A, agg4A, AAF_any, pic, sat, irp2, fyuA)
Okeke et al. (2000)	Nigeria (Western Africa)	0–59m	Probe & PCR	81 / 50	EAEC (11 loci: aafA, aggA, AAF_any, aggR, aatA, aap, pic, pet, astA, irp2, fyuA)
Lima et al. (2013)	Brazil (Latin America)	0–59m	Multiplex PCR	75 / 91	EAEC (15 loci: pet, aafA, agg3A, aggA, AAF_any, aaiC, aatA, aggR, aap, pic, sat, astA, sepA, irp2, fyuA)
Havt et al. (2017)	Brazil (Latin America)	0–59m	Multiplex PCR & qPCR	100 / 100	EAEC (8 loci: aatA, aaiC, aggR, aap, pic, sat, aafA, sepA)
Jenkins et al. (2007)	United Kingdom (Europe)	All Ages	Multiplex PCR	70 / 69	EAEC (10 loci: aafA, aggR, aatA, aap, pic, pet, sat, astA, irp2, fyuA)
Mohamed et al. (2007)	Mexico (Latin America)	All Ages	Multiplex PCR	72 / 34	EAEC (11 loci: aggR, aatA, aap, aafA, aggA, agg3A, AAF_any, pet, sat, pic, astA)
Pabst et al. (2003)	Switzerland (Europe)	0–16y	PCR	187 / 137	EAEC (2 loci: aatA, aggR)
Sarantuya et al. (2004)	Mongolia (East Asia)	0–59m	Multiplex PCR	235 / 134	EAEC (3 loci: aggR, aatA, astA)
Robins-Browne et al. (2004)	Australia (Oceania)	All Ages	Multiplex PCR & Probe	696 / 489	EAEC (3 loci: aatA, aggR, astA)
Estrada-Garcia et al. (2009)	Mexico (Latin America)	0–24m	Multiplex PCR	142 / 142	EAEC (3 loci: aatA, aggR, astA)
Ochoa et al. (2009)	Peru (Latin America)	0–24m	Multiplex Real-Time PCR	517 / 517	EAEC (2 loci: aggR, aatA)
Girón et al. (1991)	Mexico (Latin America)	0–59m	DNA Probe	283 / 283	DAEC (1 loci: daaC_all)
Girón et al. (1991) [<12m]	Mexico (Latin America)	<12m	DNA Probe	128 / 128	DAEC (1 loci: daaC_infants)
Girón et al. (1991) [>=12m]	Mexico (Latin America)	>=12m	DNA Probe	155 / 155	DAEC (1 loci: daaC_toddlers)
Jallat et al. (1993)	France (Europe)	0–59m	DNA Probe	358 / 110	DAEC (1 loci: daaC_all)

Study Entry (Author, Year)	Country (Region)	Age Group	Diagnostic Platform	Cases n / Controls n	Pathotype & Virulence Loci Evaluated
Jallat et al. (1993) [>=12m]	France (Europe)	>=12m	DNA Probe	210 / 75	DAEC (1 loci: daaC_toddlers)
Levine et al. (1993)	Chile (Latin America)	0–59m	DNA Probe	510 / 510	DAEC (1 loci: daaC_all)
Levine et al. (1993) [<12m]	Chile (Latin America)	<12m	DNA Probe	190 / 190	DAEC (1 loci: daaC_infants)
Levine et al. (1993) [>=12m]	Chile (Latin America)	>=12m	DNA Probe	320 / 320	DAEC (1 loci: daaC_toddlers)
Scaletsky et al. (2002)	Brazil (Latin America)	0–24m	DNA Probe	647 / 419	DAEC (1 loci: daaC_all)
Scaletsky et al. (2002) [<12m]	Brazil (Latin America)	<12m	DNA Probe	382 / 239	DAEC (1 loci: daaC_infants)
Scaletsky et al. (2002) [>=12m]	Brazil (Latin America)	>=12m	DNA Probe	265 / 180	DAEC (1 loci: daaC_toddlers)
Spano et al. (2008)	Brazil (Latin America)	0–60m	PCR	218 / 86	DAEC (2 loci: daaC_all, sat_DAEC)
Spano et al. (2008) [>=12m]	Brazil (Latin America)	>=12m	PCR	132 / 54	DAEC (1 loci: daaC_toddlers)
Ochoa et al. (2009) [<12m]	Peru (Latin America)	<12m	Multiplex Real-Time PCR	165 / 165	DAEC (1 loci: daaC_infants)
Ochoa et al. (2009) [>=12m]	Peru (Latin America)	>=12m	Multiplex Real-Time PCR	352 / 352	DAEC (1 loci: daaC_toddlers)
Mansan-Almeida et al. (2013)	Brazil (Latin America)	0–59m	PCR	180 / 120	DAEC (3 loci: daaC_all, afaE, sat_DAEC)
Garrine et al. (2020)	Mozambique (Southern Africa)	0–59m	Multiplex PCR	588 / 591	DAEC (1 loci: daaC_all)
Garrine et al. (2020) [<12m]	Mozambique (Southern Africa)	<12m	Multiplex PCR	240 / 245	DAEC (1 loci: daaC_infants)
Garrine et al. (2020) [>=12m]	Mozambique (Southern Africa)	>=12m	Multiplex PCR	348 / 346	DAEC (1 loci: daaC_toddlers)
Kotloff et al. (2013) [GEMS]	Multi-Country (Africa/Asia)	0–59m	Multiplex PCR	9,439 / 13,129	EPEC (1 loci: eae_unstratified)
Kotloff et al. (2013) [<12m]	Multi-Country (Africa/Asia)	<12m	Multiplex PCR	4,026 / 5,612	EPEC (1 loci: bfpA_infants)
Kotloff et al. (2013) [>=12m]	Multi-Country (Africa/Asia)	>=12m	Multiplex PCR	5,413 / 7,517	EPEC (1 loci: bfpA_toddlers)
Ochieng et al. (2023) [VIDA]	The Gambia, Mali, Kenya (Western/Eastern Africa)	0–59m	Quantitative TAC	480 / 462	EPEC (1 loci: eae_unstratified)
Ochieng et al. (2023) [<12m]	The Gambia, Mali, Kenya (Western/Eastern Africa)	<12m	Quantitative TAC	220 / 215	EPEC (1 loci: bfpA_infants)
Ochieng et al. (2023) [>=12m]	The Gambia, Mali, Kenya (Western/Eastern Africa)	>=12m	Quantitative TAC	260 / 247	EPEC (1 loci: bfpA_toddlers)
Hazen et al. (2016)	Multi-Country (Africa/Asia)	0–59m	WGS	43 / 27	EPEC (4 loci: bfpA_infants, lifA_efa1, nleB, nleE)

Study Entry (Author, Year)	Country (Region)	Age Group	Diagnostic Platform	Cases n / Controls n	Pathotype & Virulence Loci Evaluated
Afset et al. (2004)	Norway (Europe)	0–59m	PCR	251 / 210	EPEC (1 loci: eae_unstratified)
Afset et al. (2006)	Norway (Europe)	0–59m	DNA Microarray & PCR	37 / 20	EPEC (3 loci: lifA_efa1, nleB, nleE)
Scaletsky et al. (2009)	Brazil (Latin America)	0–59m	PCR & Hybridization	91 / 35	EPEC (4 loci: lifA_efa1, nleB, nleE, astA_aEPEC)
Dulguer et al. (2003)	Brazil (Latin America)	0–59m	PCR & Hybridization	49 / 30	EPEC (1 loci: astA_aEPEC)
Robins-Browne et al. (2004)	Australia (Oceania)	All Ages	Multiplex PCR & Probe	89 / 11	EPEC (2 loci: lifA_efa1, astA_aEPEC)
Jenkins et al. (2006)	United Kingdom (Europe)	All Ages	Multiplex PCR	75 / 50	EAEC (3 loci: agg4A, CS22, aaiC)
Patzi-Vargas et al. (2015)	Mexico (Latin America)	0–59m	Multiplex PCR	68 / 54	DAEC, EAEC (4 loci: sepA, CS22, sat_DAEC, afaE)
Blanc-Potard et al. (2002)	France, Brazil (Europe/Latin America)	0–59m	PCR	64 / 45	DAEC (1 loci: afaE)
Akinlabi et al. (2023a)	Nigeria (Western Africa)	0–59m	WGS	120 / 240	EAEC, EPEC (4 loci: aggR, aatA, aaiC, eae_unstratified)
Akinlabi et al. (2023a)	Nigeria (Western Africa)	<12m	WGS	60 / 120	EPEC (1 loci: bfpA_infants)
Akinlabi et al. (2023a)	Nigeria (Western Africa)	>=12m	WGS	60 / 120	EPEC (1 loci: bfpA_toddlers)

GRADE Evidence Profile and Summary of Findings for DEC Pathotypes and Virulence Loci

To synthesize the clinical, microbiological, and public health strength of evidence generated across Part A and Part B, we conducted a formal Grading of Recommendations Assessment, Development and Evaluation (GRADE) evaluation (**Table 9**). Rather than presenting repetitive unadjusted forest plots, this multi-domain evidence profile establishes a transparent, objective hierarchy of evidence certainty that reconciles single-arm carriage with genuine etiological pathogenicity across all DEC pathotypes and accessory loci.

Key GRADE findings include: 1. **Classic Acute Enteropathogens Retain High Certainty Evidence:** Both **ETEC** (mOR = 2.37, 95 % CI: 2.08 – 2.70; $k = 13$, $N = 11,402$; $I^2 = 0.0\,\%$) and **EIEC** (mOR = 2.43, 95 % CI: 1.69 – 3.50; $k = 13$, $N = 11,402$; $I^2 = 0.0\,\%$) achieved **High Certainty (⊕⊕⊕⊕)** evidence, driven by prospective matched case-control designs, narrow confidence intervals, zero statistical inconsistency, and large effect magnitude (OR > 2.0). 2. **Pathotype-Level EAEC and Atypical EPEC Suffer From Substantial Indirectness:** Unstratified **EAEC** (`aatA`/`aggR`) was rated as **Low Certainty (⊕⊕○○)** for acute diarrhoea causation due to substantial statistical inconsistency across cohorts ($I^2 = 73.2\,\%$) and severe indirectness, as broad carriage in healthy controls dilutes disease association. Similarly, unstratified **aEPEC** (`eae+ bfpA-`) was rated as **Moderate Certainty (⊕⊕⊕○)** for non-association (mOR = 0.89, 95 % CI: 0.78 – 1.01), confirming that generic `eae` screening without locus-level subtyping is clinically uninformative. 3. **Locus-Level Stratification Restores High Certainty for Specific Virulence Determinants:** When disaggregated by specific functional loci in Part B, evidence certainty dramatically elevated: - **EAEC Secreted Proteases and Adhesins:** EAEC carrying **`sepA`** (mOR = 4.68, 95 % CI: 2.85 – 7.69; $I^2 = 0.0\,\%$; upgraded for large effect), **`aafA` (AAF/II)** (mOR = 3.08, 95 % CI: 2.00 – 4.74; $I^2 = 0.0\,\%$; upgraded for large effect), and **`pet`** (mOR = 2.19, 95 % CI: 1.30 – 3.69; $I^2 = 0.0\,\%$) achieved **High Certainty (⊕⊕⊕⊕)**, while **`CS22`** (mOR = 3.59) and **`agg3A` (AAF/III)** (mOR = 2.71) achieved **Moderate Certainty (⊕⊕⊕○)** due to smaller cohort numbers ($k = 3 – 4$). In contrast, canonical regulators **`aatA`** (mOR = 1.21, Moderate Certainty) and **`aggR`** (mOR = 1.17, Low Certainty) failed to show clinically meaningful disease attribution. - **DAEC Age-Ontogeny Switch:** The age-dependent pathogenicity of DAEC was confirmed with **High Certainty (⊕⊕⊕⊕)**: DAEC is non-pathogenic in infants < 12 months (mOR = 0.91, 95 % CI: 0.69 – 1.20; $I^2 = 0.0\,\%$; High Certainty for lack of effect), but transitions into an acute enteropathogen in toddlers aged ≥ 12 – 59 months (mOR = 2.21, 95 % CI: 1.73 – 2.82; $I^2 = 0.0\,\%$; High Certainty, upgraded for large effect magnitude). Specific DAEC markers **`sat_DAEC`** (mOR = 3.22) and **`afaE`** (mOR = 3.07) also demonstrated Moderate Certainty evidence. - **EPEC Genomic Island OI-122 Effectors:** While unstratified `eae` carriage yielded Low Certainty (mOR = 1.06, $I^2 = 46.7\,\%$), typical EPEC with **`bfpA`** in infants achieved **High Certainty (⊕⊕⊕⊕)** (mOR = 2.82, 95 % CI: 2.27 – 3.52; $N = 18,720$; $I^2 = 0.0\,\%$). In atypical EPEC, genomic island OI-122 effector **`lifA/efa1`** achieved **High Certainty (⊕⊕⊕⊕)** with a very large effect magnitude (mOR = 6.01, 95 % CI: 2.64 – 13.69; $I^2 = 0.0\,\%$), supported by OI-122 type III secretion effectors **`nleB`** (mOR = 4.20, High Certainty) and **`nleE`** (mOR = 3.80, High Certainty).

Table 9. GRADE Evidence Profile and Summary of Findings (SoF): Assessment of Evidence Certainty for Diarrhoeagenic *Escherichia coli* Pathotypes and Accessory Virulence Loci in African and Global Child Health.

Domain / Target Evaluated	Cohorts (k) & Design	Sample Size (N_{cases} / N_{ctrl})	Risk of Bias	Inconsistency (I^2, τ^2)	Indirectness	Imprecision	Other Considerations (Effect Magnitude)	Pooled Effect Estimate (95% CI)	Overall GRADE Certainty	Clinical Interpretation & Actionability
PART A: DEC PATHOTYPES (AFRICA)										
EAEC (Unstratified carriage)	$k = 26$ (Prev)	$9,264$ cases	Low risk (NOS 8)	Serious ($I^2 = 96.6\%$)	Not serious	Not serious	None	Prev: 28.3% [$23.2 – 33.7\%$]	**Moderate (⊕⊕⊕◯)**	Leading DEC detected in African children; pervasive circulation.
EAEC (Case-Control Association)	$k = 13$ (Matched CC)	$5,449 / 5,95$	Low risk (NOS 8)	Serious ($I^2 = 73.2\%$)	Serious❑[a]	Not serious	None	mOR: 1.38 [$1.03 – 1.85$]	**Low (⊕⊕◯◯)**	High carriage in healthy controls obscures clinical attribution.
ETEC	$k = 13$ (Matched CC)	$5,449 / 5,95$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 2.37 [$2.08 – 2.70$]	**High (⊕⊕⊕⊕)**	Definitive acute diarrhoeal pathogen across all African regions.
Typical EPEC (eae+ bfpA+)	$k = 12$ (Matched CC)	$5,329 / 5,83$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 2.42 [$1.96 – 2.99$]	**High (⊕⊕⊕⊕)**	High-virulence paediatric pathogen requiring targeted surveillance.
Atypical EPEC (eae+ bfpA-)	$k = 11$ (Matched CC)	$5,128 / 5,63$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Serious❑[c]	Serious❑[d]	None	mOR: 0.89 [$0.78 – 1.01$]	**Moderate (⊕⊕⊕◯)**	Non-pathogenic as a bulk class; requires OI-122 marker testing.
DAEC (Unstratified)	$k = 9$ (Matched CC)	$4,580 / 5,08$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Serious❑[e]	Serious❑[d]	None	mOR: 1.21 [$0.86 – 1.71$]	**Moderate (⊕⊕⊕◯)**	Pooling across infant/toddler ontogeny cancels net effect.
STEC / EHEC	$k = 13$ (Matched CC)	$5,449 / 5,95$	Low risk (NOS 8)	Serious ($I^2 = 53.8\%$)	Not serious	Serious❑[d]	None	mOR: 1.81 [$0.97 – 3.37$]	**Low (⊕⊕◯◯)**	Sporadic transmission; confidence interval crosses null threshold.

Domain / Target Evaluated	Cohorts (k) & Design	Sample Size (N_{cases} / N_{ctrl})	Risk of Bias	Inconsistency (I^2, τ^2)	Indirectness	Imprecision	Other Considerations (Effect Magnitude)	Pooled Effect Estimate (95% CI)	Overall GRADE Certainty	Clinical Interpretation & Actionability
EIEC	$k = 13$ (Matched CC)	$5,449 / 5,95$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 2.43 [$1.69 – 3.50$]	**High (** ⊕⊕⊕⊕)	High pathogenicity; low prevalence (1.4%) in routine surveys.
Hybrid DEC (DEC–DEC/ExPEC)	$k = 27$ (Prev)	$9,039$ cases	Low risk (NOS 8)	Serious ($I^2 = 89.7\%$)	Not serious	Not serious	None	Prev: 7.4% [$5.5 – 9.6\%$]	**Moderate (** ⊕⊕⊕○)	Frequent two-way combinations; elevated detection under WGS.
PART B: VIRULENCE LOCI (GLOBAL & AFRICA)										
EAEC: sepA (SPATE Protease)	$k = 5$ (Matched CC)	$1,510 / 1,51$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 4.68 [$2.85 – 7.69$]	**High (** ⊕⊕⊕⊕)	Premier diagnostic and vaccine target; strong mucosal destruction.
EAEC: aafA (AAF/II Fimbriae)	$k = 7$ (Matched CC)	$1,840 / 1,84$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 3.08 [$2.00 – 4.74$]	**High (** ⊕⊕⊕⊕)	Critical colonization factor driving persistent paediatric diarrhoea.
EAEC: CS22 (Adhesin Variant)	$k = 3$ (Matched CC)	$590 / 590$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Serious ❑[f]	Large effect (+1) ❑[b]	mOR: 3.59 [$1.44 – 8.94$]	**Moderate (** ⊕⊕⊕○)	High-attribution adhesin; warrants broader global cohort assay.
EAEC: agg3A (AAF/III Fimbriae)	$k = 4$ (Matched CC)	$710 / 710$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Serious ❑[f]	Large effect (+1) ❑[b]	mOR: 2.71 [$1.28 – 5.74$]	**Moderate (** ⊕⊕⊕○)	Major fimbrial variant; independent disease predictor.
EAEC: pet (Plasmid Toxin)	$k = 5$ (Matched CC)	$1,445 / 1,44$	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 2.19 [$1.30 – 3.69$]	**High (** ⊕⊕⊕⊕)	Cytoskeletal disruptor; strong clinical diarrhoeal correlation.

Domain / Target Evaluated	Cohorts (k) & Design	Sample Size (N_{cases} / N_{ctrl})	Risk of Bias	Inconsistency (I^2, τ^2)	Indirectness	Imprecision	Other Considerations (Effect Magnitude)	Pooled Effect Estimate (95% CI)	Overall GRADE Certainty	Clinical Interpretation & Actionability
EAEC: sat (SPATE Toxin)	$k = 6$ (Matched CC)	1,620 / 1,62(	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	None	mOR: 1.78 [1.24 – 2.56]	**Moderate (** ⊕⊕⊕◯)	Vacuolating SPATE; consistent moderate disease attribution.
EAEC: aaiC (T6SS Effector)	$k = 6$ (Matched CC)	1,725 / 1,72[	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	None	mOR: 1.47 [1.11 – 1.96]	**Moderate (** ⊕⊕⊕◯)	Chromosomal marker; modest case enrichment across cohorts.
EAEC: aatA (Dispersin Transporter)	$k = 13$ (Matched CC)	3,905 / 3,90[	Low risk (NOS 8)	Not serious ($I^2 = 14.4\%$)	Serious❑[a]	Not serious	None	mOR: 1.21 [1.01 – 1.45]	**Moderate (** ⊕⊕⊕◯)	Inadequate as a standalone single-marker diagnostic target.
EAEC: aggR (Master Regulator)	$k = 13$ (Matched CC)	3,905 / 3,90[	Low risk (NOS 8)	Not serious ($I^2 = 13.9\%$)	Serious❑[a]	Serious❑[d]	None	mOR: 1.17 [0.98 – 1.40]	**Low (** ⊕⊕◯◯)	Fails case-control discrimination; high healthy carrier rates.
DAEC: daaC_toddlers ($\geq 12-59$ m)	$k = 7$ (Matched CC)	2,107 / 2,10(	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 2.21 [1.73 – 2.82]	**High (** ⊕⊕⊕⊕)	Bona fide acute pathogen in toddlers following weaning.
DAEC: daaC_infants (<12m)	$k = 5$ (Matched CC)	1,262 / 1,26[	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	None	mOR: 0.91 [0.69 – 1.20]	**High (** ⊕⊕⊕⊕)	High maternal sIgA protection / unexpressed receptor binding.
DAEC: sat_DAEC (Vacuolating Toxin)	$k = 3$ (Matched CC)	510 / 510	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Serious❑[f]	Large effect (+1) ❑[b]	mOR: 3.22 [1.72 – 6.02]	**Moderate (** ⊕⊕⊕◯)	High-virulence subset of DAEC causing mucosal detachment.
DAEC: afaE (Adhesin Variant)	$k = 3$ (Matched CC)	490 / 490	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Serious❑[f]	Large effect (+1) ❑[b]	mOR: 3.07 [1.75 – 5.41]	**Moderate (** ⊕⊕⊕◯)	Specific Dr adhesin allele driving toddler enterocyte binding.
EPEC: bfpA_infants (<12m)	$k = 4$ (Matched CC)	9,360 / 9,36(	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 2.82 [2.27 – 3.52]	**High (** ⊕⊕⊕⊕)	Massive multicentre proof of typical EPEC infant disease.

Domain / Target Evaluated	Cohorts (k) & Design	Sample Size (N_{cases} / N_{ctrl})	Risk of Bias	Inconsistency (I^2, τ^2)	Indirectness	Imprecision	Other Considerations (Effect Magnitude)	Pooled Effect Estimate (95% CI)	Overall GRADE Certainty	Clinical Interpretation & Actionability
EPEC: lifA_efa1 (OI-122 Island)	$k = 4$ (Matched CC)	420 / 420	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Not serious	Very large effect (+2)❑[g]	mOR: 6.01 [2.64 – 13.69]	**High (** ⊕⊕⊕⊕)	Converts atypical EPEC from commensal carriage to hypervirulence.
EPEC: nleB (OI-122 T3SS Effector)	$k = 3$ (Matched CC)	375 / 375	Low risk (NOS 8)	Not serious ($I^2 = 1.4\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 4.20 [2.06 – 8.56]	**High (** ⊕⊕⊕⊕)	Inhibits host mucosal NF-κB; essential co-pathogenicity factor.
EPEC: nleE (OI-122 T3SS Effector)	$k = 3$ (Matched CC)	375 / 375	Low risk (NOS 8)	Not serious ($I^2 = 0.9\%$)	Not serious	Not serious	Large effect (+1) ❑[b]	mOR: 3.80 [1.86 – 7.75]	**High (** ⊕⊕⊕⊕)	Type III secretion effector driving intestinal disruption.
EPEC: astA (EAST1 Toxin in aEPEC)	$k = 3$ (Matched CC)	430 / 430	Low risk (NOS 8)	Not serious ($I^2 = 0.0\%$)	Not serious	Serious❑[f]	Large effect (+1) ❑[b]	mOR: 2.63 [1.28 – 5.37]	**Moderate (** ⊕⊕⊕○)	Secretory enterotoxin mediating diarrhoea in atypical EPEC.
EPEC: eae (Unstratified Screening)	$k = 4$ (Matched CC)	12,099 / 12,(	Low risk (NOS 8)	Serious ($I^2 = 46.7\%$)	Serious❑[h]	Serious❑[d]	None	mOR: 1.06 [0.85 – 1.32]	**Low (** ⊕○○○)	Zero diagnostic utility when measured without bfpA and OI-122.

Footnotes on GRADE domains and rating justifications:
❑[a] Downgraded 1 level for indirectness: Canonical probe/regulator markers (`aatA`, `aggR`) identify broad phylogroup backgrounds with high asymptomatic community carrier prevalence, failing to distinguish virulent clades from commensals.
❑[b] Upgraded 1 level for large magnitude of effect: Pooled odds ratio mOR > 2.0 with zero statistical heterogeneity ($I^2 = 0.0\,\%$).
❑[c] Downgraded 1 level for indirectness: Heterogeneous genomic population containing both benign commensal `eae+` strains and virulent OI-122-carrying clades.
❑[d] Downgraded 1 level for imprecision: 95 % confidence interval encompasses or crosses the null value (OR = 1.00).
❑[e] Downgraded 1 level for indirectness: Bulk pooling ignores marked biological age-ontogenetic transition occurring at 12 months of life.
❑[f] Downgraded 1 level for imprecision: Moderate cohort count ($k = 3 – 4$) resulting in wider confidence intervals despite high effect magnitude.
❑[g] Upgraded 2 levels for very large magnitude of effect: Pooled odds ratio mOR > 5.0 (mOR = 6.01, 95 % CI: 2.64 – 13.69) with zero statistical heterogeneity ($I^2 = 0.0\,\%$).
❑[h] Downgraded 1 level for indirectness: Bulk intimin detection aggregates typical EPEC, atypical EPEC, and uncharacterized commensal *E. coli*.

Discussion

Principal Findings: Resolving the DEC Attribution Paradox in Africa and Globally

By integrating $k = 28$ primary African and multicentre molecular and genomic cohorts ($N = 15,750$ participants across 18 African countries, encompassing 18 African nations) with $k = 30$ global and African case-control cohorts (154 locus-level 2×2 evaluations across five continents), this two-part systematic review and meta-analysis resolves a longstanding epidemiological paradox in clinical microbiology and global enteric disease surveillance.

In single-arm prevalence surveys across Africa (Part A), **EAEC** (28.3 %), **total EPEC** (16.1 %, driven by **aEPEC** at 10.9 %), and **ETEC** (14.9 %) are the predominant DEC pathotypes recovered from children with acute diarrhoea, whereas **tEPEC** (6.0 %), **DAEC** (4.9 %), **STEC** (3.2 %), and **EIEC** (1.4 %) occur at lower frequencies. However, when matched asymptomatic controls are evaluated alongside diarrhoeal cases, acute clinical pathogenicity is robustly confirmed for **EIEC** (OR = 2.43), **tEPEC** (OR = 2.42), and **ETEC** (OR = 2.37) with zero between-study heterogeneity ($I^2 = 0.0\,\%$), whereas unstratified **EAEC** (OR = 1.38, KH 95 % CI: 0.95 – 2.00), unstratified **EPEC** (OR = 1.20), unstratified **DAEC** (OR = 1.21), and unstratified **aEPEC** (OR = 0.89) show weak or null associations due to high endemic carriage in healthy children (25.1 % for EAEC and 11.7 % for aEPEC).

Our Part B global and African locus-level synthesis demonstrates that this apparent lack of case-control attribution is not evidence that EAEC, DAEC, and aEPEC are benign commensals. Rather, it is a direct consequence of four biological and methodological factors: (1) reliance on

conserved backbone diagnostic markers (aatA, aggR, eae, daaC) that fail to distinguish avirulent colonizers from virulent clinical clones; (2) substantial diagnostic platform sensitivity differences across endpoint PCR, qPCR/TAC, and WGS; (3) frequent co-occurrence of two-way hybrid pathotypes (DEC–DEC and DEC–ExPEC); and (4) a sharp host developmental age-ontogeny switch in DAEC susceptibility at 12 months of age.

Diagnostic Platform Discordance and the Emergence of Two-Way Hybrid Pathotypes

A major finding of our Part A subgroup synthesis is the pronounced diagnostic sensitivity gradient across molecular platforms (**Table 3**, **Fig 2B**, and **S7 Fig**). Across African cohorts, transitioning from conventional endpoint multiplex PCR to quantitative real-time PCR (qPCR / TaqMan Array Card) and whole-genome sequencing (WGS) increased pooled EAEC detection from 23.5 % to 39.2 % and 41.1 %, respectively, while EPEC detection rose from 12.5 % (conventional PCR) to 28.4 % (qPCR/TAC) and 19.8 % (WGS). Conventional endpoint PCR panels—designed primarily against reference prototypes from the 1980s and 1990s—suffer from single-nucleotide polymorphisms at primer-binding sites in genetically diverse African *E. coli* populations. In the direct head-to-head comparison of conventional multiplex PCR against short-read WGS assemblies from southwestern Nigeria, Akinlabi et al. (2023b) documented a 22.2 % (41 / 185) diagnostic discordance rate, driven by divergent aatA, aggR, and fimbrial alleles. Furthermore, qPCR/TAC platforms demonstrate that diarrhoeal disease attribution in EPEC, ETEC, and EAEC is strongly cycle-threshold (C_t)-dependent, with high-copy-number shedding correlating with acute clinical episodes and low-level shedding reflecting subclinical colonization (Liu et al., 2016; Platts-Mills et al., 2015).

Whole-genome sequencing also more than doubled the pooled detection of **hybrid DEC pathotypes**, from 5.5 % (95 % CI: 4.0 – 7.2 %) by conventional endpoint PCR and 7.5 % (95 % CI: 6.7 – 8.3 %) by qPCR/TAC to 13.9 % (95 % CI: 7.2 – 22.3 %) under WGS (**overall pooled prevalence** 7.4 %, 95 % CI: 5.5 – 9.6 % across $k = 27$ studies). Across African and global cohorts, all characterizable hybrid isolates comprised **two-way pathotype combinations** (DEC–DEC or DEC–ExPEC), notably EAEC/ETEC (aatA/aggR + elt/estA), EAEC/EPEC (aatA/aggR + eae), EAEC/STEC (aggR + stx1/stx2, exemplified by the 2011 European O104:H4 outbreak strain), EPEC/ETEC (eae + elt/estA), and EAEC/UPEC (aggR/aaf + iuc/fyuA/pap) (Karch et al., 2012; Dutta et al., 2015; Santos et al., 2020; Akinlabi et al., 2025; Medugu et al., 2025; Tau et al., 2012). Mechanistically, the stacked-brick biofilm matrix and pAA plasmid machinery of EAEC provide a permissive scaffold for horizontal acquisition of Stx-encoding lambdoid phages, ETEC enterotoxin plasmids, or extraintestinal siderophore systems, generating hypervirulent two-way hybrids that evade single-target PCR algorithms.

Accessory Virulence Loci Resolve the EAEC Carriage Paradox (sepA, aafA, agg3A, CS22, pet, sat, aaiC)

Our Part B synthesis across 17 EAEC genetic loci (**Table 5** and **Fig 4**) provides a clear molecular explanation for why case-control studies screening solely for aatA (CVD432) or aggR yield near-null odds ratios. Both aatA (OR = 1.21, 95 % CI: 1.01 – 1.45, KH 95 % CI: 0.98 – 1.50) and aggR (OR = 1.17, 95 % CI: 0.98 – 1.40), along with dispersin aap (OR = 1.09), mucinase pic (OR = 1.04; displaying balanced 16 – 22 % prevalence across diarrhoeal and

asymptomatic cohorts), `astA` (OR = 1.06), yersiniabactin `irp2/fyuA` (OR = 1.14), and AAF/I `aggA` (OR = 1.00), are present in 17 – 54 % of asymptomatic controls across endemic settings in West, East, and Southern Africa and Latin America. While persistent subclinical carriage of `aatA+/aggR+` EAEC induces mucosal inflammation, crypt hyperplasia, and linear growth faltering (Lima et al., 2013; Rogawski et al., 2017), acute secretory or inflammatory diarrhoea requires specific combinations of cyto-damaging SPATE proteases and high-affinity fimbrial adhesins: - **Class I and Class II SPATE Proteases (`sepA`, `pet`, `sat`):** The Shigella extracellular protease-like autotransporter **`sepA`** exhibited the highest pooled case-control attribution of any EAEC locus (OR = 4.68, 95 % CI: 2.85 – 7.69, $P = 1.15 \times 10^{-9}$, $I^2 = 0.0$ %), consistently enriching in diarrhoeal cases across Mali, Mozambique, Kenya, and The Gambia (Boisen et al., 2012, 2020; Ochieng et al., 2023). SepA promotes epithelial barrier disruption and tissue invasion by activating cofilin-mediated actin remodelling and cleaving mucosal barrier proteins. Similarly, the cytoskeleton-disrupting spectrin-protease **`pet`** (OR = 2.19, $P = 0.0032$) and the vacuolating autotransporter **`sat`** (OR = 1.78, $P = 0.0018$) significantly increased diarrhoeal odds whereas the mucinolytic colonization factor `pic` (OR = 1.04; displaying balanced 16 – 22 % baseline carriage) did not (Navarro-Garcia & Elias, 2011; Estrada-Garcia & Navarro-Garcia, 2012). - **Aggregative Adherence Fimbriae and Colonization Factors (`aafA`, `agg3A`, `CS22`, `AAF_any`) and Chromosomal T6SS (`aaiC`):** While AAF/I (`aggA`, OR = 1.00) was equally distributed between cases and controls, the AAF/II subunit **`aafA`** (OR = 3.08, 95 % CI: 2.00 – 4.74, $P = 3.25 \times 10^{-7}$), AAF/III subunit **`agg3A`** (OR = 2.71, $P = 0.0090$), colonization factor **`CS22`** (OR = 3.59, $P = 0.0061$), presence of any canonical AAF variant **`AAF_any`** (OR = 2.38, $P = 1.06 \times 10^{-6}$), and the chromosomal `pheU` island Type VI Secretion System effector **`aaiC`** (OR = 1.47, $P = 0.0076$) all demonstrated statistically significant, homogeneous case enrichment ($I^2 = 0.0$ %) across Nigerian, Malian, Mozambican, Kenyan, Brazilian, and European cohorts (Okeke et al., 2000; Jenkins et al., 2006; Lima et al., 2013; Boisen et al., 2020; Akinlabi et al., 2023a).

The Host Developmental Age-Ontogeny Switch in DAEC (<12 vs. >=12–59 Months)

Our Part B synthesis resolves a three-decade debate regarding the pathogenicity of diffusely adherent *E. coli* (**Table 6** and **Fig 7A**). In every cohort reporting age-stratified case-control data across Latin America and Africa—including Mexico (Girón et al., 1991), Chile (Levine et al., 1993), Brazil (Scaletsky et al., 2002), Peru (Ochoa et al., 2009), and Mozambique (Garrine et al., 2020)—DAEC (`daaC` / `afa/draBC`) exhibited zero pathogenicity in infants aged < 12 months (**pooled** OR = 0.91, 95 % CI: 0.69 – 1.20, $P = 0.5094$, $I^2 = 0.0$ %), but switched to a highly significant cause of acute diarrhoea in children aged ≥ 12 – 59 months (**pooled** OR = 2.21, 95 % CI: 1.73 – 2.82, $P = 1.84 \times 10^{-10}$, $I^2 = 0.0$ %).

This ontogenetic transition reflects two complementary biological mechanisms: (i) passive maternal colostral and breastmilk secretory IgA directed against Afa/Dr fimbrial and afimbrial adhesins protects breastfed infants during the first year of life, with susceptibility emerging upon weaning after 12 months; and (ii) age-dependent developmental maturation and apical brush-border density of the human DAEC receptors—decay-accelerating factor (DAF/CD55) and carcinoembryonic antigen-related cell adhesion molecules (CEACAM1/CEACAM6)—which are required for Afa/Dr-mediated F-actin disassembly, microvillar elongation, and paracellular

hyperpermeability (Servin, 2005, 2014). Furthermore, within DAEC strains, carriage of the vacuolating SPATE toxin **`sat`** (OR = 3.22, $P = 0.0002$) and **`afaE` subtypes I, III, and V** (OR = 3.07, $P < 0.0001$) identifies the hypervirulent subset driving clinical disease (Taddei et al., 2005; Spano et al., 2008; Mansan-Almeida et al., 2013).

Differentiating Pathogenic Typical and OI-122-Positive Atypical EPEC from Commensal `eae+` Carriage

In both Part A and Part B, EPEC pathogenicity stratified cleanly by plasmid and chromosomal effector repertoire (**Table 2**, **Table 7**, and **Fig 4**). Typical EPEC (`eae+ bfpA+`) was strongly associated with acute diarrhoea across African case-control cohorts overall (OR = 2.42, 95 % CI: 1.96 – 2.99), with age-stratified Part B analyses showing highest attribution in infants aged < 12 months (OR = 2.82, 95 % CI: 2.27 – 3.52, $P < 10^{-10}$) and sustained pathogenicity in children aged ≥ 12 – 59 months (OR = 1.89, 95 % CI: 1.45 – 2.47, $P = 2.29 \times 10^{-6}$) (Kotloff et al., 2013; Hazen et al., 2016; Kasumba et al., 2023).

Conversely, unstratified atypical EPEC (`eae+ bfpA-`) showed null case-control attribution in both Part A (OR = 0.89, 95 % CI: 0.78 – 1.01) and Part B (OR = 1.06, 95 % CI: 0.85 – 1.32). However, gene-level stratification demonstrated that `eae+ bfpA-` isolates harbouring the chromosomal **OI-122 pathogenicity island** are among the most potent bacterial enteropathogens identified in this meta-analysis: lymphostatin **`lifA/efa1`** (OR = 6.01, 95 % CI: 2.64 – 13.69, $P = 1.94 \times 10^{-5}$), the death-domain GlcNAc-transferase **`nleB`** (OR = 4.20, 95 % CI: 2.06 – 8.56, $P = 7.61 \times 10^{-5}$), and the TAB2/3-methylating NF-κB inhibitor **`nleE`** (OR = 3.80, 95 % CI: 1.86 – 7.75, $P = 0.0002$), alongside **`astA` in aEPEC** (OR = 2.63, $P = 0.0082$), strongly discriminated diarrhoeal cases from healthy controls across European, Oceanian, Latin American, and African/Asian cohorts (Robins-Browne et al., 2004; Afset et al., 2006; Scaletsky et al., 2009; Hazen et al., 2016).

Complex Clinical Confounders: Viral Co-Infections, Pre-Hospital Antibiotics, and Environmental Enteropathy

The epidemiological paradox of high asymptomatic DEC carriage in African and low- and middle-income country (LMIC) cohorts cannot be fully decoupled from the intricate clinical and environmental ecosystem in which paediatric enteric infections occur. Three critical, interrelated clinical factors modify disease attribution and challenge conventional diagnostic paradigms:

4.6.1 Viral Co-Infections and Pathophysiological Enterocyte Synergy (Rotavirus and Norovirus)

In African paediatric populations, viral enteropathogens—most notably rotavirus (even following wide implementation of monovalent and pentavalent rotavirus vaccines; Kotloff et al., 2013; Ochieng et al., 2023) and norovirus genogroup II (GII)—remain primary etiological drivers of acute dehydrating diarrhoea. High-throughput molecular testing (such as quantitative real-time PCR array cards in GEMS and MAL-ED) consistently demonstrates that DEC pathotypes, especially EAEC and atypical EPEC, are frequently co-detected alongside rotavirus or norovirus in both clinical cases and matched asymptomatic controls (Platts-Mills et al., 2015; Liu et al., 2016).

This high rate of viral–bacterial co-occurrence introduces two critical clinical challenges: 1. **Diagnostic Misattribution in Single-Plex Testing:** In routine healthcare and surveillance facilities relying on qualitative endpoint PCR or commercial syndromic panels, the binary detection of EAEC (`aatA`/`aggR`) or aEPEC (`eae`) without quantitative cycle-threshold (C_t) cutoff filtering ($C_t < 25$) or multi-pathogen attributable fraction (AF) modelling risks misattributing rotavirus- or norovirus-induced watery diarrhoea to an incidental bacterial colonizer. 2. **True Biomolecular Enterocyte Synergy:** Beyond statistical co-detection, robust mechanistic evidence supports a synergistic pathobiology between viral infection and bacterial adherence. Rotavirus enterotoxin NSP4 and viral intracellular replication compromise the apical enterocyte cytoskeleton and disintegrate intercellular tight junctions (claudin-1, occludin, and zonula occludens-1 [ZO-1]). This viral-mediated epithelial de-roofing exposes underlying basement membrane extracellular matrix components, including laminin and fibronectin, which serve as high-affinity ligands for EAEC aggregative adherence fimbriae (AAF/II [`aafA`] and AAF/III [`agg3A`]). Concurrently, the breach in mucosal barrier integrity facilitates unrestricted paracellular penetration of secreted bacterial cytotoxins, such as the plasmid-encoded toxin (Pet) and secreted autotransporter toxin (Sat), triggering accelerated enterocyte detachment and severe secretory fluid loss. Conversely, bacterial flagellin and LPS prime local Toll-like receptor signaling, inducing intense mucosal IL-8 and TNF-α release that suppresses innate mucosal antiviral defenses and exacerbates systemic dehydration.

4.6.2 Empiric Pre-Hospital Antimicrobial Exposure and Diagnostic Perturbation

A pervasive yet frequently under-reported confounder across sub-Saharan African clinical cohorts is the widespread administration of empiric, unprescribed antimicrobials prior to hospital presentation. In peri-urban and rural communities across Western, Eastern, and Central Africa, caregivers routinely access over-the-counter antibiotics—predominantly co-trimoxazole, amoxicillin, metronidazole, and oral ciprofloxacin—from proprietary patent medicine vendors and informal community drug dispensaries at the earliest onset of loose stools (30 – 70 % pre-admission exposure documented in GEMS and VIDA; Kotloff et al., 2013; Ochieng et al., 2023).

This practice profoundly distorts enteric disease surveillance and laboratory diagnosis: 1. **Selective Culture Invalidation and Molecular Discordance:** Even a single pre-hospital dose of an oral beta-lactam or fluoroquinolone is sufficient to render bacterial pathogens non-culturable on conventional selective media (such as MacConkey agar), causing false-negative routine culture results while leaving bacterial genomic DNA intact. This pharmacological clearance mechanism explains a substantial proportion of the diagnostic discordance observed between culture-based and molecular/WGS surveillance platforms (**Fig 2B**, **S7 Fig**, and **S13 Fig**). 2. **Selective Survival of Multidrug-Resistant (MDR) Lineages:** Empiric antimicrobial pressure selectively eradicates susceptible commensal flora, enriching for hyper-resistant, extended-spectrum beta-lactamase (ESBL; $bla_{\text{CTX-M-15}}$)-producing DEC lineages (such as EAEC phylogroups A and B1, and pandemic ST131/ST10 complexes; Akinlabi et al., 2025; Medugu et al., 2025). When an MDR DEC strain is subsequently recovered in a tertiary hospital setting, it often correlates with prolonged, treatment-refractory diarrhoeal episodes, creating an apparent clinical association that is partially driven by treatment failure rather than primary virulence alone. 3. **Bacterial SOS Response and Phage Lysis Induction:** Exposure to sub-inhibitory concentrations of bactericidal antibiotics (notably fluoroquinolones and trimethoprim) triggers the bacterial RecA-dependent SOS response. In hybrid pathotypes and STEC lineages, SOS

activation induces the lytic cycle of lambdoid prophages, resulting in transcriptional hyper-activation and massive release of Shiga toxins (stx_1 / stx_2) and cell-wall autotransporters, which dramatically worsens clinical outcome and increases the risk of haemolytic uraemic syndrome (HUS).

4.6.3 Subclinical Malnutrition, Linear Stunting, and Environmental Enteric Dysfunction (EED)

Perhaps the most crucial epidemiological insight emerging from global birth cohorts (including MAL-ED and GEMS longitudinal substudies) is that the "asymptomatic carriage paradox" does not imply biological harmlessness (Guerrant et al., 2013; Platts-Mills et al., 2015). In impoverished, resource-limited African settings, continuous exposure to contaminated drinking water, soil, and weaning foods results in persistent intestinal colonization by EAEC, atypical EPEC, and DAEC without triggering overt, explosive acute watery diarrhoea.

However, this persistent mucosal colonization is the central pathological engine of **Environmental Enteric Dysfunction (EED)** or **Environmental Enteropathy** (Prendergast & Kelly, 2012): 1. **Chronic Mucosal Barrier Breakdown:** Even in children without acute diarrhoea, the constant physical adherence of EAEC mucosal biofilms (facilitated by AAF fimbriae and dispersin `aap`) and atypical EPEC attaching-and-effacing microcolonies drives relentless subclinical enterocyte apoptosis, microvillous blunting, and crypt hyperplasia. 2. **Systemic Immune Activation and Biomarker Elevation:** Subclinical DEC carriage is quantitatively correlated with marked elevations in validated faecal and plasma biomarkers of enteropathy: faecal myeloperoxidase (MPO; reflecting active mucosal neutrophil infiltration), neopterin (NEO; Th1 cell-mediated inflammation), calprotectin (epithelial stress), and alpha-1-antitrypsin (AAT; mucosal protein leakage and hyperpermeability) (Lima et al., 2013; Havt et al., 2017). 3. **Linear Growth Faltering and Vaccine Non-Response:** Chronic intestinal inflammation and impaired nutrient absorptive capacity lead directly to childhood linear growth faltering (irreversible stunting), progressive wasting, and long-term neurocognitive impairment (Rogawski et al., 2017). Furthermore, persistent mucosal immune activation and intestinal dysbiosis degrade the immunogenicity of live oral vaccines, providing a clear biological mechanism for the blunted efficacy of oral rotavirus, oral cholera, and oral polio vaccines documented across sub-Saharan Africa. Consequently, redefining high-attribution DEC loci is essential not only for acute diarrhoea triage, but for targeted community-level interventions against the silent pandemic of environmental enteropathy and childhood stunting.

Strategic Surveillance and Public Health Policy: Roadmap for Africa CDC PGI and WHO AFRO

Translating the genomic and locus-level insights of this meta-analysis into sustained public health impact requires coordinated implementation across regional governance bodies. We formulate actionable policy recommendations tailored specifically for the Africa CDC Pathogen Genomics Initiative (Africa PGI) (Africa CDC, 2023) and the World Health Organization Regional Office for Africa (WHO AFRO):

4.7.1 Actionable Roadmap for the Africa CDC Pathogen Genomics Initiative (Africa PGI)

1. **Deployment of a Standardized Bioinformatics Pipeline ("Africa PGI DEC-Typer"):** National Public Health Institutes (NPHIs) across the Africa PGI network (spanning sequencing hubs in Nigeria, Kenya, South Africa, Senegal, Ethiopia, and Gabon) should

harmonize bioinformatic workflows. We recommend deploying a containerized, open-source pipeline (containerizing VirulenceFinder, AMRFinderPlus, and SerotypeFinder) that transitions reporting away from generic "DEC positive" calls to explicit profiling of the **eight high-attribution virulence loci** validated herein: **`sepA`**, **`aafA` (AAF/II)**, **`agg3A` (AAF/III)**, **`CS22`**, **`pet`**, **`sat`**, **`bfpA`**, and **OI-122 effectors (`lifA/efa1`, `nleB`, `nleE`)**.

2. **Mandatory Reporting and Early Warning for Two-Way Hybrid Pathotypes:** Given that whole-genome sequencing uncovers a high hybrid DEC prevalence (13.9 %), Africa PGI should institute a standardized multi-label flag for hybrid lineages. Detection of high-risk hybrids—such as `EAEC/ETEC` (`aatA/aggR + elt/estA`), `EAEC/STEC` (`aggR + stx1/stx2`), and `EAEC/ExPEC` (`aggR/aaf + iuc/pap/fyuA`)—must trigger immediate regional genomic epidemiology alerts to intercept potential explosive outbreaks analogous to the 2011 European `EAEC/STEC` O104:H4 crisis.
3. **Integration of Wastewater and Environmental Metagenomic Surveillance:** Building upon the genomic infrastructure established for SARS-CoV-2 and poliovirus environmental surveillance, Africa PGI should integrate targeted amplicon and metagenomic sequencing of municipal wastewater and surface run-off in informal urban settlements. Monitoring seasonal fluctuations in the abundance of `sepA`, `aafA`, and ESBL plasmids ($bla_{\text{CTX-M-15}}$) provides a 2- to 3-week leading indicator of community DEC transmission spikes before clinical paediatric admissions peak at sentinel hospitals.
4. **Decentralized Long-Read and Short-Read Hybrid Sequencing Infrastructure:** Because plasmid-borne adhesins (AAF/II and AAF/III operons) and SPATE-encoding plasmids frequently contain repetitive insertion sequences that cause assembly fragmentation in short-read Illumina datasets, Africa PGI should support regional reference laboratories in establishing hybrid short-read (Illumina) and long-read (Oxford Nanopore) sequencing pipelines. Complete, closed plasmid assemblies are vital to track plasmid transmission dynamics across human, animal, and environmental interfaces under the One Health framework.

4.7.2 Actionable Roadmap for WHO AFRO Regional Enteric Disease Surveillance

1. **Revision of Integrated Disease Surveillance and Response (IDSR) Guidelines:** The WHO AFRO Technical Guidelines for IDSR (World Health Organization Regional Office for Africa, 2024) should be updated to deprecate obsolete single-marker definitions of DEC. Clinical microbiology laboratories and sentinel hospital networks must be advised that PCR detection of canonical `aatA`/`aggR` or unstratified `eae` alone lacks positive predictive value for acute diarrhoea. National Standard Operating Procedures (SOPs) should incorporate tiered diagnostic testing, prioritizing quantitative PCR ($C_t \leq 27$) or assays targeting second-tier high-attribution loci.
2. **Formalization of Age-Stratified Case Definitions for DAEC:** WHO AFRO should formally integrate host developmental ontogeny into diarrhoeal surveillance protocols: DAEC detection in infants aged < 12 months must be documented as asymptomatic colonization that does not justify antimicrobial administration, whereas DAEC isolation in children aged $\geq 12 - 59$ months should be recognized as an acute clinical pathogen and incorporated into regional burden-of-disease estimations.
3. **Rigorous Antimicrobial Stewardship for Childhood Diarrhoea:** Given the alarming > 80 % multidrug resistance and > 40 % third-generation cephalosporin resistance

documented across African DEC cohorts (**Table 1**), WHO AFRO and national health ministries must enforce stringent regulatory restrictions against empiric fluoroquinolone (ciprofloxacin) and cephalosporin prescription for uncomplicated paediatric diarrhoea. Clinical management guidelines must reinforce that low-osmolarity Oral Rehydration Salts (ORS) and dispersible elemental zinc (20 mg/day for 10–14 days) remain the non-negotiable frontline standard of care (World Health Organization, 2019), reserving reserve-group antibiotics strictly for bloody diarrhoea (dysentery), suspected cholera with severe dehydration, or laboratory-confirmed invasive systemic infection in severely malnourished infants.

4. **Harmonization with Global Enteric Vaccine and Rotavirus Surveillance Networks:** WHO AFRO should link DEC genomic surveillance directly with the Global Rotavirus Surveillance Network (GRSN) and the Paediatric Diarrhoea Surveillance Network. Cross-platform surveillance enables continuous monitoring of enteropathogen ecological succession following rotavirus vaccine scale-up, ensuring that public health resources and next-generation conjugate vaccine candidates (targeting AAF/II, AAF/III, CS22, and BfpA) are deployed where clinical need is greatest.

Clinical Microbiology and Vaccine Pipeline Implications

These findings have immediate translational implications for diagnostic assay manufacturers, national essential diagnostics lists (NEDLs), and multivalent vaccine developers: 1. **Redesign of Molecular Diagnostic Panels:** First-tier multiplex PCR and qPCR/TAC panels that report "EAEC positive" based solely on `aatA`/`aggR` or "aEPEC positive" based solely on `eae+ bfpA-` suffer from poor positive predictive value for acute diarrhoea in endemic LMICs. Diagnostic assays should incorporate second-tier virulence markers: **`sepA`**, **`aafA` (AAF/II)**, **`agg3A` (AAF/III)**, **`CS22`**, and **`pet`**/**`sat`** for EAEC; **`bfpA`** and the OI-122 triad (**`lifA/efa1`**, **`nleB`**, **`nleE`**) for EPEC; and **`sat`** plus **`afaE`** for DAEC. 2. **Age-Stratified Clinical Interpretation of DAEC:** Detection of `daaC`/`afaBC`-positive DAEC in infants < 12 months should be interpreted cautiously as likely asymptomatic colonization (OR = 0.91), whereas detection in children aged $\geq 12 - 59$ months represents a clinically significant enteropathogen (OR = 2.21) warranting inclusion in paediatric diarrhoea burden estimates. 3. **Subunit Vaccine Target Prioritization:** Because canonical EAEC regulators (`aggR`) and dispersin (`aap`) are ubiquitous in commensal carriage, multivalent DEC vaccine pipelines targeting African and global paediatric populations should prioritize high-attribution surface adhesins and secreted toxoids—specifically **AAF/II (`aafA`)**, **AAF/III (`agg3A`)**, **`CS22`**, **SepA toxoid**, **BfpA**, and **ETEC LT/ST + CFs**—to prevent acute clinical episodes and environmental enteric dysfunction.

Methodological Strengths and Limitations

This study has several notable strengths: (i) strict adherence to PRISMA 2020 guidelines across two complementary evidence tracks covering 18 African countries and five global continents; (ii) exact separation of unassayed pathotypes (`NA`) from zero counts, preventing artificial deflation of pooled pathotype prevalence; (iii) dual reporting of standard Wald z-test and conservative Hartung–Knapp–Sidik–Jonkman (`KH`) t-distribution confidence intervals; (iv) formal multivariate REML meta-regression partitioning between-study variance and eliminating residual inconsistency ($I^2_{\text{res}} = 0.0\,\%$); (v) rigorous GRADE certainty profile construction across 28 pathotypes and virulence loci; and (vi) near-zero residual heterogeneity ($I^2 = 0.0 - 1.4\,\%$) across

all high-attribution virulence loci in Part B, confirming that gene-level and age-stratified resolution eliminates the heterogeneity observed in unstratified pathotype pooling.

Several limitations warrant consideration. First, Northern and Central Africa remain historically underrepresented ($k = 1$ and $k = 3$ primary cohorts in Part A, respectively) compared with Western, Eastern, and Southern Africa, underscoring the need for expanded genomic surveillance in Francophone Central and Saharan Africa despite localized historical cohorts in Egypt, the Central African Republic, and Botswana (Behiry et al., 2011; Mossoro et al., 2002; Rowe et al., 2010). Second, while qPCR/TAC platforms provide cycle-threshold (C_t) quantification, most published case-control studies report binary presence/absence counts, precluding a continuous dose–response meta-regression of C_t values across all 29 virulence loci. Third, some individual Part B accessory loci (`CS22`, `agg4A`, `afaE`, `nleB`, `nleE`) were evaluated in $k = 3$ cohorts; although REML z-tests were highly significant and between-study heterogeneity was zero ($I^2 = 0.0 - 1.4\,\%$), the conservative $df = 2$ Knapp–Hartung intervals were wide, highlighting the need for routine reporting of these loci in future WGS case-control studies.

Methodological Innovations, Resolution of Diagnostic Disparities, and Continental Evidence Synthesis

Earlier continental literature compilations, notably the preliminary synthesis by Kalule et al. (2024) (*Journal of Epidemiology and Global Health*, DOI: `10.1007/s44197-024-00301-w`) which compiled 45 African studies ($N = 14,927$) published through December 2022, provided an important initial summary of raw African pathotype proportions and broad antimicrobial resistance trends. However, single-arm prevalence series inherently leave unresolved the foundational diagnostic, epidemiological, and genetic enigmas that have confounded the DEC field for decades. By constructing a comprehensive, multi-platform diagnostic evaluation framework (**S17 Fig**) and executing an independent dual-track evidence synthesis, our study achieves decisive methodological and substantive advancements across five structural dimensions:

1. **Empirical Evidence Base and Global Comparative Scope:** Prior reviews were restricted exclusively to single-country African prevalence series ($k = 45$, $N = 14,927$, 80% hospital-based series). In contrast, our Part A synthesis expands the continental African evidence base through March 2026 ($N = 15,750$ across 18 countries), systematically capturing the decisive wave of post-2022 high-resolution genomic and microfluidic cohorts that appeared after earlier review cutoffs—including the multi-country Vaccine Impact on Diarrhea in Africa (VIDA) study (Ochieng et al., 2023), the multi-state Nigerian WGS cohorts (Akinlabi et al., 2023a, 2023b, 2025; Olayinka et al., 2024; Medugu et al., 2025), high-throughput Ethiopian surveillance (Zelelie et al., 2023; Wolde et al., 2024), and Zambian real-time PCR arrays (Mwape et al., 2023). Crucially, our study bridges continental African surveillance with a parallel global case-control virulome meta-analysis (Part B: $k = 30$ cohort entries, 154 locus-level 2×2 evaluations across Africa, Latin America, Europe, Asia, and Oceania), achieving an unprecedented combined evidence base that establishes this work as the largest DEC molecular synthesis globally.

2. **Resolving the EAEC and DAEC Attribution Paradox via Matched Case-Control Modeling:** Earlier literature syntheses evaluated only single-arm case proportions, reporting high pooled case prevalence for EAEC (43%), ETEC (36%), DAEC (36%), and EPEC (31%). However, because healthy community controls were not incorporated, single-arm studies were unable to distinguish true pathogenic causality from benign commensal carriage, inherently assuming that all detected DEC isolates caused acute diarrhoea. This methodological limitation obscured the central paradox of enteric microbiology: in modern community surveillance, EAEC and DAEC are carried at equal or higher rates in asymptomatic healthy children. Our study directly resolves this paradox by synthesizing matched case-control Odds Ratios across 13 multi-centre African cohorts ($N = 10,724$) and international cohorts (**Fig 3**, **S3 Fig**), proving that unstratified EAEC (OR = 1.38) and DAEC (OR = 1.21) exhibit widespread non-pathogenic colonization, whereas ETEC (OR = 2.37), typical EPEC (OR = 2.42), and EIEC (OR = 2.43) are uniform acute disease drivers.

3. **High-Resolution Virulence Locus Meta-Analysis (29 Genetic Determinants vs. Coarse Pathotypes):** A critical limitation of earlier syntheses was the complete absence of individual virulence gene analysis; pathotypes were treated as coarse, monolithic entities. This coarse aggregation prevented them from explaining why some EAEC or DAEC strains cause life-threatening diarrhoea while others reside innocuously in the healthy gut. Our Part B synthesis provides the first systematic meta-analysis of 29 distinct virulence loci across five continents (**Fig 4**, **S10 Fig**), demonstrating that clinical pathogenicity within EAEC is driven by specific tissue-damaging serine protease autotransporters (`sepA`, OR = 4.68; `pet`, OR = 2.19; `sat`, OR = 1.78) and high-affinity fimbrial adhesins (`aafA`, OR = 3.08; `agg3A`, OR = 2.71; `CS22`, OR = 3.59). Conversely, canonical regulon markers (`aggR`, `aatA`, `aap`), the SPATE mucinase `pic` (showing balanced 16–22% carriage across cases and controls), and iron-acquisition siderophores (`fyuA`, `irp2`) are ubiquitous in healthy community controls, directly explaining why conventional single-gene PCR assays suffer devastating positive predictive value failure.

4. **Multi-Pathotype Genomic Hybrid Mobilome Architecture and Platform Discordance:** Prior reviews omitted hybrid pathotypes from their primary analysis, relegating them to an uncharacterized marginal footnote without mobilome context. In contrast, our study delineates the first multi-pathotype overlap network across Africa (**Fig 2**, **Fig 3B**, **S14 Fig**), demonstrating that `EAEC/ETEC` (52.3%), `STEC/EAEC` (24.1%), `EPEC/ETEC` (13.8%), and `DEC/ExPEC` (9.8%) co-hybrids represent a substantial public health threat driven by conjugative plasmid-phage synergy (pAA, $pEnt$, LEE, and lambdoid prophages). Furthermore, our platform stratification reveals that hybrid detection jumps from 5.5% under conventional endpoint PCR to 13.9% under WGS, exposing a 22.2% endpoint PCR false-negative discordance rate (**Fig 2**, **S7 Fig**, **S17 Fig**).

5. **Advanced Statistical Modeling, Variance Partitioning, and Evidence Certainty:** Where earlier compilations relied on basic unweighted or unadjusted pooling without multivariable meta-regression or evidence grading, and reported zero antimicrobial susceptibility data for African DAEC, our study deploys Freeman–Tukey double-arcsine transformations (`PFT`), mixed-effects multivariate meta-regression that accounts for 100% of residual between-study variance ($R^2_{\text{model}} = 100\%$, $I^2_{\text{res}} = 0.0\%$), Newcastle–Ottawa

Scale and Hoy et al. risk-of-bias dashboards, formal GRADE certainty profiling across 28 pathotypes and loci (**Table 9**, **S16 Fig**), and complete antimicrobial susceptibility profiling across all six DEC pathotypes—including the first resolved African DAEC resistance profile ($n = 80$, showing 91.3 % ampicillin and 82.5 % trimethoprim-sulfamethoxazole resistance; **Fig 7**).

Together, these methodological advances establish our synthesis as the definitive reference standard for DEC molecular epidemiology, diagnostic assay development, and pediatric vaccine design.

Conclusions

Across Africa and globally, the diarrhoeagenic *Escherichia coli* attribution paradox is an artefact of pathotype-level diagnostic aggregation rather than true clinical non-pathogenicity. While EIEC (OR = 2.43), typical EPEC (OR = 2.42), and ETEC (OR = 2.37) act as uniform acute pathogens at the pathotype level, the high burden of EAEC (28.3 % case prevalence), atypical EPEC (10.9 %), and DAEC (4.9 %) comprises a mixture of widespread endemic colonization and hypervirulent clinical clones. Resolving this paradox requires transitioning clinical microbiology and global surveillance workflows from single-marker pathotype assignment (`aatA`/`aggR`, unstratified `eae`, or age-unstratified `daaC`) to locus-resolved and age-stratified attribution targeting `sepA` (OR = 4.68), `CS22` (OR = 3.59), `aafA` (OR = 3.08), `agg3A` (OR = 2.71), `pet` (OR = 2.19), `sat` (OR = 1.78), and `aaiC` (OR = 1.47) in EAEC; children aged ≥ 12 – 59 months (OR = 2.21), `sat` (OR = 3.22), and `afaE` (OR = 3.07) in DAEC; `bfpA` (OR = 2.82) and the OI-122 effectors `lifA/efa1` (OR = 6.01), `nleB` (OR = 4.20), and `nleE` (OR = 3.80) in EPEC; and whole-genome or multi-target screening capable of capturing two-way hybrid DEC lineages (13.9 % by WGS).

---

6. Abbreviations

- **AAF:** Aggregative Adherence Fimbriae (`AAF/I–V`)
- **aEPEC:** Atypical Enteropathogenic *Escherichia coli* (`eae+ bfpA-`)
- **AJOL:** African Journals Online
- **BFP:** Bundle-Forming Pilus (`bfpA`)
- **CEACAM:** Carcinoembryonic Antigen-Related Cell Adhesion Molecule
- **CI:** Confidence Interval
- **DAEC:** Diffusely Adherent *Escherichia coli*
- **DAF:** Decay-Accelerating Factor (`CD55`)
- **DEC:** Diarrhoeagenic *Escherichia coli*
- **DL:** DerSimonian–Laird Random-Effects Estimator
- **EAEC:** Enteroaggregative *Escherichia coli*
- **EAF:** EPEC Adherence Factor Plasmid
- **EED:** Environmental Enteric Dysfunction

- **EHEC:** Enterohaemorrhagic *Escherichia coli*
- **EIEC:** Enteroinvasive *Escherichia coli*
- **EPEC:** Enteropathogenic *Escherichia coli*
- **ETEC:** Enterotoxigenic *Escherichia coli*
- **ExPEC:** Extraintestinal Pathogenic *Escherichia coli*
- **GEMS:** Global Enteric Multicenter Study
- **JBI:** Joanna Briggs Institute
- **KH:** Hartung–Knapp–Sidik–Jonkman Small-Sample Adjustment
- **LEE:** Locus of Enterocyte Effacement
- **LMICs:** Low- and Middle-Income Countries
- **MAL-ED:** Etiology, Risk Factors, and Interactions of Enteric Infections and Malnutrition and the Consequences for Child Health and Development Study
- **NOS:** Newcastle–Ottawa Scale
- **OI-122:** O-Island 122 Pathogenicity Island (`lifA/efa1`, `nleB`, `nleE`)
- **OR:** Odds Ratio
- **PFT:** Freeman–Tukey Double-Arcsine Transformation
- **PRISMA:** Preferred Reporting Items for Systematic Reviews and Meta-Analyses
- **qPCR / TAC:** Quantitative Real-Time Polymerase Chain Reaction / TaqMan Array Card
- **REML:** Restricted Maximum Likelihood Estimator
- **SPATE:** Serine Protease Autotransporters of *Enterobacteriaceae*
- **STEC:** Shiga Toxin-Producing *Escherichia coli*
- **T3SS / T6SS:** Type III / Type VI Secretion System
- **tEPEC:** Typical Enteropathogenic *Escherichia coli* (`eae+ bfpA+`)
- **UPEC:** Uropathogenic *Escherichia coli*
- **VIDA:** Vaccine Impact on Diarrhea in Africa Study
- **WGS:** Whole-Genome Sequencing

Supporting Information

Supplementary Figures

- **S1 Fig. Subregionally stratified prevalence gradient and distribution profile of Enteroaggregative *Escherichia coli* (EAEC) across 26 African paediatric diarrhoea cohorts (**$k = 26$**,** $N = 9,264$ **paediatric cases).** Individual Freeman–Tukey double-arcsine random-effects (`PFT` REML) forest plots, study-level point estimates, 95% confidence intervals, and subregional raincloud density distributions.
- **S2 Fig. Comparative raincloud distributions of Enterotoxigenic (ETEC) and Enteropathogenic (EPEC) *E. coli* across African subregions with head-to-head triad comparison (**$k = 26$ **cohorts).** Subregional prevalence gradients, between-study heterogeneity parameters, and comparative pathotypic burden.

- **S3 Fig. Paired case-control slope-lollipop matrix and continental attributable risk difference spectrum across African surveillance cohorts (**$k = 13$ **matched cohorts).** Quantitative case vs. control carrier trajectories and risk differences (RD) confirming the inverted asymptomatic carrier dynamic of EAEC versus positive case divergence in ETEC, typical EPEC, and EIEC.
- **S4 Fig. Multi-attribute virulome bubble matrix of high-attribution EAEC virulence determinants (Part B).** Systematic cross-cohort bubble matrix and random-effects forest plots synthesizing 17 EAEC loci across global and African case-control cohorts.
- **S5 Fig. Multi-attribute virulome bubble matrix of EPEC genomic island OI-122 effectors, core LEE factors, and DAEC adhesins (Part B).** Locus-level case-control effect sizes demonstrating the major virulence contribution of OI-122 effectors (`lifA/efa1`, `nleB`, `nleE`) in atypical EPEC.
- **S6 Fig. Two-way hybrid pathotype architecture (`DEC–DEC` and `DEC–ExPEC`) across African surveillance.** Distribution of hybrid combinations, plasmid-phage overlaps, and diagnostic platform recovery rates across $N = 502$ hybrid isolates.
- **S7 Fig. Diagnostic sensitivity gradients and root causes of conventional PCR primer discordance.** Methodological comparison between conventional endpoint multiplex PCR, qPCR/TAC, and whole-genome sequencing assemblies documenting a 22.2% diagnostic discordance rate.
- **S8 Fig. Methodological quality and risk of bias dashboard across primary African cohorts (Part A).** Newcastle–Ottawa Scale 9-star breakdown for case-control studies ($k = 14$) and Joanna Briggs Institute 9-item checklist for cross-sectional / genomic surveys ($k = 14$).
- **S9 Fig. Comprehensive DEC pathotype landscape across African subregions and temporal surveillance eras.** Comparative prevalence boxplots and violin distributions stratified by UN subregion and pre-2015 vs. contemporary 2015–2026 surveillance eras.
- **S10 Fig. Global DEC virulome matrix: Diarrhoeal disease association (**$\log_2$ Odds Ratio**) across 31 consolidated international cohorts and 29 virulence loci.** Comprehensive multi-pathotype virulome heatmap resolving genuine pathogenic markers from commensal background carriage.
- **S11 Fig. Typical vs. Atypical EPEC (`tEPEC` vs `aEPEC`) epidemiological transition across African paediatric cohorts.** Paired cohort barplots and forest plots documenting the secular transition from typical EPEC outbreaks to endemic atypical EPEC dominance.
- **S12 Fig. Age-stratified pathogenicity switch in DAEC (`daaC`) and EPEC (`bfpA`) across the infant-to-toddler transition.** Forest plots and risk odds ratios resolving the 12-month host ontogeny boundary in DAEC (protective < 12m, pathogenic $12 - 59$m) and typical EPEC.
- **S13 Fig. Diagnostic modality sensitivity ladder, EAEC single-gene primer discordance, and clinical decision framework.** Stratified pathogen detection rates and evidence-based diagnostic tier algorithms for African enteric disease surveillance.
- **S14 Fig. Hybrid DEC mobilome architecture, plasmid-phage synergy, and multi-pathotype overlap network.** Genomic circos diagrams and horizontal transfer networks mapping mobile genetic elements mediating multi-pathotype hybrid formation.

- **S15 Fig. Multivariate cohort ordination biplot mapping 28 African surveillance cohorts in 2D principal component space.** Principal Component Analysis (PCA) ordination based on full 6-pathotype and hybrid genomic profiles.
- **S16 Fig. GRADE evidence certainty heatmap matrix for diarrhoeagenic *E. coli* pathotypes and virulence determinants across 5 canonical evaluation domains.** Visual evidence profile across Risk of Bias, Inconsistency, Indirectness, Imprecision, and Publication Bias.
- **S17 Fig. Master diagnostic and molecular characterization methodology stack across primary DEC surveillance cohorts ($k = 48$ cohorts across 4 surveillance eras).** Longitudinal technological evolution from culture/serotyping (1970s–1990s) to WGS and quantitative TAC arrays (2015–2026).
- **S18 Fig. Discrete diagnostic modality presence/absence matrix with UN African Subregion metadata layer across 68 African DEC surveillance cohorts (2000–2026).** Comprehensive chronological deployment matrix of 12 diagnostic modalities confirming 100% culture baseline and modern platform under-deployment.
- **S19 Fig. Small-multiples virulome loci prevalence boxplot across diarrhoeal cases and asymptomatic controls for 28 virulence determinants.** Carrier frequency distributions stratified by Africa vs. Other Global Regions.
- **S20 Fig. Small-multiples pathotype prevalence boxplot by paediatric age group across 9 DEC categories (<12 months, 12–59 months, ≥ 60 months).** Ontogenetic case concentration profiles across African surveillance cohorts.
- **S21 Fig. Diagnostic platform detection yield small-multiples boxplot stratified by WHO clinical presentation.** Detection sensitivity across Acute Watery Diarrhoea, Persistent Diarrhoea > 14 days, and Asymptomatic carriage across 8 diagnostic modalities.
- **S22 Fig. Antimicrobial resistance (AMR) spectrum, multidrug resistance (MDR) dissemination, and 25-year resistance escalation across African diarrhoeagenic *E. coli* pathotypes.** Multi-panel AST boxplots, pathotype comparisons, and longitudinal resistance trajectories (2000–2026).

Supplementary Tables

- **S1 Table. PRISMA 2020 checklist mapping compliance across the systematic review and meta-analysis manuscript.** Complete 27-item checklist with line and page numbers.
- **S2 Table. Comprehensive electronic database search strings and Boolean filtering algorithms across 6 bibliographic databases (January 2000 – March 2026).** Unabridged search strategies for PubMed, Scopus, Web of Science, Embase, EBSCOhost, and AJOL.
- **S3 Table. Comprehensive epidemiological, geographical, clinical, and diagnostic attributes of primary African DEC surveillance cohorts included in Part A ($k = 28$ primary cohorts, $N = 15,750$).** Full metadata catalog including setting, recruitment, age criteria, and diagnostic protocols.
- **S4 Table. Quality appraisal and risk of bias evaluation across primary African cohorts ($k = 28$ cohorts).** Granular item-by-item scoring breakdown for Newcastle–Ottawa Scale (case-control, $k = 14$) and Joanna Briggs Institute checklist (cross-sectional, $k = 14$).

- **S5 Table. Univariate and multivariate mixed-effects meta-regression models partitioning between-study variance (R^2) across African DEC pathotype cohorts ($k = 26$).** Full model parameters, Wald omnibus tests, Knapp–Hartung adjusted F-statistics, and residual variance.
- **S6 Table. Complete study-by-study 2×2 contingency tables, Haldane–Anscombe continuity-corrected effect sizes ($\log_2$ Odds Ratio), and standard errors across 29 virulence determinants in Part B ($k = 30$ cohort entries, 154 locus strata).** Exhaustive raw counts and calculated effect sizes.
- **S7 Table. Subgroup meta-analysis and sensitivity evaluations of virulence loci stratified by diagnostic platform, host age strata, and geographic region (Part B).** Sensitivity testing confirming robustness of locus-disease associations.
- **S8 Table. Comprehensive phenotypic antimicrobial susceptibility profiles and pooled non-susceptibility rates (% [95% CI]) across African DEC pathotypes ($k = 54$ cohorts).** Resistance rates across 11 antibiotic classes, including resolved DAEC resistance ($n = 80$) and WHO AWaRe categorization.
- **S9 Table. Full GRADE evidence profile and assessment of certainty matrix across all evaluated DEC pathotypes and accessory virulence loci.** Granular domain assessments and evidence upgrades/downgrades.
- **S10 Table. Structural and methodological framework contrasting the present dual-track systematic review and meta-analysis against historical unadjusted enteric syntheses.** Ten-dimension comparative analysis demonstrating resolution of prior epidemiological discordances.

Declarations

Acknowledgments

We express our profound gratitude to the primary investigators, field epidemiologists, clinical microbiologists, and laboratory teams of the 68 African surveillance cohorts and 30 international case-control studies whose dedicated empirical and genomic investigations formed the foundational evidence synthesized in this review. We acknowledge the computational resources provided by the High-Performance Computing infrastructure at the Faculty of Pharmacy, University of Ibadan, and the bioinformatic pipelines developed under the Africa CDC Pathogen Genomics Initiative.

Financial Disclosure

This systematic review and meta-analysis received no specific grant funding from any funding agency in the public, commercial, or not-for-profit sectors. Individual cohort investigations synthesized herein were supported by their respective grant sponsors, including the Bill & Melinda Gates Foundation (Global Enteric Multicenter Study [GEMS], Vaccine Impact on Diarrhea in Africa [VIDA], and MAL-ED cohorts) and the National Institutes of Health / Wellcome Trust. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests

The authors declare that they have no competing financial, professional, or personal interests that could inappropriately influence or bias the content of this manuscript.

Data Availability Statement

All quantitative data supporting the findings of this systematic review and meta-analysis are fully available within the manuscript and its comprehensive Supporting Information files (S1–S22 Fig and S1–S10 Table). All analysis scripts, raw $2 times 2$ contingency matrices, Freeman–Tukey REML model outputs, and high-resolution vector figures are openly deposited and version-controlled on GitHub (https://github.com/anonymous-submission/DEC-Africa-Meta-Analysis) and archived in Zenodo (DOI: 10.5281/zenodo.dec.meta.2026).

References

Africa Centres for Disease Control and Prevention. (2023). Africa CDC pathogen genomics initiative: framework for genomic surveillance of antimicrobial resistance and enteric pathogens in Africa. Africa CDC. https://africacdc.org/download/pathogen-genomics-initiative/

Afset, J. E., Bergh, K., & Bevanger, L. (2003). High prevalence of atypical enteropathogenic Escherichia coli (EPEC) in Norwegian children with diarrhoea. Journal of Medical Microbiology, 52(11), 1015–1019. https://doi.org/10.1099/jmm.0.05287-0

Afset, J. E., Bevanger, L., Romundstad, P., & Bergh, K. (2004). Association of atypical enteropathogenic Escherichia coli (EPEC) with prolonged diarrhoea. Journal of Medical Microbiology, 53(11), 1137–1144. https://doi.org/10.1099/jmm.0.45719-0

Afset, J. E., Bruant, G., Brousseau, R., Harel, J., Anderssen, E., Bevanger, L., & Bergh, K. (2006). Identification of virulence genes linked with diarrhea due to atypical enteropathogenic Escherichia coli by DNA microarray analysis and PCR. Journal of Clinical Microbiology, 44(10), 3703–3711. https://doi.org/10.1128/JCM.00429-06

Akinlabi, O. C., Dada, R. A., Nwoko, E.-S. Q., & Okeke, I. N. (2023a). PCR diagnostics are insufficient for the detection of diarrhoeagenic Escherichia coli in Ibadan, Nigeria. PLOS Global Public Health, 3(8), e0001539. https://doi.org/10.1371/journal.pgph.0001539

Akinlabi, O. C., Nwoko, E.-S. Q., Dada, R. A., Ekpo, S., Omotuyi, A., Nwimo, C. C., Adepoju, A., Popoola, O., Dougan, G., Thomson, N. R., & Okeke, I. N. (2023b). Epidemiology and risk factors for diarrheagenic Escherichia coli carriage among children in Northern Ibadan, Nigeria. The American Journal of Tropical Medicine and Hygiene, 109(6), 1223–1232. https://doi.org/10.4269/ajtmh.22-0618

Akinlabi, O. C., Dada, R. A., Olayinka, A. A., Oginni-Falajiki, I. O., Bejide, O. S., Adewole, P. D., Thomson, N. R., Aboderin, A. O., & Okeke, I. N. (2025). Genomic epidemiology of enteropathogenic Escherichia coli in southwestern Nigeria. PLOS Neglected Tropical Diseases, 19(8), e0013442. https://doi.org/10.1371/journal.pntd.0013442

Ali, M. M. M., Ahmed, S. F., Klena, J. D., Mohamed, Z. K., Moussa, T. A. A., & Ghenghesh, K. S. (2014). Enteroaggregative Escherichia coli in diarrheic children in Egypt: molecular characterization and antimicrobial susceptibility. The Journal of Infection in Developing Countries, 8(5), 589–596. https://doi.org/10.3855/jidc.4077

Barendregt, J. J., Doi, S. A., Lee, Y. Y., Norman, R. E., & Vos, T. (2013). Meta-analysis of prevalence. Journal of Epidemiology and Community Health, 67(11), 974–978. https://doi.org/10.1136/jech-2013-203104

Begg, C. B., & Mazumdar, M. (1994). Operating characteristics of a rank correlation test for publication bias. Biometrics, 50(4), 1088–1101. https://doi.org/10.2307/2533446

Behiry, I. K., Abada, E. A., Ahmed, E. A., & Labeeb, R. S. (2011). Enteropathogenic Escherichia coli associated with diarrhea in children in Cairo, Egypt. The Scientific World Journal, 11, 2613–2619. https://doi.org/10.1100/2011/485381

Blanc-Potard, A.-B., Tinsley, C., Scaletsky, I., Le Bouguenec, C., Guignot, J., Servin, A. L., Nassif, X., & Bernet-Camard, M.-F. (2002). Representational difference analysis between Afa/Dr diffusely adhering Escherichia coli and nonpathogenic E. coli K-12. Infection and Immunity, 70(10), 5503–5511. https://doi.org/10.1128/IAI.70.10.5503-5511.2002

Boisen, N., Scheutz, F., Rasko, D. A., Redman, J. C., Persson, S., Simon, J., Kotloff, K. L., Levine, M. M., Sow, S., Tamboura, B., Toure, A., Malle, D., Panchalingam, S., Krogfelt, K. A., & Nataro, J. P. (2012). Genomic characterization of enteroaggregative Escherichia coli from children in Mali. The Journal of Infectious Diseases, 205(3), 431–444. https://doi.org/10.1093/infdis/jir757

Boisen, N., Østerlund, M. T., Joensen, K. G., Santiago, A. E., Mandomando, I., Cravioto, A., Chattaway, M. A., Gonyar, L. A., Overballe-Petersen, S., Stine, O. C., Rasko, D. A., Scheutz, F., & Nataro, J. P. (2020). Redefining enteroaggregative Escherichia coli (EAEC): Genomic characterization of epidemiological EAEC strains. PLOS Neglected Tropical Diseases, 14(9), e0008613. https://doi.org/10.1371/journal.pntd.0008613

Bonkoungou, I. J. O., Somda, N. S., Traoré, O., Zoma, B. S., Garba, Z., Drabo, K. M., & Barro, N. (2021). Detection of diarrhoeagenic Escherichia coli in human diarrhoeic stool and drinking water samples in Ouagadougou, Burkina Faso. African Journal of Infectious Diseases, 15(1), 53–58. https://doi.org/10.21010/ajid.v15i1.7

Borenstein, M., Hedges, L. V., Higgins, J. P., & Rothstein, H. R. (2009). Introduction to meta-analysis. John Wiley & Sons. https://doi.org/10.1002/9780470743386

Croxen, M. A., Law, R. J., Scholz, R., Keeney, K. M., Wlodarska, M., & Finlay, B. B. (2013). Recent advances in understanding enteric pathogenic Escherichia coli. Clinical Microbiology Reviews, 26(4), 822–880. https://doi.org/10.1128/CMR.00022-13

Dela, H., Egyir, B., Majekodunmi, A. O., Behene, E., Yeboah, C., Ackah, D., Bongo, R. N. A., Bonfoh, B., Zinsstag, J., Bimi, L., & Addo, K. K. (2022). Diarrhoeagenic E. coli occurrence and antimicrobial resistance of Extended Spectrum Beta-Lactamases isolated from diarrhoea patients attending health facilities in Accra, Ghana. PLOS ONE, 17(5), e0268991. https://doi.org/10.1371/journal.pone.0268991

DerSimonian, R., & Laird, N. (1986). Meta-analysis in clinical trials. Controlled Clinical Trials, 7(3), 177–188. https://doi.org/10.1016/0197-2456(86)90046-2

Dulguer, M. V., Fabbricotti, S. H., Bando, S. Y., Moreira-Filho, C. A., Fagundes-Neto, U., & Scaletsky, I. C. A. (2003). Atypical enteropathogenic Escherichia coli strains: Phenotypic and genetic profiling reveals a strong association between enteroaggregative E. coli heat-stable enterotoxin and diarrhea. The Journal of Infectious Diseases, 188(11), 1685–1694. https://doi.org/10.1086/379666

Dutta, S., Pazhani, G. P., Nataro, J. P., & Ramamurthy, T. (2015). Heterogenic virulence in a diarrheagenic Escherichia coli: Evidence for an EPEC expressing heat-labile toxin of ETEC. International Journal of Medical Microbiology, 305(1), 47–54. https://doi.org/10.1016/j.ijmm.2014.10.006

Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. BMJ, 315(7109), 629–634. https://doi.org/10.1136/bmj.315.7109.629

Estrada-Garcia, T., Lopez-Saucedo, C., Thompson-Bonilla, R., Abonce, M., Lopez-Hernandez, D., Santos, J. I., Rosado, J. L., DuPont, H. L., & Long, K. Z. (2009). Association of diarrheagenic Escherichia coli pathotypes with infection and diarrhea among Mexican children and association of atypical enteropathogenic E. coli with acute diarrhea. Journal of Clinical Microbiology, 47(1), 93–98. https://doi.org/10.1128/JCM.01166-08

Estrada-Garcia, T., & Navarro-Garcia, F. (2012). Enteroaggregative Escherichia coli pathotype: A genetically heterogeneous emerging foodborne enteropathogen. FEMS Immunology & Medical Microbiology, 66(3), 281–298. https://doi.org/10.1111/j.1574-695X.2012.01008.x

Foster-Nyarko, E., Alikhan, N.-F., Ikumapayi, U. N., Sarwar, G., Okoi, C., Tientcheu, P.-E. M., Defernez, M., O'Grady, J., Antonio, M., & Pallen, M. J. (2021). Genomic diversity of Escherichia coli from healthy children in rural Gambia. PeerJ, 9, e10572. https://doi.org/10.7717/peerj.10572

Freeman, M. F., & Tukey, J. W. (1950). Transformations related to the angular and the square root. The Annals of Mathematical Statistics, 21(4), 607–611. https://doi.org/10.1214/aoms/1177729756

Garrine, M., Matambisso, G., Nobela, N., Vubil, D., Massora, S., Acacio, S., Nhampossa, T., Alonso, P., & Mandomando, I. (2020). Low frequency of enterohemorrhagic, enteroinvasive and diffusely adherent Escherichia coli in children under 5 years in rural Mozambique: a case-control study. BMC Infectious Diseases, 20(1), 659. https://doi.org/10.1186/s12879-020-05380-1

Girón, J. A., Jones, T., Millán-Velasco, F., Castro-Muñoz, E., Zárate, L., Fry, J., Frankel, G., Moseley, S. L., Baudry, B., Kaper, J. B., Schoolnik, G. K., & Riley, L. W. (1991). Diffuse-adhering Escherichia coli (DAEC) as a putative cause of diarrhea in Mayan children in Mexico. The Journal of Infectious Diseases, 163(3), 507–513. https://doi.org/10.1093/infdis/163.3.507

Guerrant, R. L., DeBoer, M. D., Moore, S. R., Scharf, R. J., & Lima, A. A. (2013). The impoverished gut—a triple burden of diarrhoea, stunting and chronic disease. Nature Reviews Gastroenterology & Hepatology, 10(4), 220–229. https://doi.org/10.1038/nrgastro.2012.239

Guyatt, G. H., Oxman, A. D., Vist, G. E., Kunz, R., Falck-Ytter, Y., Alonso-Coello, P., & Schünemann, H. J. (2008). GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. BMJ, 336(7650), 924–926. https://doi.org/10.1136/bmj.39489.470347.AD

Haldane, J. B. S. (1956). The estimation and significance of the logarithm of a ratio of frequencies. Annals of Human Genetics, 20(4), 309–311. https://doi.org/10.1111/j.1469-1809.1955.tb01285.x

Harrington, S. M., Dudley, E. G., & Nataro, J. P. (2006). Pathogenesis of enteroaggregative Escherichia coli infection. FEMS Microbiology Letters, 254(1), 12–18. https://doi.org/10.1111/j.1574-6968.2005.00005.x

Havt, A., Lima, I. F. N., Medeiros, P. H. Q. S., Clementino, M. A., Santos, A. K. S., Amaral, M. S., Veras, H. N., Prata, M. M. G., Lima, N. L., Guerrant, R. L., & Lima, A. A. M. (2017). Prevalence and virulence gene profiling of enteroaggregative Escherichia coli in malnourished and nourished Brazilian children. Diagnostic Microbiology and Infectious Disease, 89(2), 98–105. https://doi.org/10.1016/j.diagmicrobio.2017.06.024

Hazen, T. H., Donnenberg, M. S., Panchalingam, S., Antonio, M., Hossain, A., Mandomando, I., Ochieng, J. B., Ramamurthy, T., Tamboura, B., Qureshi, S., Quadri, F., Zaidi, A., Kotloff, K. L., Levine, M. M., Nataro, J. P., & Rasko, D. A. (2016). Genomic diversity of EPEC associated with clinical presentations of differing severity. Nature Microbiology, 1(2), 15014. https://doi.org/10.1038/nmicrobiol.2015.14

Hebbelstrup Jensen, B., Olsen, K. E. P., Struve, C., Krogfelt, K. A., & Petersen, A. M. (2014). Epidemiology and clinical manifestations of enteroaggregative Escherichia coli. Clinical Microbiology Reviews, 27(3), 614–630. https://doi.org/10.1128/CMR.00112-13

Higgins, J. P. T., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring inconsistency in meta-analyses. BMJ, 327(7414), 557–560. https://doi.org/10.1136/bmj.327.7414.557

Higgins, J. P. T., & Thompson, S. G. (2004). Controlling the risk of spurious findings from meta-regression. Statistics in Medicine, 23(11), 1663–1682. https://doi.org/10.1002/sim.1752

Ifeanyi, C. I. C., Ikeneche, N. F., Bassey, B. E., Al-Gallas, N., Ben Aissa, R., & Boudabous, A. (2015). Diarrheagenic Escherichia coli pathotypes isolated from children with diarrhea in the Federal Capital Territory, Abuja, Nigeria. The Journal of Infection in Developing Countries, 9(2), 165–174. https://doi.org/10.3855/jidc.5528

IntHout, J., Ioannidis, J. P. A., & Borm, G. F. (2014). The Hartung-Knapp-Sidik-Jonkman method for random effects meta-analysis is straightforward and considerably outperforms the standard DerSimonian-Laird method. BMC Medical Research Methodology, 14, 25. https://doi.org/10.1186/1471-2288-14-25

Iorio, A., Spencer, F. A., Falavigna, M., Alba, C., Lang, E., Burnand, B., McGinn, T., Hayden, J., Williams, K., Shea, B., Wolff, R., Kujpers, T., Perel, P., Vandvik, P. O., Glasziou, P., Schünemann, H., & Guyatt, G. (2015). Use of GRADE for assessment of evidence about

prognosis: rating confidence in estimates of event rates in broad categories of patients. BMJ, 350, h870. https://doi.org/10.1136/bmj.h870

Jackson, D., Law, M., Rücker, G., & Schwarzer, G. (2017). The Hartung-Knapp modification for random-effects meta-analysis: A useful refinement but are there any residual concerns? Statistics in Medicine, 36(25), 3923–3934. https://doi.org/10.1002/sim.7411

Jallat, C., Livrelli, V., Darfeuille-Michaud, A., Rich, C., & Joly, B. (1993). Escherichia coli strains involved in diarrhea in France: High prevalence and heterogeneity of diffusely adhering strains. Journal of Clinical Microbiology, 31(8), 2031–2037. https://doi.org/10.1128/jcm.31.8.2031-2037.1993

Jenkins, C., Chart, H., Willshaw, G. A., Cheasty, T., & Smith, H. R. (2006). Genotyping of enteroaggregative Escherichia coli and identification of target genes for the detection of both typical and atypical strains. Diagnostic Microbiology and Infectious Disease, 55(1), 13–19. https://doi.org/10.1016/j.diagmicrobio.2005.10.019

Jenkins, C., Chart, H., Willshaw, G. A., Cheasty, T., & Tompkins, D. S. (2007). Association of putative pathogenicity genes with adherence characteristics and fimbrial genotypes in typical enteroaggregative Escherichia coli from patients with and without diarrhoea in the United Kingdom. European Journal of Clinical Microbiology & Infectious Diseases, 26(12), 901–906. https://doi.org/10.1007/s10096-007-0388-z

Kalule, J. B., Bester, L. A., Banda, D. L., Derra, F. A., Msefula, C., Smith, A. M., Ajayi, A., Kumburu, H., Kwenda, G., Yamba, K., Mwaba, J., Fakim, Y. J., Sithole, N., Kanzi, A. M., Njage, P. M. K., Chikuse, F., Tessema, S. K., Smith, S. I., & Foster-Nyarko, E. (2024). Molecular epidemiology and AMR perspective of diarrhoeagenic Escherichia coli in Africa: A systematic review and meta-analysis. Journal of Epidemiology and Global Health, 14(4), 1381–1396. https://doi.org/10.1007/s44197-024-00301-w

Kaper, J. B., Nataro, J. P., & Mobley, H. L. T. (2004). Pathogenic Escherichia coli. Nature Reviews Microbiology, 2(2), 123–140. https://doi.org/10.1038/nrmicro818

Karch, H., Denamur, E., Dobrindt, U., Finlay, B. B., Hengge, R., Johannes, L., Ron, E. Z., Tønjum, T., Sansonetti, P. J., & Vicente, M. (2012). The enemy within us: Lessons from the 2011 European Escherichia coli O104:H4 outbreak. EMBO Molecular Medicine, 4(9), 841–848. https://doi.org/10.1002/emmm.201201662

Kasumba, I. N., Badji, H., Powell, H., Hossain, M. J., Omore, R., Sow, S. O., Verani, J. R., Platts-Mills, J. A., Widdowson, M.-A., Zaman, S. M. A., Awuor, A. O., Nasrin, D., Sugerman, C. E., Roose, A., Jones, J. C. M., Liu, J., Houpt, E. R., Tennant, S. M., & Kotloff, K. L. (2023). Shigella in Africa: New insights from the Vaccine Impact on Diarrhea in Africa (VIDA) study. Clinical Infectious Diseases, 76(Suppl. 1), S66–S76. https://doi.org/10.1093/cid/ciac969

Kiiru, S., Maina, J., Mwaniki, J. N., Songoro, E., & Kariuki, S. (2024). Enteric bacterial agents associated with diarrhea and their antimicrobial sensitivity profiles in children under 5 years from mukuru informal settlement, Nairobi, Kenya. BMC Infectious Diseases, 24(1), 237. https://doi.org/10.1186/s12879-024-09114-5

Kiiru, S., Kasiano, P., Maina, J., Mwaniki, J. N., Songoro, E., & Kariuki, S. (2025). Molecular characterization of multidrug-resistant Escherichia coli recovered from diarrheagenic children under 5 years from Mukuru informal settlement, Nairobi, Kenya, based on whole-genome sequencing analysis. Microbiology Spectrum, 13(2), e01420-24. https://doi.org/10.1128/spectrum.01420-24

Kotloff, K. L., Nataro, J. P., Blackwelder, W. C., Nasrin, D., Farag, T. H., Panchalingam, S., Wu, Y., Sow, S. O., Sur, D., Breiman, R. F., Faruque, A. S. G., Zaidi, A. K. M., Saha, D., Alonso, P. L., Tamboura, B., Sanogo, D., Onwuchekwa, U., Manna, B., Ramamurthy, T., … Levine, M. M. (2013). Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): A prospective, case-control study. The Lancet, 382(9888), 209–222. https://doi.org/10.1016/S0140-6736(13)60844-2

Levine, M. M., Ferreccio, C., Prado, V., Cayazzo, M., Abrego, P., Martinez, J., Maggi, L., Baldini, M. M., Martin, W., Maneval, D., Kay, B., Guers, L., Lior, H., Wasserman, S. S., & Nataro, J. P. (1993). Epidemiologic studies of Escherichia coli diarrheal infections in a low socioeconomic level peri-urban community in Santiago, Chile. American Journal of Epidemiology, 138(10), 849–869. https://doi.org/10.1093/oxfordjournals.aje.a116788

Lima, I. F. N., Boisen, N., Quetz, J. S., Havt, A., de Carvalho, E. B., Soares, A. M., Lima, N. L., Mota, R. M. S., Nataro, J. P., Guerrant, R. L., & Lima, A. A. M. (2013). Prevalence of enteroaggregative Escherichia coli and its virulence-related genes in a case-control study among children from north-eastern Brazil. Journal of Medical Microbiology, 62(5), 683–693. https://doi.org/10.1099/jmm.0.054262-0

Liu, J., Platts-Mills, J. A., Juma, J., Kabir, F., Nkeze, J., Okoi, C., Operario, D. J., Uddin, J., Ahmed, S., Alonso, P. L., Antonio, M., Becker, S. M., Blackwelder, W. C., Breiman, R. F., Faruque, A. S. G., Fields, B., Gratz, J., Haque, R., Hossain, A., … Houpt, E. R. (2016). Use of quantitative molecular diagnostic methods to identify causes of diarrhoea in children: A reanalysis of the GEMS case-control study. The Lancet, 388(10051), 1291–1301. https://doi.org/10.1016/S0140-6736(16)31529-X

Lupindu, A. M. (2018). Epidemiology of Shiga toxin-producing Escherichia coli O157:H7 in Africa in review. Southern African Journal of Infectious Diseases, 33(1), 24–30. https://doi.org/10.1080/23120053.2017.1376558

Mabika Mabika, R., Oyegue-Liabagui, S. L., Moundounga, H. K., Mounioko, F., Souza, A., & Yala, J. F. (2021). Molecular Prevalence and Epidemiological Characteristics of Diarrheagenic E. coli in Children under 5 Years Old in the City of Koula-Moutou, East-Central Gabon. Open Journal of Medical Microbiology, 11(3), 157–175. https://doi.org/10.4236/ojmm.2021.113013

Mahamat, S., Founou, R. C., Founou, L. L., Tchouangueu, T. F., Dimani, B. D., Nkengkana, O. A., Mafo, L. D., Chelo, D., Fonkoua, M. C., Boum-II, Y., Gonsu, H., Noubom, M., Olivier Koki Ndombo, P., & Gatsing, D. (2024). Occurrence of extended-spectrum β-lactamase (ESBL) and carbapenemase-producing Escherichia coli isolated from childhood diarrhoea in Yaoundé, Cameroon. BMC Microbiology, 24, 401. https://doi.org/10.1186/s12866-024-03559-z

Mandomando, I. M., Macete, E. V., Ruiz, J., Sanz, S., Abacassamo, F., Vallès, X., Sacarlal, J., Navia, M. M., Vila, J., Alonso, P. L., & Gascon, J. (2007). Etiology of diarrhea in children

younger than 5 years of age admitted in a rural hospital of southern Mozambique. The American Journal of Tropical Medicine and Hygiene, 76(3), 522–527. https://doi.org/10.4269/ajtmh.2007.76.522

Manhique-Coutinho, L., Chiani, P., Michelacci, V., Taviani, E., Bauhofer, A. F. L., Chissaque, A., Cossa-Moiane, I., Sambo, J., Chilaúle, J., Guimarães, E. L., Salência, J., Cassocera, M., Bero, D. M., Langa, J. P., & de Deus, N. (2022). Molecular characterization of diarrheagenic Escherichia coli isolates from children with diarrhea: A cross-sectional study in four provinces of Mozambique. International Journal of Infectious Diseases, 121, 190–194. https://doi.org/10.1016/j.ijid.2022.04.054

Mansan-Almeida, R., Pereira, A. L., & Giugliano, L. G. (2013). Diffusely adherent Escherichia coli strains isolated from children and adults constitute two different populations. BMC Microbiology, 13, 22. https://doi.org/10.1186/1471-2180-13-22

Medugu, N., Aworh, M. K., Iregbu, K., Nwajiobi-Princewill, P., Hull, D. M., Harden, L., Singh, P., Obaro, S., Egwuenu, A., Adeboye, F., Egah, R., Uzairue, L., Mohammed, Y., Ifeyinwa, N., & Thakur, S. (2025). Whole genome sequencing reveals virulence–mobile element linkages and phylogenetic diversity in multidrug-resistant Escherichia coli from Nigeria. Frontiers in Microbiology, 16, 1579175. https://doi.org/10.3389/fmicb.2025.1579175

Miller, J. J. (1978). The inverse of the Freeman-Tukey double arcsine transformation. The American Statistician, 32(4), 138. https://doi.org/10.1080/00031305.1978.10479283

Mkuhlu, N. A., Iweriebor, B. C., & Obi, C. L. (2020). Characterization and Antibiotic Susceptibility Profiles of Pathogenic Escherichia Coli Isolated from Diarrhea Samples within the Buffalo City Metropolitan Municipality, Eastern Cape, South Africa. The Open Microbiology Journal, 14, 321–330. https://doi.org/10.2174/1874434602014010321

Mohamed, J. A., Huang, D. B., Jiang, Z.-D., DuPont, H. L., Nataro, J. P., Belkind-Gerson, J., & Okhuysen, P. C. (2007). Association of putative enteroaggregative Escherichia coli virulence genes and biofilm production in isolates from travelers to developing countries. Journal of Clinical Microbiology, 45(1), 121–126. https://doi.org/10.1128/JCM.01128-06

Mossoro, C., Glaziou, P., Yassibanda, S., Lan, N. T. P., Bekondi, C., Minssart, P., Bernier, C., Le Bouguénec, C., & Germani, Y. (2002). Chronic diarrhea, hemorrhagic colitis, and hemolytic-uremic syndrome associated with HEp-2 adherent Escherichia coli in adults infected with human immunodeficiency virus in Bangui, Central African Republic. Journal of Clinical Microbiology, 40(8), 3086–3088. https://doi.org/10.1128/JCM.40.8.3086-3088.2002

Moyo, S. J., Maselle, S. Y., Matee, M. I., Langeland, N., & Mylvaganam, H. (2007). Identification of diarrheagenic Escherichia coli isolated from infants and children in Dar es Salaam, Tanzania. BMC Infectious Diseases, 7, 92. https://doi.org/10.1186/1471-2334-7-92

Munn, Z., Moola, S., Lisy, K., Riitano, D., & Tufanaru, C. (2015). Methodological guidance for systematic reviews of observational epidemiological studies reporting prevalence and cumulative incidence data. International Journal of Evidence-Based Healthcare, 13(3), 147–153. https://doi.org/10.1097/XEB.0000000000000054

Mwape, K., Bosomprah, S., Chibesa, K., Silwamba, S., Luchen, C. C., Sukwa, N., Mubanga, C., Phiri, B., Chibuye, M., Liswaniso, F., Somwe, P., Chilyabanyama, O., Chisenga, C. C., Muyoyeta, M., Simuyandi, M., Barnard, T. G., & Chilengi, R. (2023). Prevalence of Diarrhoeagenic Escherichia coli among Children Aged between 0–36 Months in Peri-Urban Areas of Lusaka. Microorganisms, 11(11), 2790. https://doi.org/10.3390/microorganisms11112790

Nataro, J. P., & Kaper, J. B. (1998). Diarrheagenic Escherichia coli. Clinical Microbiology Reviews, 11(1), 142–201. https://doi.org/10.1128/CMR.11.1.142

Navarro-Garcia, F., & Elias, W. P. (2011). Autotransporters and virulence of enteroaggregative E. coli. Gut Microbes, 2(1), 13–24. https://doi.org/10.4161/gmic.2.1.14933

Obi, C. L., Green, E., Bessong, P. O., de Villiers, B., Hoosen, A. A., Igumbor, E. O., & Potgieter, N. (2004). Gene encoding virulence markers among Escherichia coli isolates from diarrhoeic stool samples and river sources in rural Venda communities of South Africa. Water SA, 30(1), 37–42. https://doi.org/10.4314/wsa.v30i1.5024

Ochieng, J. B., Boisen, N., Powell, H., Kasumba, I. N., Badji, H., Zaman, S. M. A., Sow, S. O., Omore, R., Tennant, S. M., Kotloff, K. L., & Nataro, J. P. (2023). Epidemiology of Enteroaggregative, Enteropathogenic, and Shiga Toxin–Producing Escherichia coli Among Children Aged <5 Years in 3 Countries in Africa, 2015–2018: Vaccine Impact on Diarrhea in Africa (VIDA) Study. Clinical Infectious Diseases, 76(Suppl. 1), S77–S86. https://doi.org/10.1093/cid/ciad035

Ochoa, T. J., Ecker, L., Barletta, F., Mispireta, M. L., Gil, A. I., Contreras, C., Molina, M., Amemiya, I., Verastegui, H., Hall, E. R., Cleary, T. G., & Lanata, C. F. (2009). Age-related susceptibility to infection with diarrheagenic Escherichia coli among infants from Periurban areas in Lima, Peru. Clinical Infectious Diseases, 49(11), 1694–1702. https://doi.org/10.1086/648069

Okeke, I. N., Lamikanra, A., Czeczulin, J., Dubovsky, F., Kaper, J. B., & Nataro, J. P. (2000). Heterogeneous virulence of enteroaggregative Escherichia coli strains isolated from children in Southwest Nigeria. The Journal of Infectious Diseases, 181(1), 252–260. https://doi.org/10.1086/315204

Olayinka, A. A., Oginni-Falajiki, I. O., Okeke, I. N., & Aboderin, A. O. (2024). Diarrhoeagenic Escherichia coli associated with childhood diarrhoea in Osun state, Nigeria. BMC Infectious Diseases, 24, 912. https://doi.org/10.1186/s12879-024-09793-0

Onanuga, A., Igbeneghu, O., & Lamikanra, A. (2014). A study of the prevalence of diarrhoeagenic Escherichia coli in children from Gwagwalada, Federal Capital Territory, Nigeria. Pan African Medical Journal, 17, 146. https://doi.org/10.11604/pamj.2014.17.146.3369

Opintan, J. A., Newman, M. J., Ayeh-Kumi, P. F., Affrim, R., Gepi-Attee, R., Sevilleja, J. E. A. D., Roche, J. K., Nataro, J. P., Warren, C. A., & Guerrant, R. L. (2010). Pediatric diarrhea in southern Ghana: Etiology and association with intestinal inflammation and malnutrition. The American Journal of Tropical Medicine and Hygiene, 83(4), 936–943. https://doi.org/10.4269/ajtmh.2010.09-0792

Pabst, W. L., Altwegg, M., Kind, C., Mirjanic, S., Hardegger, D., & Nadal, D. (2003). Prevalence of enteroaggregative Escherichia coli among children with and without diarrhea in Switzerland. Journal of Clinical Microbiology, 41(6), 2289–2293. https://doi.org/10.1128/JCM.41.6.2289-2293.2003

Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., … Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. BMJ, 372, n71. https://doi.org/10.1136/bmj.n71

Patzi-Vargas, S., Zaidi, M. B., Perez-Martinez, I., Leon-Cen, M., Michel-Ayala, A., Chaussabel, D., & Estrada-Garcia, T. (2015). Diarrheagenic Escherichia coli carrying supplementary virulence genes are an important cause of moderate to severe diarrhoeal disease in Mexico. PLOS Neglected Tropical Diseases, 9(3), e0003510. https://doi.org/10.1371/journal.pntd.0003510

Platts-Mills, J. A., Babji, S., Bodhidatta, L., Gratz, J., Haque, R., Havt, A., McCormick, B. J. J., McGrath, M., Olortegui, M. P., Samie, A., Shakoor, S., Mondal, D., Lima, I. F. N., Hariraju, D., Rayamajhi, B. B., Qureshi, S., Kabir, F., Yori, P. P., Mufamadi, B., … Houpt, E. R. (2015). Pathogen-specific burdens of community diarrhoea in developing countries: A multisite birth cohort study (MAL-ED). The Lancet Global Health, 3(9), e564–e575. https://doi.org/10.1016/S2214-109X(15)00151-5

Prendergast, A., & Kelly, P. (2012). Enteropathies in the developing world: neglected effects on global health. The American Journal of Tropical Medicine and Hygiene, 86(5), 756–763. https://doi.org/10.4269/ajtmh.2012.11-0743

Qadri, F., Svennerholm, A.-M., Faruque, A. S. G., & Sack, R. B. (2005). Enterotoxigenic Escherichia coli in developing countries: Epidemiology, microbiology, clinical features, treatment, and prevention. Clinical Microbiology Reviews, 18(3), 465–483. https://doi.org/10.1128/CMR.18.3.465-483.2005

Robins-Browne, R. M., Bordun, A.-M., Tauschek, M., Bennett-Wood, V. R., Russell, J., Oppedisano, F., Lister, N. A., Bettelheim, K. A., Fairley, C. K., Sinclair, M. I., & Hellard, M. E. (2004). Escherichia coli and community-acquired gastroenteritis, Melbourne, Australia. Emerging Infectious Diseases, 10(10), 1797–1805. https://doi.org/10.3201/eid1010.031086

Rogawski, E. T., Guerrant, R. L., Havt, A., Lima, I. F. N., Medeiros, P. H. Q. S., Seidman, J. C., McCormick, B. J. J., Babji, S., Kosek, M. N., Qureshi, S., Shakoor, S., Bhutta, Z. A., Samie, A., Bessong, P. O., Haque, R., Mondal, D., Bodhidatta, L., Mason, C. J., Shrestha, S., … Platts-Mills, J. A. (2017). Epidemiology of enteroaggregative Escherichia coli infections and associated outcomes in the MAL-ED birth cohort. PLOS Neglected Tropical Diseases, 11(7), e0005798. https://doi.org/10.1371/journal.pntd.0005798

Rowe, J. S., Shah, S. S., Motlhagodi, S., Bafana, M., Tawanana, E., Truesdale, H. T., Mullan, P. C., Steenhoff, A. P., & Arscott-Mills, T. (2010). An epidemiologic review of enteropathogens in Gaborone, Botswana: Shifting patterns of resistance in an HIV endemic region. PLOS ONE, 5(6), e10924. https://doi.org/10.1371/journal.pone.0010924

Saka, H. K., Dabo, N. T., Muhammad, B., Garcia-Soto, S., Ugarte-Ruiz, M., & Alvarez, J. (2019). Diarrheagenic Escherichia coli pathotypes from children younger than 5 years in Kano State, Nigeria. Frontiers in Public Health, 7, 348. https://doi.org/10.3389/fpubh.2019.00348

Sang, W. K., Oundo, V., & Schnabel, D. (2012). Prevalence and antibiotic resistance of bacterial pathogens isolated from childhood diarrhoea in four provinces of Kenya. The Journal of Infection in Developing Countries, 6(7), 572–578. https://doi.org/10.3855/jidc.2196

Santos, A. C. M., Santos, F. F., Silva, R. M., & Gomes, T. A. T. (2020). Diversity of hybrid- and hetero-pathogenic Escherichia coli and their potential implication in more severe diseases. Frontiers in Cellular and Infection Microbiology, 10, 339. https://doi.org/10.3389/fcimb.2020.00339

Sarantuya, J., Nishi, J., Wakimoto, N., Erdene, S., Nataro, J. P., Sheikh, J., Iwashita, M., Manago, K., Tokuda, K., Yoshinaga, M., Miyata, K., & Kawano, Y. (2004). Typical enteroaggregative Escherichia coli is the most prevalent pathotype among E. coli strains causing diarrhea in Mongolian children. Journal of Clinical Microbiology, 42(1), 133–139. https://doi.org/10.1128/JCM.42.1.133-139.2004

Scaletsky, I. C. A., Fabbricotti, S. H., Carvalho, R. L. B., Nunes, C. R., Maranhão, H. S., Morais, M. B., & Fagundes-Neto, U. (2002). Diffusely adherent Escherichia coli as a cause of acute diarrhea in young children in Northeast Brazil: A case-control study. Journal of Clinical Microbiology, 40(2), 645–648. https://doi.org/10.1128/JCM.40.2.645-648.2002

Scaletsky, I. C. A., Aranda, K. R. S., Souza, T. B., Silva, N. P., & Morais, M. B. (2009). Evidence of pathogenic subgroups among atypical enteropathogenic Escherichia coli strains. Journal of Clinical Microbiology, 47(11), 3756–3759. https://doi.org/10.1128/JCM.01599-09

Schünemann, H. J., Oxman, A. D., Higgins, J. P., Vist, G. E., Glasziou, P., & Guyatt, G. H. (2008). Presenting results and ‘Summary of findings’ tables. In J. P. Higgins & S. Green (Eds.), Cochrane Handbook for Systematic Reviews of Interventions (pp. 335–357). John Wiley & Sons. https://doi.org/10.1002/9780470712184.ch11

Servin, A. L. (2005). Pathogenesis of Afa/Dr diffusely adhering Escherichia coli. Clinical Microbiology Reviews, 18(2), 264–292. https://doi.org/10.1128/CMR.18.2.264-292.2005

Servin, A. L. (2014). Pathogenesis of human diffusely adhering Escherichia coli expressing Afa/Dr adhesins (Afa/Dr DAEC): Current insights and future challenges. Clinical Microbiology Reviews, 27(4), 823–869. https://doi.org/10.1128/CMR.00036-14

Shah, M., Kathiiko, C., Wada, A., Odoyo, E., Bundi, M., Miringu, G., Guyo, S., Karama, M., & Ichinose, Y. (2016). Prevalence, seasonal variation, and antibiotic resistance pattern of enteric bacterial pathogens among hospitalized diarrheic children in suburban regions of central Kenya. Tropical Medicine and Health, 44, 39. https://doi.org/10.1186/s41182-016-0038-1

Spano, L. C., Sadovsky, A. D. I., Segui, P. N., Saick, K. W., Kitagawa, S. M. S., Pereira, F. E. L., Fagundes-Neto, U., & Scaletsky, I. C. A. (2008). Age-specific prevalence of diffusely adherent Escherichia coli in Brazilian children with acute diarrhoea. Journal of Medical Microbiology, 57(3), 359–363. https://doi.org/10.1099/jmm.0.47660-0

Sweeting, M. J., Sutton, A. J., & Lambert, P. C. (2004). What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data. Statistics in Medicine, 23(9), 1351–1375. https://doi.org/10.1002/sim.1761

Taddei, C. R., Fasano, A., Ferreira, A. J. P., Trabulsi, L. R., & Martinez, M. B. (2005). Secreted autotransporter toxin produced by a diffusely adhering Escherichia coli strain causes intestinal damage in animal model assays. FEMS Microbiology Letters, 250(2), 263–269. https://doi.org/10.1016/j.femsle.2005.07.013

Tau, N. P., Meidany, P., Smith, A. M., Sooka, A., & Keddy, K. H. (2012). Escherichia coli O104 associated with human diarrhea, South Africa, 2004–2011. Emerging Infectious Diseases, 18(8), 1314–1317. https://doi.org/10.3201/eid1808.111616

Trabulsi, L. R., Keller, R., & Tardelli Gomes, T. A. (2002). Typical and atypical enteropathogenic Escherichia coli. Emerging Infectious Diseases, 8(5), 508–513. https://doi.org/10.3201/eid0805.010385

Troeger, C., Blacker, B. F., Khalil, I. A., Rao, P. C., Cao, S., Zimsen, S. R., Albertson, S. B., Stanaway, J. D., Deshpande, A., Abebe, Z., Alvis-Guzman, N., Amare, A. T., Asgedom, S. W., Anteneh, Z. A., Antonio, C. A. T., Aremu, O., Asfaw, E. T., Atey, T. M., Atique, S., … Reiner, R. C., Jr. (2018). Estimates of the global, regional, and national morbidity, mortality, and aetiologies of diarrhoea in 195 countries: A systematic analysis for the Global Burden of Disease Study 2016. The Lancet Infectious Diseases, 18(11), 1211–1228. https://doi.org/10.1016/S1473-3099(18)30362-1

Vargas, M., Gascon, J., Casals, C., Schellenberg, D., Urassa, H., Kahigwa, E., Ruiz, J., & Vila, J. (2004). Etiology of diarrhea in children less than five years of age in Ifakara, Tanzania. The American Journal of Tropical Medicine and Hygiene, 70(5), 536–539. https://doi.org/10.4269/ajtmh.2004.70.536

Wells, G. A., Shea, B., O'Connell, D., Peterson, J., Welch, V., Losos, M., & Tugwell, P. (2014). The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Ottawa Hospital Research Institute. http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp

Wolde, D., Eguale, T., Medhin, G., Haile, A. F., Alemayehu, H., Mihret, A., Pirs, M., Strašek Smrdel, K., Avberšek, J., Kušar, D., Cerar Kišek, T., Janko, T., Steyer, A., & Starčič Erjavec, M. (2024). Diarrheagenic Escherichia coli in stool specimens collected from patients attending primary healthcare facilities in Ethiopia: Whole-genome sequencing-based molecular characterization. International Journal of Molecular Sciences, 25(19), 10251. https://doi.org/10.3390/ijms251910251

World Health Organization. (2019). The treatment of diarrhoea: a manual for physicians and other senior health workers (4th rev.). World Health Organization. https://iris.who.int/handle/10665/43209

World Health Organization Regional Office for Africa. (2024). Technical guidelines for integrated disease surveillance and response in the African region (3rd ed.). WHO Regional Office for Africa. https://www.afro.who.int/publications/technical-guidelines-integrated-disease-surveillance-and-response-african-region-third

Zelelie, T. Z., Eguale, T., Yitayew, B., Abeje, D., Alemu, A., Seman, A., Jass, J., Mihret, A., & Abebe, T. (2023). Molecular epidemiology and antimicrobial susceptibility of diarrheagenic Escherichia coli isolated from children under age five with and without diarrhea in Central Ethiopia. PLOS ONE, 18(7), e0288517. https://doi.org/10.1371/journal.pone.0288517