

ACUTE TOXICITY IN MICE

COMPOUND 3-Hydroxy-3-methylbutanoic acid (Tur 13)SOURCE Lorillard - Organic Chemistry LOT NO. OR39-23DATE RECEIVED Unk. TESTED 12/28/78 REPORTED 5/3/79
10/6/80, UpdateINVESTIGATOR(S) H. S. Tong & M. S. Forte NOTEBOOK PAGE BIO14-23SIGNATURE(S) H. S. Tong M. S. Forte (by A. Poole)STRAIN OF MICE Swiss-Webster MALE ☒ FEMALE ☐ DATE RECEIVED Unk.AVERAGE WEIGHT(RANGE) (GM) _____ SOURCE Camu ResearchROUTE OF COMPOUND ADMINISTRATION ☒ P.O. ☐ I.P. ☐ I.V. ☐ INHALATIONCOMPOUND VEHICLE ☒ 0.5 % METHYL CELLULOSE ☐ CORN OIL ☐ SALINE ☐ OTHER _____

GROUP NO.	% SOLUTION	DOSE (mg/kg BODY WEIGHT)	RESULTS (NO DEAD/NO TESTED)
1	5	1800	1/6
2	10	2160	0/6
3	10	2592	0/6
4	10	3732	3/6
5	10	4479	6/6

REFERENCE FOR CALCULATION Litchfield, J. T. and Wilcoxin, F., J. of Pharmacol.
and Exper. Ther., 90:99, 1948.LD50 (95% CONFIDENCE LIMITS) 3.5 (3.1 to 3.9)g/kgCONCLUSION This compound appears to act as a CNS depressant with symptoms
of respiratory depression, constriction of blood vessels, and in-
activity. Survivors recovered in 48 hours. The recommended safe
dose for a single trial by inhalation in man is 0.3 mg.Copies to the Following: Dr. H. J. Minnemeyer
Ms. L. B. Gray

00040534