

XVIII.

Methodus Testaceorum D. D'ARGENTVILLE.

		Species.	Varietates.
Classis I. Conchæ Univalves.	Familia XIV. <i>Concha Globosa.</i> Concha globosa est univalvis, ventre instar globi subrotundo, apertura laxiore, dentata, edentula, clavícula mediocriter umbonata, depressa, columella aspera, lævi.	1. Dolium rotundum & umbilicatum 2. Dolium oblongum & læve. 3. Dolium oblongum costatum, umbonatum. 4. Dolium cauda elongata & arcuata. 5. Dolium Bulla dictum. 6. Dolium, concha sphaerica fasciata, cœrulea, intus flavida, Vitta cœrulea appellata, rarissima.	1. — 6. 2. — 6. 3. — 8. 4. — 5. 5. — 7. 6. — 1.
	Familia XV. <i>Porcellana seu Veneræ.</i> Porcellana seu Venera est Concha univalvis, a rimula sic nominata, ore dentato ex utraque parte, conglobata, oblonga, gibbosa, umbonata.	1. Porcellana conglobata & crassa. 2. Porcellana pyriformis & tenuis. 3. Porcellana figura oblonga & crassa. 4. Porcellana in aliqua parte gibbosa.	1. — 17. 2. — 4. 3. — 16. 4. — 5.
Classis II. Conchæ Bivalves.	Familia I. <i>Ostrea.</i> Ostreum est Concha bivalvis, testis extus sordidis, luto obductis, e crustis multis seu laminis compositis, lævis, aculeis, striis, tuberculis instructa, plana, plicata, globosa, valva inferiori elatior quam superiore.	1. Ostreum planum & læve. 2. Ostreum plicatum læve. 3. Ostreum globosum & spinosum. 4. Ostreum globosum, laminis eminentibus laciniatum. 5. Ostreum globosum, foliaceum. 6. Ostreum oblongum & umbonatum.	1. — 15. 2. — 6. 3. — 10. 4. — 2. 5. — 4. 6. — 5.