

XVIII.

Methodus Testaceorum D. D'ARGENTVILLE.

Classis II. Conchæ Bivalves.	Familia II.		
	<i>Chama.</i>		
	Chama est Concha bivalvis, lævis, aliquantulum rugosa, spinosa, valvis æqualibus clatis, convexis, ore patulo & hiantē.	Species.	Varietates.
		1. Chama oblonga, plana, transverse striata.	1. — 4.
		2. Chama oblonga, elata, ab apice striata.	2. — 6.
		3. Chama rotunda æquilateralis.	3. — 25.
		4. Chama Venerea, inæquilateralis.	4. — 15.
		5. Chama striis diversimode exaratis.	5. — 1.
	Familia III.		
	<i>Musculi</i> seu <i>Mytuli</i> .		
	Musculus seu Mytulus est Concha bivalvis, oblonga, in acumen definens æqualibus extremitatibus, lævis, aspera, plana, elata, aliquando rostro paulisper elato.	1. Musculus planus in acumen definens.	1. — 21.
		2. Musculus elatus, æquilateralis.	2. — 8.
		3. Tellina oblonga, plana, æqualibus extremitatibus.	3. — 11.
		4. Tellina oblonga, dissimilibus extremitatibus.	4. — 10.
		5. Tellina mutilata.	5. — 3.
	Familia IV.		
	<i>Cordiformes.</i>		
	Concha Cordiformis est bivalvis, globosa, elata, canaliculata, inaurita, spinosa vel imbricata.	1. Bucardium globosum.	1. — 7.
		2. Cordiforme triangulare.	2. — 6.
		3. Cordiforme, Cor hominis, seu Veneris.	3. — 3.
		4. Cordiforme imbricatum.	4. — 4.
		5. Cordiforme, Archa Noëni.	5. — 2.
		6. Concha exotica margine in mucronem emisso, intus fistulosa, tota candida & tenuis F. Columnæ.	
	Familia V.		
	<i>Pectines.</i>		
	Pecten est Concha bivalvis, ex omni parte clusilis & striata in formam Pectinis, quo capilli pectuntur, plana, elata, aurita, semiaurita, inaurita.	1. Pecten utrinque auritus.	1. — 17.
		2. Pecten semiauritus.	2. — 6.
		3. Pecten inauritus.	3. — 7.