

XVIII.

Methodus Testaceorum D. D'ARGENTVILLE.

Class. II. Conchæ Bivalves.	Familia VI.		Species.	Varietates.
	<i>Solenes five Ungues.</i>			
	Solēn est Concha bi- valvis, corpore longo, ex utraque extremitate patente, recto vel arcuato.		1. Solen corpore recto.	1. — 12.
			2. Solen corpore arcuato, seu cur- vus.	2. — 2.
Class. III. Conchæ Multivalves seu Polyvalves.	Familia I.			
	<i>Echini.</i>			
	Echinus marinus est	1. Echinus orbicularis.	1. — 9.	
	Concha multivalvis,	2. Echinus ovalis.	2. — 2.	
	orbicularis, ovalis, an- gulosa, irregularis, pla- na, aculeis, tuberculis insignita, aliquando spinis denudata.	3. Echinus angulosus.	3. — 3.	
		4. Echinus irregularis.	4. — 9.	
		5. Echinus planus, stellatus.	5. — 2.	
	Familia II.			
	<i>Vermiculi marini.</i>			
	Vermiculi marini sunt	1. Vermiculi directe dispositi.	1. — 5.	
	Conchæ multivalves,	2. Vermiculi incurvati.	2. — 4.	
	canaliformes, subro- tundæ, rugosæ, incur- vatæ, rectæ, intortæ.	3. Vermiculi circulariter dispositi.	3. — 9.	
	Familia III.			
	<i>Balani.</i>			
	Balanus est Concha	1. Balanus ore ampliore, major latus.	1. — 6.	
	multivalvis, glandi- formis, duodecim lami- nas continens, ore ampliore, angustiore.	2. Balanus ore angustiore, striatus parvus, seu minor.	2. — 4.	