


# **DIC and Fibrin Related Markers**

**A comprehensive offer**




# What is DIC?

(Disseminated Intravascular Coagulation)




➔ DIC involves a dynamic situation due to an imbalance between procoagulant and anticoagulant proteins

# Disorders usually associated with DIC




- ➔ DIC is not a disease itself but is always secondary to an underlying disease
- ➔ DIC is a life-threatening condition

# Pathogenesis of DIC



# Fibrin-Related Markers


Markers of plasmin & thrombin action (1)






# Fibrin-Related Markers

Markers of plasmin & thrombin action (2)



# Fibrin-Related Markers


Markers of plasmin & thrombin action (3)





# Fibrin-Related Markers

Markers of plasmin & thrombin action



- Concomitant generation of Thrombin & Plasmin
- Balance between FM and FDP/D-Di



# DIC Diagnosis

## *Key aspects*

---

- Encompass **both clinical and laboratory** information
- The ISTH DIC Scoring system correlates key clinical observations and outcomes
- Repeat tests to monitor the dynamically changing scenario

*"Guidelines for the diagnosis and management of disseminated intravascular coagulation",  
British Society of Haematology, British Journal of Haematology, 2009*

# Composite scoring systems for DIC

## *ISTH diagnostic algorithm for overt DIC*

1. **Risk assessment: does the patient have an underlying disorder known to be associated with overt DIC?**
  - If yes: **proceed**; if no: **do not use this algorithm**
2. **Order global coagulation tests**  
*platelet count, PT, fibrinogen, soluble fibrin monomers or fibrin degradation products*
3. **Score global coagulation test results:**
  - platelet count (G/L):  $>100 = 0$  ■  $<100 = 1$  ■  $<50 = 2$
  - prolonged PT:  $<3 \text{ s} = 0$  ■  $>3 \text{ s but } <6 \text{ s} = 1$  ■  $>6 \text{ s} = 2$
  - fibrinogen level:  $>1.0 \text{ g/L} = 0$  ■  $<1.0 \text{ g/L} = 1$
  - elevated fibrin-related markers (*e.g. FM, FDP*):  
no increase = **0** ■ moderate increase = **2** ■ strong increase = **3**
4. **Calculate score**
  - If  $\geq 5$ : **compatible with overt DIC**; repeat scoring daily
  - If  $< 5$ : suggestive (*not affirmative*) **for non-overt DIC**; repeat next 1-2 days.

➔ a DIC score encompassing clinical and laboratory evidence showed higher diagnostic performances than any test alone

# DIC

## Which lab tests are most useful?

### Natural Coagulation Inhibitors

Antithrombin (AT),  
Protein C (PC):  
Low levels in 90%  
of DIC patients

### Markers of Fibrinolysis

### Fibrinogen

### Coagulation Factors

### Global Clotting times

Prolonged PT and APTT:  
> 95% of DIC patients

### Platelet Count

Incidence in DIC patients

< 100G/L:

50 - 60% (critically ill medical patients)

> 80% (surgical and trauma patients)

FRMs



➔ No single test is sufficiently specific to diagnose or rule-out DIC  
DIC score encompass multiple parameters assessment and follow-up

# DIC

## Focus on FRMs

---

- **Fibrin monomers (FM)**

- Helpful to diagnose intra-vascular fibrin formation in DIC
- Above a defined threshold a diagnosis of DIC can be made
- Only generated intravascularly  
Not influenced by extravascular fibrin formation  
(*local inflammation or trauma*)

- **Fibrin(ogen) degradation products (FDP)**

- Detectable in 99% of patients with sepsis and DIC

- **D-dimer (DDi)**


- High levels in patients with DIC
- Poorly distinguish patients with DIC from patients with VTE, recent surgery and inflammatory conditions



➔ All FRMs are elevated in hypercoagulable states and may be useful for the diagnosis of DIC



# Theoretical course of FRMs



In normal conditions, activated thrombin will cleave Fibrinogen into Fibrin Monomers (FM) and fibrinopeptides. FM will form non covalent bindings to form Soluble fibrin polymer.

The plasma FM level will decrease 2 or 3 days after the onset of thrombosis but the D-dimer level will not increase immediately.

## All FRMs do not have the same time course


- **D-Dimer** and **FDP** reflect secondary fibrinolysis after clot formation
- FM reflect the early phase of thrombosis; may be an earlier predictive marker

# Study on 105 plasma samples from patients with sepsis, SIRS\* and healthy individuals

Toh JMH, Ken-Dror G., Downey C.

*Fibrin-Related Marker comparison in Sepsis: Correlation to global and molecular parameters of haemostatic dysfunction.*

58th Annual meeting of SSC ISTH, LIVERPOOL, UK , 2012.



→ Equivalence of the 3 FRM as markers of intravascular fibrininformation with FM performing better in relation to global hemostatic tests (*PT, PLT, FIB*) used in DIC scoring.

\*SIRS : Systemic Inflammatory Response Syndrome

1

Wada H, Sakuragawa N,  
*Are Fibrin –Related Markers Useful for the Diagnosis of Thrombosis?*  
*Seminar in Thrombosis and Hemostasis, 2008; Volume 34; Number 1.*

### Objectives

Comparison the pathogenesis and course of elevation of FRMs in hypercoagulable states and evaluate their usefulness in the diagnosis of thrombosis.

### Outcomes

Timecourse of FRMs after the onset of thrombosis

### Findings

Elevated levels of FRMs indicate a high risk of thrombosis providing useful information for the diagnosis; SF may reflect the early phase of thrombosis whereas D-Dimer reflect the secondary fibrinolysis after clot formation.

As they don't have the same dynamic of elevation and decrease relatively to the onset of thrombosis, **both D-dimer and SF measurements may be recommended in patients with DIC.**

2

*Gris JC, Faille JL, Cochery-Nouvellon E, Lissalde G, Lefrant JY, J Thromb Haemost, 2011; 9: 1252-5.*

## Objectives

Prospective study for further assess early FM, DDi and related ISTH overt DIC score values and prognostic values.

## Methods

350 patients > 18 years entering ICU with acute septic shock.

## Outcomes

Monitoring of survival at 90 days

## Findings

A positive overt DIC score using DDi or FM as FRMs were associated with a poor 90-Days outcome; FM being a better prognosis marker.



3

*Dempfe CE, Wurst M, Smolinski M, Lorenz S, Osika A, Olenik D, Fiedler F, Borggrefe M. Thromb Haemost, 2004; 91: 812-8.*

## Objectives

Comparison of the predictive value of the ISTH overt DIC score in a surgical intensive care cohort using either DDi or FM as FRM.

## Methods

331 patients - day1 blood sample

## Outcomes

Monitoring of 28-day mortality

## Findings

FM-DIC score displayed the highest prognostic power.

## FM, DDi and FDP:

- Are elevated in hypercoagulable states
- Have different time course (FM prethrombotic, FDP/D-Di post-thrombotic)
- May be useful as part of DIC composite score
- May be useful prognosis markers in DIC patients

# Take home messages

## FM, DDi and FDP:

- Are elevated in hypercoagulable states
- Have different time courses  
(*FM prethrombotic, FDP/DDi post-thrombotic*)
- May be useful as part of DIC composite score
- May be useful prognosis markers in DIC patients

