Notations and conventions

We close this section with some notational conventions. We let $[m]$ denote the set $\{1, 2, \ldots, m\}$ for every positive integer m. $\mathbb{R}, \mathbb{C}, \mathbb{Z}$ and $\mathbb{Z}_+$ stand for the set of all real, complex, integer and non-negative integer values, respectively. Tr denotes the trace operator and $\mathrm{tr} = 2^{-\frac{n}{2}}\,\mathrm{Tr}$ is the normalized trace (so that $\mathrm{tr}(\mathbb{I}) = 1$ for the identity operator $\mathbb{I}$ in $\mathbb{C}^{2^{\frac{n}{2}}}$).

Next we introduce some graph-theoretic notations. A simple q-uniform hypergraph, or just simple hypergraph is a pair (V, E) where V is the set of nodes and E is a subset of the set of all $\binom{V}{q}$ cardinality q subsets of V. For two hyperedges $e_1, e_2 \in E$ we write $e_1 \sim e_2$ when $e_1 \cap e_2 \neq \emptyset$. With some abuse of notation we let $\binom{V}{q}$ denote the set of all q-size subsets of V. A multi q-uniform hypergraph, or just multi hypergraph, is a pair $(V, \bar{r})$ where $\bar{r} \in \mathbb{Z}_+^{\binom{|V|}{q}}$. For each subset $e \in \binom{V}{q}$ $r_e \in \mathbb{Z}_+$ is called the multiplicity of the hyperedge e. $m(V, E) = \sum_{e \in \binom{V}{q}} r_e$ denotes the total multiplicity of the multi hypergraph (V, E). Each multi hypergraph naturally induces a simple hypergraph where hyperedges are included iff $r_e \geq 1$. Let $\mathcal{H}^{\mathrm{sim}}([m])$ denote the set of all simple hypergraphs on the node set $[m]$. $\mathcal{H}([m])$ denotes the set of all multi hypergraphs on the node set $[m]$ with even multiplicities, namely $r_e \in 2\mathbb{Z}_+$ for all e. Note that while $\mathcal{H}^{\mathrm{sim}}([m])$ is a finite set for every m, the set $\mathcal{H}([m])$ is infinite.

A simple hypergraph (V, E) is connected if for every $u, v \in V$ there exists a path from u to v. Namely, there exists a sequence of hyperedges $e_1, \ldots, e_r$ such that $u \in e_1, v \in e_r$ and $e_i \cap e_{i+1} \neq \emptyset, i \in [r-1]$. A multi hypergraph is connected if its induced simple hypergraph is connected. Every simple and multi hypergraph H admits a unique decomposition into connected components $H_1, \ldots, H_C$ so that H is the disjoint union of H_i and each H_i is a connected hypergraph. A simple hypergraph (V, E) is called a hypertree or simply a tree if it is a connected hypergraph with precisely $(|V|-1)/(q-1)$ hyperedges. It is easy to verify that every two hyperedges of a hypertree intersect in at most one node. The set of all simple hypertrees on the node set $[m]$ is denoted by $\mathcal{T}_m^{\mathrm{sim}}$. A singleton, namely a graph on one node with no hyperedges is defined to be connected by default, so that $\mathcal{T}_1^{\mathrm{sim}} = \{1\}$. A special role in our derivation will be played by multi hypertrees where all multiplicities are $r_e = 2$, so that the total number of hyperedges is $2(m-1)/(q-1)$. We denote the set of such hypertrees by $\mathcal{T}_{m,2}$. Given $m \geq 1, \Delta \geq 0$, $\mathcal{H}_{m,\Delta} \subset \mathcal{H}([m])$ denotes the set of all connected multi hypergraphs with precisely $2(m-1+\Delta)/(q-1)$ hyperedges. Note that $\mathcal{H}_{v,0} = \mathcal{T}_{v,2}$. We call Δ the excess of H for $H \in \mathcal{H}_{m,\Delta}$.

1 Main result. Free energy limit

The SYK model is defined on the Hilbert space $\mathbb{C}^{2^{\frac{n}{2}}}$ via the so-called Majorana operators $\psi_i, i \in [n]$ on $\mathbb{C}^{2^{\frac{n}{2}}}$ satisfying

$$\{\psi_i, \psi_j\} \triangleq \psi_i\psi_j + \psi_j\psi_i = \delta_{ij}. \tag{1}$$

Here δ_{ij} is the identity operator $\mathbb{I}$ on $\mathbb{C}^{2^{\frac{n}{2}}}$, when $i = j$, and 0 otherwise, so that $\psi_i^2 = \frac{1}{2}\mathbb{I}$. These operators can be explicitly constructed using standard Pauli operators X, Y, Z by setting $\psi_{2i+1} = \frac{1}{\sqrt{2}}(\otimes_{1\leq k\leq i} Z) \otimes X_{i+1} \otimes \mathbb{I}$ and $\psi_{2i+2} = \frac{1}{\sqrt{2}}(\otimes_{1\leq k\leq i} Z) \otimes Y_{i+1} \otimes \mathbb{I}$, for $0 \leq i \leq n/2 - 1$. This explicit construction though will not be necessary for us. We fix a positive even integer q, the interaction parameter. For every $I = (i_1, \ldots, i_q) \in \binom{[n]}{q}, 1 \leq i_1 < \cdots < i_q \leq n$ we define

$\psi_I \triangleq i^{q/2}\psi_{i_1}\cdots\psi_{i_q}$. In particular $\psi_I^2 = 2^{-q}\mathbb{I}$, which is an easy implication of (1). The Hamiltonian of the SYK model with q-body interaction is then defined as the Hermitian operator

$$H_{\mathrm{SYK}} \triangleq \sum_{I\in\binom{[n]}{q}} J_I\psi_I, \tag{2}$$

where $(J_I, I \in \binom{[n]}{q})$ is a sequence of i.i.d. random variables defined by $J_I = \sigma\bar{J}_I$ where $\sigma^2 = \frac{(q-1)!}{n^{q-1}}$ and $\bar{J}_I$ are Rademacher random variables, independent across $I \in \binom{[n]}{q}$. I.e. $\bar{J}_I = \pm 1$ with equal probabilities $1/2$. By the universality property established in [1] the free energy limit (defined below) will be the same for the case when $\bar{J}_I$ are i.i.d. standard normal random variables, which is a more common choice in defining the SYK Hamiltonian. For every fixed inverse temperature $\beta \in \mathbb{R}$ we define the associated (random) partition function $Z_{n,q}(\beta) \triangleq \mathrm{tr}\left(e^{\beta H_{\mathrm{SYK}}}\right)$, where we recall that $\mathrm{tr} = 2^{-\frac{n}{2}}\,\mathrm{Tr}$ stands for the normalized trace operator. The normalization by $2^{\frac{n}{2}}$ is chosen for convenience as the trace of the identity operator $\mathbb{I}$ in $\mathbb{C}^{2^{\frac{n}{2}}}$ is $2^{\frac{n}{2}}$. We define the associated free energy as

$$F_{n,q}(\beta) \triangleq \frac{1}{\beta}\log Z_{n,q}(\beta).$$

Our main result is the computation of the free energy limit

$$F_q(\beta) \triangleq \lim_{n\to\infty}\frac{F_{n,q}(\beta)}{n},$$

for small enough constant inverse temperature β. In particular, we prove the existence of this limit (see the discussion in the introduction), and establish that it holds w.h.p. as $n \to \infty$. In light of the "quenched $\approx$ annealed" result (**??**) mentioned earlier and in light of the concentration of the quenched free energy around its expectation, also verified in [1], we instead focus on

$$f_{n,q}(\beta) \triangleq \frac{1}{\beta}\log \mathbb{E}[Z_{n,q}(\beta)].$$

Thus our goal is proving the existence and computing the limit

$$f_q(\beta) \triangleq \lim_{n\to\infty}\frac{f_{n,q}(\beta)}{n},$$

as $f_q(\beta) = F_q(\beta)$ for all β.

In order to state our result formally, we begin by introducing some necessary technical preliminaries. Let $\Omega = [0,1]^2$. Elements $x = (u_1, u_2) \in \Omega$ will be referred to as "chords". For every $x = (u_1, u_2), y = (v_1, v_2) \in \Omega$, define the chord intersection function

$$K(x,y) = \begin{cases} -1 & \text{if } u_1 < v_1 < u_2 < v_2 \text{ or } v_1 < u_1 < v_2 < u_2, \\ 1 & \text{otherwise.} \end{cases}$$

Namely $K(x,y) = -1$ when the chords x and y cross and $= 1$ otherwise. Let ν be the probability law of $(\min(U_1,U_2), \max(U_1,U_2)) \in \Omega$ when $U_j, j = 1,2$ are generated uniformly at random from $[0,1]$, independently.




Figure 1: Chord intersection sign for two chords $x = (u_1, u_2), y = (v_1, v_2) \in \Omega$.

On the space of continuous functions from $\{z \in \mathbb{C} : |z| \le 1/e^2\} \times \Omega$ to $\mathbb{C}$ define a norm $\|G\|_{e,\Omega} = \sup_{|z|\le 1/e^2, x\in\Omega} |G(z,x)|$. The restriction $|z| \le 1/e^2$ can be relaxed a little and is chosen for convenience. Let $\mathcal{F}_{e,\Omega}$ be the subset of these functions G satisfying $\|G\|_{e,\Omega} \le 1$. On $\mathcal{F}_{e,\Omega}$ we introduce a function H, which we call the Chord-Cavity-Generator (CCG), defined implicitly via the following fixed point equation

$$H(z,x) = z + z \sum_{d\ge 1} \frac{1}{d!} \int_{\Omega^d} \prod_{i=1}^{d} \left[K(x,y_i) \frac{H(z,y_i)^{q-1}}{(q-1)!} \right] \prod_{1\le i<j\le d} K(y_i,y_j)\, \nu^{\otimes d}(dy_1 \cdots dy_d). \tag{3}$$

The validity of this definition is justified by the following claim.

Proposition 1 (Existence of CCG)**.** *For every even $q \ge 2$ there exists $H \in \mathcal{F}_{e,\Omega}$ which is the unique solution of the equation (3). Furthermore, H is analytic on the domain $\{z \in \mathbb{C} : |z| \le 1/e^2\} \times \Omega$.*

The proof of this proposition is found in Section 3. Next we provide a more intuitive and equivalent description of CCG via the chord crossing diagrams on trees. Consider any simple q-uniform hypergraph $H = (V, E)$ on the node set $V = V(H)$ and edge set $E = E(H)$. Suppose we have a chord mapping $X : E \to \Omega$. Define

$$B_H(X) = \prod_{e_1,e_2\in E:|e_1\cap e_2| \text{ odd}} K(X(e_1), X(e_2)) \in \{\pm 1\}.$$

Namely, $B_H(X)$ is $(-1)^R$, where R is the number of pairs of hyperedges which share an odd number of nodes and such that their corresponding chords cross. Let

$$b_H = \mathbb{E}[B_H(X)], \tag{4}$$

when X is generated according to the law $\nu^{\otimes|E|}$. Note that $|b_H| \le 1$.

Similarly, suppose we have a simple rooted hypergraph H with a distinguished node $r \in V$ called a root. Fix $x \in \Omega$ and $X : E \to \Omega$. Define

$$B_{H,r}(X,x) = \prod_{e\in E: r\in e} K(x, X(e)) \prod_{e_1,e_2\in E:|e_1\cap e_2| \text{ odd}} K(X(e_1), X(e_2)) \in \{\pm 1\}. \tag{5}$$

We can think of $B_{H,r}(X,x)$ as $B_{\tilde{H}}(X)$ of a modified graph $\tilde{H}$ obtained from H by adding a virtual hyperedge $\tilde{e}$ containing root node r and $q-1$ new nodes, and such that the chord at $\tilde{e}$ is fixed at value x. Let $b_{H,r}(x) = \mathbb{E}[B_{H,r}(X,x)]$ again when X is generated according to the law $\nu^{\otimes|E|}$. We have $|b_{H,r}(x)| \le 1$ as well. We now provide an alternative characterization of the CCG function $H(z,x)$ through the expected crossing functions b on trees. Recall that $\mathcal{T}_v^{\text{sim}}$ is the set of simple hypertrees on v nodes.

Proposition 2 (CCG graph theoretically)**.** *The function H defined in Proposition 1 satisfies*

$$H(z,x) \triangleq \sum_{v=1}^{\infty} \frac{z^v}{v!} \sum_{T\in\mathcal{T}_v^{\text{sim}}} \sum_{r\in V(T)} b_{T,r}(x) \tag{6}$$

on $\{z : |z| \le 1/e^2\} \times \Omega$.

Here the double sum $\sum_{T\in\mathcal{T}_v^{\text{sim}}} \sum_{r\in V(T)}$ is set to zero when the set $\mathcal{T}_v^{\text{sim}}$ is empty (which is the case when $q-1$ does not divide $|V|-1$). The proof of this proposition is found in Section 3.

Next we introduce two H-related functions, $\mathcal{R}$ and $\mathcal{E}$ via

$$\mathcal{R}(z) = z + z\sum_{d\ge1} \frac{1}{d!} \int_{\Omega^d} \prod_{i=1}^{d} \left[\frac{H(z,y_i)^{q-1}}{(q-1)!}\right] \prod_{1\le i<j\le d} K(y_i,y_j)\, \nu^{\otimes d}(dy_1\cdots dy_d), \tag{7}$$

$$\mathcal{E}(z) = \frac{1}{q!}\int_{\Omega} H(z,x)^q\, \nu(dx). \tag{8}$$

Both are well-defined and analytic on $\{z : |z| \le 1/e^2\}$ since $\sup_{z:|z|\le1/e^2} \sup_{x\in\Omega} |H(z,x)| < \infty$ by Proposition 1. Let

$$\rho = \left(\frac{(q-1)!\beta^2}{2^{q+1}}\right)^{1/(q-1)},$$

and let

$$\mathcal{F}(z) = \rho^{-1}\left[\mathcal{R}(\rho z) - (q-1)\mathcal{E}(\rho z)\right],$$

defined whenever both $\mathcal{R}$ and $\mathcal{E}$ are defined. As for the CCG function H, there is a graph-theoretic equivalent representation of $\mathcal{F}$, which is as follows. For every $v \ge 1$ and β define

$$\mu_v = \frac{\sum_{T\in\mathcal{T}_v^{\text{sim}}} b_T}{v!}\rho^{v-1}. \tag{9}$$

We set $\mu_v = 0$ when $\mathcal{T}_v$ is an empty set. We note that the set $\mathcal{T}_1^{\text{sim}}$ is not empty and consists of a singleton. As it turns out $\mathcal{F}$ is the generating function for the sequence μ_v. Specifically, define

$$\Phi(z) = \sum_{v\ge1} \mu_v z^v. \tag{10}$$

Our third technical preliminary is as follows.

Proposition 3. *There exists small enough $\beta^* > 0$ such that for all $\beta \in \mathbb{R}$ satisfying $|\beta| \le \beta^*$, Φ is analytic on $\{z : |z| \le 2\}$. For all such β, there exists a unique solution $t^* = t^*(\beta) \in \mathbb{R}$ of*

$$t^* \dot{\Phi}(t^*) = 1, \tag{11}$$

in the range $(1/2, 3/2)$.

The constant 2 in $\{z : |z| \le 2\}$ is a bit arbitrary and can be made as large as desired by making β^* sufficiently small. The importance of the property $t^* \in (1/2, 3/2)$ will be revealed only in proof details. We now relate $\mathcal{F}$ to Φ.

Proposition 4. *For $\beta^* > 0$ sufficiently small and $|\beta| \le \beta^*$, $\mathcal{F}(z)$ is analytic on $\{z : |z| \le 2\}$ and*

$$\Phi(z) = \mathcal{F}(z). \tag{12}$$

Note that provided β is small enough $|\rho z| \le 1/e^2$ when $|z| \le 2$ so that $\mathcal{F}$ is indeed analytic on $|z| \le 2$. We are now in a position to state our main result.

Theorem 5. *For every even q there exists small enough $\beta^* = \beta^*(q) > 0$ such that for all β satisfying $|\beta| \le \beta^*$, the following convergence holds:*

$$\lim_{n\to\infty} \frac{f_{n,q}(\beta)}{n} = \frac{\mathcal{F}(t^*) - 1 - \log t^*}{\beta} \triangleq f_q(\beta), \tag{13}$$

where t^* is defined in (11).

We compute f_q numerically and we find that $f_q(\beta) > 0$ for all $0 < \beta \le \beta^*$. We leave it as an open question to verify positivity analytically. A simple consequence of the fact $f_q(\beta) > 0$ for some $\beta > 0$ is that the ground state energy is of the order $\Theta(n)$ w.h.p. Indeed, let $\lambda_{\max}$ be the ground state energy of the Hamiltonian H_{SYK}, namely the largest eigenvalue of H_{SYK}. As the dimension of the space is $2^{\frac{n}{2}}$ we have $Z_{n,q} \le 2^{-\frac{n}{2}} 2^{\frac{n}{2}} \exp(\beta\lambda_{\max}) = \exp(\beta\lambda_{\max})$ leading to

$$F_{n,q}(\beta) \le \frac{1}{\beta} \log\left(\exp(\beta\lambda_{\max})\right) \le \lambda_{\max}.$$

The convergence of $F_{n,q}$ w.h.p. to $f_q(\beta) > 0$ implies $\lambda_{\max} = \Theta(n)$ w.h.p.

2 Proof of the main result Theorem 5

2.1 Power series expansion of the free energy

Given any two superindices $I, J \in \binom{[n]}{q}$, applying (1) we have $\psi_I \psi_J = (-1)^{|I\cap J|} \psi_J \psi_I$. Suppose we have any ordered sequence $(I_1, \dots, I_m) \in \binom{[n]}{q}^m$, such that each superindex I appears an even number of times in this sequence. That is, for every $k \in [m]$ there exists an odd number of indices $[m] \ni r \ne k$ such that $I_k = I_r$. Then from $\psi_i^2 = \frac{1}{2}\mathbb{I}$ we have $\psi_I^2 = \frac{1}{2^q}\mathbb{I}$ (recall that q is

even) and $\prod_{k\in[m]} \psi_{I_k} = \pm 2^{-qm/2}\mathbb{I}$, where the sign of the expression is determined by the ordering $(I_1, \ldots, I_m)$. We denote this multiplier ± 1 by $B(I_1, \ldots, I_m)$ so that

$$\prod_{k\in[m]} \psi_{I_k} = B(I_1, \ldots, I_m) 2^{-qm/2}\mathbb{I}. \tag{14}$$

Anticipating a forthcoming derivation, the function $B(I_1, \ldots, I_m)$ will be related to $B_H(X)$ introduced earlier, when the multiplicity of each superindex I is exactly two.

Consider any q-uniform multi-hypergraph $H = ([n], \bar{m})$ on the node set $V = [n]$. For each $I \in \binom{[n]}{q}$ its multiplicity is denoted by m_I and $\bar{m} = (m_I, I \in \binom{[n]}{q})$. Let $w(H)$ be the set of all ordered sequences $(I_1, \ldots, I_m)$ which induce precisely the multi-hypergraph H. That is $w(H)$ is simply the set of all ordered sequences $(I_1, \ldots, I_m)$ such that each I appears exactly m_I times in this sequence. In particular, $m = \sum_I m_I$. Let

$$b(H) = \sum_{(I_1,\ldots,I_m)\in w(H)} B(I_1, \ldots, I_m). \tag{15}$$

The values $b(H)$ will be the key building block in our proof. We note that they factorize over connected components. That is, let $H_1, \ldots, H_R$ denote the connected components of a multi hypergraph H. Recall that $m = m(H)$ denotes the sum of edge multiplicities of a multi hypergraph H. Then

$$b(H) = \binom{m(H)}{m(H_k), k \in [R]} \prod_{1\le k\le R} b(H_k). \tag{16}$$

This holds since operators ψ_I commute over disjoint superindices I. Returning to the partition function, we have

$$\mathbb{E}[Z_{n,q}] = \mathbb{E}[\mathrm{tr}\big(e^{\beta H}\big)] = \sum_{r\ge 0} \frac{\beta^{2r}}{(2r)!} \mathbb{E}[\mathrm{tr}\big(H^{2r}\big)].$$

Here we use the fact that the expectation for odd r vanishes, since odd moments of $\bar{J}_I$ vanish. We expand $\mathbb{E}[H^{2r}]$ using (14) and the fact that $\mathbb{E}[\bar{J}_I^m] = 1$ for even m and $\mathbb{E}[\bar{J}_I^m] = 0$ for odd m. We obtain

$$\begin{aligned}
\mathbb{E}[\mathrm{tr}\big(H^{2r}\big)] &= \sum_{I_1,\ldots,I_{2r}\in\binom{[n]}{q}} \mathbb{E}[\prod_{k\in[2r]} J_{I_k}] \,\mathrm{tr}\left(\prod_{k\in[2r]} \psi_{I_k}\right) \\
&= \sum_{I_1,\ldots,I_{2r}\in\binom{[n]}{q}} \mathbb{E}[\prod_{k\in[2r]} J_{I_k}] B(I_1, \ldots, I_{2r}) 2^{-rq} \\
&= \sum_H 2^{-qm(H)/2} \sigma^{m(H)} b(H),
\end{aligned}$$

where the sum is over all multi hypergraphs H such that $m(H) = 2r$ and all multiplicities $m_I(H)$ are even. In light of (16) we obtain

$$\mathbb{E}[\mathrm{tr}\big(H^{2r}\big)] = (2r)! \sum_H \prod_{1\le k\le C} \frac{1}{m(H_k)!} 2^{-qm(H_k)/2} \sigma^{m(H_k)} b(H_k),$$

where $H_1, \ldots, H_C$ is the connected component decomposition of H. Here we used $m(H) = \sum_k m(H_k)$. Recall that given a subset $C \subset [n]$, $\mathcal{H}(C)$ denotes the set of all connected multi-hypergraphs H with the node set C with even multiplicities. Then we can rewrite the above as

$$\mathbb{E}[\mathrm{tr}\big(H^{2r}\big)] = \sum_{C_0, C_1, \ldots, C_R} \sum_{H_k \in \mathcal{H}(C_k), k \in [R]} (2r)! \prod_{k \in [R]} \frac{1}{m(H_k)!} 2^{-qm(H_k)/2} \sigma^{m(H_k)} b(H_k),$$

where the first sum is over all partitions of the node set $[n]$ into subsets $C_0, C_1, \ldots, C_R$, and the second sum is over all tuples $H_1, \ldots, H_R$ with $H_k \in \mathcal{H}(C_k)$ such that $\sum_{k \in [R]} m(H_k) = 2r$. In particular $\cup_{1 \le k \le R} C_k$ is the union of the supports of the multi hypergraphs H_k, and $C_0 = [n] \setminus \cup_{1 \le k \le R} C_k$. Combining and incorporating β into the product terms, we obtain

$$\mathbb{E}[Z_{n,q}] = \sum_{C_0, C_1, \ldots, C_R} \sum_{H_k, k \in [R]} \prod_{k \in [R]} \frac{1}{m(H_k)!} (\beta 2^{-q/2} \sigma)^{m(H_k)} b(H_k).$$

Given $v \ge 1, \Delta \ge 0$ recall that $\mathcal{H}_{v,\Delta} \subset \mathcal{H}([v])$ is the subset of $\mathcal{H}([v])$ consisting of multi hypergraphs H with precisely $m(H) = 2(v - 1 + \Delta)/(q-1)$ hyperedges and Δ is the excess of H. We then rewrite the above as

$$\mathbb{E}[Z_{n,q}] = \sum_{C_0, C_1, \ldots, C_R} \sum_{\Delta_1, \ldots, \Delta_R \ge 0} \prod_{k \in [R]} \left(\sum_{H \in \mathcal{H}_{|C_k|, \Delta_k}} \frac{1}{m(H)!} (\beta 2^{-q/2} \sigma)^{m(H)} b(H) \right).$$

Introduce

$$\mu_{v,\Delta} = \frac{n^{v-1}}{v!} \sum_{H \in \mathcal{H}_{v,\Delta}} \frac{1}{m(H)!} (\beta 2^{-q/2} \sigma)^{m(H)} b(H). \tag{17}$$

The extra factor $n^{v-1}/v!$ is introduced for later convenience. We stress a distinction between $\mu_{v,\Delta}$ and μ_v defined in (9). As will be established later, $\mu_v = \mu_{v,0}$. We also define $\mu_{v,\Delta} = 0$ when $\mathcal{H}_{v,\Delta}$ is an empty set. For example this is the case when $2 \le v < q$ since the graph H cannot contain any hyperedges and thus is not connected. When $v = 1, \Delta = 0$, which corresponds to a (nominally) connected graph on a single node, we set $\mu_{1,0} = 1$. Then

$$\mathbb{E}[Z_{n,q}] = \sum_{C_0, C_1, \ldots, C_R} \sum_{\Delta_1, \ldots, \Delta_R \ge 0} \prod_{k \in [R]} (|C_k|!) n^{-(|C_k|-1)} \mu_{|C_k|, \Delta_k}.$$

Fix any set of integers $(k_{v,\Delta}, v \ge 1, \Delta \ge 0)$ satisfying

$$\sum_{v,\Delta} v k_{v,\Delta} = n. \tag{18}$$

Consider the set of partitions $(C_{v,\Delta,j}, v \ge 1, \Delta \ge 0, j \in [k_{v,\Delta}])$ of $[n]$ such that $|C_{v,\Delta,j}| = v$. The total number of such partitions is

$$\frac{n!}{\prod_{v,\Delta} (v!)^{k_{v,\Delta}} k_{v,\Delta}!}. \tag{19}$$

These partitions represent partitions $C_0, C_1, \ldots, C_R$ where $C_{v,\Delta,j}, j \in [k_{v,\Delta}]$ support graphs in $\mathcal{H}_{v,\Delta}$ and C_0 represent nodes in $[n]$ not supporting any graphs. Then after cancellation of $v!$ and $|C_k|!$ we obtain

$$\begin{aligned}\mathbb{E}[Z_{n,q}] &= n! \sum_{k_{v,\Delta}} \prod_{v,\Delta} \frac{1}{n^{(v-1)k_{v,\Delta}}} \frac{1}{k_{v,\Delta}!} \mu_{v,\Delta}^{k_{v,\Delta}} \\ &= n! \sum_{k_{v,\Delta}} \frac{1}{n^{\sum_{v,\Delta}(v-1)k_{v,\Delta}}} \prod_{v,\Delta} \frac{1}{k_{v,\Delta}!} \mu_{v,\Delta}^{k_{v,\Delta}} \end{aligned} \tag{20}$$

where the sum is over all $(k_{v,\Delta}, v \geq 1, \Delta \geq 0)$ satisfying (18). In particular, $k_{1,0}$ represents the number of nodes not supporting graphs, hence the choice $\mu_{1,0} = 1$.

Our next goal is to represent this expression as coefficients of a generating function. Introduce functions $M_0, \Delta M : \mathbb{C} \to \mathbb{C} \cup \{\infty\}$ defined by

$$M_0(z) = \exp\left(n \sum_{v \geq 1} \mu_{v,0} z^v\right),$$

$$\Delta M(z) = \exp\left(n \sum_{1 \leq v \leq n, \Delta \geq 1} \mu_{v,\Delta} z^v\right),$$

We note that while in the first expression the range of $v \geq 1$ is unrestricted, in the second expression it is restricted to $v \leq n$. This will be dictated by the fact that the first will be analytic for the infinite sum $\sum_{v \geq 1}$ for sufficiently bounded $|z|$, while for the second expression analyticity will hold only for the final sum $\sum_{1 \leq v \leq n}$. Let $M(z) = M_0(z)\Delta M(z)$.

Proposition 6. *There exists $\beta^* > 0$ small enough such that for all β satisfying $|\beta| \leq \beta^*$, $M_0, \Delta M$ are analytic on $\{z : |z| \leq 2\}$. Furthermore, let M_n be the coefficient of M in front of z^n, so that $M(z) = \sum_{n \geq 0} M_n z^n$. Then $\mathbb{E}[Z_{n,q}] = \frac{n!}{n^n} M_n$.*

Proof. The proof of analyticity is delayed and will be subsumed by stronger claims, in particular Lemma 8 covering the analyticity of M_0 and Lemma 9 covering the analyticity of ΔM. We now establish the second part. We have

$$M(z) = \prod_{v \geq 1, \Delta \geq 0} \exp\left(n \mu_{v,\Delta} z^v\right),$$

where for $\Delta \geq 1$ the product is restricted to $v \leq n$. Then

$$\begin{aligned} M(z) &= \prod_{v \geq 1, \Delta \geq 0} \left(\sum_{k \geq 0} \frac{\left(n\mu_{v,\Delta} z^v\right)^k}{k!}\right) \\ &= \sum_{k_{v,\Delta}} \prod_{v \geq 1, \Delta \geq 0} \frac{\left(n\mu_{v,\Delta} z^v\right)^{k_{v,\Delta}}}{k_{v,\Delta}!}, \end{aligned}$$

where as before, terms with $\Delta \geq 1$ are restricted to $v \leq n$. The coefficient of z^n is the sum above set to $z = 1$ and restricted to tuples $(k_{v,\Delta}, v \geq 1, \Delta \geq 0)$ satisfying (18). With this restriction

$$\prod_{v \geq 1, \Delta \geq 0} n^{k_{v,\Delta}} = \prod_{v \geq 1, \Delta \geq 0} n^{(v-(v-1))k_{v,\Delta}} = n^n \prod_{v \geq 1, \Delta \geq 0} \frac{1}{n^{(v-1)k_{v,\Delta}}}.$$

Multiplying this restricted sum by $\frac{n!}{n^n}$ we obtain (20). □

Our next goals are (a) simplifying M_0 and analyzing its scaling limit, and (b) establishing that the contribution of ΔM is vanishing.

2.2 Contribution from trees

In light of Proposition 6, we focus on estimating $M(z)$. In this section we estimate the contribution from M_0 and show how it leads to the final expression (13) in our main result. In the following subsection we combine this estimation with the vanishing contribution from ΔM to complete the main result.

Recall that $\mathcal{T}_v^{\text{sim}}$, (respectively $\mathcal{T}_{v,2}$) is the set of all simple hypertrees (respectively multi hypertrees with multiplicity 2) on v nodes. Recall the definition (4) of b_H in terms of uniformly generated chords. Recall also the definition (15) of $b(H)$ which is in terms of permutations of superindices. We now relate b_H and $b(H)$.

Lemma 7. *Given $H \in \mathcal{H}_v^{\text{sim}}$ consider the corresponding multi-hypergraph in $\mathcal{H}_{v,2}$ with multiplicity 2 for every hyperedge, which we denote by H as well with some abuse of notation. Then*

$$b(H) = b_H \frac{m(H)!}{2^{\frac{m(H)}{2}}}. \tag{21}$$

Proof. $H \in \mathcal{H}_v^{\text{sim}}$ let E^{sim} be its set of (distinct) hyperedges, so that for H viewed as an element of $\mathcal{H}_{v,2}$ its set of hyperedges E is E^{sim} each repeated twice. Consider any ordered sequence $\tau = (I_1, \dots, I_{2|E|})$ of $2|E|$ hyperedges of the multi hypergraph version of H. The total number of such sequences is $|E|!/2^{|E|/2}$. Define $X_\tau : E^{\text{sim}} \to [|E|]^2$ as follows. For each $e \in E^{\text{sim}}$, $X(e) = (u^1, u^2)$ if the first copy of the edge e in the permutation τ appears in position u^1 and the second copy in position u^2. Given X_τ generate a random $X : E \to \Omega$ where each u^j is mapped to a point in $[\frac{u^j-1}{|E|}, \frac{u^j}{|E|}]$ chosen uniformly at random. Suppose τ is also chosen u.a.r. The distribution of X is then $\nu^{\otimes |E|}$. The relation (21) then follows since

$$b(H) \left(\frac{m(H)!}{2^{\frac{m(H)}{2}}} \right)^{-1}$$

is the expectation of $B(I_1, \dots, I_{2|E|})$ with respect to uniformly random τ. □

Next we validate $\mu_{v,0} = \mu_v$ for the case of multi hypertrees $T \in \mathcal{T}_{v,2}$ where μ_v was defined in (9). By definition we have $\mu_{1,0} = \mu_1$. For the remaining cases, applying Lemma 7 we have

$$\begin{aligned} \mu_{v,0} &= \frac{n^{v-1}}{v!} \sum_{T \in \mathcal{T}_{v,2}} \frac{1}{m(T)!} (\beta^2 2^{-q} \sigma^2)^{m(T)/2} b(H) \\ &= \frac{n^{v-1}}{v!} \sum_{T \in \mathcal{T}_v^{\text{sim}}} (\beta^2 2^{-q-1} \sigma^2)^{m(T)} b_T. \end{aligned}$$

For every $T \in \mathcal{T}_v^{\text{sim}}$ we have $m(T) = (v-1)/(q-1)$. Then

$$n^{v-1} (\sigma^2)^{m(T)} = n^{v-1} \left(\frac{(q-1)!}{n^{q-1}} \right)^{\frac{v-1}{q-1}} = ((q-1)!)^{\frac{v-1}{q-1}}.$$

We obtain:

$$\begin{aligned}\mu_{v,0} &= \frac{(\beta^2 2^{-q-1}(q-1)!)^{\frac{v-1}{q-1}}}{v!} \sum_{H\in\mathcal{T}_v^{\text{sim}}} b_H \\ &= \frac{\rho^{v-1}}{v!} \sum_{H\in\mathcal{T}_v^{\text{sim}}} b_H \\ &= \mu_v.\end{aligned}$$

Our next goal is proving Proposition 3.

Proof of Proposition 3. We begin by establishing some bounds for Φ.

Lemma 8. *For every $c>0$ there exists small enough $\beta^*>0$ such that for all β with $|\beta|\le\beta^*$ the function $\Phi(z)$ (defined by (10)) is analytic on $|z|\le 2$. Furthermore*

$$\sup_{|z|\le 2} |\Phi(z)-z| \le c, \tag{22}$$

$$\sup_{|z|\le 2} \left|\dot\Phi(z)-1\right| \le c, \tag{23}$$

$$\sup_{|z|\le 2} |z|^2\left|\ddot\Phi(z)\right| \le c. \tag{24}$$

Proof. Since $|b_H|\le 1$ and $\left|\mathcal{T}_v^{\text{sim}}\right| = \frac{(v-1)!v^{s-1}}{s!((q-1)!)^s}$, where $s=(v-1)/(q-1)$ is the number of hyperedges. By Stirling's approximation we have

$$\begin{aligned}|\mu_v| &\le (1+o(1))e^v\left(\frac{(q-1)!\beta^2}{2^{q+1}}\right)^{(v-1)/(q-1)} \\ &\le (1+o(1))e\left(e\left(\frac{(q-1)!\beta^2}{2^{q+1}}\right)^{1/(q-1)}\right)^{v-1} \\ &\triangleq (1+o(1))eg^{v-1}(\beta).\end{aligned} \tag{25}$$

We choose β^* small enough so that when $|\beta|\le\beta^*$, $\sum_v g^v(\beta)2^v<\infty$. Then, recalling $\mu_1=1$ and $\mu_v=0$ when $v\ge 2$ and $\mathcal{T}_v$ is empty, we obtain $|\Phi(z)| = |z+\sum_{v\ge2}\mu_v z^v| \le 2+\sum_v g^v(\beta)2^v<\infty$, and thus Φ is analytic on $|z|\le 2$.

We now establish the claimed bounds. We have $\Phi(z)-z=\sum_{v\ge2}\mu_v z^v$. Given $c>0$, we can find β^* small enough so that $|g(\beta^*)|$ in (25) satisfies $\sum_{v\ge2}(1+o(1))eg^{v-1}(\beta^*)2^v\le c$ for every $v\ge 2$. Namely (22) holds.

Next, $\dot\Phi(z)-1=\sum_{v\ge2} v\mu_v z^{v-1}$. Using a very similar argument we find β^* small enough so that (23) holds. Finally, $\ddot\Phi(z)=\sum_{v\ge2} v(v-1)\mu_v z^{v-2}$. A similar argument is used to find β^* small enough so that (24) holds as well. □

We return to the proof of the proposition and show that for β^* sufficiently small $F(t)\triangleq t\dot\Phi(t)-1$ has a unique zero t^* in the interval $(1/2,3/2)$. Choose $c<1/3$. We find β^* small enough so that by Lemma 8 $\frac12(\dot\Phi\left(\frac12\right)-1)\le c/2$, implying $\frac12\dot\Phi\left(\frac12\right)-1\le c/2-1/2<0$. Similarly,

$\frac{3}{2}(\dot{\Phi}\left(\frac{3}{2}\right)-1) \geq -3c/2$, implying $\frac{3}{2}\dot{\Phi}\left(\frac{3}{2}\right)-1 \geq 1/2-3c/2 > 0$. Thus F has zero in the interval $[1/2, 3/2]$. To show uniqueness, again using Lemma 8, we note that for $t \in [1/2, 3/2]$

$$\dot{F}(t) = \dot{\Phi}(t) + t\ddot{\Phi}(t) \geq 1 - c - \frac{1}{(1/2)}c = 1 - 3c > 0.$$

Therefore F is strictly increasing and the root is unique. □

2.3 Contribution from non-trees

We turn to analyzing ΔM. Define

$$B(z) = \log \Delta M(z) = n \sum_{1 \leq v \leq n, \Delta \geq 1} \mu_{v,\Delta} z^v.$$

Lemma 9. *There exists small enough $\beta^* > 0$ so that when $|\beta| \leq \beta^*$*

$$\sup_{n \geq 1} \sup_{|z| \leq 2} |B(z)| \leq 1, \quad \sup_{n \geq 1} \sup_{|z| \leq 2} |\dot{B}(z)| \leq 1. \tag{26}$$

Proof. We begin by establishing the following bound. For all $v \geq 1$ and $\Delta \geq 1$ we claim

$$\sum_{H \in \mathcal{H}_{v,\Delta}} \frac{|b(H)|}{m(H)!} \leq (2e)^{v+\Delta} v^{v+\Delta}. \tag{27}$$

Recall that $\mathcal{H}_{v,\Delta} = \emptyset$ when $\Delta \geq 1$ and $v < q$, and thus we may assume $v \geq q$. To show the claimed bound it suffices to show

$$|\mathcal{H}_{v,\Delta}| \leq (qe)^{v+\Delta} v^{v+\Delta},$$

since $|b(H)| \leq m(H)!$. Recall that the total number of hyperedges in a graph $H \in \mathcal{H}_{v,\Delta}$ is $E \triangleq 2\frac{v-1+\Delta}{q-1}$ by definition. The number of distinct possible hyperedges is $\binom{v}{q}$. Let $2m_1, \ldots, 2m_{\binom{v}{q}}$ be their multiplicities (including the possibility $m_j = 0$), which have to be even by definition of $\mathcal{H}_{v,\Delta}$. Then $2m_1 + 2m_2 + \cdots + 2m_{\binom{v}{q}} = E$, implying

$$|\mathcal{H}_{v,\Delta}| = \binom{\binom{v}{q} + \frac{E}{2} - 1}{\binom{v}{q} - 1}.$$

Let $\bar{v} = \binom{v}{q}$. We have

$$\begin{aligned}
\binom{\bar{v} + \frac{E}{2} - 1}{\bar{v} - 1} &= \binom{\bar{v} + \frac{E}{2} - 1}{\frac{E}{2}} \\
&\leq \left(\frac{e(\bar{v} + \frac{E}{2})}{\frac{E}{2}} \right)^{\frac{E}{2}} \\
&= e^{\frac{E}{2}} \left(1 + \frac{\bar{v}}{\frac{E}{2}} \right)^{\frac{E}{2}}.
\end{aligned}$$

We have

$$\frac{\bar{v}}{\frac{E}{2}} = \frac{\binom{v}{q}}{\frac{v-1+\Delta}{q-1}} = \frac{v(v-1)\cdots(v-q+1)}{q!}\frac{q-1}{v-1+\Delta}$$

Since $\Delta \geq 1$ we have $v-1+\Delta \geq v$ and thus we obtain a bound

$$\frac{v(v-1)\cdots(v-q+1)}{q!}\frac{q}{v} \leq v^{q-1}.$$

Then

$$\begin{aligned} e^{\frac{E}{2}}\left(1+\frac{\bar{v}}{\frac{E}{2}}\right)^{\frac{E}{2}} &\leq e^{\frac{E}{2}}\left(1+v^{q-1}\right)^{\frac{E}{2}} \\ &\leq (2e)^{\frac{E}{2}}\left(v^{q-1}\right)^{\frac{v-1+\Delta}{q-1}} \\ &= (2e)^{\frac{E}{2}}v^{(v-1+\Delta)} \\ &\leq (2e)^{v+\Delta}v^{v+\Delta}. \end{aligned}$$

This completes the proof of (27).

We now return to the proof of the lemma. Recall that $\mu_{v,\Delta} = 0$ when $\Delta \geq 1$ and $v < q$. We have

$$\mu_{v,\Delta} = \sum_{H\in\mathcal{H}_{v,\Delta}} \frac{1}{\left(2\frac{v-1+\Delta}{q-1}\right)!}\frac{n^{v-1}}{v!}(\beta^2 2^{-q}\sigma^2)^{\frac{v-1+\Delta}{q-1}}b(H).$$

Then

$$|\mu_{v,\Delta}| \leq \frac{\sum_{H\in\mathcal{H}_{v,\Delta}}|b(H)|}{\left(2\frac{v-1+\Delta}{q-1}\right)!}\frac{n^{v-1}}{v!}(\beta^2 2^{-q}\sigma^2)^{\frac{v-1+\Delta}{q-1}}.$$

By (27) it is at most

$$\begin{aligned} |\mu_{v,\Delta}| &\leq (2e)^{v+\Delta}v^{v+\Delta}\frac{n^{v-1}}{v!}(\beta^2 2^{-q}\sigma^2)^{\frac{v-1+\Delta}{q-1}} \\ &= (2e)^{v+\Delta}v^{v+\Delta}\frac{n^{v-1}}{v!}(\beta^2 2^{-q}(q-1)!n^{-(q-1)})^{\frac{v-1+\Delta}{q-1}} \\ &= (2e)^{v+\Delta}v^{v+\Delta}\frac{n^{v-1}}{v!}(\beta^2 2^{-q}(q-1)!)^{\frac{v-1+\Delta}{q-1}}\frac{1}{n^{v-1+\Delta}} \\ &= (2e)^{v+\Delta}v^{v+\Delta}\frac{1}{v!}(\beta^2 2^{-q}(q-1)!)^{\frac{v-1+\Delta}{q-1}}\frac{1}{n^{\Delta}} \end{aligned}$$

Then

$$\max_{|z|\leq 2}|B(z)| \leq n\sum_{q\leq v\leq n,\Delta\geq 1} 2^v|\mu_{v,\Delta}| \leq n(4e(\beta^2 2^{-q}(q-1)!)^{\frac{1}{q-1}})^{v+\Delta}v^{v+\Delta}\frac{1}{v!}\frac{1}{n^{\Delta}}$$

We find universal constant $c \geq 1$ so that $\frac{v^v}{v!} \leq c^v$ for all $v \geq 1$. Denote $(4e)c(\beta^2 2^{-q}(q-1)!)^{\frac{1}{q-1}}$ by $g(\beta)$. We obtain a bound

$$\max_{|z|\leq 2} |B(z)| \leq (g(\beta))^{v+\Delta} v^{\Delta} \frac{1}{n^{\Delta-1}}$$

For every fixed $v \leq n$ consider the sum

$$\sum_{\Delta \geq 1} \left(\frac{g(\beta)v}{n}\right)^{\Delta-1} = \frac{1}{1-\frac{g(\beta)v}{n}} \leq 2,$$

provided β small enough so that $g(\beta)v/n \leq g(\beta) \leq 1/2$. We obtain a bound

$$\max_{|z|\leq 2} |B(z)| \leq 2 \sum_{q \leq v \leq n} (g(\beta))^{v+1} v$$

which we arrange to be less than 1 by taking β sufficiently small.

The derivation for $\dot{B}(z) = n \sum_{1 \leq v \leq n, \Delta \geq 1} \mu_{v,\Delta} v z^{v-1}$ is similar. □

2.4 Proof completion

Recalling Proposition 6, we now focus on asymptotics of M_n. We fix β^* small enough and find the corresponding t^* as in Proposition 3. By Cauchy's integral formula on $z = t^* e^{i\theta}$, we can write M_n as

$$\begin{aligned} M_n &= \frac{1}{2\pi i} \oint_{|z|=t^*} \frac{M(z)}{z^{n+1}} \, dz \\ &= \frac{1}{2\pi i} \oint_{|z|=t^*} \frac{\exp(n\Phi(z)) \Delta M(z)}{z^{n+1}} \, dz \\ &= \frac{(t^*)^{-n}}{2\pi} \int_{-\pi}^{\pi} \exp\left(n\Psi(\theta) + B(t^* e^{i\theta})\right) d\theta, \end{aligned}$$

where we define $\Psi(\theta) = \Phi(t^* e^{i\theta}) - i\theta$ and recall that $B(z) = \log \Delta M(z)$. The remainder of the section is devoted to establishing the following proposition, from which the proof of Theorem 5 will be an easy consequence.

Proposition 10. *The following holds*

$$\lim_n \frac{1}{n} \log M_n = -\log t^* + \Psi(0) = -\log t^* + \Phi(t^*).$$

Proof. The proof is based on standard saddle point method. In particular, we will show that $\theta = 0$ is the unique maximizer of Ψ, and show that the contribution from B is negligible, due to Lemma 9.

2.4.1 Proof of upper bound.

We first establish

$$\limsup_n \frac{1}{n} \log M_n \le -\log t^* + \Psi(0). \tag{28}$$

Since $t^*\dot{\Phi}(t^*) = 1$, we have $\dot{\Psi}(0) = it^*\dot{\Phi}(t^*) - i = 0$. Next

$$\begin{aligned}\ddot{\Psi}(\theta) &= -t^* e^{i\theta}\dot{\Phi}(t^* e^{i\theta}) - (t^* e^{i\theta})^2 \ddot{\Phi}(t^* e^{i\theta}) \\ &= -t^* e^{i\theta} + t^* e^{i\theta}\left(1 - \dot{\Phi}(t^* e^{i\theta})\right) - (t^* e^{i\theta})^2 \ddot{\Phi}(t^* e^{i\theta}).\end{aligned} \tag{29}$$

Using bounds from Lemma 8, which apply since $|t^* e^{i\theta}| = t^* \in [1/2, 3/2]$, we obtain

$$\begin{aligned}|1 - \dot{\Phi}(t^* e^{i\theta})| &\le c, \\ |t^* e^{i\theta}|^2 |\ddot{\Phi}(t^* e^{i\theta})| &\le c.\end{aligned}$$

Thus

$$\operatorname{Re} \ddot{\Psi}(\theta) \le -\operatorname{Re}(t^* e^{i\theta}) + (3/2)c + c = -t^* \cos(\theta) + (5/2)c.$$

Assume $|\theta| \le 1$. Then $t^* \cos(\theta) \ge (1/2)\cos(1)$. We take c small enough so that $-(1/2)\cos(1) + (5/2)c \le -1/4$ and obtain that $\operatorname{Re} \ddot{\Psi}(\theta) \le -1/4$ in the range $|\theta| \le 1$. Then in this range by Taylor expansion we obtain

$$\operatorname{Re} \Psi(\theta) \le \Psi(0) - \theta^2/8 = \Phi(t^*) - \theta^2/8. \tag{30}$$

Next we bound $\operatorname{Re} \Psi(\theta)$ in the remaining range $|\theta| \ge 1$. We have

$$\begin{aligned}\operatorname{Re} \Psi(\theta) &= \operatorname{Re} \Phi(t^* e^{i\theta}) \\ &= \Phi(t^*) - t^*(1 - \cos\theta) - (\Phi(t^*) - t^*) + \operatorname{Re}\left(\Phi(t^* e^{i\theta}) - t^* e^{i\theta}\right)\end{aligned}$$

Applying Lemma 8

$$|\Phi(t^*) - t^*|, |\Phi(t^* e^{i\theta}) - t^* e^{i\theta}| \le c.$$

Using $t^* \ge 1/2$, we obtain an upper bound

$$\operatorname{Re} \Psi(\theta) \le \Phi(t^*) - (1/2)(1 - \cos(1)) + 2c.$$

We assume $c > 0$ is small enough so that the right-hand side is at most $\Phi(t^*) - 1/5$. Thus when $|\theta| \ge 1$ we obtain a bound

$$\operatorname{Re} \Psi(\theta) \le \Psi(0) - 1/5. \tag{31}$$

We now complete the proof of (28). We assume β^* small enough so that bounds (26) from Lemma 9 apply. Then

$$\begin{aligned} M_n &= \frac{(t^*)^{-n}}{2\pi}\int_{-\pi}^{\pi} \mathrm{Re}\left(\exp\left(n\Psi(\theta)+B(t^*e^{i\theta})\right)\right)d\theta \\ &\le \frac{(t^*)^{-n}}{2\pi}\int_{-\pi}^{\pi} \exp\left(\mathrm{Re}\left(n\Psi(\theta)\right)+B(t^*e^{i\theta})\right)\right)d\theta \\ &\le \frac{(t^*)^{-n}}{2\pi}\left(\int_{|\theta|\le 1} e^{n\Phi(t^*)-n\theta^2/8+1}d\theta + \int_{|\theta|\ge 1,\theta\in[-\pi,\pi]} e^{n\Phi(t^*)-n/5+1}d\theta\right) \\ &\le (t^*)^{-n}e^{n\Phi(t^*)}\frac{e}{2\pi}\left(\sqrt{\frac{8\pi}{n}}+2\pi e^{-n/5}\right), \end{aligned}$$

and (28) follows.

2.4.2 Proof of lower bound.

We now establish

$$\liminf_n \frac{1}{n}\log M_n \ge -\log t^* + \Psi(0). \tag{32}$$

For this we partition the integrated region $[-\pi,\pi]$ into three regions. Introduce constant $\eta \in (0,1], L \ge 1$ to be specified later. The choice of η will depend on β^*. The choice of L will be universal. We assume that n is sufficiently large so that $1/\sqrt{n} \le \eta$. Our regions will be $|\theta| \le L/\sqrt{n}, L/\sqrt{n} \le \theta \le 1$ and $|\theta| \ge 1$. The choice of L in particular will be only relevant to the regime $L/\sqrt{n} \le \theta \le 1$.

1. $|\theta| \le L/\sqrt{n}$. We have from (29)

$$\ddot{\Psi}(0) = -1 - (t^*)^2\ddot{\Phi}(t^*) \in \mathbb{R}.$$

Applying Lemma 8 $1+\ddot{\Psi}(0) \in [-c,c]$. We find sufficiently small $\eta > 0$ such that when $|\theta| \le \eta$ we have $1+\mathrm{Re}(\ddot{\Psi}(\eta)) \in [-2c,2c]$. Recall that $\dot{\Psi}(0) = 0$. Then by Taylor expansion,

$$\Phi(t^*) - \frac{1}{2}(1+2c)\theta^2 \le \mathrm{Re}\,\Psi(\theta) \le \Phi(t^*) - \frac{1}{2}(1-2c)\theta^2.$$

We assume c is small enough so that

$$\Phi(t^*) - \theta^2 \le \mathrm{Re}\,\Psi(\theta) \le \Phi(t^*) - \frac{1}{8}\theta^2. \tag{33}$$

Next we control the imaginary part of Ψ. Since $\dot{\Psi}(0) = 0$ and $\ddot{\Psi}(0)$ does not have an imaginary part we have

$$|\mathrm{Im}(\Psi(\theta) - \Phi(t^*))| \le |\theta|^3 \cdot \frac{1}{6}\sup_{|w|\le\eta}\left|\frac{d^3\Psi(w)}{dw^3}\right| \triangleq C|\theta|^3.$$

Here C depends only on the choice of β^* and η. Applying Lemma 9

$$\left|\mathrm{Im}(B(t^*e^{i\theta}) - B(t^*))\right| \le \left|t^*e^{i\theta} - t^*\right| \sup_\theta |\dot{B}(t^*e^{i\theta})| \le \frac{3}{2}2|\theta| = 3|\theta|.$$

Assume n sufficiently large such that $L/\sqrt{n} \le \eta$ and

$$\frac{CL^3 + 3L}{\sqrt{n}} \le \frac{\pi}{3}, \tag{34}$$

Then for $|\theta| \le L/\sqrt{n}$

$$\left|\mathrm{Im}\left(n\left(\Psi(\theta) - \Phi(t^*)\right) + B(t^*e^{i\theta}) - B(t^*)\right)\right| \le Cn|\theta|^3 + 3|\theta| \le \frac{\pi}{3}.$$

Using $\cos(\pi/3) = 1/2$ we have

$$\mathrm{Re}\int_{|\theta|\le L/\sqrt{n}} e^{n\Psi(\theta)+B(t^*e^{i\theta})}d\theta \ge \frac{1}{2}\int_{|\theta|\le L/\sqrt{n}} e^{\mathrm{Re}\, n\Psi(\theta)+\mathrm{Re}\, B(t^*e^{i\theta})}d\theta.$$

By Lemma 9 $|B(t^*) - B(t^*e^{i\theta})| \le |\theta| < 1$. Then $|\,\mathrm{Re}(B(t^*e^{i\theta})) - \mathrm{Re}(B(t^*))| < 1$, and since $B(t^*)$ is real valued

$$\mathrm{Re}(B(t^*e^{i\theta})) \ge B(t^*) - 1 \ge -2,$$

where in the last inequality Lemma 9 is invoked again. Then

$$\int_{|\theta|\le L/\sqrt{n}} e^{\mathrm{Re}\, n\Psi(\theta)+\mathrm{Re}\, B(t^*e^{i\theta})}d\theta \ge e^{-2}\int_{|\theta|\le L/\sqrt{n}} e^{\mathrm{Re}\, n\Psi(\theta)}d\theta.$$

Applying the left part of (33) the right-hand side above is at least

$$e^{-2}e^{n\Phi(t^*)}\int_{|\theta|\le L/\sqrt{n}} e^{-n\theta^2}d\theta = e^{n\Phi(t^*)}\frac{e^{-2}}{\sqrt{n}}\int_{-L}^{L} e^{-u^2}du \triangleq K_n. \tag{35}$$

We note

$$\lim_n \frac{1}{n}\log K_n = \Phi(t^*). \tag{36}$$

2. $L/\sqrt{n} \le \theta \le 1$. For this regime, we have by (30) and Lemma 9

$$\begin{aligned}\int_{L/\sqrt{n}\le|\theta|\le 1} \left|e^{n\Psi(\theta)+B(t^*e^{i\theta})}\right|d\theta &\le e^{1+n\Phi(t^*)}\int_{L/\sqrt{n}\le|\theta|\le 1} e^{-n\theta^2/8}d\theta \\ &\le \frac{e}{\sqrt{n}}e^{n\Phi(t^*)}\int_{|u|\ge L} e^{-u^2/8}du.\end{aligned}$$

We choose L large enough so that this upper bound is at most $(1/3)K_n$. Namely

$$\frac{e}{\sqrt{n}}e^{n\Phi(t^*)}\int_{|u|\ge L} e^{-u^2/8}du \le \frac{1}{3}e^{n\Phi(t^*)}\frac{e^{-2}}{\sqrt{n}}\int_{-L}^{L} e^{-u^2}du.$$

The choice of L here is universal.

3. $\theta \geq 1$. For the outer integral, we have by Lemma 9 and (31) that

$$\int_{|\theta|\geq 1} \mathrm{Re}(\exp\big(n\Psi(\theta) + B(t^* e^{i\theta})\big))d\theta \leq 2\pi e e^{n\Phi(t^*)-n/5}.$$

We choose n large enough so that again this is at most $(1/3)K_n$. Combining all these bounds, the sum of three integrals is at least $(1/3)K_n$. Applying (36) we complete the proof of (32) and thus the proof of Proposition 10,

□

Proof of Theorem 5. We have completed the proof of the main result when Φ is defined as in (10). In order to complete the proof we need to validate Propositions 1 and 4. We will also prove Proposition 2, which will be a step in proving Proposition 4. These propositions are established in the next section. □

3 Loose ends. Proof of Propositions 1,2,4

Proof of Proposition 1. Consider an operator $\mathcal{T}$ acting on functions $G \in \mathcal{F}_{e,\Omega}$ defined by

$$\mathcal{T}(G)(z,x) = z + z\sum_{d\geq 1}\frac{1}{d!}\int_{\Omega^d}\prod_{i=1}^{d}\left[K(x,y_i)\,\frac{G^{q-1}(z,y_i)}{(q-1)!}\right]\prod_{1\leq i<j\leq d}K(y_i,y_j)\,\nu^{\otimes d}(dy_1\cdots dy_d).$$

While the range of $\mathcal{T}(G)$ could potentially include ∞ if the infinite sum is not absolutely convergent, we show below that this is not the case.

Lemma 11. *For every $G \in \mathcal{F}_{e,\Omega}$, we have $\mathcal{T}(G) \in \mathcal{F}_{e,\Omega}$. Furthermore, for every $G_1, G_2 \in \mathcal{F}_{e,\Omega}$,*

$$\|\mathcal{T}(G_1) - \mathcal{T}(G_2)\|_{e,\Omega} \leq e^{-1}\|G_1 - G_2\|_{e,\Omega}. \tag{37}$$

Namely, $\mathcal{T}$ is a contraction on the space $\mathcal{F}_{e,\Omega}$ with respect to the metric induced by the norm $\|\cdot\|_{e,\Omega}$.

Proof. Let

$$D_e = \{z \in \mathbb{C} : |z| \leq e^{-2}\}.$$

Fix $G \in \mathcal{F}_{e,\Omega}$. We first show that $\mathcal{T}(G)$ is well-defined and belongs to $\mathcal{F}_{e,\Omega}$.

For $z \in D_e$ and $x \in \Omega$, using $|K| = 1$, $\nu(\Omega) = 1$, and $\|G\|_{e,\Omega} \leq 1$, we have

$$\begin{aligned}
|\mathcal{T}(G)(z,x)| &\leq |z| + |z|\sum_{d\geq 1}\frac{1}{d!}\int_{\Omega^d}\prod_{i=1}^{d}\frac{|G(z,y_i)|^{q-1}}{(q-1)!}\,\nu^{\otimes d}(dy_1\cdots dy_d)\\
&\leq |z| + |z|\sum_{d\geq 1}\frac{1}{d!}\\
&= |z|e \leq e^{-1} \leq 1.
\end{aligned}$$

The same estimate shows that the defining series for $\mathcal{T}(G)$ converges absolutely and uniformly on $D_e \times \Omega$.

It remains to check continuity. For each fixed d, the corresponding integral is continuous in (z, x) by dominated convergence. Indeed, G is continuous, the integrand is uniformly bounded by an integrable function, and for every fixed $x \in \Omega$ the discontinuity set of the factors involving $K(x, y_i)$ has $\nu^{\otimes d}$-measure zero. Since the series converges uniformly on $D_e \times \Omega$, $\mathcal{T}(G)$ is continuous on $D_e \times \Omega$. Therefore $\mathcal{T}(G) \in \mathcal{F}_{e,\Omega}$.

Now fix $G_1, G_2 \in \mathcal{F}_{e,\Omega}$ and set

$$\Delta = \|G_1 - G_2\|_{e,\Omega}.$$

For fixed $z \in D_e$, $x \in \Omega$, $d \geq 1$, and $y_1, \ldots, y_d \in \Omega$, define

$$A_i = K(x, y_i)\frac{G_1(z, y_i)^{q-1}}{(q-1)!}, \qquad B_i = K(x, y_i)\frac{G_2(z, y_i)^{q-1}}{(q-1)!}.$$

Since $G_1, G_2 \in \mathcal{F}_{e,\Omega}$, we have $|A_i| \leq 1$ and $|B_i| \leq 1$. By the telescoping identity,

$$\prod_{i=1}^{d} A_i - \prod_{i=1}^{d} B_i = \sum_{k=1}^{d} \left(\prod_{i<k} A_i\right)(A_k - B_k)\left(\prod_{i>k} B_i\right).$$

Therefore,

$$\left|\prod_{i=1}^{d} A_i - \prod_{i=1}^{d} B_i\right| \leq \sum_{k=1}^{d} |A_k - B_k|.$$

For each k,

$$\begin{aligned} |A_k - B_k| &= \frac{|G_1(z, y_k)^{q-1} - G_2(z, y_k)^{q-1}|}{(q-1)!} \\ &\leq \frac{q-1}{(q-1)!}|G_1(z, y_k) - G_2(z, y_k)| \\ &\leq \Delta. \end{aligned}$$

Here we used the elementary inequality

$$|a^m - b^m| \leq m|a - b|, \qquad |a|, |b| \leq 1,$$

with $m = q - 1$, together with $(q-1)/(q-1)! \leq 1$ for $q \geq 2$.

The linear term z cancels when subtracting $\mathcal{T}(G_1)$ and $\mathcal{T}(G_2)$. Hence, using again $|K| = 1$,

$$\begin{aligned} |\mathcal{T}(G_1)(z, x) - \mathcal{T}(G_2)(z, x)| &\leq |z| \sum_{d \geq 1} \frac{1}{d!} \int_{\Omega^d} \left|\prod_{i=1}^{d} A_i - \prod_{i=1}^{d} B_i\right| \nu^{\otimes d}(dy_1 \cdots dy_d) \\ &\leq |z| \sum_{d \geq 1} \frac{d}{d!}\Delta \\ &= |z| e \, \Delta \\ &\leq e^{-1}\Delta. \end{aligned}$$

Taking the supremum over $z \in D_e$ and $x \in \Omega$ gives

$$\|\mathcal{T}(G_1) - \mathcal{T}(G_2)\|_{e,\Omega} \leq e^{-1}\|G_1 - G_2\|_{e,\Omega}.$$

This proves (37). □

Returning to the proof of the proposition, the existence and uniqueness of the solution to the fixed point equation (3) now follow from the Banach fixed point theorem, applied to the complete metric space $\mathcal{F}_{e,\Omega}$ with the metric induced by $\|\cdot\|_{e,\Omega}$. Equivalently, the unique fixed point is the uniform limit of the sequence

$$H_0 \equiv 1, \qquad H_{n+1} = \mathcal{T}(H_n), \quad n \geq 0.$$

□

Next we prove Proposition 2.

Proof of Proposition 2. Let $\hat{H}(z,x)$ be defined as the right-hand side of (6). It is finite and analytic on $\{z : |z| \leq 1-\delta\}$ for any $\delta > 0$ since $\left|\mathcal{T}_v^{\text{sim}}\right| = \frac{(v-1)!v^{s-1}}{s!((q-1)!)^s}$ with $s = (v-1)/(q-1)$, which by Stirling's approximation is $e^{O(v)}v!$. We claim that $\hat{H} = H$. Recall the definition of $B_{H,r}(X,x)$ from (5). Suppose T is an r-rooted tree with at least one hyperedge. Let $e_1, \ldots, e_d$ be the hyperedges containing r. Let $T_{e_i,k}, i \in [d], k \in [q-1]$ be the subtrees generated by the k-th node of hyperedge e_i, where we assume that the node r is the last node q in the hyperedge e_i. The hyperedge e_i itself is included into the tree $T_{e_i,k}$. Note that for any two hyperedges e, f of a tree we have $|e \cap f| \in \{0,1\}$. Thus $|e \cap f| = 1$ iff $e \sim f$. Then we can expand (5) further as

$$\begin{aligned} B_{T,r}(x) = &\prod_{i\in[d]} K(x, X(e_i)) \prod_{e_i,e_j,1\leq i<j\leq d} K(X(e_i), X(e_j)) \\ &\times \prod_{i\in[d],k\in[q-1]} \prod_{f\in T_{e_i,k},(i,k)\in f} K(X(e_i), X(f)) \prod_{f_1\sim f_2\in T_{e_i,k}, f_1,f_2\neq e_i} K(X(f_1), X(f_2)). \end{aligned}$$

Here (i,k) denotes the k-th node of e_i. We can simplify this as

$$B_{T,r}(x) = \prod_{i\in[d]} K(x, X(e_i)) \prod_{e_i,e_j,1\leq i<j\leq d} K(X(e_i), X(e_j)) \prod_{i\in[d],k\in[q-1]} B_{T_{e_i,k}}(X(e_i)).$$

Consider the expectation of $B_{T,r}(x)$ conditioned on $X(e_i) = y_i, i \in [d]$. Then

$$\mathbb{E}[B_{T,r}(x) | X(e_i) = y_i, i \in [d]] = \prod_{i\in[d]} K(x, y_i) \prod_{e_i,e_j,1\leq i<j\leq d} K(y_i, y_j) \prod_{i\in[d],k\in[q-1]} b_{T_{e_i,k}}(y_i)$$

Then

$$b_{T,r}(x) = \int_{\Omega^v} \prod_{i\in[d]} K(x, y_i) \prod_{e_i,e_j,1\leq i<j\leq d} K(y_i, y_j) \prod_{i\in[d],k\in[q-1]} b_{T_{e_i,k}}(y_i) \nu(dy_i).$$

From this we obtain that $\hat{H}$ is a fixed point of (3), where normalizations $1/d!$ and $1/(q-1)!$ are included since the hyperedges $e_i, i \in [d]$ and nodes $i \in [q-1]$ in each hyperedge $e_i, i \in [d]$ are unordered. Therefore, by Proposition 1 $\hat{H} = H$. □

Our final step is proving Proposition 4.

Proof of Proposition 4. Introduce $\hat{\mathcal{R}}(z)$ and $\hat{\mathcal{E}}(z)$ as follows.

$$\hat{\mathcal{R}}(z) = \sum_{v=1}^{\infty} \frac{z^v}{v!} \sum_{T \in \mathcal{T}_v^{\text{sim}}} |V(T)| b_T.$$

$$\hat{\mathcal{E}}(z) = \sum_{v=1}^{\infty} \frac{z^v}{v!} \sum_{T \in \mathcal{T}_v^{\text{sim}}} |E(T)| b_T.$$

As above, it is easy to see that $\hat{\mathcal{R}}, \hat{\mathcal{E}}$ are analytic on $\{z : |z| \le 1/e^2\}$.

Lemma 12. *We claim that* $\hat{\mathcal{R}} = \mathcal{R}, \hat{\mathcal{E}} = \mathcal{E}$.

Proof. Fix a tree $T \in \mathcal{T}_v^{\text{sim}}$ with a distinguished root r. Let $e_1, \dots, e_d$ hyperedges containing r. We have

$$B_T(X) = \prod_{i,j \in [d]} K(X(e_i), X(e_j)) \prod_{i \in [d], k \in [q-1]} B_{T_{i,k}}(X),$$

where the trees $T_{i,k}$ are defined as in the proof of Proposition 2. Then

$$b_T = \int_{\Omega^d} \prod_{i,j \in [d]} K(y_i, y_j) \prod_{i,k \in [q-1]} b_{T_{i,k}}(y_i) \nu(dy_i).$$

The proof of $\hat{\mathcal{R}} = \mathcal{R}$ is then completed similarly to the proof of Proposition 2 where the multiplier v appears in front of b_T due to v choices of the root r.

The proof of $\hat{\mathcal{E}} = \mathcal{E}$ is obtained similarly. Fix a tree T and an edge e in this tree. Consider q subtrees rooted at q nodes of e. Fixing the value $X(e)$ to x we obtain q copies of $H(x, z)$. The multiplier $|E(T)|$ appears due to that many choices of e. Dividing by $q!$ is due to symmetry of q nodes in e. □

We now complete the proof of the proposition. Applying Lemma 12 and $E(T) = (v-1)/(q-1)$ for every $T \in \mathcal{T}_v^{\text{sim}}$ we have

$$\begin{aligned}
\mathcal{R}(z) - (q-1)\mathcal{E}(z) &= z + \sum_{v=1}^{\infty} \frac{z^v}{v!} \sum_{T \in \mathcal{T}_v^{\text{sim}}} |V(T)| b_T - (q-1) \sum_{v=1}^{\infty} \frac{z^v}{v!} \sum_{T \in \mathcal{T}_v^{\text{sim}}} (v-1)/(q-1) b_T \\
&= z + \sum_{v=1}^{\infty} \frac{z^v}{v!} \sum_{T \in \mathcal{T}_v^{\text{sim}}} b_T.
\end{aligned}$$

Since $\mathcal{R}$ and $\mathcal{E}$ are analytic on $|z| \le 1/e^2$, then $\mathcal{R}(\rho z), \mathcal{E}(\rho z)$ are analytic on $|z| \le 2$ when $|\beta| \le \beta^*$

and β^* is sufficiently small. Then

$$\begin{aligned}\mathcal{F}(z) &= \rho^{-1}\left(\rho z + \sum_{v=1}^{\infty} \frac{(\rho z)^v}{v!} \sum_{T \in \mathcal{T}_v^{\mathrm{sim}}} b_T\right) \\ &= z + \sum_{v=1}^{\infty} \frac{\rho^{v-1} z^v}{v!} \sum_{T \in \mathcal{T}_v^{\mathrm{sim}}} b_T \\ &= z + \sum_{v=1}^{\infty} z^v \mu_v \\ &= \Phi(z).\end{aligned}$$

This completes the proof of the proposition. □

References

[1] Eric R Anschuetz, David Gamarnik, and Bobak T Kiani, *Bounds on the ground state energy of quantum p-spin hamiltonians*, Communications in Mathematical Physics **406** (2025), no. 10, 232.