Sequence-to-Expression Models Fail to Generalise to New Genomic Neighbourhoods

Anonymous Author(s)
Affilation, Address
anon.email@example.org

Abstract

Machine learning for synthetic biology is moving beyond interpreting DNA towards designing it. This shift requires verifiers, i.e. models that predict how designed DNA sequences will be expressed before they are constructed, but it remains unclear whether current sequence-to-expression (S2E) models can perform this role outside native genomic contexts. Baker's yeast provides a uniquely powerful eukaryotic testbed for this question: its genetic tractability enables complex genetic perturbations such as genome-wide reporter integrations and large structural rearrangements that remain difficult to construct at comparable scale in mammalian genomes. Here, we benchmark two S2E models zero-shot across eleven yeast tasks. Both models approximately recover the experimental ordering of short, gene-proximal variants, but perform at chance level on genome-wide reporter integrations and poorly on structural rearrangements. For native-promoter swaps, their predicted transcript-abundance ranges are only approximately two-fold, whereas measured fluorescence spans one to two orders of magnitude more for two reporter genes. These results identify contextual generalisation and dynamic-range compression as major obstacles to using current S2E models as verifiers for genome design. An open-source implementation of our benchmark can be found online.[1]

1 Introduction

The next frontier for machine learning in genomics is in designing DNA and not merely interpreting it. Models intended to guide genome engineering must therefore predict how genes behave under DNA sequence perturbations beyond the native genomic contexts that make up their training data. Baker's yeast offers a powerful eukaryotic testbed for this problem because its genome is far easier to engineer systematically and at scale than the human genome.

Foundational insights in genomics where established in yeast first, such as centromere function, DNA replication, and cell-cycle control [1, 2, 3]. Its experimental advantage now extends to synthetic genomics. Sc2.0, the synthetic yeast genome project, enables deletions, duplications, inversions, and translocations to be generated throughout synthetic yeast chromosomes. Using these rearrangements, *Brooks et al.* [4] showed that transcriptional units are not context-independent parts: neighbouring transcription alters expression levels and transcript isoform boundaries. Yeast therefore makes it possible to test whether sequence-to-expression models learn regulatory principles that remain valid when genomes are engineered rather than merely observed.

Genomic sequence-to-expression (S2E) models predict RNA-seq and related expression readouts directly from DNA sequence. If reliable under cis-perturbations, they could become core tools for synthetic biology: tuning the expression of engineered pathways and serving as predictive oracles

[1] https://github.com/anon-schmoo/neurips-icbinb-2026

Submitted to the I Can't Believe It's Not Better (ICBINB): Failure Modes of AI in Biology workshop at NeurIPS 2026.

that screen or steer sequences generated by DNA design models before they are constructed [5]. Yet, in the native genomic data that dominate current training sets, most genes occur in only one neighbourhood. Large receptive fields make the surrounding sequence available to a model without forcing it to learn its causal contribution.

Here, we exploit yeast's genetic tractability to benchmark *Shorkie* [6] and *Yorzoi* [7] zero-shot across eleven tasks, ranging from single-nucleotide variants and massively parallel reporter assays (MPRAs) to promoter swaps, genome-wide reporter relocations, structural rearrangements, and exogenous bacterial chromosomes. We find that models rank short, gene-proximal variants, particularly in promoters and terminators, correctly, but predict severely compressed expression ranges (promoter swaps). Both perform at approximately chance level when identical reporter cassettes are integrated at new genomic sites and remain weak on structural rearrangements. We show that predictive features of position effects, such as reporter distance to Micro-C defined boundaries, are linearly decodable by *Yorzoi* but are nevertheless not converted to accurate expression predictions. Together, these results identify generalisation to distant sequence neighbourhoods and dynamic range compression as central obstacles to using current S2E models for genome design.

2 Related Work

S2E model benchmarks *Karollus et al.* [8] benchmark mammalian S2E models across a variety of tasks, mainly population studies and deep perturbation assays. Their key finding is that mammalian S2E models at the time, such as *Enformer* [9], were substantially more accurate at predicting the effects of promoter perturbations but systematically underestimated the effects of distal enhancers. In contrast, we leverage the greater variety of genetic perturbations possible in yeast and benchmark more recent models that explore novel training methods or datasets.

Yeast S2E models for RNA-seq *Chao et al.* [6] and *Schneider et al.* [7] introduce some of the benchmark tasks presented here to evaluate the generalisation of their models, *Shorkie* and *Yorzoi*, to downstream tasks, including expression quantitative trait loci (eQTL) studies and promoter massively parallel reporter assays. However, with the exception of one task, they do not benchmark their models on the same tasks and do not cover the variety of genetic perturbations found in the yeast literature, such as perturbations of other regulatory regions, engineered structural variants, or genome-wide reporter translocations. Our benchmark fills this gap.

Assay-specific S2E models For many of the benchmark tasks we present, the authors trained small convolutional, decision-tree-based, or linear models to predict the outcomes of their reporter assays. For example, *Hong et al.* [10] train a small XGBoost model to predict the fluorescence of a reporter gene integrated at different genomic locations. *Rafi et al.* [11] evaluate a variety of deep-learning models, some contributed by the wider community, to predict the outcome of their massively parallel promoter assay. Although sometimes useful for uncovering explanatory features, these models are usually constrained to the narrow experimental datasets on which they were trained and are generally not applicable to other assays. Here, we benchmark S2E models trained on RNA-seq and additional sequencing-based assays that, in theory, should be able to generalise to other assays.

3 A benchmark for genomic sequence-to-expression models in baker's yeast

We introduce eleven benchmark tasks that probe different capabilities of S2E models in a zero-shot setting. Before explaining our results, we first introduce two important concepts: a formal definition of sequence-to-expression models and an overall explanation of how zero-shot evaluations work across benchmark tasks.

S2E models map a DNA sequence $x \in \{A, C, G, T\}^L$ to non-negative coverage predictions

$$\hat{y} = f(x) \in \mathbb{R}_{\geq 0}^{T \times C}, \tag{1}$$

where f is the model, T is the number of tracks, and C is the number of predicted positions. Rows correspond to tracks, and columns correspond to positions. We primarily evaluate RNA-seq tracks: *Shorkie* [6] predicts one unstranded track per experiment, whereas *Yorzoi* [7] predicts separate

forward- and reverse-strand tracks. Typically, $C < L$ because predictions are restricted to the central region of the input, reducing boundary effects from unobserved flanking sequences.

Zero-shot evaluation Both models predict coverage profiles with one scalar per position. In contrast, most assays that we evaluate report a single scalar expression value per gene. Thus, we transform our predictions into per-gene scalars as well.

Let $\tilde{y}(x) \in \mathbb{R}^{C}_{\geq 0}$ denote the processed coverage profile for input sequence x. For a target gene g, let $\mathcal{R}_g(x)$ denote the set of output positions used to read out that gene. Unless stated otherwise, this is its coding sequence, i.e., $\mathcal{R}_g(x) = \mathrm{CDS}_g(x)$. We define

$$\varphi_g(x) := \sum_{j \in \mathcal{R}_g(x)} \tilde{y}_j(x). \tag{2}$$

Thus, $\varphi_g(x)$ is the model's scalar prediction of transcript abundance for gene g. When the target gene is unambiguous, we write $\varphi(x)$.

For all benchmark tasks, we use S2E-models in zero-shot fashion. This means that we derive scalar predictions of transcript abundance for the experimentally measured sequences and compute our metrics as pairwise comparisons of a scalar prediction and a scalar experimental readout which are mostly fluorescence or RNA-seq-derived gene expression measurements.

3.1 S2E models fail to predict gene expression of genome-wide integrated reporter genes

Motivation The core result from [12], [13], and [10] is that the same gene expression cassette, complete with a promoter, coding sequence, and terminator, is differentially expressed when integrated at different positions in the genome. Hong et al. [10] found that the distance from the integration site to the nearest boundary of a self-associating chromatin domain [14] explains part of this variation. Although the models are not trained directly on Micro-C data to predict such boundaries, gene-expression neighbourhood effects could be learned from RNA-seq alone, which lies squarely within the existing models' training data.

Task We task the models with predicting the expression of a gene expression cassette containing a promoter, reporter CDS, and terminator at two types of integration site. Expression is experimentally measured as fluorescence. We correlate the predicted transcript abundance of the reporter with the experimental fluorescence measurement.

Data *Hong et al.* [10] integrate a red fluorescent protein expression cassette at 150 intergenic regions (one integration per genome) and measure fluorescence by flow cytometry. In contrast, *Wu et al.* [13] choose a different type of integration site and insert their reporter cassette into different strains of the yeast single-knockout collection at the positions of previously deleted non-essential genes. We note that some of the variation in this assay might be better explained by variation in the trans-regulatory environment (e.g., changes in transcription-factor abundance) than by the local position effect.

Evaluation For each insertion site $i \in I$, we reconstruct sequence x_i as the concatenation of the upstream genomic neighbourhood, the fixed reporter cassette, and the downstream genomic neighbourhood:

$$x_i := \mathrm{concat}\Big(\mathrm{genome}_{i-\frac{L-R}{2}:i}, \mathrm{reporter}, \mathrm{genome}_{i:i+\frac{L-R}{2}}, \Big), \quad i \in I. \tag{3}$$

We predict reporter expression as

$$\hat{y}_i := \varphi_{\mathrm{reporter}}(x_i), \quad i \in I. \tag{4}$$

Here, I is the set of surveyed integration sites, $\mathrm{genome}_{m:n}$ denotes the half-open genomic interval from m inclusive to n exclusive, R is the length of the reporter cassette, and L is the model input length. We report Pearson correlation r and Spearman rank correlation ρ across the paired values

$$\{(\hat{y}_i, y_i) \mid i \in I\}, \tag{5}$$

where y_i is the measured reporter fluorescence at integration site i.

Table 1: Zero-shot performance on the chromosomal position-effect benchmarks. Wu evaluates 1,043 scorable ORF-replacement loci. Hong reports the held-out IntProp set as the primary test set ($n = 52$) and IntTrain as a secondary split ($n = 98$). All correlations are close to zero or negative, indicating that neither model captures genomic position effects.

Assay	n	Metric	Shorkie	Yorzoi
Wu et al.	1,043	Pearson r	–0.039	0.012
		Spearman ρ	–0.029	0.016
Hong et al.	52	IntProp Spearman ρ	–0.148	0.063
Hong et al.	98	IntTrain Spearman ρ	–0.269	–0.073

Results In both assays, Yorzoi and Shorkie have approximately random performance and fail to predict the effect of inserting the reporter cassette at a new site (Table 1).

Hong et al. [10] fit an XGBoost model to part of their data and determine that, among their hand-crafted features, the distance from the insertion site to the nearest Micro-C-determined boundary is the most predictive (single-feature $r^2 \approx 0.1$). Here, a Micro-C boundary denotes a local position with the minimum number of crossing Micro-C interactions considered within a 10 kbp window (Section A.6). We wondered to what extent the evaluated models capture this feature. Shorkie and Yorzoi both broadly follow a U-Net architecture in which the sequence is first downsampled to a shorter length, processed through transformer blocks with bidirectional self-attention, and then upsampled to sequence length while the number of channels is expanded to produce an embedding. Finally, a linear head transforms this embedding to match the target number of channels.

To investigate the presence of the aforementioned feature, we trained linear probes on outputs of *Yorzoi* at three layers: the output of the downsampling convolutions, the output of the bottleneck transformer blocks, and the output of the upsampling convolutions before the output head [6, 7]. The target was the distance to the nearest boundary at each position in the sequence. Interestingly, the downsampling layers had no linearly decodable notion of the boundary distances, with a low Pearson correlation of $r = 0.16$ on held-out sequences. The output of the transformer layers, however, achieved a higher correlation of 0.46 on held-out sequences, demonstrating that some version of this feature is learned during training. Right before the head, the feature is less decodable ($r = 0.31$).

Our hypothesis is that this feature is not essential for predicting training windows and is thus only weakly learned during training. In follow-up experiments, we intend to retrain one of the models with Micro-C boundary distances or contact maps as an additional output, which would force the model to retain the feature and potentially improve performance on integration assays such as that of *Hong et al.*.

3.2 S2E models poorly predict expression changes after structural rearrangements

Motivation Although the native yeast genome offers limited diversity for testing how the expression of the same coding sequence changes across structurally distinct genetic neighbourhoods, engineered strains from the Sc2.0 project provide extensive structural variation. In contrast to other benchmark tasks that use a fluorescent marker, a directly matched transcriptomic readout allows model failures to be attributed more clearly to the model rather than to inherent differences between the biological mechanisms captured by the assay and the model.

Task Models are benchmarked on their ability to predict the log-fold change in expression, quantified by long-read direct RNA-seq, for a gene in two different genetic neighbourhoods: the reference sequence ref on an unrecombined chromosome 9 (right arm) and the same gene in a recombined and therefore structurally rearranged genetic neighbourhood (alt).

Data *Brooks et al.* [4] profile transcription in approximately 70 yeast strains with a structurally rearranged chromosome 9 (right arm) using long-read Nanopore RNA-seq. These rearrangements are produced by inserting recombination sites between the stop codon and 3′ UTR and then inducing recombination. This produces various structural variants, including deletions, duplications, inversions, and translocations. Each structural variant necessarily changes the 3′ UTR of the gene of

Table 2: Results for *Brooks et al.* [4], log-fold-change (LFC) prediction on the shared cohort ($n = 327$). Metrics are reported separately for each of the three parental sequencing runs. *Ceiling* is the leave-one-out reproducibility bound for that run, computed on the samples where the other two runs are available ($n = 198/80/169$).

Model	Run 1 ($n = 205$)	Run 2 ($n = 80$)	Run 3 ($n = 291$)
LFC direction balanced accuracy			
Shorkie	0.51	0.61	0.53
Yorzoi	**0.62**	**0.65**	**0.63**
Ceiling	*0.76*	*0.96*	*0.69*
LFC Spearman ρ			
Shorkie	–0.06	–0.16	0.11
Yorzoi	**0.44**	**0.28**	**0.22**
Ceiling	*0.78*	*0.95*	*0.59*
LFC Pearson r			
Shorkie	–0.09	0.00	0.02
Yorzoi	**0.30**	**0.21**	**0.15**
Ceiling	*0.79*	*0.95*	*0.67*

interest. This dataset yields 215 unique sequences from 30 genes across approximately 50 strains, totalling 327 samples.

Metric The *ref* strain was sequenced three times. Thus, a gene can be compared against up to three independent reference measurements. A run serves as a reference for a gene only if that gene reached at least ten reads in it. Because the threshold is applied per gene and per run, the three runs cover different sets of genes of the 327 samples. We therefore report every metric separately for each run. For run $k \in \{\mathrm{Run\ 1}, \mathrm{Run\ 2}, \mathrm{Run\ 3}\}$, gene g, and a sequence i containing g in the original (ref) or rearranged (alt) neighborhood:

$$\mathrm{LFC}_{i,g,k} = \log_2 \varphi_g(x_{\mathrm{alt},i}) - \log_2 \varphi_g(x_{\mathrm{ref}})_k \tag{6}$$

where $x_{\mathrm{alt},i}$ is and the predicted LFC is formed the same way, by predicting coverage for the ref and alt sequences and summing over the coding-sequence bins. Within each run we report balanced accuracy, Spearman ρ and Pearson r across the samples that run supports. Each run is compared against its own leave-one-out ceiling which is the correlation between that run's measured LFCs and the mean of the other two runs.

Results Neither Shorkie nor Yorzoi can accurately predict the effect of structural variation on gene expression. Yorzoi beats Shorkie on every metric in every run, but its balanced accuracy in predicting the direction of the effect reaches only 62–65%, and its correlations stay far below the leave-one-out ceiling in all three runs Table 2. Shorkie is at chance throughout. Unlike Shorkie, Yorzoi was trained on data from [4], although the benchmark sequences scored here were held out from both models. These results generally point to the failure of current models to generalise to new sequence neighbourhoods that influence expression. Some of this performance difference could be attributable to the different RNA-seq assays used to train Shorkie and Yorzoi: while the benchmark uses long-read Nanopore RNA-seq, Shorkie is trained only on short-read Illumina RNA-seq.

3.3 S2E models rank promoter swaps but predict a compressed expression range

Motivation The *Yeast Toolkit* (YTK) [15] is a frequently used modular set of building blocks for yeast synthetic biology. Its characterisation and engineering make it possible to construct gene expression cassettes by combining genetic parts, most importantly promoters, coding sequences, and terminators. By selecting promoters of different "strengths", characterised by fluorescence, the authors show that gene expression can be tuned across a very wide range, from no change above background to a thousand-fold increase in fluorescence. We investigated the models zero-shot for two reasons: (1) S2E models almost never see the same gene with different promoters during training

(except for a handful of homologous genes), and (2) we previously observed that the dynamic range predicted for MPRAs seemed suspiciously narrow.

Task Models are tasked with predicting the expression of two fluorescent genes (mRuby and Venus), each driven by a set of 19 promoters obtained from the native yeast genome (e.g., pTDH3).

Data We obtain plate-reader fluorescence measurements for 19 promoters and two fluorescent genes (the median across four replicates per promoter-gene pair) from *Lee et al.* [15]. Each data point is fluorescence normalised by biomass (OD600) and recorded as fold change over background fluorescence.

Evaluation For each promoter $p \in \mathcal{P}$ and reporter $r \in \{\mathrm{mRuby2}, \mathrm{Venus}\}$, we reconstruct the sequence

$$x_{p,r} := \mathrm{concat}(\mathrm{upstream_DNA}, p, r, \mathrm{downstream_DNA},). \tag{7}$$

The upstream and downstream flanking sequences are fixed across all constructs. We obtain a scalar transcript abundance prediction for the reporter using (2):

$$S_{p,r} := \varphi_r(x_{p,r}). \tag{8}$$

We compare $S_{p,r}$ with the corresponding experimental fluorescence measurement using Pearson correlation r and Spearman rank correlation ρ.

Let $L_{p,r}$ denote the median fluorescence for promoter p and reporter r, expressed as fold change over the non-fluorescent-cell background. To assess how well the predictions capture the observed dynamic range, we calculate the observed and predicted max-to-min ratios separately for each reporter:

$$F_{\mathrm{obs},r} = \frac{\max_{p\in\mathcal{P}} L_{p,r}}{\min_{p\in\mathcal{P}} L_{p,r}}, \quad F_{\mathrm{pred},r} = \frac{\max_{p\in\mathcal{P}} S_{p,r}}{\min_{p\in\mathcal{P}} S_{p,r}}. \tag{9}$$

The observed and predicted extrema are selected independently. Thus, the promoter producing the predicted maximum or minimum need not be the promoter producing the corresponding experimental extremum. We convert the range ratios into base-10 logarithmic spans:

$$D_{\mathrm{obs},r} = \log_{10}(F_{\mathrm{obs},r}), \quad D_{\mathrm{pred},r} = \log_{10}(F_{\mathrm{pred},r}). \tag{10}$$

Results As with other gene-proximal genetic perturbations, both models approximately recover the ranking of promoter strengths, and Shorkie outperforms Yorzoi for both reporter genes (Table 3). However, their predicted dynamic expression ranges are many times smaller than the observed gene expression ranges. The observed fluorescence ranges are 66-fold and 481-fold for mRuby and Venus, respectively, while the predicted transcript-abundance range is around two-fold for both models and both reporter genes. *Lee et al.* did not collect the experimental data necessary to determine the true dynamic range of transcript abundance. Thus, an unconfounded comparison between the dynamic ranges is not possible because switching promoters also changes the 5′ untranslated region, which is known to modify translation efficiency and could therefore modulate the observed fluorescence range. However, the difference between the dynamic ranges is so large that it seems implausible that differences in protein abundance caused by 5′ UTR changes alone would explain the gap. An alternative explanation is that the models indeed predict a much narrower range of gene expression. This may be partly because the models never see the same gene with different promoters during training.

3.4 S2E models rank MPRA variants best in promoters and terminators, worst in 5′ UTRs

Motivation Massively parallel reporter assays (MPRAs) screen thousands to millions of DNA sequences at one specific position with a fixed reporter protein. Models that have learned the appropriate regulatory logic of gene expression in those regions should be able to rank the different sequences correctly.

Task Here, we examine four different datasets, each probing a different part of regulatory logic and benchmarked in the same way. We predict expression for a sequence construct i and compute gene

Table 3: Results on the *Lee et al.* [15] YTK promoter benchmark, reported separately for each reporter CDS ($n = 19$ promoters per reporter). The observed range is the max/min ratio of median fold-over-background fluorescence; the predicted range is the max/min ratio of raw reporter-CDS coverage sums. Pearson r is calculated after $\log_{10}$ transformation; ρ denotes Spearman rank correlation. Bold denotes the better model based on unrounded values.

Metric	**mRuby2**		**Venus**	
	Shorkie	Yorzoi	Shorkie	Yorzoi
Observed range[2]	66×	66×	481×	481×
Predicted range	2.6×	1.9×	2.2×	2.2×
Pearson r	**0.50**	0.41	**0.78**	0.49
Spearman ρ	**0.52**	0.47	**0.76**	0.46

expression as the sum of the coding-sequence bins. This is then correlated with the true fluorescence measurements. The datasets investigate promoter, 5′ untranslated region, coding sequence, and terminator variants.

Data We use four different datasets from [11], [16], [17], and [18]. *Rafi et al.* [11] provide a dataset of approximately 70,000 80-bp promoter sequences, each embedded in a plasmid-based promoter scaffold that drives the transcription of a fluorescent protein. Cells are sorted into 18 bins by fluorescence-activated cell sorting (FACS). The cell-count-weighted mean bin is z-normalised and then reported as a proxy for *reporter expression*. *Cuperus et al.* [16] engineer 500,000 5′-untranslated-region (5′ UTR) variants and measure gene expression in a pooled growth assay. We note that variants in the 5′ UTR also alter translation efficiency, which no model currently captures. However, variants can, for example, alter mRNA stability, which is within the theoretical capabilities of current models. *Chen et al.* [17] perturb small stretches (up to 20 bp) of the coding sequences of two genes, GFP and TDH3, while keeping their amino acid sequences unchanged. Their proxy for gene expression is the ratio of RNA-seq reads to DNA-seq reads for each variant in the pool. *Shalem et al.* [18] screen a collection of thousands of terminator variants in a similar manner to [11]. Terminator variants are known to alter transcription stop sites and transcript isoform distributions, resulting, for example, in different mRNA stabilities.

Evaluation For Rafi, Shalem, and Chen, we use the host-marginalised log-expression-difference score m_v defined in Section A.4. Its per-context terms are computed from the scalar transcript-abundance readout $\varphi_g(x)$ defined in (2). For Cuperus, we score the fixed reporter construct directly using $\varphi_{\mathrm{HIS3}}(x_v)$. We correlate the resulting scalar scores with the corresponding experimental reporter measurements.

Results Broadly speaking, we observe the strongest correlations with experimentally measured gene expression for the promoter and terminator assays (r and ρ from $0.61 - 0.77$), slightly weaker correlations for synonymous codon sequences, and the weakest for the 5′ UTR MPRA. *Shorkie* outperforms *Yorzoi* on the promoter MPRA. We speculate that this is mostly attributable to its masked-language-modelling pretraining procedure, since promoter motifs are well conserved across yeast species. *DREAM-RNN* [11], a supervised model is evaluated on the same test set but trained on a much larger set of sequences from the same experiment achieves a Pearson r of 0.98. This baseline demonstrates a substantial margin over both zero-shot approaches (Δr =0.21 compared with Shorkie).

In contrast, terminator motifs are more cryptic and less conserved. We speculate that *Yorzoi's* advantage over *Shorkie* on the terminator MPRA stems from its training on genomically rearranged sequences, in which some coding sequences are seen multiple times with different terminator sequences[3].

Cuperus et al. [16] is the least predictable MPRA of the four tested ones. We suspect that even perfect RNA-seq coverage prediction would yield limited correlation as the 5′ UTR sequence, especially around the start codon, has a direct influence on translation efficiency, which current models cannot learn from their transcription and chromatin focussed target tracks. Secondly, instead of relying

[2]The observed range is derived from experimental fluorescence measurements and is model-independent.

[3]The *Brooks* benchmark tasks only evaluates on held-out sequences not used during training.

Table 4: Results on MPRA benchmark tasks. Paired values report Pearson correlation / Spearman rank correlation (r / ρ); the single Cuperus values report ρ. R1 and R2 denote GFP replicates 1 and 2. Bold indicates the better-performing model.

Benchmark	Assay	Shorkie	Yorzoi
Rafi / deBoer promoter	Overall	**0.76 / 0.77**	0.61 / 0.63
Shalem terminator	Overall	0.64 / 0.65	**0.71 / 0.71**
Chen synonymous	GFP R1	**0.30 / 0.39**	0.28 / 0.36
	GFP R2	**0.57 / 0.59**	0.56 / 0.57
	TDH3	0.53 / **0.30**	**0.61** / 0.23
Cuperus 5′-UTR	Overall	**0.25**	0.12

on a reporter gene [16] use a pooled growth experiment in HIS3-knockout strains, for which it is unclear that enrichment in the pool remains correlated with protein expression above a certain protein abundance. Future models that predict protein abundance in addition to transcript abundance might improve upon the results shown here.

4 Discussion

Across eleven benchmark tasks, Shorkie and Yorzoi generalise best to short, gene-proximal variants, particularly in promoters and terminators. Both models recover weaker but meaningful signals for synonymous variants and eQTLs. However, accurate ranking does not ensure quantitative calibration. In the YTK promoter assay, both models approximately recover promoter ordering while predicting only a two-fold expression range, compared with observed fluorescence ranges of 66-fold and 481-fold. Although fluorescence is also affected by translation, this discrepancy suggests that the models substantially underestimate effect magnitude.

The largest failures occur when fixed expression cassettes or native genes are moved into new genomic neighbourhoods. Performance is near random in both position-effect assays and remains weak for structural rearrangements. Large receptive fields therefore make contextual sequence available without ensuring that models learn its causal contribution. One possible explanation lies in the observational training data: most native genes occur in only one neighbourhood, allowing models to rely on correlated proximal features instead. Our linear-probe results provide a clue: Micro-C boundary distance is linearly decodable from Yorzoi's intermediate representations. However, this does not conclusively establish that the model uses this information when predicting expression or that its failure to do so causes the observed position-effect errors.

Neither model is consistently superior. Shorkie generally performs better on promoter variants and eQTLs, whereas Yorzoi leads on structural rearrangements, terminators, and foreign bacterial chromosomes. Because the models differ in architecture, training data, receptive field, and output assays, and because we evaluate only two models, controlled ablations would be required to attribute these differences to particular design choices.

Individual benchmark tasks also impose limits on how specifically the failures can be interpreted. The Hong and YTK assays measure fluorescence rather than RNA abundance, while the Wu strains potentially alter the trans-regulatory background as well as the reporter's integration site. Moreover, although the evaluated Brooks sequences were held out from Yorzoi's training set, other data from the same study were used during training. These factors complicate comparisons of absolute performance and prevent us from attributing every error solely to genomic context. Nevertheless, the recurrence of weak contextual generalisation across independent datasets and perturbation types indicates a practical limitation for DNA sequence design.

Addressing this limitation will require training data in which identical transcriptional units are measured across multiple engineered genomic neighbourhoods. Future evaluations should also assess effect magnitude alongside ranking accuracy, since models can recover an experimental ordering while severely compressing the predicted response. Overall, current models are useful for prioritising local variants in matched contexts but are not yet reliable verifiers for genome engineering, where accurate predictions under structural and contextual changes are essential.

5 References

[1] L. Clarke and J. Carbon, 'Isolation of a yeast centromere and construction of functional small circular chromosomes', *Nature*, vol. 287, no. 5782, pp. 504–509, Oct. 1980, doi: 10.1038/287504a0.

[2] Stinchcomb, 'Isolation and characterisation of a yeast chromosomal replicator', 1979.

[3] B. J. Reid, J. G. Culotti, R. S. Nash, and J. R. Pringle, 'Forty-five years of cell-cycle genetics', *Molecular Biology of the Cell*, vol. 26, no. 24, pp. 4307–4312, Dec. 2015, doi: 10.1091/mbc.E14-10-1484.

[4] A. N. Brooks, A. L. Hughes, S. Clauder-Münster, L. A. Mitchell, J. D. Boeke, and L. M. Steinmetz, 'Transcriptional neighborhoods regulate transcript isoform lengths and expression levels', *Science*, vol. 375, no. 6584, pp. 1000–1005, Mar. 2022, doi: 10.1126/science.abg0162.

[5] G. Brixi *et al.*, 'Genome modelling and design across all domains of life with Evo 2', *Nature*, vol. 652, no. 8112, pp. 1349–1361, Apr. 2026, doi: 10.1038/s41586-026-10176-5.

[6] K.-H. Chao, M. M. Magzoub, E. Stoops, S. Hackett, J. Linder, and D. R. Kelley, 'Predicting dynamic expression patterns in budding yeast with a fungal DNA language model'. Accessed: May 22, 2026. [Online]. Available: https://www.biorxiv.org/content/10.1101/2025.09.19.677475v1

[7] T. Schneider *et al.*, 'Yorzoi: Predicting RNA-seq coverage from DNA sequence in yeast'. Accessed: May 22, 2026. [Online]. Available: https://www.biorxiv.org/content/10.1101/2025.09.20.677345v1

[8] A. Karollus, T. Mauermeier, and J. Gagneur, 'Current sequence-based models capture gene expression determinants in promoters but mostly ignore distal enhancers', *Genome Biology*, vol. 24, no. 1, p. 56, Mar. 2023, doi: 10.1186/s13059-023-02899-9.

[9] Ž. Avsec *et al.*, 'Effective gene expression prediction from sequence by integrating long-range interactions', *Nature Methods*, vol. 18, no. 10, pp. 1196–1203, Oct. 2021, doi: 10.1038/s41592-021-01252-x.

[10] H. Hong, Y. Cai, J. Wang, C. Dong, Q. Zhang, and J. Lian, 'Exploring chromosomal position effects for predictable tuning of metabolic pathways in yeast', *Nucleic Acids Research*, vol. 54, no. 8, p. gkag401, May 2026, doi: 10.1093/nar/gkag401.

[11] A. M. Rafi *et al.*, 'A community effort to optimize sequence-based deep learning models of gene regulation', *Nature Biotechnology*, pp. 1–11, Oct. 2024, doi: 10.1038/s41587-024-02414-w.

[12] X. Chen and J. Zhang, 'The Genomic Landscape of Position Effects on Protein Expression Level and Noise in Yeast', *Cell Systems*, vol. 2, no. 5, pp. 347–354, May 2016, doi: 10.1016/j.cels.2016.03.009.

[13] X.-L. Wu *et al.*, 'Genome-wide landscape of position effects on heterogeneous gene expression in Saccharomyces cerevisiae', *Biotechnology for Biofuels*, vol. 10, no. 1, p. 189, July 2017, doi: 10.1186/s13068-017-0872-3.

[14] T.-H. S. Hsieh, A. Weiner, B. Lajoie, J. Dekker, N. Friedman, and O. J. Rando, 'Mapping nucleosome resolution chromosome folding in yeast by Micro-C', *Cell*, vol. 162, no. 1, pp. 108–119, July 2015, doi: 10.1016/j.cell.2015.05.048.

[15] M. E. Lee, W. C. DeLoache, B. Cervantes, and J. E. Dueber, 'A Highly Characterized Yeast Toolkit for Modular, Multipart Assembly', *ACS Synthetic Biology*, vol. 4, no. 9, pp. 975–986, Sept. 2015, doi: 10.1021/sb500366v.

[16] J. T. Cuperus *et al.*, 'Deep learning of the regulatory grammar of yeast 5′ untranslated regions from 500,000 random sequences', *Genome Research*, vol. 27, no. 12, pp. 2015–2024, Dec. 2017, doi: 10.1101/gr.224964.117.

[17] S. Chen *et al.*, 'Codon-Resolution Analysis Reveals a Direct and Context-Dependent Impact of Individual Synonymous Mutations on mRNA Level', *Molecular Biology and Evolution*, vol. 34, no. 11, pp. 2944–2958, Nov. 2017, doi: 10.1093/molbev/msx229.

[18] O. Shalem *et al.*, 'Systematic Dissection of the Sequence Determinants of Gene 3' End Mediated Expression Control', *PLOS Genetics*, vol. 11, no. 4, p. e1005147, Apr. 2015, doi: 10.1371/journal.pgen.1005147.

[19] L. Meneu *et al.*, 'Sequence-dependent activity and compartmentalization of foreign DNA in a eukaryotic nucleus', *Science*, vol. 387, no. 6734, p. eadm9466, Feb. 2025, doi: 10.1126/science.adm9466.

[20] É. Caudal *et al.*, 'Pan-transcriptome reveals a large accessory genome contribution to gene expression variation in yeast', *Nature Genetics*, vol. 56, no. 6, pp. 1278–1287, June 2024, doi: 10.1038/s41588-024-01769-9.

[21] R. Kita, S. Venkataram, Y. Zhou, and H. B. Fraser, 'High-resolution mapping of cis-regulatory variation in budding yeast', *Proceedings of the National Academy of Sciences*, vol. 114, no. 50, pp. E10736–E10744, Dec. 2017, doi: 10.1073/pnas.1717421114.

A Appendix

A.1 Additional Result: S2E model predictions weakly correlate with true coverage of exogenous bacterial DNA

Motivation None of the other benchmark tasks tests the models on the task for which they were trained: RNA-seq track prediction. Testing on the native genome is difficult because Shorkie and Yorzoi have different, potentially overlapping training and test sets. *Meneu et al.* [19] transform two bacterial chromosomes into yeast and profile their expression using RNA-seq, providing a clean held-out test set comprising 2 Mbp of DNA sequence. Because one of the bacterial chromosomes has a substantially lower GC content, this experiment further tests the models' ability to generalise to out-of-distribution sequences.

Task Both models predict RNA-seq coverage profiles in a zero-shot setting for the two bacterial chromosomes, which are then compared with the measured RNA-seq coverage profiles.

Data *Meneu et al.* build two yeast strains, each with one additional linearised chromosome. One chromosome comes from *M. pneumoniae*, is roughly 818 kbp long, has a yeast-matched GC content of 40%, and is largely transcribed. The other, longer chromosome comes from the bacterium *M. mycoides*, is approximately 1.2 Mbp long, and has a lower GC content of 24%, which also substantially decreases its average expression.

Evaluation Because the input sequence window of neither Shorkie nor Yorzoi spans the full genomic sequence, we tile each bacterial genome according to the output-window length of the respective model (Yorzoi: 3 kbp; Shorkie: approximately 14 kbp) and predict coverage profiles using the full input window (i.e., the sequence corresponding to the respective tile plus flanking sequences extending to the boundaries of the input window). The resulting coverage profile for the full chromosome is then split into the same tiles for both models and evaluated against ground truth with two metrics: bin-wise Pearson correlation and Jensen-Shannon divergence of the normalised profiles. Here, normalising a profile means dividing each predicted value by the sum of the bins in a given window so that the bins sum to 1. Formally,

$$\boldsymbol{y} \in \mathbb{R}^{L}_{\geq 0}, \quad \hat{\boldsymbol{y}} \coloneqq f(x) \in \mathbb{R}^{L}_{\geq 0}, \tag{11}$$

where x is the input sequence and f is the respective model. Furthermore,

$$\boldsymbol{p} \coloneqq \frac{\boldsymbol{y}}{\|\boldsymbol{y}\|_1}, \quad \hat{\boldsymbol{p}} \coloneqq \frac{\hat{\boldsymbol{y}}}{\|\hat{\boldsymbol{y}}\|_1}, \quad \boldsymbol{p}, \hat{\boldsymbol{p}} \in \Delta^{L-1}. \tag{12}$$

where $\boldsymbol{y} \neq \mathbf{0}$, $\hat{\boldsymbol{y}} \neq \mathbf{0}$, and Δ^{L-1} denotes the $(L-1)$-dimensional probability simplex.

$$r_{\text{bin}} \coloneqq r(\boldsymbol{y}, \hat{\boldsymbol{y}}), \quad D_{\text{JS}} \coloneqq \text{JSD}(\boldsymbol{p}, \hat{\boldsymbol{p}}). \tag{13}$$

where r is the usual bin-wise Pearson correlation.

Results Yorzoi consistently outperforms Shorkie across both bacterial chromosomes and all evaluated dimensions (Table 5). Its median per-window Pearson correlation is higher and its Jensen–Shannon divergence is lower in every comparison. Nevertheless, the absolute correlations remain low: even Yorzoi reaches only $r_{\text{bin}} = 0.38$ on *M. mycoides* and $r_{\text{bin}} = 0.11$ on *M. pneumoniae*. *M. mycoides* has approximately three-fold higher Pearson correlation than *M. pneumoniae* for both models, despite its substantially more out-of-distribution GC content (24% compared with 40% for *M. pneumoniae* and approximately 38% for yeast). In contrast, *M. mycoides* has higher Jensen-Shannon divergence, indicating that the models place transcriptional mass less accurately within each window.

A.2 Additional result: S2E models weakly distinguish eQTLs from matched variants

Table 5: Results for *M. mycoides* (Mmmyco) and *M. pneumoniae* (Mpneumo). Bold denotes the best-performing model. JSD refers to Jensen–Shannon divergence; ↓ indicates that lower values are better.

Benchmark	Metric	Shorkie	Yorzoi
***M. mycoides* (Mmmyco)**			
Overall	Bin-wise Pearson	0.2638	**0.3843**
Overall	Shape JSD ↓	0.2844	**0.2382**
***M. pneumoniae* (Mpneumo)**			
Overall	Bin-wise Pearson	0.0912	**0.1103**
Overall	Shape JSD ↓	0.0982	**0.0826**

Table 6: Results on eQTL benchmark tasks. Values are means across four negative sets; standard errors (in parentheses) are in units of 10^{-3}.

Model	Kita (AUROC)	Kita (AUPRC)	Caudal (AUROC)	Caudal (AUPRC)
Shorkie	**0.67 (8.7)**	**0.64 (9.7)**	**0.57 (1.5)**	**0.58 (2.2)**
Yorzoi	0.60 (2.8)	0.57 (4.8)	0.53 (0.7)	0.54 (1.5)

Motivation A particularly difficult test of how sequence regulates gene expression is predicting the effect of a single-nucleotide change in the input sequence on the expression of a gene. This task is challenging because none of the evaluated models sees sequences with only one or a few nucleotides changed during training.

Task We use a collection of expression quantitative loci (eQTLs) and ask our models to classify single-nucleotide variants into two classes: eQTL or non-eQTL. Here, an eQTL is a (gene, variant) pair in which the presence of the variant is associated with a significant change in the expression of the selected gene. This benchmark task was adapted from [6].

Data We derive eQTLs from two studies, *Caudal et al.* [20] and *Kita et al.* [21], which use two yeast collections. *Caudal et al.* test gene–variant associations in the 1,011-yeast-genome collection, while *Kita et al.* use a smaller set of 86 yeast isolates. Variant–gene pairs are further filtered so that the variant and associated gene fit within the coverage windows of both models. This yields 1,850 and 605 SNP–eQTLs for [20] and [21], respectively. As described in [6], we build four sets of distance-matched negative *non-eQTL* variant–gene pairs by matching each eQTL pair with a distance-matched variant from the same studies that is not an eQTL.

Evaluation For each (variant v, gene g) pair (a true eQTL or a matched negative), we make two predictions: one for the reference sequence (x_{ref}) and one for the sequence containing variant v (x_{alt}). Under the assumption that eQTLs should produce larger changes in gene expression than non-QTL variants, we compute the log-fold change for gene g between x_{ref} and x_{alt}:

$$s(v, g) := \left| \log_2\big(\varphi_g(x_{\mathrm{alt}}) + 1\big) - \log_2\big(\varphi_g(x_{\mathrm{ref}}) + 1\big) \right| \tag{14}$$

where i and j are the start and end positions of gene g's coding sequence in the model output. We use $s(v, g)$ to discriminate between the classes (with $s(v, g) > \mathrm{threshold}$ classified as eQTL). We then compute AUROC and AUPRC independently for the four negative sets and report the mean AUROC/AUPRC and standard error.

Results Shorkie and Yorzoi both show slightly better-than-random performance at distinguishing eQTL (variant, gene) pairs from non-eQTL pairs (Table 6). On *Kita et al.* in particular, Shorkie shows some ability to distinguish non-eQTLs from eQTLs. Additional statistical fine-mapping might help move beyond effect direction and determine the models' ability to predict effect sizes as well.

A.3 Model training, data and architecture details (*Shorkie* and *Yorzoi*)

The two sequence-to-expression models that we investigate in the benchmark are *Shorkie* (*Chao et al.* [6]) and *Yorzoi* (*Schneider et al.* [7]). Both are trained on sequences from the model organism *S.*

cerevisiae (baker's yeast). Although both models predict RNA-seq tracks from yeast DNA sequences, there are a few differences that make them worth comparing. First, Shorkie fine-tunes ShorkieLM, a masked language model trained on 165 yeast species, to predict expression and ChIP-seq tracks, and has an input sequence window of approximately 16 kilobase pairs (kbp). Its predicted RNA-seq tracks are unstranded. Yorzoi, on the other hand, is trained on long-read RNA-sequencing data; its training data supplement native yeast DNA sequences with human DNA profiled in yeast and structurally rearranged yeast sequences. It has a shorter input sequence window of approximately 5 kbp. Its predicted RNA-seq tracks are stranded, i.e., one track for the forward strand and one for the reverse strand in each RNA-seq experiment. For Yorzoi, a pair of tracks represents the *coverage* of the two DNA strands, whereas Shorkie predicts only unstranded RNA-seq tracks, and thus one track represents one experiment.

A.4 MPRA metrics calculation and marginalisation procedure

The Rafi, Shalem, and Chen assays measure synthetic sequences in reporter constructs whose sequence context differs substantially from the native genomic contexts on which the sequence-to-expression models were trained. We therefore evaluate each assayed sequence across multiple native host loci.

For an assayed sequence v and host locus $h \in H$, let $x_{v,h}^{\mathrm{alt}}$ denote the reconstructed sequence containing v, and let $x_{v,h}^{\mathrm{ref}}$ denote its assay-specific reference sequence in the same host context. Let $g_{v,h}$ denote the gene whose predicted expression is read out. Using the scalar transcript-abundance readout defined in (2), we calculate the context-specific log-expression difference

$$d(v,h) \coloneqq \log_2\Big(\varphi_{g_{v,h}}\Big(x_{v,h}^{\mathrm{alt}}\Big)+1\Big) - \log_2\Big(\varphi_{g_{v,h}}\Big(x_{v,h}^{\mathrm{ref}}\Big)+1\Big). \tag{15}$$

We then average the context-specific differences over the fixed set of host loci:

$$m_v \coloneqq \frac{1}{|H|}\sum_{h\in H} d(v,h). \tag{16}$$

Thus, m_v is one scalar prediction for assayed sequence v. We compare these predictions with the corresponding experimental measurements using Pearson correlation r and Spearman rank correlation ρ. The host panels are fixed before evaluation and contain expressed genes spanning different expression levels and both genomic strands.

Rafi promoter assay For Rafi, v is the 110-base-pair assayed sequence comprising the 80-base-pair variable promoter and its fixed flanks. We insert it 180 base pairs upstream of the transcription start site of each of 22 native host genes. The alternative sequence contains the insert, whereas the reference sequence is the corresponding unmodified native locus. This is the procedure used in [6].

Shalem terminator assay For Shalem, v is the 150-base-pair assayed 3′-end sequence. At each of 22 native host genes, we replace the 450-base-pair region immediately downstream of the stop codon with v followed by a fixed 300-base-pair non-terminating CYC1-derived sequence. The reference is the corresponding unmodified native locus.

Chen coding-sequence assay For Chen, each synonymous variant is evaluated at 20 host genes that are active in yeast-extract–peptone–dextrose conditions. At each locus, the host coding sequence is replaced by the GFP or TDH3 reporter coding sequence and its downstream cassette while retaining the native promoter and genomic context. The alternative and reference constructs differ only in the assayed 36-nucleotide coding-sequence block. Here, $g_{v,h}$ denotes the inserted GFP or TDH3 reporter rather than the host gene, and the reporter coding sequence is used as the readout region.

Cuperus 5′-UTR assay We do not marginalise the Cuperus sequences. Each variable 5′ untranslated region is evaluated directly in the fixed CYC1-promoter–HIS3–CYC1-terminator construct used by the benchmark. For sequence v, the prediction is

$$\varphi_{\mathrm{HIS3}}(x_v). \tag{17}$$

The marginalisation procedure applies only to the genomic sequence-to-expression models. The supervised DREAM-RNN reference for Rafi predicts expression directly from the original reporter-assay sequence and is therefore not scored using (16).

A.5 *Brooks et al.* benchmark preprocessing and calculation

We use three replicate RNA-seq experiments of unscrambled SynIXR JS94 strains from *Brooks et al.*. This means that while strains have no induced genetic rearrangements, the right arm of chromosome 9 is synthetic with loxPsym sites (used for recombination) installed after approximately every non-essential coding sequence before the native 3′ UTR sequence. We compute LFCs between the rearranged sequence and the parental control separately for each replicate yielding three true LFCs. To account for potential dropouts, we only keep true LFCs where the parental strain's gene has at least 10 reads. This results in:

- 205 samples for replicate r0
- 80 samples for replicate r1
- 291 samples for replicate r2

We compute correlations for each of the replicates separately. Additionally, we compute a leave-one-out ceiling which simply uses the average LFC of two replicates to "predict" the third replicate.

A.6 *Hong et al.* boundary-distance definition

Hsieh et al. [14] measure Micro-C interactions across the yeast genome and define *chromosomal interaction domains* delineated by boundaries. *Hong et al.* find that the distance from an integration site to the nearest such boundary predicts integrated-reporter expression. Boundaries are defined as local minima in the number of Micro-C interactions crossing a genomic position:

$$\text{boundary_distance}_i = \min_{b \in B} |\text{insertion_site}_i - b|, \tag{18}$$

where

$$B = \left\{ i \in [L] \mid C(i) = \min_{k \in \mathcal{N}(i)} C(k) \right\} \tag{19}$$

and

$$C(i) = \sum_{(x,y) \in \mathcal{E}} \mathbb{1}[x < i < y] \mathbb{1}[500 < y - x < 10,000]. \tag{20}$$

Here, $\mathcal{E}$ is the set of Micro-C interaction pairs with ordered endpoints $x < y$, $[L]$ is the set of positions on the chromosome, and $\mathcal{N}(i)$ is a local neighborhood around i.

B Use of LLMs for creating this work

LLMs were involved in creating this work through coding agents that helped to write, debug, and test the code produced for this benchmark. LLMs were also used to write parts of the code documentation (`docs/` in the repository). Additionally, they were used to improve the fluency and correct the grammar and spelling of the manuscript.