

1 Solving Equations

Solving Equations

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Solving any equation

$$f(x) = g(x)$$

Solving Equations

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Solving any equation

$$f(x) = g(x)$$

can be recast as a **root-finding exercise**

$$f(x) - g(x) = 0.$$

Solving Equations

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Solving any equation

$$f(x) = g(x)$$

can be recast as a **root-finding exercise**

$$f(x) - g(x) = 0.$$

Sometimes **closed-form** solutions exist, e.g. for polynomials with degree ≤ 4

$$ax^2 + bx + c = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Solving Equations

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Solving any equation

$$f(x) = g(x)$$

can be recast as a **root-finding exercise**

$$f(x) - g(x) = 0.$$

Sometimes **closed-form** solutions exist, e.g. for polynomials with degree ≤ 4

$$ax^2 + bx + c = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

but for most equations, including almost all polynomials degree ≥ 5 , **no closed-form solution exists**

$$x^5 - x - 1 = 0 \implies x = 1.1673\dots$$

Solving Equations

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Solving any equation

$$f(x) = g(x)$$

can be recast as a **root-finding exercise**

$$f(x) - g(x) = 0.$$

Sometimes **closed-form** solutions exist, e.g. for polynomials with degree ≤ 4

$$ax^2 + bx + c = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

but for most equations, including almost all polynomials degree ≥ 5 , **no closed-form solution exists**

$$x^5 - x - 1 = 0 \implies x = 1.1673\dots$$

Q. How do we solve such equations?

Solving Equations

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Solving any equation

$$f(x) = g(x)$$

can be recast as a **root-finding exercise**

$$f(x) - g(x) = 0.$$

Sometimes **closed-form** solutions exist, e.g. for polynomials with degree ≤ 4

$$ax^2 + bx + c = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

but for most equations, including almost all polynomials degree ≥ 5 , **no closed-form solution exists**

$$x^5 - x - 1 = 0 \implies x = 1.1673\dots$$

Q. How do we solve such equations?

A. Make a **guess**, and try to **improve** it.

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Consider that deceptively difficult quintic,

$$x^5 - x - 1 = 0$$

which has a solution when $x = \xi$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Consider that deceptively difficult quintic,

$$x^5 - x - 1 = 0$$

which has a solution when $x = \xi$

We can rearrange to express x as a **slowly-varying** function of x ,

$$x = \phi(x) = \sqrt[5]{x+1}$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Consider that deceptively difficult quintic,

$$x^5 - x - 1 = 0$$

which has a solution when $x = \xi$

We can rearrange to express x as a **slowly-varying** function of x ,

$$x = \phi(x) = \sqrt[5]{x + 1}$$

REARRANGEMENT METHOD

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Consider that deceptively difficult quintic,

$$x^5 - x - 1 = 0$$

which has a solution when $x = \xi$

We can rearrange to express x as a **slowly-varying** function of x ,

$$x = \phi(x) = \sqrt[5]{x + 1}$$

REARRANGEMENT METHOD

At the **solution** to the quintic, **ϕ is equal to x** ,

$$\phi(\xi) = \xi$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Consider that deceptively difficult quintic,

$$x^5 - x - 1 = 0$$

which has a solution when $x = \xi$

We can rearrange to express x as a **slowly-varying** function of x ,

$$x = \phi(x) = \sqrt[5]{x + 1}$$

REARRANGEMENT METHOD

At the **solution** to the quintic, **ϕ is equal to x** ,

$$\phi(\xi) = \xi$$

but at other points x_i there is an **error**, **ϵ_i** ,

$$\phi(x_i) = \xi + \epsilon_i$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Idea:

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Idea:

- 1 start with some guess, x_0 ;

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Idea:

- 1 start with some guess, x_0 ;
- 2 compute a new guess, $x_1 = \phi(x_0)$, and hope its closer to ξ ;

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Idea:

- 1 start with some guess, x_0 ;
- 2 compute a new guess, $x_1 = \phi(x_0)$, and hope its closer to ξ ;
- 3 iterate until $|x_{k+1} - x_k|$ is less than some error tolerance.

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Idea:

- 1 start with some guess, x_0 ;
- 2 compute a new guess, $x_1 = \phi(x_0)$, and hope its closer to ξ ;
- 3 iterate until $|x_{k+1} - x_k|$ is less than some error tolerance.

Problem: how do we know if the sequence $x_0, x_1, \dots, x_k, x_{k+1}$ **converges** to ξ ?

$$\lim_{k \rightarrow \infty} \phi(x_k) = \xi$$

Convergence

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

In order for this sequence to converge, we require the **absolute error**

$$|\epsilon_k| = |\xi - x_k|$$

to **decrease after each iteration**.

In order for this sequence to converge, we require the **absolute error**

$$|\epsilon_k| = |\xi - x_k|$$

to **decrease after each iteration**.

Starting with,

$$\xi + \epsilon_{k+1} = x_{k+1} = \phi(x_k) = \phi(\xi + \epsilon_k),$$

In order for this sequence to converge, we require the **absolute error**

$$|\epsilon_k| = |\xi - x_k|$$

to **decrease after each iteration**.

Starting with,

$$\xi + \epsilon_{k+1} = x_{k+1} = \phi(x_k) = \phi(\xi + \epsilon_k),$$

we expand the right-most term as a Taylor series,

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \phi'(\xi) + \mathcal{O}(\epsilon_k^2)$$

$$\epsilon_{k+1} = \epsilon_k \phi'(\xi) + \mathcal{O}(\epsilon_k^2)$$

In order for this sequence to converge, we require the **absolute error**

$$|\epsilon_k| = |\xi - x_k|$$

to **decrease after each iteration**.

Starting with,

$$\xi + \epsilon_{k+1} = x_{k+1} = \phi(x_k) = \phi(\xi + \epsilon_k),$$

we expand the right-most term as a Taylor series,

$$\begin{aligned}\xi + \epsilon_{k+1} &= \phi(\xi) + \epsilon_k \phi'(\xi) + \mathcal{O}(\epsilon_k^2) \\ \epsilon_{k+1} &= \epsilon_k \phi'(\xi) + \mathcal{O}(\epsilon_k^2)\end{aligned}$$

therefore a **necessary** (but not sufficient) condition for the sequence to converge to ξ is

$$\|\phi'(\xi)\| < 1$$

Q-Convergence

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

How quickly does $\phi(x)$ converge to ξ ?

How quickly does $\phi(x)$ converge to ξ ?

In general we can write,

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|^q} = \mu \quad (1)$$

Q-CONVERGENCE

How quickly does $\phi(x)$ converge to ξ ?

In general we can write,

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|^q} = \mu \quad (1)$$

Q-CONVERGENCE

where q is the **order of convergence** and μ is the **rate of convergence**

How quickly does $\phi(x)$ converge to ξ ?

In general we can write,

$$\boxed{\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|^q} = \mu} \quad (1)$$

Q-CONVERGENCE

where q is the **order of convergence** and μ is the **rate of convergence**

In the example,

$$\epsilon_{k+1} = \epsilon_k \phi'(\xi) + \mathcal{O}(\epsilon_k^2)$$

How quickly does $\phi(x)$ converge to ξ ?

In general we can write,

$$\boxed{\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|^q} = \mu} \quad (1)$$

Q-CONVERGENCE

where q is the **order of convergence** and μ is the **rate of convergence**

In the example,

$$\epsilon_{k+1} = \epsilon_k \phi'(\xi) + \mathcal{O}(\epsilon_k^2)$$

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|} = |\phi'(\xi)|$$

$$q = 1 \text{ and } \mu = |\phi'(\xi)|$$

How quickly does $\phi(x)$ converge to ξ ?

In general we can write,

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|^q} = \mu \quad (1)$$

Q-CONVERGENCE

where q is the **order of convergence** and μ is the **rate of convergence**

In the example,

$$\epsilon_{k+1} = \epsilon_k \phi'(\xi) + \mathcal{O}(\epsilon_k^2)$$

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|} = |\phi'(\xi)|$$

$$q = 1 \text{ and } \mu = |\phi'(\xi)|$$

we say that the sequence converges **linearly** to ξ with rate $|\phi'(\xi)|$.

If ϕ' **vanishes at the solution**, but ϕ'' is finite and non-zero,

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \phi'(\xi) + \frac{1}{2} \epsilon_k^2 \phi''(\xi) + \mathcal{O}(\epsilon_k^3)$$

If ϕ' **vanishes at the solution**, but ϕ'' is finite and non-zero,

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \phi'(\xi) + \frac{1}{2} \epsilon_k^2 \phi''(\xi) + \mathcal{O}(\epsilon_k^3)$$

If ϕ' vanishes at the solution, but ϕ'' is finite and non-zero,

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \phi'(\xi) + \frac{1}{2} \epsilon_k^2 \phi''(\xi) + \mathcal{O}(\epsilon_k^3)$$

$$\epsilon_{k+1} = \epsilon_k^2 \phi''(\xi) + \mathcal{O}(\epsilon_k^3)$$

If ϕ' vanishes at the solution, but ϕ'' is finite and non-zero,

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \phi'(\xi) + \frac{1}{2} \epsilon_k^2 \phi''(\xi) + \mathcal{O}(\epsilon_k^3)$$

$$\frac{\epsilon_{k+1}}{\epsilon_k^2} = \frac{1}{2} \phi''(\xi) + \mathcal{O}(\epsilon_k)$$

If ϕ' **vanishes at the solution**, but ϕ'' is finite and non-zero,

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \phi'(\xi) + \frac{1}{2} \epsilon_k^2 \phi''(\xi) + \mathcal{O}(\epsilon_k^3)$$

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|^2} = \frac{1}{2} \phi''(\xi)$$

If ϕ' **vanishes at the solution**, but ϕ'' is finite and non-zero,

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \phi'(\xi) + \frac{1}{2} \epsilon_k^2 \phi''(\xi) + \mathcal{O}(\epsilon_k^3)$$

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|^2} = \frac{1}{2} \phi''(\xi)$$

$q = 2$, $\mu = \frac{1}{2} \phi''(\xi)$, and the sequence converges **quadratically** to the limit ξ .

If ϕ' **vanishes at the solution**, but ϕ'' is finite and non-zero,

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \phi'(\xi) + \frac{1}{2} \epsilon_k^2 \phi''(\xi) + \mathcal{O}(\epsilon_k^3)$$

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|^2} = \frac{1}{2} \phi''(\xi)$$

$q = 2$, $\mu = \frac{1}{2} \phi''(\xi)$, and the sequence converges **quadratically** to the limit ξ .

A heuristic to estimate q is given by,

$$q \approx \frac{\log \left| \frac{x_{k+1} - x_k}{x_k - x_{k-1}} \right|}{\log \left| \frac{x_k - x_{k-1}}{x_{k-1} - x_{k-2}} \right|} \quad (2)$$

Q-CONVERGENCE RATE ESTIMATE

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\sqrt[5]{x_k + 1}}{x_k + 1}$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$x_{k+1} = \phi(x_k)$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$\xi + \epsilon_{k+1} = \phi(\xi + \epsilon_k)$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \phi'(\xi) + \mathcal{O}(\epsilon_k^2)$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$\xi + \epsilon_{k+1} = \phi(\xi) + \epsilon_k \frac{1}{5} \frac{\phi(\xi)}{\xi + 1} + \mathcal{O}(\epsilon_k^2)$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$\xi + \epsilon_{k+1} = \xi + \epsilon_k \frac{1}{5} \frac{\xi}{\xi + 1} + \mathcal{O}(\epsilon_k^2)$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$\epsilon_{k+1} = \epsilon_k \frac{1}{5} \frac{\xi}{\xi + 1} + \mathcal{O}(\epsilon_k^2)$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$\frac{\epsilon_{k+1}}{\epsilon_k} = \frac{1}{5} \frac{\xi}{\xi + 1} + \mathcal{O}(\epsilon_k)$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|} = \left| \frac{1}{5} \frac{\xi}{\xi + 1} \right|$$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|} = \left| \frac{1}{5} \frac{\xi}{\xi + 1} \right|$$

A. $q = 1$, $\mu \approx 0.11 < 0$

Solving the Quintic

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

Q. Does the quintic sequence converge?

$$x^5 - x - 1 = 0$$

$$x_{k+1} = \phi(x_k) = \sqrt[5]{x_k + 1}$$

$$\phi'(x) = \frac{1}{5} \frac{\phi(x)}{x_k + 1}$$

$$\lim_{k \rightarrow \infty} \frac{|\epsilon_{k+1}|}{|\epsilon_k|} = \left| \frac{1}{5} \frac{\xi}{\xi + 1} \right|$$

A. $q = 1$, $\mu \approx 0.11 < 1$

the sequence **converges** to the solution for **all** starting values x_k .

Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 find an interval (a, b) that contains a zero;

Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 find an interval (a, b) that contains a zero;
- 2 find $c = (a + b)/2$;

Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 find an interval (a, b) that contains a zero;
- 2 find $c = (a + b)/2$;
- 3 set $(a, b) \leftarrow (a, c)$ or $(a, b) \leftarrow (c, b)$ such that (a, b) still contains a zero

Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 find an interval (a, b) that contains a zero;
- 2 find $c = (a + b)/2$;
- 3 set $(a, b) \leftarrow (a, c)$ or $(a, b) \leftarrow (c, b)$ such that (a, b) still contains a zero
- 4 continue until $|a - b| < \delta$

- 1 find an interval (a, b) that contains a zero;
- 2 find $c = (a + b)/2$;
- 3 set $(a, b) \leftarrow (a, c)$ or $(a, b) \leftarrow (c, b)$ such that (a, b) still contains a zero
- 4 continue until $|a - b| < \delta$

$$a_{k+1}, b_{k+1} = \phi(a_k, b_k) = \begin{cases} \left(a_k, \frac{a_k + b_k}{2}\right) & \text{if } f\left(\frac{a_k + b_k}{2}\right) > 0; \\ \left(\frac{a_k + b_k}{2}, b_k\right) & \text{if } f\left(\frac{a_k + b_k}{2}\right) < 0. \end{cases} \quad (3)$$


BISECTION UPDATE

Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Bisection

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations



Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose an interval (x_0, x_1) , ideally containing a zero;

Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose an interval (x_0, x_1) , ideally containing a zero;
- 2 find the secant line that passes through $(x_0, f_0), (x_1, f_1)$

$$y = \frac{f_1 - f_0}{x_1 - x_0}(x - x_1) + f_1$$

Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose an interval (x_0, x_1) , ideally containing a zero;
- 2 find the secant line that passes through $(x_0, f_0), (x_1, f_1)$

$$y = \frac{f_1 - f_0}{x_1 - x_0}(x - x_1) + f_1$$

- 3 find the zero of the secant line, x_2

$$x_2 = x_1 - f_1 \frac{x_1 - x_0}{f_1 - f_0}$$

Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose an interval (x_0, x_1) , ideally containing a zero;
- 2 find the secant line that passes through $(x_0, f_0), (x_1, f_1)$

$$y = \frac{f_1 - f_0}{x_1 - x_0}(x - x_1) + f_1$$

- 3 find the zero of the secant line, x_2

$$x_2 = x_1 - f_1 \frac{x_1 - x_0}{f_1 - f_0}$$

- 4 $(x_0, x_1) \leftarrow (x_1, x_2)$

Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose an interval (x_0, x_1) , ideally containing a zero;
- 2 find the secant line that passes through $(x_0, f_0), (x_1, f_1)$

$$y = \frac{f_1 - f_0}{x_1 - x_0}(x - x_1) + f_1$$

- 3 find the zero of the secant line, x_2

$$x_2 = x_1 - f_1 \frac{x_1 - x_0}{f_1 - f_0}$$

- 4 $(x_0, x_1) \leftarrow (x_1, x_2)$
- 5 continue until $|x_0 - x_1| < \delta$

Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose an interval (x_0, x_1) , ideally containing a zero;
- 2 find the secant line that passes through $(x_0, f_0), (x_1, f_1)$

$$y = \frac{f_1 - f_0}{x_1 - x_0}(x - x_1) + f_1$$

- 3 find the zero of the secant line, x_2

$$x_2 = x_1 - f_1 \frac{x_1 - x_0}{f_1 - f_0}$$

- 4 $(x_0, x_1) \leftarrow (x_1, x_2)$
- 5 continue until $|x_0 - x_1| < \delta$

$$(x_{k+1}, x_{k+2}) = \left(x_{k+1}, \left\{ x_{k+1} - f_{k+1} \frac{x_{k+1} - x_k}{f_{k+1} - f_k} \right\} \right) \quad (4)$$


SECANT UPDATE

Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Secant Method

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations



Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose a point x_0 , ideally near a zero;

Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose a point x_0 , ideally near a zero;
- 2 find the tangent to the curve at x_0

$$y = f'_0(x - x_0) + f_0$$

Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose a point x_0 , ideally near a zero;
- 2 find the tangent to the curve at x_0

$$y = f'_0(x - x_0) + f_0$$

- 3 find the zero of the tangent line, x_1

$$x_1 = x_0 - \frac{f_0}{f'_0}$$

Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

1 choose a point x_0 , ideally near a zero;

2 find the tangent to the curve at x_0

$$y = f'_0(x - x_0) + f_0$$

3 find the zero of the tangent line, x_1

$$x_1 = x_0 - \frac{f_0}{f'_0}$$

4 $x_0 \leftarrow x_1$

Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

1 choose a point x_0 , ideally near a zero;

2 find the tangent to the curve at x_0

$$y = f'_0(x - x_0) + f_0$$

3 find the zero of the tangent line, x_1

$$x_1 = x_0 - \frac{f_0}{f'_0}$$

4 $x_0 \leftarrow x_1$

5 continue until $|x_0 - x_1| < \delta$

Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

1 choose a point x_0 , ideally near a zero;

2 find the tangent to the curve at x_0

$$y = f'_0(x - x_0) + f_0$$

3 find the zero of the tangent line, x_1

$$x_1 = x_0 - \frac{f_0}{f'_0}$$

4 $x_0 \leftarrow x_1$

5 continue until $|x_0 - x_1| < \delta$

$$x_{k+1} = x_k - \frac{f_k}{f'_k}$$

(5)


NEWTON–RAPHSOIN UPDATE

Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Newton–Raphson

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations



Inverse Quadratic Interpolation

- 1 choose three non-collinear points (x_0, x_1, x_2) , ideally near a zero;

Inverse Quadratic Interpolation

- 1 choose three non-collinear points (x_0, x_1, x_2) , ideally near a zero;
- 2 find the quadratic function of y that passes through f at those points,

$$f^{-1}(y) = \sum_{i=0}^2 x_i \prod_{i \neq j} \frac{y - f_j}{f_i - f_j}$$

Inverse Quadratic Interpolation

- 1 choose three non-collinear points (x_0, x_1, x_2) , ideally near a zero;
- 2 find the quadratic function of y that passes through f at those points,

$$f^{-1}(y) = \sum_{i=0}^2 x_i \prod_{i \neq j} \frac{y - f_j}{f_i - f_j}$$

- 3 find the zero of that quadratic function, x_3

$$x_3 = f^{-1}(0) = \sum_{i=0}^2 x_i \prod_{i \neq j} \frac{f_j}{f_i - f_j}$$

Inverse Quadratic Interpolation

- 1 choose three non-collinear points (x_0, x_1, x_2) , ideally near a zero;
- 2 find the quadratic function of y that passes through f at those points,

$$f^{-1}(y) = \sum_{i=0}^2 x_i \prod_{i \neq j} \frac{y - f_j}{f_i - f_j}$$

- 3 find the zero of that quadratic function, x_3

$$x_3 = f^{-1}(0) = \sum_{i=0}^2 x_i \prod_{i \neq j} \frac{f_j}{f_i - f_j}$$

- 4 $(x_0, x_1, x_2) \leftarrow (x_1, x_2, x_3)$

Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose three non-collinear points (x_0, x_1, x_2) , ideally near a zero;
- 2 find the quadratic function of y that passes through f at those points,

$$f^{-1}(y) = \sum_{i=0}^2 x_i \prod_{i \neq j} \frac{y - f_j}{f_i - f_j}$$

- 3 find the zero of that quadratic function, x_3

$$x_3 = f^{-1}(0) = \sum_{i=0}^2 x_i \prod_{i \neq j} \frac{f_j}{f_i - f_j}$$

- 4 $(x_0, x_1, x_2) \leftarrow (x_1, x_2, x_3)$
- 5 continue until $|x_3 - x_2| < \delta$

Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations

- 1 choose three non-collinear points (x_0, x_1, x_2) , ideally near a zero;
- 2 find the quadratic function of y that passes through f at those points,

$$f^{-1}(y) = \sum_{i=0}^2 x_i \prod_{i \neq j} \frac{y - f_j}{f_i - f_j}$$

- 3 find the zero of that quadratic function, x_3

$$x_3 = f^{-1}(0) = \sum_{i=0}^2 x_i \prod_{i \neq j} \frac{f_j}{f_i - f_j}$$

- 4 $(x_0, x_1, x_2) \leftarrow (x_1, x_2, x_3)$
- 5 continue until $|x_3 - x_2| < \delta$

$$\begin{aligned} x_{k+1} = & \frac{f_{k-1} f_k}{(f_{k-2} - f_{k-1})(f_{k-2} - f_k)} x_{k-2} \\ & + \frac{f_{k-2} f_k}{(f_{k-1} - f_{k-2})(f_{k-1} - f_k)} x_{k-1} \\ & + \frac{f_{k-2} f_{k-1}}{(f_k - f_{k-2})(f_k - f_{k-1})} x_k \end{aligned}$$

(6)


IQR UPDATE

Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations




Inverse Quadratic Interpolation

Intro to
Comp. Phys.

James
Farrell &
Jure
Dobnikar

Solving
Equations



Pathological Cases

Implement the five root-finding methods we have studied, and use them to find the roots of,

1 $x^5 - x - 1 = 0;$

2 $16x^4 - 8x + 3 = 0;$

3 $x^3 - 2x^2 - 11x + 12 = (x - 4)(x - 1)(x + 3) = 0,$

for a range of starting values (plot the curves to get an idea of what a sensible range of values might be). What do you notice? Do all of the methods find all of the roots? Estimate the order of convergence using the formula given above. Is the order of convergence always the theoretical maximum?