

JAMA | Original Investigation | CARING FOR THE CRITICALLY ILL PATIENT

# High-Flow Nasal Oxygen vs Noninvasive Ventilation in Patients With Acute Respiratory Failure

## The RENOVATE Randomized Clinical Trial

RENOVATE Investigators and the BRICNet Authors

**IMPORTANCE** High-flow nasal oxygen (HFNO) and noninvasive ventilation (NIV) are commonly used respiratory support therapies for patients with acute respiratory failure (ARF).

**OBJECTIVE** To assess whether HFNO is noninferior to NIV on the rates of endotracheal intubation or death at 7 days in 5 patient groups with ARF.

**DESIGN, SETTING, AND PARTICIPANTS** This noninferiority, randomized clinical trial enrolled hospitalized adults (aged  $\geq 18$  years; classified as 5 patient groups with ARF: nonimmunocompromised with hypoxemia, immunocompromised with hypoxemia, chronic obstructive pulmonary disease [COPD] exacerbation with respiratory acidosis, acute cardiogenic pulmonary edema [ACPE], or hypoxemic COVID-19, which was added as a separate group on June 26, 2023) at 33 hospitals in Brazil between November 2019 and November 2023 (final follow-up: April 26, 2024).

**INTERVENTIONS** High-flow nasal oxygen (n = 883) or NIV (n = 883).

**MAIN OUTCOMES AND MEASURES** The primary outcome was endotracheal intubation or death within 7 days assessed using a bayesian hierarchical model with dynamic borrowing across patient groups. Noninferiority was defined by a posterior probability of 0.992 or greater for an odds ratio (OR) less than 1.55.

**RESULTS** Among 1800 patients, 1766 completed the study (mean age, 64 [SD, 17] years; 707 [40%] were women). The primary outcome of endotracheal intubation or death at 7 days occurred in 39% (344/883) in the HFNO group vs 38% (336/883) in the NIV group. In the immunocompromised with hypoxemia patient group, the primary outcome occurred in 57.1% (16/28) in the HFNO group vs 36.4% (8/22) in the NIV group; enrollment was stopped for futility (final OR, 1.07; 95% credible interval [CrI], 0.81-1.39; noninferiority posterior probability [NPP], 0.989). In the nonimmunocompromised with hypoxemia group, the primary outcome occurred in 32.5% (81/249) in the HFNO group vs 33.1% (78/236) in the NIV group (OR, 1.02 [95% CrI, 0.81-1.26]; NPP, 0.999). In the ACPE group, the primary outcome occurred in 10.3% (14/136) in the HFNO group vs 21.3% (29/136) in the NIV group (OR, 0.97 [95% CrI, 0.73-1.23]; NPP, 0.997). In the hypoxemic COVID-19 group, the primary outcome occurred in 51.3% (223/435) in the HFNO group vs 47.0% (210/447) in the NIV group (OR, 1.13 [95% CrI, 0.94-1.38]; NPP, 0.997). In the COPD exacerbation with respiratory acidosis group, the primary outcome occurred in 28.6% (10/35) in the HFNO group vs 26.2% (11/42) in the NIV group (OR, 1.05 [95% CrI, 0.79-1.36]; NPP, 0.992). However, a post hoc analysis without dynamic borrowing across the 5 ARF patient groups revealed some qualitatively different results in patients with COPD, immunocompromised patients, and patients with ACPE. The incidence of serious adverse events was similar (9.4% of patients in HFNO group vs 9.9% in NIV group).

**CONCLUSIONS AND RELEVANCE** Compared with NIV, HFNO met prespecified criteria for noninferiority for the primary outcome of endotracheal intubation or death within 7 days in 4 of the 5 patient groups with ARF. However, the small sample sizes in some patient groups and the sensitivity of the findings to the choice of analysis model suggests the need for further study in patients with COPD, immunocompromised patients, and patients with ACPE.

**TRIAL REGISTRATION** ClinicalTrials.gov Identifier: [NCT03643939](https://clinicaltrials.gov/ct2/show/study/NCT03643939)

JAMA. 2025;333(10):875-890. doi:[10.1001/jama.2024.26244](https://doi.org/10.1001/jama.2024.26244)  
Published online December 10, 2024.

◀ Editorial pages 848 and 850

+ Multimedia

+ Supplemental content

**Group Information:** The RENOVATE Investigators and BRICNet Authors appear at the end of this article. The RENOVATE Investigators and BRICNet nonauthors appear in Supplement 4.

**Corresponding Author:** Alexandre Biasi Cavalcanti, MD, PhD, Hcor Research Institute, Rua Desembargador Eliseu Guilherme, 200 Paraíso, São Paulo 04004-030, Brazil ([abiasi@hcor.com.br](mailto:abiasi@hcor.com.br)).

**Section Editor:** Christopher Seymour, MD, Associate Editor, JAMA ([christopher.seymour@jamanetwork.org](mailto:christopher.seymour@jamanetwork.org)).

Clinicians use high-flow nasal oxygen and noninvasive ventilation (pressure support ventilation with positive end-expiratory pressure) to treat acute respiratory failure associated with several conditions, including hypoxemic acute respiratory failure in both nonimmunocompromised and immunocompromised patients, exacerbation of chronic obstructive pulmonary disease (COPD), acute cardiogenic pulmonary edema, and COVID-19.<sup>1</sup> However, patients may not always tolerate noninvasive ventilation due to discomfort.<sup>2</sup> Compared with low-flow oxygen, high-flow nasal oxygen offers physiological advantages, including improved oxygenation, reduced dead space, alveolar recruitment, humidification, heating, and enhanced secretion clearance.<sup>3,4</sup> Moreover, high-flow nasal oxygen has been shown to effectively reduce  $\text{Paco}_2$  levels in patients experiencing an exacerbation of COPD,<sup>5,6</sup> decrease cardiac preload, and improve signs of respiratory failure for patients with acute heart failure.<sup>7,8</sup>

Compared with noninvasive ventilation, high-flow nasal oxygen is easier to use and more comfortable for the patient.<sup>9</sup> Patients using a high-flow nasal cannula can eat, drink, and talk more easily than with noninvasive ventilation. However, high-flow nasal oxygen may be less effective than noninvasive ventilation in reducing the workload on respiratory muscles during acute respiratory failure.<sup>10</sup> Evidence on the most appropriate form of noninvasive respiratory support across different causes of acute respiratory failure is limited. Guidelines recommend noninvasive ventilation for acute respiratory failure caused by COPD exacerbation and acute cardiogenic pulmonary edema; however, the evidence<sup>11,12</sup> supporting the use of noninvasive ventilation is based on comparison with low-flow oxygen not high-flow nasal oxygen. High-flow nasal oxygen is preferred over low-flow oxygen for treating patients with hypoxemic acute respiratory failure, including those with immunosuppression or COVID-19.<sup>13,14</sup> However, considerable uncertainty remains regarding the comparative effectiveness of high-flow nasal oxygen vs noninvasive ventilation.<sup>9,15-19</sup>

We conducted a multicenter, adaptive, randomized clinical trial to evaluate the noninferiority of high-flow nasal oxygen compared with noninvasive ventilation and assess potential superiority in reducing the rates of endotracheal intubation or death across 5 patient groups hospitalized with acute respiratory failure (nonimmunocompromised with hypoxemia, immunocompromised with hypoxemia, COPD exacerbation with respiratory acidosis, acute cardiogenic pulmonary edema, or hypoxemic COVID-19).

## Methods

### Study Design and Oversight

The High-Flow Nasal Oxygen Cannula Compared to Non-Invasive Ventilation in Adult Patients With Acute Respiratory Failure (RENOVATE) was a multicenter, adaptive, noninferiority randomized clinical trial conducted at 33 hospitals in Brazil. The main adaptation in this trial was the use of frequent interim analyses that allowed for the stopping of individual patient groups with acute respiratory failure based on futility, noninferiority, or superiority (additional details

## Key Points

**Question** Is high-flow nasal oxygen noninferior to noninvasive ventilation for rate of endotracheal intubation or death at 7 days in patients with acute respiratory failure due to a variety of causes?

**Findings** In this randomized clinical trial ( $n = 1766$  patients), a bayesian hierarchical model with dynamic borrowing across 5 patient groups found high-flow nasal oxygen was noninferior (defined by a posterior probability  $\geq 0.992$  for an odds ratio  $< 1.55$ ) to noninvasive ventilation for rates of endotracheal intubation or death at 7 days in 4 of the patient groups (nonimmunocompromised with hypoxemic acute respiratory failure, chronic obstructive pulmonary disease exacerbation with respiratory acidosis, acute cardiogenic pulmonary edema, and hypoxemic COVID-19). One patient group (immunocompromised with hypoxemic acute respiratory failure) was stopped for futility.

**Meaning** Compared with noninvasive ventilation, high-flow nasal oxygen met prespecified criteria for noninferiority for the outcome of endotracheal intubation or death within 7 days in 4 of the 5 patient groups with acute respiratory failure.

appear in the eResults in [Supplement 1](#)). The trial protocol ([Supplement 2](#)) and statistical analysis plan ([Supplement 3](#)) were published.<sup>20,21</sup> The trial was overseen by an international steering committee and an independent data and safety monitoring committee whose members were unaware of the treatment group assignment (additional details appear in the eMethods in [Supplement 1](#)).

The bayesian adaptive trial design, analysis of the primary outcome, and 28-day mortality assessments were performed by Berry Consultants, LLC. The Hcor Research Institute coordinated the trial and conducted all other statistical analyses. Additional details regarding trial governance and responsibilities appear in the eMethods in [Supplement 1](#). The trial was conducted in accordance with principles of the Declaration of Helsinki,<sup>22</sup> and the trial protocol ([Supplement 2](#)) was approved by the institutional review boards at each site and a national ethics review board. Due to the urgency of clinical decisions regarding respiratory support and based on the recommendation of the national ethics review board, written consent was sought from all patients or their surrogate decision-makers after enrollment in the trial.

### Patients

Adult patients (aged  $\geq 18$  years) were eligible if they were admitted to intensive care units (ICUs), emergency departments, or hospital wards because of acute respiratory failure, which was defined by presence of hypoxemia (oxygen saturation as measured by pulse oximetry [ $\text{SpO}_2$ ]  $< 90\%$  or  $\text{Pao}_2$   $< 60$  mm Hg at room air) and either respiratory effort (use of accessory musculature, paradoxical breathing, or thoracoabdominal asynchrony) or tachypnea (respiratory rate  $> 25$  breaths/min). Patients were initially classified into 1 of the following 4 mutually exclusive patient groups with acute respiratory failure: nonimmunocompromised with hypoxemia, immunocompromised with hypoxemia, COPD exacerbation with respiratory acidosis, or acute cardiogenic pulmonary edema

(additional eligibility criteria appear in the eMethods in [Supplement 1](#)). Hypoxemic COVID-19 was added later and is further described below. Before randomization, the use of low-flow oxygen was allowed for all patients; the use of noninvasive ventilation was allowed for most patients, but not for those with acute cardiogenic pulmonary edema.

Immunocompromised patients were defined as those with (1) regular use of immunosuppressive drugs for more than 3 months or high doses of corticosteroids ( $>0.5$  mg/kg/d); (2) presence of solid organ transplants, solid tumor treated with chemotherapy within the past 5 years, or hematologic malignancy treated within the past 5 years; or (3) diagnosed with AIDS or a primary immunodeficiency.<sup>23</sup> A COPD exacerbation with respiratory acidosis was defined as being present in patients with (1) high clinical suspicion of COPD or previous COPD diagnosis; (2) a respiratory rate greater than 25 breaths/min or use of accessory musculature, paradoxical breathing, or thoracoabdominal asynchrony; or (3) an arterial blood pH level lower than 7.35 and a  $\text{PaCO}_2$  level greater than 45 mm Hg. Acute cardiogenic pulmonary edema was defined as being present in patients (1) with sudden-onset dyspnea and diffuse crackling rales with or without the third audible heart sound; (2) without a history of pulmonary aspiration, infection, or pulmonary fibrosis; (3) with acute cardiogenic pulmonary edema as the primary clinical hypothesis; or (4) with a chest x-ray with bilateral alveolar infiltrates suggestive of pulmonary edema, respiratory rate greater than 25 breaths/min, and  $\text{SpO}_2$  level less than 95%.

The main exclusion criteria were (1) an urgent need for endotracheal intubation (prolonged respiratory pauses, cardiorespiratory arrest, heart rate  $<50$  beats/min with decreased level of consciousness, arterial blood pH level  $<7.15$ , irrespective of the cause), (2) hemodynamic instability, and (3) contraindications to noninvasive ventilation (eg, uncontrolled vomiting, copious oral secretions, facial deformities, a Glasgow Coma Scale score  $\leq 12$  points, presence of pneumothorax, or a do-not-intubate order).

After March 2020, patients with COVID-19 inevitably entered the trial in 1 of the 4 patient groups with acute respiratory failure described above. All analyses were conducted without separating patients with COVID-19 into a distinct group until the fifth interim analysis was conducted in March 2023. After March 2023, patients with COVID-19 were separated out for the purposes of the statistical analysis (described in an amendment to the trial protocol in June 2023). This decision was made by the trial steering committee while still blinded to any outcome data and as a good faith effort to match the aims of the original design while addressing the emergence of the COVID-19 pandemic. This decision was based on several critical factors, including that COVID-19 represented a novel disease with an anticipated intubation rate that could differ significantly from other causes of acute respiratory failure.

In addition, there was a rapidly evolving learning curve regarding the optimal management of patients with COVID-19, particularly in relation to noninvasive respiratory support and endotracheal intubation. This learning curve led to substantial shifts in intubation rates and mortality over time as clinical practices adapted. Consequently, the original assump-

tions about endotracheal intubation and mortality rates in the noninvasive ventilation (control) group during the trial's design phase may have been overly optimistic to accurately represent patients with COVID-19 (eMethods in [Supplement 1](#)).

### Randomization

The randomization list was generated electronically with permuted block sizes that were unknown to the investigators. The allocation concealment was maintained via an online central automated system available 24 hours daily, stratified by center and acute respiratory failure patient group. Patients were randomly assigned in a 1:1 ratio to high-flow nasal oxygen or noninvasive ventilation ([Figure 1](#)). The blinding of patients and clinicians was not possible because of the nature of the intervention.

### Interventions


High-flow nasal oxygen was delivered continuously (Airvo-2, Fisher & Paykel Healthcare) with a flow starting at 30 L/min for patients with COPD exacerbation with respiratory acidosis and at 45 L/min for those in the 4 other patient groups with acute respiratory failure, titrated gradually toward 60 L/min or the highest flow tolerated. Fraction of inspired oxygen ( $\text{FiO}_2$ ) started at 50% and was titrated to maintain  $\text{SpO}_2$  within 88% to 92% in the COPD exacerbation with respiratory acidosis patient group and within 92% to 98% in the other patient groups. After 24 hours, weaning from high-flow nasal oxygen therapy could begin if clinical improvement had been achieved (eMethods in [Supplement 1](#)). Noninvasive ventilation rescue therapy was allowed for the COPD exacerbation with respiratory acidosis and the acute cardiogenic pulmonary edema patient groups at the discretion of the treating physician (eFigure 1 in [Supplement 1](#)).

In the noninvasive ventilation group, the therapy was delivered through a face mask using either a ventilator designed primarily for invasive or noninvasive ventilation. In the COPD exacerbation with respiratory acidosis patient group, the settings for inspiratory positive airway pressure were between 12 and 16 cm of  $\text{H}_2\text{O}$ ; for all the other patient groups, the settings were between 12 and 14 cm of  $\text{H}_2\text{O}$ . In the COPD exacerbation patient group, the expiratory positive airway pressure setting was 4 cm of  $\text{H}_2\text{O}$ ; for all the other patient groups, the setting was 8 cm of  $\text{H}_2\text{O}$ . In the COPD exacerbation patient group, the  $\text{FiO}_2$  level was titrated to maintain an  $\text{SpO}_2$  within 88% and 92%; for all the other patient groups,  $\text{FiO}_2$  level was titrated to maintain an  $\text{SpO}_2$  within 92% and 98%.

The inspiratory positive airway pressure or the expiratory positive airway pressure could be increased independently by 1 or 2 cm of  $\text{H}_2\text{O}$  until (1) a maximum of 20 cm of  $\text{H}_2\text{O}$  for inspiratory positive airway pressure and a maximum of 12 cm of  $\text{H}_2\text{O}$  for expiratory positive airway pressure or (2) patient tolerability and signs of clinical improvement (defined as a respiratory rate  $\leq 25$  breaths/min and no signs of accessory respiratory muscle activity). The tidal volume target was 6 to 9 mL/kg of ideal body weight.

On the first day, 24 hours of noninvasive ventilation use was encouraged. Thereafter, weaning from noninvasive ventilation could be started in patients without signs of respiratory

Figure 1. Enrollment, Randomization, and Follow-Up in the Study





distress and included decreasing the duration of the noninvasive ventilation sessions, decreasing  $\text{FIO}_2$  level, and decreasing pressure levels each day at the discretion of the clinical team. A patient was considered weaned when  $\text{FIO}_2$  level was less than or equal to 30%, and both the expiratory positive airway pressure and pressure support were less than or equal to 6 cm of  $\text{H}_2\text{O}$  (eMethods in Supplement 1). The criteria for endotracheal intubation appear in the eMethods in Supplement 1. The final decision about endotracheal intubation was made by the attending physician.

### Primary Outcome

The primary outcome was endotracheal intubation or death within 7 days of randomization. The primary outcome was initially defined as endotracheal intubation within 7 days. However, we expanded the definition in the fifth version of the protocol (April 2021) to include both endotracheal intubation and death within 7 days to also consider eventual patients with acute respiratory failure who died without being intubated.

To maintain consistency in the indication for endotracheal intubation across sites and to minimize the risk of delayed intubation, predefined criteria were applied: (1) respiratory or cardiac arrest; (2) hemodynamic instability with mean arterial pressure lower than 65 mm Hg, systolic arterial pressure lower than 90 mm Hg after proper fluid resuscitation, or need for increasing doses of vasopressors ( $>0.3$  g/kg/min of norepinephrine); (3) cognitive impairment and agitation that prevents medical or nursing care without full sedation; (4) a Glasgow Coma Scale score of less than 11; (5) failure to maintain an  $\text{SpO}_2$  level greater than 92% (or  $>88\%$  in patients with COPD) despite having an  $\text{FIO}_2$  level of 60%; (6) a progressive increase in  $\text{Paco}_2$  that is greater than 10 mm Hg and a concurrent drop in arterial blood pH level despite attempts of improving ventilation; (7) an intolerability to high-flow nasal oxygen or noninvasive ventilation; (8) development of hypersecretion and inability to eliminate such secretion; (9) require frequent discontinuation of noninvasive ventilation therapy; (10) severe arrhythmia with hemodynamic instability; (11) persistent respiratory acidosis with arterial blood pH level lower than 7.2 after 60 minutes of optimal treatment; or (12) physician discretion.

### Secondary and Tertiary Outcomes

The secondary outcomes were 28-day mortality, 90-day mortality, mechanical ventilation-free days at 28 days, and ICU-free days at 28 days. The tertiary outcomes were hospital and ICU length of stay within 90 days, vasopressor-free days within 28 days, the proportion of patients who received a do-not-intubate order within 7 days after randomization, and patient comfort score.

### Statistical Analysis

The trial used bayesian adaptive statistical methods to assess the primary outcome (statistical analysis plan appears in Supplement 3). We hypothesized that high-flow nasal oxygen would be noninferior or superior to noninvasive ventilation in all 5 patient groups with acute respiratory failure. The

noninferiority margin was established based on an estimated absolute effect of 36% for noninvasive ventilation on the rate of endotracheal intubation.<sup>24</sup> Considering that the noninferiority margin should not exceed half of this effect, we selected an absolute difference of 10%.<sup>25</sup> Next, we proportionately extrapolated the 10% absolute difference on a 30% event rate (derived from a population with hypoxemic acute respiratory failure) to the other patient groups through the conversion to an odds ratio (OR) of 1.55 (the noninferiority margin).

The noninferiority and superiority hypotheses were tested in order with gatekeeping, thus without the need for a multiplicity correction. Noninferiority was declared if the noninferiority posterior probability was higher than 0.992. If noninferiority was demonstrated, then superiority was declared if the superiority posterior probability was also higher than 0.992 for an OR less than 1. If noninferiority was not demonstrated, the final result was futility. Stopping for futility, superiority, and noninferiority was allowed and conditioned on thresholds defined for each interim analysis (eMethods in Supplement 1 and Supplements 2-3). The interim and final values for the posterior probability thresholds for noninferiority and superiority were set to control for type I error at less than 0.025 in all 5 patient groups with acute respiratory failure (eTable 1 in Supplement 1). Up to 6 interim analyses were planned, starting after the 500th patient was randomized, with a maximum total sample size of 2000 patients. Simulations showed that, in a scenario of similar treatment effects, the power of our trial to demonstrate noninferiority would be 84.9% for nonimmunocompromised patients with hypoxemia, 76.1% for immunocompromised patients with hypoxemia, 68.5% for patients with COPD exacerbation with respiratory acidosis, 5.5% for patients with acute cardiogenic pulmonary edema, and 92.5% for patients with hypoxemic COVID-19 (eTable 2 in Supplement 1).

The primary analysis considered an intention-to-treat population that consisted of all randomized patients with informed consent. The primary outcome was analyzed using bayesian hierarchical modeling with dynamic borrowing across the 5 patient groups with acute respiratory failure, which shrinks the posterior distributions of the effects in each group toward the overall estimate. We used modestly informative prior distribution for the primary outcome rate for the noninvasive ventilation (control) group for each of the 5 patient groups with acute respiratory failure (eMethods in Supplement 1 and Supplement 3). A time-varying effect was considered for patients with hypoxemic COVID-19 to account for potential changes in the risk of the primary outcome across time.<sup>26</sup> The results are presented as rates, model-fitted median ORs, 95% credible intervals (CrIs), and posterior probabilities of noninferiority and superiority.

Mortality within 28 days was analyzed using the same bayesian model structure as the primary outcome, with adjustments to the prior distributions to account for different rates of 28-day mortality. For the other secondary outcomes, we used bayesian linear and logistic regression models to calculate the posterior probability of the intervention's superiority, using uninformative prior distributions. Specifically, we

applied flat priors for the model coefficients, allowing the data to drive the estimates without influence from prior assumptions. Posterior probabilities of superiority and the 95% CrIs for the secondary outcomes did not account for multiplicity, and, therefore, should not be used to infer definitive differences in treatment effects. There were no missing data for the primary outcome. The secondary outcomes were analyzed using only patients with available data, without imputation of missing data.

The sensitivity analyses included (1) a per-protocol analysis; (2) an analysis only including the COPD exacerbation with respiratory acidosis and the acute cardiogenic pulmonary edema patient groups and using escalation of ventilatory support as the outcome (escalation from noninvasive ventilation to endotracheal intubation and escalation from high-flow nasal oxygen therapy to noninvasive ventilation or endotracheal intubation); (3) a post hoc analysis with weaker informative priors for all 5 patient groups with acute respiratory failure; and (4) a post hoc analysis of the primary outcome using the same bayesian model structure as the main analysis, but without borrowing.

The cmdstanr version 0.6.1 (Stan Development Team) was used in conjunction with R version 4.3.1 (R Foundation for Statistical Computing).<sup>27,28</sup>

## Results

### Patients

Enrollment began in November 2019. A total of 2731 patients with acute respiratory failure were screened, and 1800 were enrolled at 33 sites in Brazil and randomized (Figure 1). Thirty-four patients were excluded after randomization because informed consent was not obtained.

In April 2021 (first interim analysis), enrollment was stopped for futility (defined as a posterior probability of noninferiority <0.30) in the immunocompromised with hypoxemia patient group (eTable 3 in Supplement 1). Enrollment was stopped when the predetermined noninferiority threshold was met in March 2023 (fifth interim analysis) for the hypoxemic COVID-19 patient group and in October 2023 (sixth interim analysis) for the nonimmunocompromised with hypoxemia and the acute cardiogenic pulmonary edema patient groups. Enrollment continued as planned until the final analysis for the COPD exacerbation with respiratory acidosis patient group.

Informed consent was not obtained from 49 patients because they died before their next of kin could provide it. However, the ethics committees at the sites permitted the use of data from 15 of these patients, resulting in a total of 1766 patients (mean age, 63.7 years [SD, 16.5 years]; 40% were women) included in the final analysis (eFigures 2-5 in Supplement 1).

The characteristics at enrollment were comparable across all 5 patient groups with acute respiratory failure (Table 1 and eTables 4-5 in Supplement 1). There were 485 nonimmunocompromised patients (27.5%) with hypoxemia, 50 immunocompromised patients (2.8%) with hypoxemia, 77 patients

(4.4%) with COPD exacerbation with respiratory acidosis, 272 patients (15.4%) with acute cardiogenic pulmonary edema, and 882 patients (49.9%) with hypoxemic COVID-19. Forty percent of the patients were randomized and treatment was initiated while they were being treated in the emergency department or hospital ward.

### Interventions

The allocated intervention was received by 93.3% of patients (824/883) in the high-flow nasal oxygen treatment group and by 91.5% of patients (808/883) in the noninvasive ventilation treatment group. There was a lower duration of therapy use within the first 48 hours in the noninvasive ventilation group (Table 2). Among patients assigned to high-flow nasal oxygen, the use of noninvasive ventilation was less than 10% in 4 of the 5 patient groups with acute respiratory failure; however, the use of noninvasive ventilation was 22.9% in the COPD exacerbation with respiratory acidosis patient group. The ventilatory settings for both high-flow nasal oxygen and noninvasive ventilation are described in eTable 6 in Supplement 1.

### Primary Outcome

The primary outcome of endotracheal intubation or death within 7 days occurred in 39.0% of patients (344 of 883) in the high-flow nasal oxygen group vs 38.1% of patients (336 of 883) in the noninvasive ventilation group. High-flow nasal oxygen was noninferior to noninvasive ventilation in the patient groups of nonimmunocompromised with hypoxemia, COPD exacerbation with respiratory acidosis, acute cardiogenic pulmonary edema, and hypoxemic COVID-19 (Figure 2 and eFigure 6 in Supplement 1). Among nonimmunocompromised patients with hypoxemia, the primary outcome occurred in 32.5% of patients in the high-flow nasal oxygen group vs 33.1% of patients in the noninvasive ventilation group (OR, 1.02 [95% CrI, 0.81-1.26]; noninferiority posterior probability, 0.999).

Among patients with COPD exacerbation with respiratory acidosis, the primary outcome occurred in 28.6% of patients in the high-flow nasal oxygen group vs 26.2% of patients in the noninvasive ventilation group (OR, 1.05 [95% CrI, 0.79-1.36]; noninferiority posterior probability, 0.992). Among patients with acute cardiogenic pulmonary edema, the primary outcome occurred in 10.3% of patients in the high-flow nasal oxygen group vs 21.3% of patients in the noninvasive ventilation group (OR, 0.97 [95% CrI, 0.73-1.23]; noninferiority posterior probability, 0.997).

Among patients with hypoxemic COVID-19, the primary outcome occurred in 51.3% of patients in the high-flow nasal oxygen group vs 47.0% of patients in the noninvasive ventilation group (OR, 1.13 [95% CrI, 0.94-1.38]; noninferiority posterior probability, 0.997). Among immunocompromised patients with hypoxemia, the primary outcome occurred in 57.1% of patients in high-flow nasal oxygen group vs 36.4% of patients in noninvasive ventilation group (OR, 1.07 [95% CrI, 0.81-1.39]; noninferiority posterior probability, 0.989), which met the criterion for futility. Summary of the individualized components of primary outcome and

Table 1. Characteristics of the Patients at Baseline

|  | Patient groups with acute respiratory failure (ARF) |  |  |   |  |   |
|--|---|--|--|---|--|---|
|  | Nonimmunocompromised with hypoxemia                 |  | Immunocompromised with hypoxemia             |   | COPD exacerbation with respiratory acidosis  |   |
|  | High-flow nasal oxygen (n = 249) <sup>a</sup>       | Noninvasive ventilation (n = 236) <sup>b</sup> | High-flow nasal oxygen (n = 28) <sup>a</sup> | Noninvasive ventilation (n = 22) <sup>b</sup> | High-flow nasal oxygen (n = 35) <sup>a</sup> | Noninvasive ventilation (n = 42) <sup>b</sup> |
| Age, mean (SD), y  | 66 (17)   | 70 (63)  | 54 (16)                                      | 53 (13)                                       | 74 (10)                                      | 71 (11)                                       |
| Sex, No. (%)   |   |  |  |   |  |   |
| Female   | 115 (46.2)  | 105 (44.7)                                     | 12 (42.9)                                    | 7 (31.9)                                      | 15 (42.9)                                    | 27 (64.3)                                     |
| Male   | 134 (53.8)  | 131 (55.3)                                     | 16 (57.1)                                    | 15 (68.1)                                     | 20 (57.1)                                    | 15 (37.7)                                     |
| Site of inclusion, No. (%) <sup>c</sup>                        |   |  |  |   |  |   |
| Intensive care unit  | 135 (54.2)  | 125 (53.0)                                     | 26 (92.9)                                    | 22 (100.0)                                    | 12 (34.3)                                    | 22 (52.4)                                     |
| Emergency department   | 100 (40.2)  | 94 (39.8)                                      | 0  | 0   | 20 (57.1)                                    | 19 (45.2)                                     |
| Hospital ward  | 14 (5.6)  | 17 (7.2)                                       | 2 (7.1)                                      | 0   | 3 (8.6)                                      | 1 (2.4)                                       |
| SAPS 3, median (IQR) <sup>d</sup>                              | 60 (54-69)  | 60 (53-66)                                     | 65 (60-71)                                   | 63 (57-68)                                    | 63 (57-67)                                   | 58 (51-64)                                    |
| ARF diagnosis to randomization, median (IQR), h                | 1 (0-4)   | 2 (0-4)  | 7 (4-14)                                     | 2 (1-11)                                      | 2 (0-4)                                      | 1 (0-3)                                       |
| Respiratory support before randomization, No. (%) <sup>e</sup> |   |  |  |   |  |   |
| Low-flow oxygen  | 230 (92.4)  | 218 (92.4)                                     | 23 (82.1)                                    | 18 (85.7)                                     | 31 (88.6)                                    | 38 (92.7)                                     |
| Noninvasive ventilation  | 96 (39.0)   | 81 (34.3)                                      | 5 (17.9)                                     | 4 (18.2)                                      | 13 (37.1)                                    | 16 (38.1)                                     |
| Heart rate, median (IQR), beats/min                            | 90 (75-107)   | 90 (79-108)                                    | 101 (84-115)                                 | 97 (87-110)                                   | 93 (84-104)                                  | 97 (79-109)                                   |
| Mean arterial pressure, median (IQR), mm Hg                    | 93 (83-103)   | 91 (81-102)                                    | 93 (85-101)                                  | 85 (80-94)                                    | 97 (84-104)                                  | 95 (88-107)                                   |
| Respiratory rate, median (IQR), breaths/min                    | 28 (26-31)  | 28 (26-30)                                     | 29 (28-33)                                   | 29 (26-31)                                    | 27 (26-29)                                   | 27 (25-30)                                    |
| SpO <sub>2</sub> , median (IQR), % <sup>f</sup>                | 86 (82-88)  | 86 (82-88)                                     | 86 (83-91)                                   | 88 (84-93)                                    | 83 (80-88)                                   | 85 (81-88)                                    |
| Arterial blood gas levels                                      |   |  |  |   |  |   |
| pH, median (IQR)   | 7.41 (7.35-7.45)                                    | 7.41 (7.37-7.46)                               | 7.42 (7.40-7.47)                             | 7.42 (7.40-7.44)                              | 7.32 (7.28-7.34)                             | 7.30 (7.24-7.36)                              |
| Pao <sub>2</sub> , median (IQR), mm Hg <sup>g</sup>            | 72 (58-90)  | 71 (60-86)                                     | 82 (55-106)                                  | 68 (62-89)                                    | 61 (52-93)                                   | 80 (60-114)                                   |
| Paco <sub>2</sub> , median (IQR), mm Hg                        | 37 (31-45)  | 38 (31-44)                                     | 31 (28-38)                                   | 32 (28-38)                                    | 55 (46-73)                                   | 64 (51-73)                                    |
| Pao <sub>2</sub> :Fio <sub>2</sub> , median (IQR) <sup>g</sup> | 191 (127-262)                                       | 194 (136-261)                                  | 181 (115-220)                                | 212 (150-281)                                 | 189 (143-290)                                | 209 (163-308)                                 |

(continued)

Table 1. Characteristics of the Patients at Baseline (continued)

|                                     | Patient groups with acute respiratory failure (ARF) |  |  |   |  |   | COPD exacerbation with respiratory acidosis   |  |   |  | Acute cardiogenic pulmonary edema             |  | Hypoxemic COVID-19                             |  |
|-------------------------------------|---|--|--|---|--|---|---|--|---|--|---|--|--|--|
|                                     | Nonimmunocompromised with hypoxemia                 |  | Immunocompromised with hypoxemia             |   | COPD exacerbation with respiratory acidosis  |   | High-flow nasal oxygen (n = 35) <sup>a</sup>  |  | Noninvasive ventilation (n = 42) <sup>b</sup> |  | High-flow nasal oxygen (n = 136) <sup>a</sup> |  | Noninvasive ventilation (n = 447) <sup>b</sup> |  |
| Comorbidities, No. (%) <sup>c</sup> | High-flow nasal oxygen (n = 249) <sup>a</sup>       | Noninvasive ventilation (n = 236) <sup>b</sup> | High-flow nasal oxygen (n = 28) <sup>a</sup> | Noninvasive ventilation (n = 22) <sup>b</sup> | High-flow nasal oxygen (n = 35) <sup>a</sup> | Noninvasive ventilation (n = 42) <sup>b</sup> | High-flow nasal oxygen (n = 136) <sup>a</sup> | Noninvasive ventilation (n = 447) <sup>b</sup> | High-flow nasal oxygen (n = 136) <sup>a</sup> | Noninvasive ventilation (n = 447) <sup>b</sup> | High-flow nasal oxygen (n = 136) <sup>a</sup> | Noninvasive ventilation (n = 447) <sup>b</sup> | High-flow nasal oxygen (n = 136) <sup>a</sup>  | Noninvasive ventilation (n = 447) <sup>b</sup> |
| Arterial hypertension               | 127 (51.0)  | 122 (51.7)                                     | 11 (39.3)                                    | 5 (22.7)                                      | 21 (60.0)                                    | 22 (52.4)                                     | 104 (76.5)                                    | 117 (86.0)                                     | 190 (43.7)                                    | 195 (43.6)                                     | 190 (43.7)                                    | 195 (43.6)                                     | 190 (43.7)                                     | 195 (43.6)                                     |
| Diabetes                            | 81 (32.5)   | 68 (28.8)                                      | 3 (10.7)                                     | 2 (9.1)                                       | 12 (34.3)                                    | 13 (31.0)                                     | 64 (47.1)                                     | 69 (50.7)                                      | 111 (25.5)                                    | 108 (24.2)                                     | 111 (25.5)                                    | 108 (24.2)                                     | 111 (25.5)                                     | 108 (24.2)                                     |
| COPD                                | 70 (28.1)   | 65 (27.5)                                      | 2 (7.1)                                      | 3 (13.6)                                      | 35 (100.0)                                   | 42 (100.0)                                    | 26 (19.1)                                     | 33 (24.3)                                      | 14 (3.2)                                      | 22 (4.9)                                       | 14 (3.2)                                      | 22 (4.9)                                       | 14 (3.2)                                       | 22 (4.9)                                       |
| Heart failure                       | 44 (17.7)   | 53 (22.5)                                      | 0  | 2 (9.1)                                       | 9 (25.7)                                     | 8 (19.0)                                      | 72 (52.9)                                     | 80 (58.8)                                      | 26 (6.0)                                      | 22 (4.9)                                       | 26 (6.0)                                      | 22 (4.9)                                       | 26 (6.0)                                       | 22 (4.9)                                       |
| Chronic kidney failure              | 29 (11.6)   | 19 (8.1)                                       | 1 (3.6)                                      | 3 (13.6)                                      | 2 (5.7)                                      | 2 (4.8)                                       | 23 (16.9)                                     | 37 (27.2)                                      | 29 (6.7)                                      | 36 (8.1)                                       | 29 (6.7)                                      | 36 (8.1)                                       | 29 (6.7)                                       | 36 (8.1)                                       |
| Solid tumors                        | 21 (8.4)  | 15 (6.4)                                       | 9 (32.1)                                     | 11 (50.0)                                     | 0  | 2 (4.8)                                       | 3 (2.2)                                       | 3 (2.2)  | 15 (3.4)                                      | 12 (2.7)                                       | 15 (3.4)                                      | 12 (2.7)                                       | 15 (3.4)                                       | 12 (2.7)                                       |
| Hematologic malignancies            | 2 (0.8)   | 1 (0.4)  | 5 (17.9)                                     | 3 (13.6)                                      | 1 (2.9)                                      | 0   | 0   | 0  | 6 (1.4)                                       | 4 (0.9)  | 6 (1.4)                                       | 4 (0.9)  | 6 (1.4)  | 4 (0.9)  |

Abbreviations: COPD, chronic obstructive pulmonary disease; FiO<sub>2</sub>, fraction of inspired oxygen; SAPS 3, Simplified Acute Physiology Score 3; SpO<sub>2</sub>, oxygen saturation as measured by pulse oximetry.

<sup>a</sup> Delivered through a nasal cannula with larger-bore tubing.

<sup>b</sup> Noninvasive positive-pressure ventilation delivered through a face mask using either a ventilator in the intensive care unit or a bilevel device.

<sup>c</sup> Percentages may not total 100 because of rounding.

<sup>d</sup> Calculated from 20 variables at enrollment, information about previous health status, and information obtained at admission. Scores range from 0 to 217 points; higher scores indicate a greater severity of illness. A score

<sup>e</sup> A patient may have received both types.

<sup>f</sup> The timing of these measurements was different. The level of SpO<sub>2</sub> was measured at room air and arterial blood gas was collected under oxygen delivered via low-flow oxygen cannula, nonbreathing face mask, or noninvasive ventilation.

<sup>g</sup> The level of FiO<sub>2</sub> was determined using the following calculation: (oxygen flow in L/min) × 0.03 + 0.21.

between 55 and 65 indicates a moderate to high severity of illness, suggesting a mortality risk between 30% and 40%.




Table 2. Adherence to Trial Therapy and Co-Interventions

| Patient groups with acute respiratory failure  |  |  |   |  |   |   |  |   |  |   |  |   |  |  |
|--|--|--|---|--|---|---|--|---|--|---|--|---|--|--|
| Nonimmunocompromised with hypoxemia  |  |  | Immunocompromised with hypoxemia              |  |   | COPD exacerbation with respiratory acidosis   |  |   | Acute cardiogenic pulmonary edema              |   |  | Hypoxemic COVID-19                            |  |  |
| High-flow nasal oxygen (n = 249) <sup>a</sup>  | Noninvasive ventilation (n = 236) <sup>b</sup> | High-flow nasal oxygen (n = 28) <sup>a</sup> | Noninvasive ventilation (n = 22) <sup>b</sup> | High-flow nasal oxygen (n = 35) <sup>a</sup> | Noninvasive ventilation (n = 42) <sup>b</sup> | High-flow nasal oxygen (n = 136) <sup>a</sup> | Noninvasive ventilation (n = 136) <sup>b</sup> | High-flow nasal oxygen (n = 435) <sup>a</sup> | Noninvasive ventilation (n = 447) <sup>b</sup> | High-flow nasal oxygen (n = 136) <sup>a</sup> | Noninvasive ventilation (n = 136) <sup>b</sup> | High-flow nasal oxygen (n = 435) <sup>a</sup> | Noninvasive ventilation (n = 447) <sup>b</sup> |  |
| Trial respiratory support during first 3 d   |  |  |   |  |   |   |  |   |  |   |  |   |  |  |
| Received assigned therapy, No. (%)   | 223 (89.6)                                     | 205 (86.9)                                   | 26 (92.9)                                     | 20 (90.9)                                    | 30 (85.7)                                     | 37 (88.1)                                     | 130 (95.6)                                     | 415 (95.4)                                    | 415 (92.8)                                     |   |  |   |  |  |
| Time receiving therapy, median (IQR), d  | 2 (1-3)  | 1 (1-3)                                      | 3 (1-3)                                       | 2 (1-3)                                      | 1 (1-3)                                       | 1.5 (1-3)                                     | 1 (1-2)  | 3 (1-3)                                       | 2 (1-3)  |   |  |   |  |  |
| Crossed over to another therapy, No. (%) <sup>c</sup>  | 23 (9.2)                                       | 5 (2.1)                                      | 1 (3.6)                                       | 0  | 8 (22.9)                                      | 0   | 7 (5.1)  | 0   | 34 (7.6)                                       |   |  |   |  |  |
| Trial respiratory support on day 1   |  |  |   |  |   |   |  |   |  |   |  |   |  |  |
| Received assigned therapy, No. (%)   | 221 (88.9)                                     | 203 (86.0)                                   | 26 (92.9)                                     | 20 (90.9)                                    | 30 (85.7)                                     | 37 (88.1)                                     | 129 (94.9)                                     | 415 (95.4)                                    | 409 (91.5)                                     |   |  |   |  |  |
| Duration of use, median (IQR), h   | 24 (9-24)                                      | 10 (4-17)                                    | 19 (12-24)                                    | 10 (3-21)                                    | 21 (7-24)                                     | 12 (6-17)                                     | 13 (7-24)                                      | 24 (12-24)                                    | 13 (6-22)                                      |   |  |   |  |  |
| Trial respiratory support on day 2   |  |  |   |  |   |   |  |   |  |   |  |   |  |  |
| Received assigned therapy, No./total (%)   | 138/191 (72.3)                                 | 114/184 (62.0)                               | 20/21 (95.2)                                  | 14/18 (77.8)                                 | 17/25 (68.0)                                  | 21/35 (60)                                    | 46/121 (38)                                    | 309/331 (93.4)                                | 289/354 (81.6)                                 |   |  |   |  |  |
| Duration of use, median (IQR), h   | 24 (22-24)                                     | 6 (2-13)                                     | 24 (23-24)                                    | 10 (4-12)                                    | 24 (18-24)                                    | 4 (2-14)                                      | 24 (10-24)                                     | 24 (22-24)                                    | 11 (5-16)                                      |   |  |   |  |  |
| Trial respiratory support on day 3   |  |  |   |  |   |   |  |   |  |   |  |   |  |  |
| Received assigned therapy, No./total (%)   | 108/178 (60.7)                                 | 82/161 (50.9)                                | 16/17 (94.1)                                  | 8/14 (57.1)                                  | 13/24 (54.2)                                  | 13/30 (43.0)                                  | 28/107 (26.2)                                  | 246/288 (85.0)                                | 212/291 (72.9)                                 |   |  |   |  |  |
| Duration of use, median (IQR), h   | 24 (20-24)                                     | 3.2 (2.0-8.1)                                | 24.0 (22.5-24.0)                              | 5.5 (2.0-7.2)                                | 21.9 (4.2-24.0)                               | 3 (1.7-6.0)                                   | 24.0 (19.2-24.0)                               | 24.0 (24.0-24.0)                              | 8.4 (3.3-13.3)                                 |   |  |   |  |  |
| Type of co-intervention within 7 d, No. (%)  |  |  |   |  |   |   |  |   |  |   |  |   |  |  |
| Vasopressors or inotropes  | 79 (31.7)                                      | 63 (26.7)                                    | 12 (42.9)                                     | 6 (27.3)                                     | 7 (20.0)                                      | 10 (23.8)                                     | 65 (47.8)                                      | 144 (33.1)                                    | 155 (34.7)                                     |   |  |   |  |  |
| Dexametomidine   | 32 (12.9)                                      | 44 (18.6)                                    | 3 (10.7)                                      | 4 (18.2)                                     | 4 (11.4)                                      | 9 (21.4)                                      | 13 (9.6)                                       | 57 (13.1)                                     | 117 (26.2)                                     |   |  |   |  |  |
| Abbreviation: COPD, chronic obstructive pulmonary disease.   |  |  |   |  |   |   |  |   |  |   |  |   |  |  |
| Delivered through a nasal cannula with larger-bore tubing.   |  |  |   |  |   |   |  |   |  |   |  |   |  |  |
| Noninvasive positive-pressure ventilation delivered through a face mask using either a ventilator in the intensive care unit or a bilevel device. If needed, standard oxygen therapy could be given through a nonrebreather face mask or low-flow nasal cannula between sessions in patients with device intolerance or agitation at a flow rate varying from 2 to 10 L/min. |  |  |   |  |   |   |  |   |  |   |  |   |  |  |
| If there was a worsening of respiratory failure, the trial protocol recommended noninvasive ventilation for patients with COPD exacerbation and for those with acute cardiogenic pulmonary edema. The use of noninvasive ventilation was not a protocol violation in these patients.   |  |  |   |  |   |   |  |   |  |   |  |   |  |  |

Abbreviation: COPD, chronic obstructive pulmonary disease.

<sup>a</sup> Delivered through a nasal cannula with larger-bore tubing.<sup>b</sup> Noninvasive positive-pressure ventilation delivered through a face mask using either a ventilator in the intensive care unit or a bilevel device. If needed, standard oxygen therapy could be given through a nonrebreather face mask or low-flow nasal cannula between sessions in patients with device intolerance or agitation at a flow rate varying from 2 to 10 L/min.<sup>c</sup> If there was a worsening of respiratory failure, the trial protocol recommended noninvasive ventilation for patients with COPD exacerbation and for those with acute cardiogenic pulmonary edema. The use of noninvasive ventilation was not a protocol violation in these patients.

Figure 2. Primary Outcome of Endotracheal Intubation or Death Within 7 Days



reasons for endotracheal intubation appear in **Table 3** and in **eTable 7** in **Supplement 1**.

### Other Outcomes

The secondary outcomes, including median 28-day mortality rates, were not different between the 2 treatment groups for any of the 5 patient groups with acute respiratory failure (**Table 3**). The tertiary outcome of comfort was superior with high-flow nasal oxygen vs noninvasive ventilation (**eTable 8** in **Supplement 1**). Other outcomes appear in **eTable 9** in **Supplement 1**.

### Sensitivity Analysis

The results for the posterior probabilities of noninferiority were similar to those of the main analysis for the per-protocol popu-

lation and the post hoc sensitivity analysis with weakly informative priors (**eTables 10-11** in **Supplement 1**).

The post hoc sensitivity analysis of the primary outcome without borrowing appears in **Figure 2**. Among nonimmunocompromised patients with hypoxemia, the OR was 0.98 (95% CrI, 0.73-1.33; noninferiority posterior probability, 0.996). Among immunocompromised patients with hypoxemia, the OR was 2.56 (95% CrI, 1.14-5.68; noninferiority posterior probability, 0.144).

Among patients with COPD exacerbation with respiratory acidosis, the OR was 1.48 (95% CrI, 0.67-3.09; noninferiority posterior probability, 0.549). Among patients with acute cardiogenic pulmonary edema, the OR was 0.52 (95% CrI, 0.29-0.91; noninferiority posterior probability, 0.999; superiority

Table 3. Primary and Secondary Outcomes

| Patient groups with acute respiratory failure                                | No. of events/total No. of patients (%) | Effect estimate (95% credible interval) |                           | Posterior probability of superiority <sup>a</sup> |
|--|---|---|---------------------------|---|
|  |   | Absolute                                | Relative                  |   |
| Nonimmunocompromised with hypoxemia  |   |   |                           |   |
| Primary outcome of endotracheal intubation or death within 7 d               | 81/250 (32.4)                           | MD, 0.5 (−4.5 to 4.6)                   | MOR, 1.02 (0.81 to 1.26)  | 0.687   |
| Individual components of primary outcome                                     |   |   |                           |   |
| Endotracheal intubation within 7 d   | 78/250 (31.2)                           |   |                           |   |
| Death within 7 d   | 30/250 (12.0)                           |   |                           |   |
| Secondary outcomes <sup>b</sup>  |   |   |                           |   |
| 28-d mortality   | 61/249 (24.5)                           | MD, −1.1 (−5.5 to 2.9) <sup>c</sup>     | MOR, 0.94 (0.74 to 1.17)  | 0.687   |
| 90-d mortality   | 74/249 (29.7)                           | MD, −4.1 (−12.3 to 3.9)                 | MHR, 0.86 (0.61 to 1.15)  | 0.826   |
| No. of ICU-free days at 28 d, median (IQR), d [No.] <sup>d</sup>             | 19 (−1 to 24) [240]                     | MD, 0 (−2 to 2.8)                       | MPOR, 1.00 (0.69 to 1.34) | 0.496   |
| No. of ventilator-free days at 28 d, median (IQR), d [No.] <sup>d</sup>      | 28 (−1 to 28) [242]                     | MD, 0.9 (−2.31 to 4.1)                  | MPOR, 1.16 (0.8 to 1.63)  | 0.795   |
| Immunocompromised with hypoxemia   |   |   |                           |   |
| Primary outcome of endotracheal intubation or death within 7 d               | 16/28 (57.1)                            | MD, 1.4 (−4.9 to 7.5)                   | MOR, 1.07 (0.81 to 1.39)  | 0.526   |
| Individual components of primary outcome                                     |   |   |                           |   |
| Endotracheal intubation within 7 d   | 14/28 (50.0)                            |   |                           |   |
| Death within 7 d   | 5/28 (17.9)                             |   |                           |   |
| Secondary outcomes <sup>b</sup>  |   |   |                           |   |
| 28-d mortality   | 15/28 (53.6)                            | MD, −0.2 (−6.3 to 6.0) <sup>c</sup>     | MOR, 0.99 (0.76 to 1.30)  | 0.523   |
| 90-d mortality   | 17/28 (60.7)                            | MD, 1.6 (24.9 to 27.1)                  | MHR, 1.29 (0.47 to 2.41)  | 0.242   |
| No. of ICU-free days at 28 d, median (IQR), d [No.] <sup>d</sup>             | −1 (−1 to 19) [27]                      | MD, −4.2 (−12.4 to 4.6)                 | MPOR, 0.48 (0.11 to 1.25) | 0.102   |
| No. of ventilator-free days at 28 d, median (IQR), d [No.] <sup>d</sup>      | −1 (−1 to 28) [28]                      | MD, −4.0 (−13.9 to 6.2)                 | MPOR, 0.56 (0.11 to 1.38) | 0.136   |
| Chronic obstructive pulmonary disease exacerbation with respiratory acidosis |   |   |                           |   |
| Primary outcome of endotracheal intubation or death within 7 d               | 10/35 (28.6)                            | MD, 0.8 (−4.0 to 5.3)                   | MOR, 1.05 (0.79 to 1.36)  | 0.546   |
| Individual components of primary outcome                                     |   |   |                           |   |
| Endotracheal intubation within 7 d   | 10/35 (28.6)                            |   |                           |   |
| Death within 7 d   | 3/35 (8.6)                              |   |                           |   |
| Secondary outcomes <sup>b</sup>  |   |   |                           |   |
| 28-d mortality   | 8/35 (22.9)                             | MD, −0.2 (−4.0 to 3.7) <sup>c</sup>     | MOR, 0.98 (0.75 to 1.29)  | 0.546   |
| 90-d mortality   | 11/35 (31.4)                            | MD, 0.5 (−20.1 to 20.8)                 | MHR, 1.09 (0.39 to 2.24)  | 0.416   |
| No. of ICU-free days at 28 d, median (IQR), d [No.] <sup>d</sup>             | 22 (0 to 28) [33]                       | MD, 0.2 (−6.6 to 7.1)                   | MPOR, 1.05 (0.33 to 2.23) | 0.541   |
| No. of ventilator-free days at 28 d, median (IQR), d [No.] <sup>d</sup>      | 28 (0 to 28) [33]                       | MD, −1.0 (−8.4 to 6.2)                  | MPOR, 0.82 (0.22 to 1.93) | 0.360   |

(continued)

Table 3. Primary and Secondary Outcomes (continued)

| Patient groups with acute respiratory failure                           | No. of events/total No. of patients (%) |                         | Effect estimate (95% credible interval) |                           | Posterior probability of superiority <sup>a</sup> |
|---|---|-------------------------|---|---------------------------|---|
|   | High-flow nasal oxygen                  | Noninvasive ventilation | Absolute                                | Relative                  |   |
| <b>Acute cardiogenic pulmonary edema</b>                                |   |                         |   |                           |   |
| Primary outcome of endotracheal intubation or death within 7 d          | 14/136 (10.3)                           | 29/136 (21.3)           | MD, -0.3 (-3.7 to 2.4)                  | MOR, 0.97 (0.73 to 1.23)  | 0.608   |
| Individual components of primary outcome                                |   |                         |   |                           |   |
| Endotracheal intubation within 7 d                                      | 12/136 (8.8)                            | 27/136 (19.9)           |   |                           |   |
| Death within 7 d  | 7/136 (5.1)                             | 16/136 (11.8)           |   |                           |   |
| <b>Secondary outcomes<sup>b</sup></b>                                   |   |                         |   |                           |   |
| 28-d mortality  | 19/136 (14.0)                           | 24/136 (17.6)           | MD, -0.5 (-4.0 to 3.2) <sup>c</sup>     | MOR, 0.97 (0.74 to 1.40)  | 0.608   |
| 90-d mortality  | 23/136 (16.9)                           | 30/136 (22.1)           | MD, -5.1 (-14.7 to 4.3)                 | MHR, 0.73 (0.38 to 1.18)  | 0.872   |
| No. of ICU-free days at 28 d, median (IQR), d [No.] <sup>d</sup>        | 28 (20 to 28) [131]                     | 28 (17 to 28) [132]     | 0.7 (-2.0 to 3.3)                       | MPOR, 1.19 (0.68 to 1.80) | 0.749   |
| No. of ventilator-free days at 28 d, median (IQR), d [No.] <sup>d</sup> | 28 (28 to 28) [132]                     | 28 (26 to 28) [132]     | 2.2 (-0.8 to 5.5)                       | MPOR, 1.85 (0.95 to 3.17) | 0.983   |
| <b>Hypoxemic COVID-19</b>   |   |                         |   |                           |   |
| Primary outcome of endotracheal intubation or death within 7 d          | 223/435 (51.3)                          | 210/437 (48.1)          | MD, 3.1 (-1.5 to 7.9)                   | MOR, 1.13 (0.94 to 1.38)  | 0.568   |
| Individual components of primary outcome                                |   |                         |   |                           |   |
| Endotracheal intubation within 7 d                                      | 219/435 (50.3)                          | 209/437 (47.8)          |   |                           |   |
| Death within 7 d  | 34/435 (7.8)                            | 17/437 (3.9)            |   |                           |   |
| <b>Secondary outcomes<sup>b</sup></b>                                   |   |                         |   |                           |   |
| 28-d mortality  | 97/435 (22.3)                           | 102/447 (22.8)          | MD, -0.4 (-4.0 to 3.3) <sup>e</sup>     | MOR, 0.98 (0.79 to 1.21)  | 0.568   |
| 90-d mortality  | 126/435 (29.0)                          | 129/447 (28.9)          | MD, 0.2 (-5.7 to 6.1)                   | MHR, 0.98 (0.76 to 1.25)  | 0.560   |
| No. of ICU-free days at 28 d, median (IQR), d [No.] <sup>d</sup>        | 15 (0 to 21) [434]                      | 17 (0 to 22) [439]      | MD, -1.0 (-3.1 to 1.1)                  | MPOR, 0.85 (0.67 to 1.06) | 0.840   |
| No. of ventilator-free days at 28 d, median (IQR), d [No.] <sup>d</sup> | 24 (1 to 28) [434]                      | 28 (0 to 28) [441]      | MD, -0.4 (-2.8 to 1.9)                  | MPOR, 0.94 (0.75 to 1.19) | 0.310   |

Abbreviations: ICU, intensive care unit; MD, median difference; MHR, median hazard ratio; MPOR, median proportional odds ratio.

<sup>a</sup> Superiority is declared if the posterior probability is greater than 0.992.

<sup>b</sup> The secondary outcomes are inferences based on a Bayesian linear and logistic regression model; however, 28-day mortality is based on a Bayesian model with the same structure as the primary outcome with some modification of the prior distributions to reflect the different rates of 28-day mortality within the groups without consideration of the noninferiority margin. Borrowing of information between groups was present.

<sup>c</sup> The model-fitted rate differences were calculated by subtracting the rate for patients who received noninvasive ventilation from the rate for those who received high-flow nasal oxygen.

<sup>d</sup> Defined as the number of days without invasive mechanical ventilation or days outside the ICU. For patients who died, a value of -1 day was assigned.

<sup>e</sup> Time-averaged rate that was weighted by the number of COVID-19 patients enrolled over time.

posterior probability, 0.970). Among patients with hypoxemic COVID-19, the OR was 1.16 (95% CrI, 0.94-1.43; noninferiority posterior probability, 0.992). The need for respiratory support escalation among patients with COPD exacerbation or acute cardiogenic pulmonary edema appears in eTable 12 in [Supplement 1](#).

### Adverse Events

The incidence of serious adverse events was similar between treatment groups (9.4% of patients in the high-flow nasal oxygen group vs 9.9% of patients in the noninvasive ventilation group; eTable 13 in [Supplement 1](#)).

## Discussion

This multicenter, adaptive, randomized clinical trial used a bayesian hierarchical model with dynamic borrowing across groups and found that high-flow nasal oxygen was noninferior to noninvasive ventilation with respect to endotracheal intubation or death within 7 days for the following patient populations with acute respiratory failure: nonimmunocompromised with hypoxemia, COPD exacerbation with respiratory acidosis, acute cardiogenic pulmonary edema, and hypoxemic COVID-19. Noninferiority of high-flow nasal oxygen compared with noninvasive ventilation was not shown among immunocompromised patients with hypoxemic acute respiratory failure.

In the FLORALI trial,<sup>16</sup> there was no significant difference in intubation rates among patients with non-hypercapnic acute hypoxemic respiratory failure who were treated with high-flow nasal oxygen (n = 106) vs noninvasive ventilation (n = 110), although 90-day mortality was lower in the high-flow nasal oxygen group. However, for patients with hypoxemic acute respiratory failure due to COVID-19, the effectiveness of high-flow nasal oxygen was uncertain compared with noninvasive ventilation.<sup>9,29</sup>

The current study included 485 nonimmunocompromised patients with acute hypoxemic respiratory failure and 881 patients with acute hypoxemic respiratory failure due to COVID-19 and showed noninferiority of high-flow nasal oxygen compared with noninvasive ventilation in both these patient groups. The results of a post hoc analysis without borrowing were also consistent with noninferiority of high-flow nasal oxygen for these patient groups (nonimmunocompromised with hypoxemia and hypoxemic COVID-19). In addition, there were no significant differences in 90-day mortality or in any secondary outcomes.

Among patients with COPD exacerbation with respiratory acidosis or acute cardiogenic pulmonary edema, previous evidence supported noninvasive ventilation as the recommended respiratory support modality.<sup>12,30,31</sup> However, these recommendations were based on studies comparing noninvasive ventilation with low-flow oxygen.<sup>12</sup> Although the results from the primary analysis in the current study demonstrated noninferiority of high-flow nasal oxygen compared with noninvasive ventilation for the COPD exacerbation with respiratory acidosis patient group, the post hoc analysis with-

out borrowing produced discrepant results; the 95% CrI included ORs as high as 3.09, suggesting potential harm with use of high-flow nasal oxygen.

Among the patients with COPD or acute cardiogenic pulmonary edema assigned to high-flow nasal oxygen who showed signs of worsening respiratory failure, the trial protocol recommended escalation to noninvasive ventilation. Notably, 23% of patients with COPD initially assigned to high-flow nasal oxygen required rescue noninvasive ventilation. Furthermore, the patients with acute cardiogenic pulmonary edema who needed noninvasive ventilation before randomization were excluded from the trial. Therefore, the results from the current trial may not be generalizable to patients with more severe forms of acute cardiogenic pulmonary edema.

There were no significant differences in 28-day or 90-day mortality between the 2 treatment groups within any of the 5 acute respiratory failure patient groups. However, overall mortality was higher compared with other studies.<sup>16,18,23,32</sup> Several factors may explain this elevated mortality. First, mortality among critically ill patients has generally been higher in Brazil than in high-income countries.<sup>33-35</sup> Second, during the COVID-19 pandemic, patients tended to avoid leaving home, so only more severe cases sought hospital care.<sup>36</sup> Third, the inclusion criteria for patients with COPD exacerbation required respiratory acidosis, thereby selecting for more severe cases.

The strengths of the current trial include its generalizability because it included a broad population of patients with acute respiratory failure. At the same time, the results are specific to each of the 5 patient groups with acute respiratory failure studied. The adaptive bayesian design allowed dynamic borrowing of information across patient groups, which leverages the information that arises from the evaluation of the treatments in multiple and related disease categories. This dynamic borrowing increases the precision of the estimates and includes information not available with simpler models without borrowing, and may be useful when the assumption of similarity of treatment effects across groups is appropriate.

### Limitations

The current trial has limitations. First, enrollment in the immunocompromised with hypoxemia patient group was stopped early for futility after the first interim analysis. The futility stopping criterion in this analysis may have been set too high (a posterior probability of noninferiority below 30%). If a lower threshold (eg, 10%) had been selected, enrollment in this group might have continued, potentially resulting in a different conclusion. As a result, the validity of effect estimates in the immunocompromised with hypoxemia patient group is limited by the small sample size. In addition, patients with COVID-19 were not classified into a separate group before March 2023, and some of the patients included in the interim analysis for the immunocompromised with hypoxemia patient group had COVID-19. Future trials comparing high-flow nasal oxygen vs noninvasive ventilation in immunocompromised patients with acute hypoxemic respiratory failure are warranted.



Second, although the COVID-19 pandemic led to a protocol amendment that categorized patients with COVID-19 as a separate group, a blinded steering committee made these adjustments. Third, the effect estimates relied on borrowing across patient groups, which assumes some degree of similarity in underlying treatment effects. A post hoc sensitivity analysis without borrowing showed substantial deviations from the main model, particularly in the COPD and the immunocompromised with hypoxemia patient groups. However, an analysis without borrowing loses information and results in a systematic overestimate of the variability of treatment effects across patient groups.

Fourth, the comparator intervention was noninvasive ventilation administered through a face mask; thus, our results do not address comparisons of high-flow nasal oxygen with con-

tinuous positive airway pressure or with helmet-delivered noninvasive ventilation.

## Conclusions

Compared with noninvasive ventilation, high-flow nasal oxygen met prespecified criteria for noninferiority for the primary outcome of endotracheal intubation or death within 7 days in 4 of the 5 patient groups with acute respiratory failure. However, the small sample sizes in some patient groups and the sensitivity of the findings to the choice of analysis model suggests the need for further study in patients with COPD, immunocompromised patients, and patients with acute cardiogenic pulmonary edema.

## ARTICLE INFORMATION

**Accepted for Publication:** November 21, 2024.

**Published Online:** December 10, 2024.

doi:10.1001/jama.2024.26244

### The RENOvATE Investigators and BRICNet

**Authors:** Israel S. Maia, MD, PhD; Letícia Kawano-Dourado, MD, PhD; Lucas Tramujas, MD; Neymar Elias de Oliveira, MD; Rafael Naoki Souza, RT; Dhaisi Faustino Signorini, RT; Mariangela Pimentel Pincelli, MD, PhD; Cássio Luis Zandonai, RT; Regiane Tamires Blasius, RT; Fabrício Freires, RT; Vanessa Marques Ferreira, RT; Marcelo Luz Pereira Romano, MD, PhD; Mieke Claudia Miura, MSc; Caroline Maschio de Censo, MSc; Eliana Bernadete Caser, MD, PhD; Betania Silva, RT; Daniela Correia Santos Bonomo, RT; Jussara Alencar Arraes, MSc; Meton Soares de Alencar Filho, MD; Jacques Gabriel Álvares Horta, MD; Débora Campos Oliveira, RT; Emerson Boschi, MD, MSc; Rafael Lessa Costa, MD; Glauco Adriano Westphal, MD, PhD; Juliano Ramos, MD; Fábio Holanda Lacerda, MD; Conrado Roberto Hoffmann Filho, MD; Bruno Valle Pinheiro, MD, PhD; Leonardo Bugarin de Andrade Neumamm, PhD; Mário Roberto Rezende Guimarães Júnior, MD; Davi Tamamaru de Souza, MD, MSc; Juliana Carvalho Ferreira, MD, PhD; Louis Nakayama Ohe, MD; Daniel Almeida Schettini, MD, MSc; Marlus Muri Thompson, MD; Maria Cristina França de Oliveira, MD; Viviane Cordeiro Veiga, MD, PhD; Karina L. Negrelli, DVM; Renato H. N. Santos, BSc(Stat); Lucas Damiani, PhD; Rodrigo M. Gurgel, Biomed Sci; Samara P. C. Gomes, Biomed Sci; Lucas M. Lima, BScIT; Tamiris A. Miranda, MSc; Lígia N. Laranjeira, PhD; Pedro Gabriel Melo de Barros e Silva, MD, PhD; Flávia R. Machado, MD, PhD; Mark Fitzgerald, PhD; Anna Bosse, MS; Joe Marion, PhD; Carlos Roberto Ribeiro Carvalho, MD, PhD; Laurent Brochard, MD, PhD; Roger J. Lewis, MD, PhD; Alexandre Biasi Cavalcanti, MD, PhD.

### Affiliations of The RENOvATE Investigators and BRICNet Authors:

Hcor Research Institute, São Paulo, Brazil (Maia, Kawano-Dourado, Tramujas, Negrelli, Santos, Damiani, Gurgel, Gomes, Lima, Miranda, Laranjeira, de Barros e Silva, Biasi Cavalcanti); Brazilian Research in Intensive Care Network (BRICNet), São Paulo, Brazil (Maia, J. C. Ferreira, Veiga, Machado, Biasi Cavalcanti); Hospital Nereu Ramos, Florianópolis, Brazil (Maia, Pincelli, Zandonai, Blasius); Universidade Federal de Santa Catarina, Florianópolis, Brazil (Maia, Pincelli); MAGIC Evidence Ecosystem Foundation, Oslo,

Norway (Kawano-Dourado); Divisão de Pneumologia, Instituto do Coracao, Hospital das Clínicas, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil (Kawano-Dourado, J. C. Ferreira, Carvalho); Fundação Faculdade Regional de Medicina de São José do Rio Preto, São José do Rio Preto, Brazil (N. E. de Oliveira, Souza, Signorini); Faculdade de Medicina de São José do Rio Preto, São José do Rio Preto, Brazil (N. E. de Oliveira); Hospital São Paulo, Escola Paulista de Medicina, Universidade Federal de São Paulo, São Paulo, Brazil (Freires, V. M. Ferreira, Machado); Intensive Care Unit, Hcor Research Institute, São Paulo, Brazil (Romano, Miura, de Censo); Hospital Unimed Vitória, Vitória, Brazil (Caser, Silva, Santos Bonomo); Hospital Maternidade São Vicente de Paulo, Barbalha, Brazil (Arraes, de Alencar Filho); Santa Casa da Misericórdia de Ouro Preto, Ouro Preto, Brazil (Álvares Horta, Oliveira); Hospital Geral de Caxias do Sul, Caxias do Sul, Brazil (Boschi, Costa); Centro Hospitalar Unimed de Joinville, Joinville, Brazil (Westphal, Ramos); Instituto Oto de Pesquisa, Ensino e Inovação-Hospital Oto Aldeota, Fortaleza, Brazil (Lacerda); Hospital Regional Hans Dieter Schmidt, Joinville, Brazil (Filho); Hospital Universitário da Universidade Federal de Juiz de Fora, Juiz de Fora, Brazil (Pinheiro); Hospital Naval Marcílio Dias, Rio de Janeiro, Brazil (de Andrade Neumamm); Santa Casa de Barretos, Barretos, Brazil (Guimarães Júnior); Hospital Universitário da Universidade Estadual de Maringá, Maringá, Brazil (de Souza); Instituto Dante Pazzanese de Cardiologia, São Paulo, Brazil (Ohe); Hospital da Luz, São Paulo, Brazil (Schettini); Hospital Evangélico de Cachoeiro de Itapemirim, Cachoeiro de Itapemirim, Brazil (Thompson); Hospital AC Camargo, São Paulo, Brazil (M. C. F. de Oliveira); Beneficência Portuguesa de São Paulo, São Paulo, Brazil (Veiga); Berry Consultants LLC, Austin, Texas (Fitzgerald, Bosse, Marion, Lewis); Keenan Research Centre, Li Ka Shing Knowledge Institute, St Michael's Hospital, Unity Health and Interdepartmental Division of Critical Care Medicine, University of Toronto, Toronto, Ontario, Canada (Brochard); Department of Emergency Medicine, Harbor-UCLA Medical Center, Los Angeles, California (Lewis); Postgraduate Program of Anesthesiology, Surgical Sciences, and Perioperative Medicine, Faculty of Medicine, Universidade de São Paulo, São Paulo, Brazil (Biasi Cavalcanti).

### Author Contributions:

Drs Maia and Biasi Cavalcanti had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Drs Maia and Kawano-Dourado contributed equally to this article.

**Concept and design:** Maia, Kawano-Dourado, Pincelli, Romano, Arraes, Alvares Horta, D. Souza, Negrelli, Damiani, Fitzgerald, Carvalho, Brochard, Lewis, Biasi Cavalcanti.

**Acquisition, analysis, or interpretation of data:** Maia, Kawano-Dourado, Tramujas, N. Oliveira, R. Souza, Signorini, Pincelli, Zandonai, Blasius, Freires, V. Ferreira, Romano, Miura, Censo, Caser, Silva, Bonomo, Alencar Filho, Arraes, D. Oliveira, Boschi, Costa, Westphal, Ramos, Lacerda, Hoffmann Filho, Pinheiro, Neumamm, Guimarães Junior, J. Ferreira, Ohe, Schettini, Thompson, M. Oliveira, Gabriel Melo de Barros e Silva, Veiga, Caser, Gomes, de Lima, Santos, Damiani, Miranda, Laranjeira, Machado, Marion, Bosse, Fitzgerald, Carvalho, Brochard, Biasi Cavalcanti.

**Drafting of the manuscript:** Maia, Kawano-Dourado, Tramujas, N. Oliveira, Pincelli, V. Ferreira, Romano, Censo, Alencar Filho, Arraes, Lacerda, Ohe, Thompson, de Lima, Santos, Bosse, Carvalho, Biasi Cavalcanti.

**Critical review of the manuscript for important intellectual content:** Maia, Kawano-Dourado, Tramujas, R. Souza, Signorini, Zandonai, Blasius, Freires, Romano, Miura, Caser, Silva, Bonomo, Arraes, Alvares Horta, D. Oliveira, Boschi, Costa, Westphal, Ramos, Hoffmann Filho, Pinheiro, Neumamm, Guimarães Junior, D. Souza, J. Ferreira, Schettini, M. Oliveira, Gabriel Melo de Barros e Silva, Veiga, Negrelli, Gurgel, Gomes, Damiani, Miranda, Laranjeira, Machado, Marion, Fitzgerald, Carvalho, Brochard, Lewis.

**Statistical analysis:** Maia, Alencar Filho, Arraes, Santos, Damiani, Marion, Bosse, Fitzgerald, Lewis.

**Obtained funding:** Maia, Arraes, Gurgel, Gomes, Biasi Cavalcanti.

**Administrative, technical, or material support:** Maia, Tramujas, N. Oliveira, Pincelli, Zandonai, Blasius, Romano, Miura, Caser, Bonomo, Alvares Horta, D. Oliveira, Costa, Neumamm, J. Ferreira, Ohe, Thompson, M. Oliveira, Veiga, Negrelli, Gurgel, Gomes, de Lima, Miranda, Laranjeira, Carvalho, Lewis.

**Supervision:** Maia, Kawano-Dourado, Romano, Caser, Bonomo, Alencar Filho, Arraes, Boschi, Costa, Lacerda, Neumamm, J. Ferreira, Thompson, Gabriel

Melo de Barros e Silva, Negrelli, Gurgel, Miranda, Machado, Carvalho, Biasi Cavalcanti.

**Conflict of Interest Disclosures:** Dr J. Ferreira reported receiving grants from the ATS Foundation. Dr Gabriel Melo de Barros e Silva reported receiving grants from Roche Diagnostics, Novartis, and Bayer. Dr Machado reported receiving personal fees from Baxter and Biomerieux. Dr Marion reported being an employee of Berry Consultants, LLC (a statistical consulting firm that specializes in the design, conduct, oversight, and analysis of adaptive and platform clinical trials). Dr Bosse reported receiving personal fees from Berry Consultants. Dr Fitzgerald reported being an employee of Berry Consultants. Dr Brochard reported receiving grants from Stimit and Vitalaire, receiving nonfinancial support from Cerebra Health, and receiving personal fees from Stimit and Fisher & Paykel Healthcare. Dr Lewis reported being an employee of Berry Consultants. No other disclosures were reported.

**Funding/Support:** Supported by grant 25000.028212/2018-10 from the Brazilian Ministry of Health (*Programa de Apoio ao Desenvolvimento Institucional do Sistema Único de Saúde*). Fisher & Paykel Healthcare provided the high-flow nasal oxygen equipment and associated disposables. The trial coordinating center and sponsor was the Hcor Research Institute.

**Role of the Funder/Sponsor:** The funders/sponsors had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Group Information:** The RENOVATE Investigators and BRICNet nonauthors appear in [Supplement 4](#).

**Disclaimer:** Dr Lewis is Statistical Editor of *JAMA* but was not involved in any of the decisions regarding review of the manuscript or its acceptance.

**Meeting Presentation:** Presented in part at the Critical Care Reviews meeting; December 10, 2024; Melbourne, Australia.

**Data Sharing Statement:** See [Supplement 5](#).

**Additional Contributions:** We thank the patients who participated in this trial and their relatives, and the medical and nursing teams at the participating sites for their contributions, whether through directly recruiting patients or providing essential care to those enrolled in the study. We also thank Mariela Lacerda, PhD (Department of Science and Technology, and Secretariat of Science, Technology, and Innovation, Health Industrial Complex, Ministry of Health), for her unwavering enthusiasm and support for the trial, technical scientific consulting, program management coordination, and general coordination of the health research funding. Dr Lacerda did not receive any compensation for her work during the trial.

## REFERENCES

1. Munshi L, Mancebo J, Brochard LJ. Noninvasive respiratory support for adults with acute respiratory failure. *N Engl J Med*. 2022;387(18):1688-1698. doi:10.1056/NEJMr2204556
2. Carron M, Freo U, BaHammam AS, et al. Complications of non-invasive ventilation techniques: a comprehensive qualitative review of randomized trials. *Br J Anaesth*. 2013;110(6):896-914. doi:10.1093/bja/aet070

3. Mauri T, Turrini C, Eronia N, et al. Physiologic effects of high-flow nasal cannula in acute hypoxemic respiratory failure. *Am J Respir Crit Care Med*. 2017;195(9):1207-1215. doi:10.1164/rccm.201605-0916OC
4. Cortegiani A, Crimi C, Noto A, et al. Effect of high-flow nasal therapy on dyspnea, comfort, and respiratory rate. *Crit Care*. 2019;23(1):201. doi:10.1186/s13054-019-2473-y
5. Cortegiani A, Longhini F, Madotto F, et al; H. F.-AECOPD study investigators. High flow nasal therapy versus noninvasive ventilation as initial ventilatory strategy in COPD exacerbation: a multicenter non-inferiority randomized trial. *Crit Care*. 2020;24(1):692. doi:10.1186/s13054-020-03409-0
6. Tan D, Wang B, Cao P, et al. High flow nasal cannula oxygen therapy versus non-invasive ventilation for acute exacerbations of chronic obstructive pulmonary disease with acute-moderate hypercapnic respiratory failure: a randomized controlled non-inferiority trial. *Crit Care*. 2024;28(1):250. doi:10.1186/s13054-024-05040-9
7. Roca O, Pérez-Terán P, Masclans JR, et al. Patients with New York Heart Association class III heart failure may benefit with high flow nasal cannula supportive therapy: high flow nasal cannula in heart failure. *J Crit Care*. 2013;28(5):741-746. doi:10.1016/j.jcrc.2013.02.007
8. Marjanovic N, Piton M, Lamarre J, et al. High-flow nasal cannula oxygen versus noninvasive ventilation for the management of acute cardiogenic pulmonary edema: a randomized controlled pilot study. *Eur J Emerg Med*. 2024;31(4):267-275. doi:10.1097/MEJ.0000000000001128
9. Oczkowski S, Ergon B, Bos L, et al. ERS clinical practice guidelines: high-flow nasal cannula in acute respiratory failure. *Eur Respir J*. 2022;59(4):2101574. doi:10.1183/13993003.01574-2021
10. Telias I, Brochard LJ, Gattarello S, et al. The physiological underpinnings of life-saving respiratory support. *Intensive Care Med*. 2022;48(10):1274-1286. doi:10.1007/s00134-022-06749-3
11. Berbenetz N, Wang Y, Brown J, et al. Non-invasive positive pressure ventilation (CPAP or bilevel NPPV) for cardiogenic pulmonary oedema. *Cochrane Database Syst Rev*. 2019;4(4):CD005351. doi:10.1002/14651858.CD005351.pub4
12. Rochwerg B, Brochard L, Elliott MW, et al. Official ERS/ATS clinical practice guidelines: noninvasive ventilation for acute respiratory failure. *Eur Respir J*. 2017;50(2):1602426. doi:10.1183/13993003.02426-2016
13. Grasselli G, Calfee CS, Camporota L, et al; European Society of Intensive Care Medicine Taskforce on ARDS. ESICM guidelines on acute respiratory distress syndrome: definition, phenotyping and respiratory support strategies. *Intensive Care Med*. 2023;49(7):727-759. doi:10.1007/s00134-023-07050-7
14. Ospina-Tascón GA, Calderón-Tapia LE, García AF, et al; HiFlo-Covid Investigators. Effect of high-flow oxygen therapy vs conventional oxygen therapy on invasive mechanical ventilation and clinical recovery in patients with severe COVID-19: a randomized clinical trial. *JAMA*. 2021;326(21):2161-2171. doi:10.1001/jama.2021.20714

15. Ferreyro BL, Angriman F, Munshi L, et al. Noninvasive oxygenation strategies in adult patients with acute respiratory failure: a protocol for a systematic review and network meta-analysis. *Syst Rev*. 2020;9(1):95. doi:10.1186/s13643-020-01363-0
16. Frat JP, Thille AW, Mercat A, et al; FLORALI Study Group; REVA Network. High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med*. 2015;372(23):2185-2196. doi:10.1056/NEJMoa1503326
17. Lewis SR, Baker PE, Parker R, Smith AF. High-flow nasal cannulae for respiratory support in adult intensive care patients. *Cochrane Database Syst Rev*. 2021;3(3):CD010172.
18. Perkins GD, Ji C, Connolly BA, et al; RECOVERY-RS Collaborators. Effect of noninvasive respiratory strategies on intubation or mortality among patients with acute hypoxemic respiratory failure and COVID-19: the RECOVERY-RS randomized clinical trial. *JAMA*. 2022;327(6):546-558. doi:10.1001/jama.2022.0028
19. Rochwerg B, Einav S, Chaudhuri D, et al. The role for high flow nasal cannula as a respiratory support strategy in adults: a clinical practice guideline. *Intensive Care Med*. 2020;46(12):2226-2237. doi:10.1007/s00134-020-06312-y
20. Maia IS, Kawano-Dourado L, Zampieri FG, et al; RENOVATE Investigators and the BRICNet. High flow nasal catheter therapy versus non-invasive positive pressure ventilation in acute respiratory failure (RENOVATE trial): protocol and statistical analysis plan. *Crit Care Resusc*. 2023;24(1):61-70.
21. Maia IS, Kawano-Dourado L, Damiani LP, Fitzgerald M, Lewis RJ, Cavalcanti AB. Update in statistical analysis plan of the RENOVATE trial. *Crit Care Resusc*. 2023;25(3):113-114. doi:10.1016/j.ccrj.2023.06.008
22. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human participants. *JAMA*. Published online October 19, 2024. doi:10.1001/jama.2024.21972
23. Azoulay E, Lemiale V, Mokart D, et al. Effect of high-flow nasal oxygen vs standard oxygen on 28-day mortality in immunocompromised patients with acute respiratory failure: the HIGH randomized clinical trial. *JAMA*. 2018;320(20):2099-2107. doi:10.1001/jama.2018.14282
24. Liu YJ, Zhao J, Tang H. Non-invasive ventilation in acute respiratory failure: a meta-analysis. *Clin Med (Lond)*. 2016;16(6):514-523. doi:10.7861/clinmedicine.16-6-514
25. Jones B, Jarvis P, Lewis JA, Ebbutt AF. Trials to assess equivalence: the importance of rigorous methods. *BMJ*. 1996;313(7048):36-39. doi:10.1136/bmj.313.7048.36
26. Saville BR, Berry DA, Berry NS, Viele K, Berry SM. The Bayesian time machine: accounting for temporal drift in multi-arm platform trials. *Clin Trials*. 2022;19(5):490-501.
27. R Foundation for Statistical Computing. A Language and Environment for Statistical Computing [computer program]. R Foundation for Statistical Computing; 2024.
28. Stan Development Team. Stan Modeling Language Users Guide and Reference Manual version 2.35 [computer program]. Accessed November 25, 2024. <https://mc-stan.org>

29. Pisciotta W, Passannante A, Arina P, Alotaibi K, Ambler G, Arulkumaran N. High-flow nasal oxygen versus conventional oxygen therapy and noninvasive ventilation in COVID-19 respiratory failure: a systematic review and network meta-analysis of randomised controlled trials. *Br J Anaesth*. 2024;132(5):936-944. doi:10.1016/j.bja.2023.12.022
30. Weng CL, Zhao YT, Liu QH, et al. Meta-analysis: noninvasive ventilation in acute cardiogenic pulmonary edema. *Ann Intern Med*. 2010;152(9):590-600. doi:10.7326/0003-4819-152-9-201005040-00009
31. Osadnik CR, Tee VS, Carson-Chahhoud KV, Picot J, Wedzicha JA, Smith BJ. Non-invasive ventilation for the management of acute hypercapnic respiratory failure due to exacerbation of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2017;7(7):CD004104. doi:10.1002/14651858.CD004104.pub4
32. Plant PK, Owen JL, Elliott MW. Early use of non-invasive ventilation for acute exacerbations of chronic obstructive pulmonary disease on general respiratory wards: a multicentre randomised controlled trial. *Lancet*. 2000;355(9219):1931-1935. doi:10.1016/S0140-6736(00)02323-0
33. Azevedo LCP, Park M, Salluh JIF, et al; ERICC (Epidemiology of Respiratory Insufficiency in Critical Care) investigators. Clinical outcomes of patients requiring ventilatory support in Brazilian intensive care units: a multicenter, prospective, cohort study. *Crit Care*. 2013;17(2):R63. doi:10.1186/cc12594
34. Santa Cruz R, Matesa A, Gómez A, et al. Mortality due to acute respiratory distress syndrome in Latin America. *Crit Care Med*. 2024;52(8):1275-1284. doi:10.1097/CCM.0000000000006312
35. Kurtz P, Bastos LSL, Dantas LF, et al. Evolving changes in mortality of 13,301 critically ill adult patients with COVID-19 over 8 months. *Intensive Care Med*. 2021;47(5):538-548. doi:10.1007/s00134-021-06388-0
36. Lau VI, Dhanoa S, Cheema H, et al. Non-COVID outcomes associated with the coronavirus disease-2019 (COVID-19) pandemic effects study (COPES): a systematic review and meta-analysis. *PLoS One*. 2022;17(6):e0269871. doi:10.1371/journal.pone.0269871