

ORIGINAL ARTICLE

Performance of Creatinine and Cystatin C-Based Equations to Estimate Glomerular Filtration Rate Among Patients With Heart Failure

John Roth¹, MD; John C. Lieske², MD; Sandra M. Herrmann, MD; A.M. Arruda-Olson, MD, PhD; Joerg Herrmann, MD; Wendy McCallum, MD; Timothy S. Larson, MD; Andrew D. Rule³, MD; Silvia M. Titan⁴, MD, PhD

BACKGROUND: The performance of estimated glomerular filtration rate (eGFR) among patients with heart failure (HF) may be worse than in the general population due to a higher prevalence of confounding factors affecting creatinine and cystatin C. Studies in this area are scarce and not stratified by type of HF. We evaluated the performance of current creatinine and cystatin C equations (eGFR creatinine-based equation [eGFR_{cr}], eGFR serum cystatin C-based, and eGFR creatinine-cystatin C equation) compared with measured GFR (mGFR) among patients with HF stratified by ejection fraction.

METHODS: We pulled data on Mayo Clinic patients with an mGFR performed for clinical indications from 2011 to 2023, with serum creatinine and cystatin C measured within 7 days and an echocardiogram performed up to 1 year before the mGFR date. HF was identified by the presence of *International Classification of Diseases* codes within 1 year before the mGFR and subgrouped into ejection fraction (EF) $\geq 50\%$ (HFEF $\geq 50\%$, n=182) or $<50\%$ (HFEF $<50\%$, n=115) and compared with no-HF controls (n=1871). CKD-EPI eGFR_{cr}, eGFR serum cystatin C-based, and eGFR creatinine-cystatin C equations were calculated, and compared for bias (mGFR minus eGFR) and accuracy (1-P30, proportion of people with $\geq 30\%$ difference between eGFR and mGFR). CIs were generated by bootstrapping.

RESULTS: The HF groups were characterized by older age, higher proportion of males, more diabetes, higher creatinine, and higher cystatin C than controls. eGFR_{cr} overestimated mGFR to a greater extent in both HF groups as compared with controls, whereas the eGFR serum cystatin C-based equation showed similar underestimation in both HF groups and controls. In the HF groups, cystatin C-based equations were more accurate than eGFR_{cr} particularly within HFEF $<50\%$ (1-P30 of 28% and 34% for eGFR serum cystatin C-based and eGFR creatinine-cystatin C equations, respectively, versus 60% for eGFR_{cr}), whereas eGFR creatinine-cystatin C equation was more accurate in controls.

CONCLUSIONS: Among patients with HF, eGFR_{cr} demonstrates inferior performance (more bias and less accuracy) compared with cystatin C-based eGFRs, with this effect being more pronounced in those with HFEF $<50\%$.

Key Words: creatinine ■ cystatin C ■ humans ■ prevalence ■ stroke volume

Accurate assessment of kidney function and diagnosis of chronic kidney disease is essential among patients with heart failure (HF) for accurate drug dosing, assessment of eligibility for treatment and trials, evaluation of risks related to treatment, safety of imaging and procedures, consideration for combined heart-kidney transplant, and prognostication.^{1–3} However, precise assessment of kidney function has posed an ongoing

challenge for decades. Most commonly, estimation of the glomerular filtration rate (GFR) relies on equations that include demographics and endogenous filtration markers such as creatinine and cystatin C (estimated GFR or eGFR). Current guidelines within the US recommend the use of a creatinine-based equation (eGFR_{cr}) for initial assessment.⁴ If there is clinical suspicion this estimate is inaccurate, providers may pursue confirmatory testing

Correspondence to: Silvia Titan, MD, PhD, Division of Nephrology and Hypertension, Mayo Clinic, 200 First St SW, Rochester, MN 55905. Email titan.silvia@mayo.edu
Supplemental Material is available at <https://www.ahajournals.org/doi/suppl/10.1161/CIRCHEARTFAILURE.125.013014>.
For Sources of Funding and Disclosures, see page XXX.

© 2025 American Heart Association, Inc.

Circulation: Heart Failure is available at www.ahajournals.org/journal/circheartfailure

WHAT IS NEW?

- Large cohort study in a major US health care system showing creatinine-based estimated glomerular filtration rate equations compared with to cystatin C-based estimated glomerular filtration rate perform worse (more bias and less accuracy) in those with heart failure.
- This effect is more pronounced in those with heart failure with reduced ejection fraction as opposed to those with preserved ejection fraction.

WHAT ARE THE CLINICAL IMPLICATIONS?

- Highlights the frequency and magnitude of errors in estimated glomerular filtration rate in patients with heart failure.
- Identifies the combined creatinine-cystatin C, as opposed to creatinine-only equations as the preferred means of estimating glomerular filtration rate in patients with heart failure.

Nonstandard Abbreviations and Acronyms

CKD-EPI	Chronic Kidney Disease Epidemiology Collaboration
CKD-EPIcr	Chronic Kidney Disease Epidemiology Collaboration creatinine equation
EF	ejection fraction
eGFR	estimated glomerular filtration rate
eGFRcr	estimated glomerular filtration rate creatinine-based equation
eGFRcrys	estimated glomerular filtration rate creatinine-cystatin C equation
eGFRcys	estimated glomerular filtration rate serum cystatin C-based
EKFC	European Kidney Failure Consortium
HF	heart failure
ICD	International Classification of Diseases
mGFR	measured glomerular filtration rate
NoHF	no HF

with the combined creatinine-cystatin C equation (eGFR-crys).⁴ In situations where accurate assessment of GFR is paramount for clinical decision-making, guidelines recommend obtaining a measured GFR (mGFR) based on urinary clearance or plasma decay of an exogenous filtration marker, considered as gold standard methods. However, clinical use of the combined eGFR equation or mGFR remains low, particularly among nonnephrologists. Patients with HF may be at increased risk for factors that confound endogenous filtration markers resulting in inaccurate GFR estimation.⁵ For instance, sarcopenia,

cachexia, and frailty are common in HF and lead to wasting of skeletal muscle, which is the primary source of creatinine, causing overestimation of true GFR with eGFRcr.^{6–8} For cystatin C, underestimation of GFR may occur as a function of changes in adipose tissue mass, smoking, thyroid diseases, endogenous and exogenous glucocorticoids,⁹ and stress states leading to increased cystatin C production.⁸ Currently, only a few studies with small sample sizes have evaluated the performance of various eGFR equations within HF populations.^{10–12} In addition, causes of HF are heterogeneous and could be associated with variable effects on creatinine and cystatin C generation. However, no studies to date have examined eGFR versus mGFR among those with reduced versus normal ejection fraction (EF). Moreover, these studies did not evaluate the race-free Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation¹³ or recent equations that are widely used in Europe and other countries.

Thus, the aim of this study was to assess the performance of the most recent CKD-EPI and European Kidney Failure Consortium (EKFC) equations compared with mGFR in people with HF stratified by HF subtype.

METHODS

This study was approved by the Mayo Clinic IRB (23-002121). Clinical data and laboratory results were extracted from the Mayo Clinic electronic health record. Briefly, we retrieved all patients >18 years old with mGFR obtained for clinical purposes across all Mayo Clinic sites from May 2011 (after creatinine and cystatin C had both been standardized) to September 2023 who also had a transthoracic echocardiogram within 1 year of the mGFR date (n=32218). Patients were then classified as having HF if there was an encounter with *International Classification of Diseases* (ICD) 9 or ICD10 code for HF (ICD10 I50 and ICD9 428)¹⁴ within 1 year before mGFR (n=2662) and subclassified according to EF as HFEF ≥50% or HFEF<50%. A control group was defined as people without any encounter for HF (NoHF) who also had an echocardiogram within 1 year of mGFR showing EF ≥50% (n=14 096). Patient with a creatinine (Creat cohort, n=14 857, being n=1467 for HFEF ≥50% and n=994 for HFEF <50%) or creatinine and cystatin C (CreatCys cohort, n=2168, being n=182 for HFEF ≥50% and n=115 for HFEF<50%) performed within 7 days of the date of the mGFR and no missing or implausible laboratory, height, or weight values were retained. If multiple mGFRs were obtained, only the first was used. In our institution, mGFRs are typically done in the outpatient setting. Height and weight were available for the same day for 95% of the sample; in those whose same-day height and weight were not available, we used the closest height and weight within 2 weeks of the date of mGFR. Patients on dialysis were excluded. Validation of HF categories was performed via chart review of 30 randomly selected patients (10 from each group) with excellent performance of 100% correct diagnosis and EF for both the HF and NoHF groups. Furthermore, 100% of patients classified as NoHF did not fulfill the Framingham Heart Failure diagnostic criteria,¹⁵

whereas 95% of those identified as HF did. [Figure S1](#) depicts the flow and criteria used to define the study population.

All laboratory tests were performed in the Mayo Clinic Central Clinical Chemistry Laboratory. Creatinine was measured using a Roche enzymatic creatinase assay standardized to international reference material using a Roche Cobas chemistry analyzer (c701 or c501; Roche Diagnostics, Indianapolis). Cystatin C was measured by an immunoturbidometric assay traceable to an international reference material (European Reference Material). Gentian reagents were used between May 2011 and May 2021, and Roche reagents from May 2021 to September 2023. We have shown previously that the residuals of the regression of cystatin C on mGFR did not change meaningfully after the introduction of the new assay.¹⁶ The following eGFR equations were evaluated: 2021 CKD-EPI creatinine equation,¹³ 2012 CKD-EPI cystatin C equation,¹⁷ 2021 CKD-EPI creatinine-cystatin C equation,¹³ 2024 EKFC creatinine equation,¹⁸ 2023 EKFC cystatin equation,¹⁹ and the EKFC creatinine-cystatin C equation, which is the average of the 2024 EKFC creatinine equation and the 2023 EKFC cystatin equation. Required demographics were extracted from the electronic health record or from a questionnaire administered at the time of the mGFR test. Diabetes was defined as self-reported diabetes or presence of an ICD code for diabetes (ICD10 E08–E13, ICD9 249 and 250), and transplantation status was defined via a verified medical history or presence of ICD codes for solid and hematologic transplants (except cornea) before mGFR.

mGFR was measured throughout the study period by non-radiolabeled iothalamate urinary clearance following a standardized protocol. Briefly, patients were asked to report for testing early in the day and in a fasting state to minimize diurnal or dietary variations. Iothalamate was administered subcutaneously followed by oral hydration to encourage urine flow, and a urinary catheter was placed in cases of incomplete bladder emptying. Plasma and urine were collected in timed intervals, and iothalamate was quantified by liquid chromatography tandem mass spectrometry at the Mayo Clinic Renal Testing Laboratory.²⁰ Measured GFR was normalized as mL/min per 1.73 m² with body surface area as estimated by the DuBois formula.²¹

Bias was defined as mGFR minus eGFR, and accuracy by 1-P30, that is, the percentage of people with an eGFR that differed from mGFR by $\geq 30\%$; root mean square error and CIs by bootstrapping ($n=500$) were also computed. The McNemar test was used to compare 1-P30% for paired samples. Results were stratified by transplant status. Another analysis used a control group for each HF group matched for key covariates (age, sex, self-reported race, body mass index, diabetes, and transplant status at the date of mGFR, and mGFR below or above 60 mL/min per 1.73 m²). Propensity score matching using the nearest neighbor was performed at a 1:1 ratio without replacement and a caliper of 0.2 (R package MatchIt). CKD staging was assessed according to the Kidney Disease Improving Global Outcomes classification²² for mGFR and each eGFR, and expressed as the percent match when staging using mGFR versus each eGFR.

All analyses and plots were performed using RStudio (2023.12.1 version). The data that support the findings of this study are available from the corresponding author upon reasonable request.

RESULTS

Table 1 contains descriptive data for the Creat and CreatCys cohorts of the 3 groups by HF status: NoHF, HFEF $\geq 50\%$, and HFEF $<50\%$. Patients with HFEF $\geq 50\%$ and HFEF $<50\%$ were older, had a higher percentage of male sex, a higher frequency of diabetes, and lower kidney function compared with those with NoHF. There was a high frequency of mGFR obtained posttransplant as part of kidney, bone marrow, and liver transplant recipient care. Most variables were similar between groups, except for mGFR, which was lower in CreatCys compared with Creat, especially among those with HF.

Table 2 contains eGFR performance metrics for the 3 groups in the Creat and CreatCys cohorts. The CKD-EPI creatinine equation (CKD-EPIcr) demonstrated minimal bias in the NoHF groups but overestimation in HFEF $\geq 50\%$ and HFEF $<50\%$ in the 2 cohorts; a similar pattern was observed using the 2024 EKFC creatinine equation. In terms of precision, in the Creat cohort, 1-P30% of 22%, 30%, and 37% were observed with CKD-EPIcr in the NoHF, HFEF $\geq 50\%$, and HFEF $<50\%$ groups, respectively, and this trend was even more pronounced in the CreatCys cohort (1-P30% of 22%, 36%, and 60% in NoHF, HFEF $\geq 50\%$, and HFEF $<50\%$, respectively). When comparing the performance of creatinine and cystatin C equations within the CreatCys cohort, eGFRcr showed the worst performance in the HF groups, with a 1-P30% of 36% in HFEF $\geq 50\%$ and 60% in HFEF $<50\%$ for the CKD-EPIcr (and 36% in HFEF $\geq 50\%$ and 56% in HFEF $<50\%$ with the 2024 EKFC creatinine equation), in comparison to 22% observed in the NoHF group. Conversely, both CKD-EPI and EKFC eGFRrcys and serum cystatin C-based (eGFRcys) equations performed better than eGFRcr in the HF groups, with eGFRrcys being the most accurate in HFEF $\geq 50\%$ and eGFRcys the most accurate in HFEF $<50\%$. Similar trends were observed when comparing the root mean square error between subgroups within each cohort. Figures 1 and 2 show the scatter plots between the mGFR and the CKD-EPI and EKFC equations, respectively, in the CreatCys cohort. [Table S1](#) shows the performance metrics stratified by transplant status, and in both populations the cystatin C-based equations performed better than eGFRcr, with eGFRcr showing larger errors in the HF groups compared with the non-HF group.

As a sensitivity analysis, the performance metrics were also evaluated using a matched control group. [Tables S2 and S3](#) show the descriptive variables before and after matching for each HF group and its matched control in the Creat and CreatCys cohorts, respectively. [Tables S4 and S5](#) depict the performance of eGFR on mGFR in the matched groups in both cohorts. While adjustments did attenuate differences, the performance was still worse in the HF groups, with larger bias and imprecision observed with eGFRcr compared with eGFRrcys or eGFRcys.

Table 1. Descriptive Data at mGFR Date of the Creat and CreatCys Cohorts

	Creat cohort				CreatCys cohort			
	NoHF	HFEF ≥50%	HFEF <50%	P	NoHF	HFEF ≥50%	HFEF <50%	P
	n=12 396	n=1 467	n=994		n=1 871	n=182	n=115	
Age at mGFR, y; mean, SD	57 (12)	59 (13)	58 (13)	<0.0001	57 (12)	61 (13)	60 (14)	<0.0001
Gender, male; n (%)	7079 (57)	849 (58)	717 (72)	<0.0001	976 (52)	110 (60)	86 (75)	<0.0001
Self-reported race, White; n (%)	10 966 (89)	1267 (86)	846 (85)	0.0004	1660 (89)	161 (89)	96 (84)	0.14
Diabetes, n (%)	2890 (23)	787 (54)	400 (40)	<0.0001	376 (20)	120 (66)	49 (43)	<0.0001
Posttransplant status, n (%)	5331 (43)	1020 (70)	275 (28)	<0.0001	616 (33)	91 (50)	32 (28)	<0.0001
BMI, kg/m ² ; mean (SD)	28.9 (6.8)	29.2 (6.7)	28.8 (5.9)	0.18	28.5 (5.7)	28.8 (6.2)	28.4 (6.1)	0.78
Creatinine, mg/dL; mean (SD)	1.2 (0.8)	1.6 (0.8)	1.5 (0.7)	<0.0001	1.3 (0.9)	1.8 (0.9)	1.7 (0.8)	<0.0001
Cystatin, mg/L; mean (SD)	1.4 (0.7)	2.2 (1.0)	2.0 (0.7)	<0.0001
BSA-adj. mGFR, mL/min per 1.73 m ² ; mean (SD)	75 (32)	53 (26)	55 (26)	<0.0001	72 (33)	42 (24)	41 (19)	<0.0001
BSA-unadj. mGFR, mL/min per 1.73; mean (SD)	84 (38)	60 (31)	63 (32)	<0.0001	80 (38)	47 (28)	48 (24)	<0.0001
CKD-EPI creatinine equation, mL/min per 1.73 m ² ; mean (SD)	74 (28)	58 (26)	59 (25)	<0.0001	72 (28)	50 (26)	50 (23)	<0.0001
CKD-EPI cystatin C equation, mL/min per 1.73 m ² ; mean (SD)	68 (33)	36 (22)	38 (18)	<0.0001
CKD-EPI creatinine-cystatin C equation, mL/min per 1.73m ² ; mean (SD)	71 (31)	42 (23)	43 (19)	<0.0001
EKFC creatinine equation, mL/min per 1.73 m ² ; mean (SD)	71 (25)	56 (24)	58 (23.1)	<0.0001	69 (26)	49(24)	49 (22)	<0.0001
EKFC cystatin equation, mL/min per 1.73 m ² ; mean (SD)	67 (28)	40 (20)	41 (17)	<0.0001
EKFC creatinine-cystatin C equation, mL/min per 1.73 m ² ; mean (SD)	68 (25)	44 (21)	45 (17)	<0.0001

P values reflect comparisons across groups within each cohort using Kruskal-Wallis and χ^2 tests for continuous and categorical variables, respectively. BMI indicates body mass index; BSA, body surface area; CKD-EPI indicates Chronic Kidney Disease Epidemiology Collaboration; Creat cohort, cohort for which creatinine was available; CreatCys cohort, cohort for which both creatinine and cystatin C were available; EKFC, European Kidney Failure Consortium; HFEF, heart failure with reduced or mildly reduced ejection fraction; HFEF, heart failure with preserved ejection fraction; mGFR, measured glomerular filtration rate; and NoHF, no heart failure.

Table 3 shows the percent concordance of CKD stage for the 3 groups for the Creat and CreatCys cohorts. Overall, the concordance of CKD staging of eGFRs compared with mGFR was lower in the HF groups than in NoHF. Both eGFR CKD-EPI and EKFC were subject to significant misclassification of CKD staging in the HFEF ≥50% and HFEF <50% groups, which was most pronounced in the latter group. Percent concordance for the NoHF, HFEF ≥50%, and HFEF<50% groups was 76%, 59%, and 55% with CKD-EPI and 75%, 60%, and 54% with EKFC, respectively. The eGFRcys demonstrated the highest percent concordance in NoHF (79% for both CKD-EPI and EKFC) and HFEF ≥50% (63% and 59% for CKD-EPI and EKFC, respectively), while eGFRcys showed the highest concordance in HFEF <50% (51% and 53% for CKD-EPI and EKFC, respectively). The concordance of eGFRcr in the HFEF <50% group was as low as 29% for both CKD-EPI and EKFC.

DISCUSSION

Our study has 2 main findings. First, bias and accuracy observed in the HFEF ≥50% and HFEF <50% groups

with all eGFR equations were generally worse than in our control group with no HF. This reduced accuracy translated into mismatched CKD staging in both HF groups, reaching concordance in staging in only 55% and 29% in the HFEF <50% group for Creat and CreatCys, respectively. This finding might be expected, since the HF population is quite different from the NoHF population due to a higher burden of cardiovascular risk factors (age, sex, diabetes, hypertension, kidney disease, smoking, cardiotoxic exposures) that may affect non-GFR determinants of eGFR. Additional contributors to the non-GFR determinants of eGFR related to HF include changes in metabolism, cachexia, inflammation, and others. Considering that many of these factors are known to interfere with creatinine and cystatin C and their performance in predicting GFR, worse performance in the HF groups was not unexpected. Regardless of the mechanism, our study demonstrates that the performance of eGFR to predict mGFR is worse in the HF population than in the control group without HF. This carries significant clinical implications and identifies those with HF as a high-risk population that may have additional health risks related to inaccuracies in GFR estimation.

Table 2. Performance of CKD-EPI and EKFC eGFR Equations in the Creat and CreatCys Cohorts Stratified by Heart Failure Diagnosis

	Creat cohort			CreatCys cohort		
	NoHF	HFEF ≥50%	HFEF <50%	NoHF	HFEF ≥50%	HFEF <50%
	n=12 396	n=1467	n=994	n=1871	n=182	n=115
Median bias (CI)						
CKD-EPI creatinine equation	0.5 (0.2 to 0.7)	−3.7 (−4.5 to −3)	−4.1 (−5 to −3.2)	0 (−0.6 to 0.8)	−6 (−8.3 to −4.4)	−8.1 (−11.4 to −3.7)
CKD-EPI cystatin C equation	4.3 (3.4 to 5.4)	4.7 (3.6 to 6.3)	2.6 (−0.4 to 4.4)
CKD-EPI creatinine-cystatin C equation	1.1 (0.4 to 1.9)	0.5 (−0.9 to 2)	−1.5 (−3.1 to 1.3)
EKFC creatinine equation	2.7 (2.4 to 3)	−2.2 (−3.2 to −1.3)	−3.7 (−4.9 to −2.5)	1.9 (0.8 to 2.5)	−4.5 (−6.5 to −2.5)	−7.7 (−11.4 to −2.9)
EKFC cystatin equation	3.4 (2.4 to 4.5)	0.9 (−1.2 to 2.7)	−0.9 (−3.6 to 0.4)
EKFC creatinine-cystatin C equation	2.6 (1.9 to 3.2)	−2 (−3.6 to −0.6)	−6.1 (−11.1 to −3.1)
1-P30 (CI)						
CKD-EPI creatinine equation	21.6 (20.8 to 22.3)	30.1 (27.8 to 32.7)	37.4 (34.3 to 40.4)	22.4 (20.7 to 24.3)	36.3 (29.4 to 43.4)	60 (50.9 to 68.7)
CKD-EPI cystatin C equation	24.6 (22.5 to 26.6)	31.9 (25 to 39)	27.8 (20 to 35.7)
CKD-EPI creatinine-cystatin C equation	15.4 (13.8 to 16.9)	25.3 (18.1 to 31.3)	33.9 (25.2 to 42.6)
EKFC creatinine equation	21.3 (20.6 to 22)	27.5 (25.2 to 29.7)	35.8 (33 to 38.7)	21.4 (19.7 to 23.2)	35.7 (28.6 to 41.8)	55.7 (46.1 to 64.3)
EKFC cystatin equation	20.8 (18.9 to 22.7)	30.8 (23.1 to 37.4)	33 (24.3 to 42.6)
EKFC creatinine-cystatin C equation	15.3 (13.6 to 16.9)	29.1 (22 to 35.2)	42.6 (33 to 50.8)
RMSE						
CKD-EPI creatinine equation	19.2	16.8	19	18.2	18.5	22.0
CKD-EPI cystatin C equation	19.2	15.8	13.4
CKD-EPI creatinine-cystatin C equation	16.1	12.9	13.5
EKFC creatinine equation	19.5	15.9	17.9	18.4	17.4	21
EKFC cystatin equation	18.3	14.4	12.8
EKFC creatinine-cystatin C equation	16.5	12.8	14.3

Bias was computed as mGFR−eGFR. Bias and RMSE are expressed as mL/min per 1.73 m². McNemar test *P* values for 1-P30 related to paired samples in the CreatCys cohort: NoHF group, *P*<0.0001 for serum creatinine-based eGFR or serum cystatin c-based eGFR compared with eGFR_{crecys}; HFEF ≥50%, *P*<0.007 and 0.03 for CKD-EPI creatinine equation, and EKFC creatinine equation in comparison to CKD-EPI creatinine-cystatin C equation and EKFC creatinine-cystatin C equation, respectively, and NS for CKD-EPI cystatin C equation and EKFC cystatin equation in comparison to CKD-EPI creatinine-cystatin C equation and EKFC creatinine-cystatin C equation, respectively; HFEF <50%, *P*<0.0001 and 0.003 for CKD-EPI creatinine equation and EKFC creatinine equation in comparison to CKD-EPI creatinine-cystatin C equation and EKFC creatinine-cystatin C equation, respectively; and NS for CKD-EPI cystatin C equation and EKFC cystatin equation in comparison to CKD-EPI creatinine-cystatin C equation and EKFC creatinine-cystatin C equation, respectively. Creat cohort indicates cohort for which creatinine was available; CreatCys cohort, cohort for which both creatinine and cystatin C were available; CKD-EPI indicates Chronic Kidney Disease Epidemiology Collaboration; eGFR, estimated glomerular filtration rate; EKFC, European Kidney Failure Consortium; HFEF, heart failure with reduced ejection fraction; HFEF, heart failure with preserved ejection fraction; mGFR, measured glomerular filtration rate; NoHF, no prior diagnosis of heart failure; P30, proportion of eGFR within 30% of measured GFR; and RMSE, root mean square error.

The second main finding refers to the comparative performance of eGFRs within the HF population. Unlike the no HF controls, eGFR_{cr} had the worst performance in comparison to both eGFR_{crecys} and eGFR_{cys} within the HF groups. Specifically, eGFR_{cr} demonstrated significant bias and was prone to overestimation of mGFR, as well as poor accuracy. This effect was more pronounced in the HFEF <50% group than in HFEF ≥50%. An eGFR within 30% of mGFR has been considered acceptable

for clinical decision-making, and Kidney Disease Improving Global Outcomes recommends eGFR equations should strive for this benchmark in >90% of patients.²² However, in the CreatCys cohort, use of eGFR_{cr} resulted in up to 60% of people falling outside this metric, with errors above 30%. Even in the Creat cohort (which is less prone to selection bias than the CreatCys cohort), despite root mean square error being similar in the NoHF and HFEF <50% groups, the frequency of larger errors




Figure 1. Scatter plots and regression lines of measured glomerular filtration rate (mGFR) and estimated glomerular filtration rate (eGFR) values for the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equations in the CreatCys cohort. Dots represent values for eGFR and mGFR expressed as mL/min per 1.73 m². Dotted line marks upper and lower 30% difference between mGFR and eGFR. Blue line shows the regression lines for the scatter plots and black line shows the reference line (perfect agreement between mGFR and eGFR). CKD-EPIcr indicates Chronic Kidney Disease Epidemiology Collaboration creatinine equation; CKD-EPIcrs, Chronic Kidney Disease Epidemiology Collaboration creatinine and cystatin C equation; CKD-EPIcys, Chronic Kidney Disease Epidemiology Collaboration cystatin C equation; HFEF, heart failure with preserved ejection fraction; HFEF, heart failure with reduced ejection fraction; and NoHF, no heart failure.

(>30%) went from 22% in the NoHF group, to 30% in the HFEF ≥50% group, and up to 37% in the HFEF <50% group. Possible contributors include sarcopenia, malnutrition, and cachexia which are associated with low muscle mass, the primary determinant of creatinine generation.¹¹ eGFRcrs demonstrated the best performance in the HFEF ≥50% group, while eGFRcys had the best performance in the HFEF <50% group followed

closely by the eGFRrcys. Regardless, eGFRcys and eGFRrcys still show only modest precision with errors >30% occurring in 25% to 42% of patients in the HF groups. Between CKD-EPI and EKFC equations, results were overall similar in numbers and patterns in agreement with prior publications suggesting these equations have found a point of convergence.²³ As there was a likely selection bias regarding which patients had cystatin




Figure 2. Scatter plots and regression lines of the measured glomerular filtration rate (mGFR) and estimated glomerular filtration rate (eGFR) values for the European Kidney Function Consortium (EKFC) equations in the CreatCys cohort.

Dots represent values for eGFR and mGFR expressed as mL/min per 1.73 m². Dotted line marks upper and lower 30% difference between mGFR and eGFR. Blue line shows the regression lines for the scatter plots and black line shows the reference line (perfect agreement between mGFR and eGFR). EKFCcr indicates European Kidney Function Consortium creatinine equation; EKFCcrs, European Kidney Function Consortium creatinine-cystatin C equation; EKFCcys, European Kidney Function Consortium cystatin C equation; HFEF, heart failure with preserved ejection fraction; HFEF, heart failure with reduced ejection fraction; and NoHF, no heart failure.

C testing due to a reason to question the eGFRcr result, it may be reasonable to use eGFRcrs over eGFRcys in patients with HF. This would also be consistent with the general eGFR literature that suggests eGFRcrs (but not eGFRcys) is more accurate than eGFRcr.^{13,16,24}

Two prior small studies have assessed the performance of eGFRs in HF. Kervella et al¹¹ evaluated the 2012 CKD-EPI creatinine and cystatin C equations on 66 people with HF as compared with urinary clearance

of inulin, showing that the CKD-EPI cystatin C equation had smaller bias but all 3 equations performed poorly, with P30% of 65% for the CKD-EPI cystatin C equation, 52% for Chronic Kidney Disease Epidemiology Collaboration creatinine and cystatin C equation, and 33% for CKD-EPIcr. In another study using plasma clearance of dextran in 50 patients, P30% of 74% for Chronic Kidney Disease Epidemiology Collaboration creatinine and cystatin C equation, 66% for CKD-EPIcr and 56% for

Table 3. Concordance of CKD Stage With eGFR in the Creat and CreatCys Cohorts Stratified by Heart Failure Diagnosis

	Creat cohort			CreatCys cohort		
	NoHF	HFEF ≥50%	HFEF <50%	NoHF	HFEF ≥50%	HFEF <50%
	n=12 396	n=1467	n=994	n=1871	n=182	n=115
CKD-EPI creatinine equation	76.0	59.6	55.3	75.8	53.3	28.7
CKD-EPI cystatin C equation	70.2	51.6	51.3
CKD-EPI creatinine-cystatin C equation	78.5	63.2	46.1
EKFC creatinine equation	75.4	59.7	54.4	74.8	52.2	29.6
EKFC cystatin equation	73.1	58.2	53.0
EKFC creatinine-cystatin C equation	79.4	59.3	40.9

CKD-EPI indicates Chronic Kidney Disease Epidemiology Collaboration; Creat cohort, cohort for which creatinine was available; CreatCys cohort, cohort for which both creatinine and cystatin C were available; EKFC, European Kidney Failure Consortium; HFEF, heart failure with reduced ejection fraction; HFEF, heart failure with preserved ejection fraction; and NoHF, no heart failure.

the CKD-EPI cystatin C equation were described.¹² Our study showed that cystatin C-based equations perform better than eGFRcr in people with HF, although still with moderate precision, and that the overestimation with eGFRcr tends to be more pronounced in those with HFEF <50% as compared with HFEF ≥50%.

Inaccurate estimates of GFR carry significant implications in the care of patients with HF. For example, GFR may affect the decision-making on the use and titration of medications, such as angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, mineralocorticoid receptor antagonists, sodium-glucose cotransporter 2 inhibitors, and glucagon-like peptides.^{25,26} It also impacts drug dosing of some direct oral anticoagulants and low-molecular-weight heparins, with the potential for increased risk of bleeding due to overdosing and ischemic or embolic events due to underdosing.²⁷ GFR estimation is part of the evaluation for mechanical circulatory support such as left ventricular assist devices,^{28,29} and is a crucial parameter to define eligibility for heart-kidney transplantation versus heart transplantation only.^{30–33} Per ACC/AHA and Kidney Disease Improving Global Outcomes guidelines, patients at elevated risk for contrast-induced nephropathy should receive adequate periprocedural hydration, and low GFR, particularly in the setting of HF, is considered a significant risk factor.^{4,34–37} It is interesting to note that HF trials may use eGFRcr as an inclusion, criterion, which may contribute to heterogeneity in the recruited population and treatment effects.

To the best of our knowledge, this is the single largest investigation into the performance of various eGFR equations in the HF population. Other strengths of our study include the use of standardized assays for serum cystatin C and creatinine, use of urinary iothalamate clearance as mGFR, availability of a control group without HF within the same population, stratification for HF, and the fact that we were able to test modern race-free equations that had not been tested in prior studies.

This study contains several limitations. First, there was likely significant selection bias in the cohorts generated.

The HF groups may have overrepresented patients with malignancy, multiple comorbidities, or organ transplants. In addition, selection bias was likely present both in patients who underwent mGFR testing and received echocardiograms, as well as in the subset with cystatin C testing. Notably, mGFR is done by protocol for certain subgroups in our institution (eg, posttransplantation), though we attempted to address this potential bias by stratified analysis by transplant status, which demonstrated similar results regardless of transplant status in the HF patients (but not in the no HF controls). Our population was also predominantly White so our results may not apply to other populations with different demographic characteristics. Notably, the race-free CKD-EPI eGFRcr equation overestimates GFR in White patients and this may have contributed to the overestimation with eGFRcr in HF seen in this study.³⁸ Since our electronic health record does not include information on medication dispensation and discontinuation systematically, we could not precisely determine if people were on chronic use of steroids at the time of mGFR date. Steroids are known to increase cystatin C production⁹ and may cause underestimation of cystatin C-based equations. In addition, we could not assess other medications that can influence creatinine or cystatin C and therefore may confound the performance of eGFRs. Creatinine and cystatin were mostly measured within 1 to 2 days of the mGFR date, but we also allowed labs from up to 1 week of mGFR testing as this introduced minimal variation in the outpatient setting.¹⁶ Our sample sizes for the HFEF ≥50% and HFEF <50% groups were relatively small. Finally, we did not have information on muscle mass, body composition, and frailty, precluding stratified analysis for variables of interest.

CONCLUSIONS

Collectively, the results demonstrate suboptimal performance of eGFRcr in HF populations, with this effect being more pronounced in those with HFEF <50%.

While significant inaccuracy still exists with eGFR_{cr} or eGFR_{cr}s, our data indicate superior performance of these in comparison to eGFR_{cr} in both HFEF $\geq 50\%$ and HFEF $< 50\%$. These findings support the use of the eGFR_{cr}s equations in HF, consistent with the general recommendation in the Kidney Disease: Improving Global Outcomes guidelines for use of eGFR_{cr}s when a more accurate estimation of GFR is needed.

ARTICLE INFORMATION

Received March 4, 2025; accepted June 30, 2025.

Affiliations

Department of Internal Medicine (J.R.), Division of Nephrology and Hypertension (J.C.L., S.M.H., T.S.L., A.D.R., S.M.T.), and Department of Cardiovascular Medicine (A.M.A.-O., J.H.), Mayo Clinic, Rochester, MN. Division of Nephrology, Tufts Medical Center, Boston, MA (W.M.C.).

Acknowledgments

All the above-mentioned authors have acknowledged and approved their inclusion in this publication and are supportive of the data and conclusions.

Sources of Funding

Mayo Clinic, Project 95070002; Mayo Clinic CcTS grant UL1TR002377.

Disclosures

None.

Supplemental Material

Tables S1–S5

Figure S1

REFERENCES

- Pinsino A, Carey MR, Husain S, Mohan S, Radhakrishnan J, Jennings DL, Ngunjiri AS, Ladanyi A, Braghieri L, Takeda K, et al. The difference between cystatin C- and creatinine-based estimated GFR in heart failure with reduced ejection fraction: insights from PARADIGM-HF. *Am J Kidney Dis*. 2023;82:521–533. doi: 10.1053/j.ajkd.2023.03.005
- Chen DC, Shlipak MG, Scherzer R, Bansal N, Potok OA, Rifkin DE, Ix JH, Muir AN, Hsu CY, Estrella MM. Association of intra-individual differences in estimated GFR by creatinine versus cystatin C with incident heart failure. *Am J Kidney Dis*. 2022;80:762–772.e1. doi: 10.1053/j.ajkd.2022.05.011
- Carrero JJ, Fu EL, Sang Y, Ballew S, Evans M, Elinder CG, Barany P, Inker LA, Levey AS, Coresh J, et al. Discordances between creatinine- and cystatin C-based estimated GFR and adverse clinical outcomes in routine clinical practice. *Am J Kidney Dis*. 2023;82:534–542. doi: 10.1053/j.ajkd.2023.04.002
- Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2024 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int*. 2024;105:S117–S314. doi: 10.1016/j.kint.2023.10.018
- Levey AS, Inker LA. Assessment of glomerular filtration rate in health and disease: a state of the art review. *Clin Pharmacol Ther*. 2017;102:405–419. doi: 10.1002/cpt.729
- Fujimoto Y, Maeda D, Kagiya N, Sunayama T, Dotare T, Jujo K, Saito K, Kamiya K, Saito H, Ogasahara Y, et al. Prevalence and prognostic impact of the coexistence of cachexia and sarcopenia in older patients with heart failure. *Int J Cardiol*. 2023;381:45–51. doi: 10.1016/j.ijcard.2023.03.035
- Konishi M, Kagiya N, Kamiya K, Saito H, Saito K, Ogasahara Y, Maekawa E, Misumi T, Kitai T, Iwata K, et al. Impact of sarcopenia on prognosis in patients with heart failure with reduced and preserved ejection fraction. *Eur J Prev Cardiol*. 2021;28:1022–1029. doi: 10.1093/eurjpc/zwaa117
- Ivey-Miranda JB, Inker LA, Griffin M, Rao V, Maulion C, Turner JM, Wilson FP, Tang WHW, Levey AS, Testani JM. Cystatin C and muscle mass in patients with heart failure. *J Card Fail*. 2021;27:48–56. doi: 10.1016/j.cardfail.2020.07.013
- Kleeman SO, Thakir TM, Demestichas B, Mourikis N, Loiero D, Ferrer M, Bankier S, Riazat-Kesh Y, Lee H, Chantzichristos D, et al. Cystatin C is glucocorticoid responsive, directs recruitment of Trem2+ macrophages, and predicts failure of cancer immunotherapy. *Cell Genom*. 2023;3:100347. doi: 10.1016/j.xgen.2023.100347
- Adingwupu OM, Barbosa ER, Palevsky PM, Vassalotti JA, Levey AS, Inker LA. Cystatin C as a GFR estimation marker in acute and chronic illness: a systematic review. *Kidney Med*. 2023;5:100727. doi: 10.1016/j.xkme.2023.100727
- Kervella D, Lemoine S, Sens F, Dubourg L, Sebbag L, Guebre-Egziabher F, Bonnefoy E, Juillard L. Cystatin C versus creatinine for GFR estimation in CKD due to heart failure. *Am J Kidney Dis*. 2017;69:321–323. doi: 10.1053/j.ajkd.2016.09.016
- Swolinsky JS, Neger NP, Leistner DM, Edelmann F, Knebel F, Tuvshinbat E, Lemke C, Roehle R, Haase M, Costanzo MR, et al. Serum creatinine and cystatin C-based estimates of glomerular filtration rate are misleading in acute heart failure. *ESC Heart Fail*. 2021;8:3070–3081. doi: 10.1002/ehf2.13404
- Inker LA, Eneanya ND, Coresh J, Tighiouart H, Wang D, Sang Y, Crews DC, Doria A, Estrella MM, Froissart M, et al. Chronic Kidney Disease Epidemiology Collaboration. New creatinine- and cystatin C-based equations to estimate GFR without race. *N Engl J Med*. 2021;385:1737–1749. doi: 10.1056/NEJMoa2102953
- Bates BA, Akhabue E, Nahass MM, Mukherjee A, Hiltner E, Rock J, Wilton B, Mittal G, Visaria A, Rua M, et al. Validity of International Classification of Diseases (ICD)-10 diagnosis codes for identification of acute heart failure hospitalization and heart failure with reduced versus preserved ejection fraction in a national medicare sample. *Circ Cardiovasc Qual Outcomes*. 2023;16:e009078. doi: 10.1161/CIRCOUTCOMES.122.009078
- McKee PA, Castelli WP, McNamara PM, Kannel WB. The natural history of congestive heart failure: the Framingham study. *N Engl J Med*. 1971;285:1441–1446. doi: 10.1056/NEJM197111232852601
- Titan SM, Lieske JC, Meeusen JW, Thorson-Spaulin Y, Nowakowski GS, Barreto JN, Barreto EF, Ruddy KJ, Leung N, et al. Performance of creatinine and cystatin-based equations on estimating measured GFR in people with hematological and solid cancers. *Clin J Am Soc Nephrol*. 2025;20:358–366. doi: 10.2215/CJN.0000000616
- Inker LA, Schmid CH, Tighiouart H, Eckfeldt JH, Feldman HI, Greene T, Kusek JW, Manzi J, Van Lente F, Zhang YL, et al. CKD-EPI Investigators. Estimating glomerular filtration rate from serum creatinine and cystatin C. *N Engl J Med*. 2012;367:20–29. doi: 10.1056/NEJMoa1114248
- Delanaye P, Rule AD, Schaeffner E, Cavalier E, Shi J, Hoofnagle AN, Nyman U, Björk J, Pottel H. Performance of the European Kidney Function Consortium (EKFC) creatinine-based equation in United States cohorts. *Kidney Int*. 2024;105:629–637. doi: 10.1016/j.kint.2023.11.024
- Pottel H, Björk J, Rule AD, Ebert N, Eriksen BO, Dubourg L, Vidal-Petiot E, Grubb A, Hansson M, Lamb EJ, et al. Cystatin C-based equation to estimate GFR without the inclusion of race and sex. *N Engl J Med*. 2023;388:333–343. doi: 10.1056/NEJMoa2203769
- Seegmiller JC, Burns BE, Fauq AH, Mukhtar N, Lieske JC, Larson TS. Iohalamate quantification by tandem mass spectrometry to measure glomerular filtration rate. *Clin Chem*. 2010;56:568–574. doi: 10.1373/clinchem.2009.133751
- Du Bois D, Du Bois EF. A formula to estimate the approximate surface area if height and weight be known. 1916. *Nutrition*. 1989;5:303–11; discussion 312.
- National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis*. 2002;39:S1–266.
- Inker LA, Tighiouart H, Adingwupu OM, Shlipak MG, Doria A, Estrella MM, Froissart M, Gudnason V, Grubb A, Kalil R, et al. CKD-EPI and EKFC GFR estimating equations: performance and other considerations for selecting equations for implementation in adults. *J Am Soc Nephrol*. 2023;34:1953–1964. doi: 10.1681/ASN.0000000000000227
- Fu EL, Levey AS, Coresh J, Elinder CG, Rotmans JI, Dekker FW, Paik JM, Barany P, Grams ME, Inker LA, et al. Accuracy of GFR estimating equations in patients with discordances between creatinine and cystatin C-based estimations. *J Am Soc Nephrol*. 2023;34:1241–1251. doi: 10.1681/ASN.0000000000000128
- Das SR, Everett BM, Birtcher KK, Brown JM, Januzzi JL Jr, Kalyani RR, Kosiborod M, Magwire M, Morris PB, Neumiller JJ, et al. 2020 expert consensus decision pathway on novel therapies for cardiovascular risk reduction in patients with type 2 diabetes: a report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol*. 2020;76:1117–1145. doi: 10.1016/j.jacc.2020.05.037
- Maddox TM, Januzzi JL, Allen LA, Breathett K, Brouse S, Butler J, Davis LL, Fonarow GC, Ibrahim NE, Lindenfeld J, et al. 2024 ACC expert consensus

- decision pathway for treatment of heart failure with reduced ejection fraction. *J Am Coll Cardiol*. 2024;83:1444–1488. doi: 10.1016/j.jacc.2023.12.024
27. Ribic C, Crowther M. Thrombosis and anticoagulation in the setting of renal or liver disease. *Hematology Am Soc Hematol Educ Program*. 2016;2016:188–195. doi: 10.1182/asheducation-2016.1.188
 28. Peura JL, Colvin-Adams M, Francis GS, Grady KL, Hoffman TM, Jessup M, John R, Kiernan MS, Mitchell JE, O'Connell JB, et al; American Heart Association Heart Failure and Transplantation Committee of the Council on Clinical Cardiology. Recommendations for the use of mechanical circulatory support: device strategies and patient selection: a scientific statement from the American Heart Association. *Circulation*. 2012;126:2648–2667. doi: 10.1161/CIR.0b013e3182769a54
 29. Pasrija C, Tran D, George P, Sorensen E, Kaczorowski DJ, Ton V-K, Kon ZN, Vorhees H, Sawan M, Griffith BP. Left ventricular assist device implantation may be feasible in appropriately selected patients with severe renal insufficiency. *J Thorac Cardiovasc Surg*. 2020;159:1307–1319.e2. doi: 10.1016/j.jtcvs.2019.03.098
 30. Kittleson MM, Sharma K, Brennan DC, Cheng XS, Chow SL, Colvin M, DeVore AD, Dunlay SM, Fraser M, Garonzik-Wang J, et al; American Heart Association Heart Failure and Transplantation Committee of the Council on Clinical Cardiology; Council on the Kidney in Cardiovascular Disease; Council on Cardiovascular Surgery and Anesthesia; Council on Cardiovascular and Stroke Nursing; Council on Quality of Care and Outcomes Research; and Council on Lifelong Congenital Heart Disease and Heart Health in the Young. Dual-organ transplantation: indications, evaluation, and outcomes for heart-kidney and heart-liver transplantation: a scientific statement from the American Heart Association. *Circulation*. 2023;148:622–636. doi: 10.1161/CIR.0000000000001155
 31. Kobashigawa J, Dadhania DM, Farr M, Tang WHW, Bhimaraj A, Czer L, Hall S, Haririan A, Formica RN, Patel J, et al; Consensus Conference Participants. Consensus conference on heart-kidney transplantation. *Am J Transplant*. 2021;21:2459–2467. doi: 10.1111/ajt.16512
 32. Delgado C, Baweja M, Crews DC, Eneanya ND, Gadegbeku CA, Inker LA, Mendu ML, Miller WG, Moxey-Mims MM, Roberts GV, et al. A unifying approach for GFR estimation: recommendations of the NKF-ASN task force on reassessing the inclusion of race in diagnosing kidney disease. *J Am Soc Nephrol*. 2021;32:2994–3015. doi: 10.1681/ASN.2021070988
 33. Kolsrud O, Ricksten SE, Holmberg E, Fellin M, Karason K, Hammarsten O, Samuelsson O, Dellgren G. Measured and not estimated glomerular filtration rate should be used to assess renal function in heart transplant recipients. *Nephrol Dial Transplant*. 2016;31:1182–1189. doi: 10.1093/ndt/gfv338
 34. Lawton JS, Tamis-Holland JE, Bangalore S, Bates ER, Beckie TM, Bischoff JM, Bittl JA, Cohen MG, DiMaio JM, Don CW, et al; Writing Committee Members. 2021 ACC/AHA/SCAI guideline for coronary artery revascularization: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2022;79:e21–e129. doi: 10.1016/j.jacc.2021.09.006
 35. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, Deswal A, Drazner MH, Dunlay SM, Evers LR, et al. 2022 AHA/ACC/HFSA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2022;79:e263–e421. doi: 10.1016/j.jacc.2021.12.012
 36. Xu T, Lin M, Shen X, Wang M, Zhang W, Zhao L, Li D, Luan Y, Zhang W. Association of the classification and severity of heart failure with the incidence of contrast-induced acute kidney injury. *Sci Rep*. 2021;11:15348. doi: 10.1038/s41598-021-94910-1
 37. Kooiman J, Seth M, Share D, Dixon S, Gurm HS. The association between contrast dose and renal complications post PCI across the continuum of procedural estimated risk. *PLoS One*. 2014;9:e90233. doi: 10.1371/journal.pone.0090233
 38. Fu EL, Coresh J, Grams ME, Clase CM, Elinder CG, Paik J, Ramspek CL, Inker LA, Levey AS, Dekker FW, et al. Removing race from the CKD-EPI equation and its impact on prognosis in a predominantly White European population. *Nephrol Dial Transplant*. 2023;38:119–128. doi: 10.1093/ndt/gfac197

Circulation: Heart Failure

FIRST PROOF ONLY