

Cardiovascular toxicities associated with novel cellular immune therapies

Malak Munir,¹ Ahmed Sayed,^{1,2} Daniel Addison,^{3,4,*} and Narendranath Epperla^{5,*}

¹Department of Medicine, Ain Shams University Faculty of Medicine, Cairo, Egypt; ²Department of Cardiology, Houston Methodist DeBakey Heart & Vascular Center, Houston, TX; ³Division of Cancer Prevention and Control, Department of Internal Medicine, College of Medicine and ⁴Division of Epidemiology, College of Public Health, The Ohio State University, Columbus, OH; and ⁵Division of Hematology and Hematologic Malignancies, Huntsman Cancer Institute, The University of Utah, Salt Lake City, UT

Over the past decade, T-cell-directed therapies, including chimeric antigen receptor T-cell (CAR-T) and bispecific T-cell engager (BTE) therapies, have reshaped the treatment of an expanding number of hematologic malignancies, whereas tumor-infiltrating lymphocytes, a recently approved cellular therapy, targets solid tumor malignancies. Emerging data suggest that these therapies may be associated with a high incidence of serious cardiovascular toxicities, including atrial fibrillation, heart failure, ventricular arrhythmias, and other cardiovascular toxicities. The development of these events is a major limitation to long-term survival after these treatments. This review examines the current state of evidence, including reported incidence rates, risk factors, mechanisms, and management strategies of cardiovascular toxicities after treatment with these novel therapies. We specifically focus on CAR-T and BTE therapies and their relation to arrhythmia, heart failure, myocarditis, bleeding, and other major cardiovascular events. Beyond the relationship between cytokine release syndrome and cardiotoxicity, we describe other potential mechanisms and highlight key unanswered questions and future directions of research.

Introduction

Over the last decade, chimeric antigen receptor T-cell (CAR-T) therapy and bispecific T-cell engagers (BTEs) have revolutionized the treatment of hematologic malignancies.¹⁻¹⁰ Commercially available CAR-Ts are ex vivo modified T cells that are engineered to target a specific tumor antigen such as CD19 in B-cell malignancies or B-cell maturation antigen in multiple myeloma.¹¹ BTE cell therapy is a shorter acting T-cell therapy in which a dual-target antibody is designed to bind on 1 end to the T-cell receptor component CD3 and on the other to tumor antigens.¹² Both CAR-T and BTE therapies are primarily used to treat hematologic malignancies, however, several ongoing CAR-T products are being investigated in early phase clinical trials for the treatment of solid tumors. Although their mechanisms of action differ, T-cell-based therapies all serve 1 goal, to facilitate T-cell attack on and eradicate tumor cells.

There are currently ~400 registered trials actively recruiting for T-cell-based therapy protocols in the United States alone.¹³ Understanding their adverse event (AE) profile becomes crucial because these novel cellular therapies expand into widespread use for a growing range of indications. In particular,

Submitted 6 June 2024; accepted 13 October 2024; prepublished online on *Blood Advances* First Edition 17 October 2024; final version published online 13 December 2024. <https://doi.org/10.1182/bloodadvances.2024013849>.

*D.A. and N.E. contributed equally to this study.

Data reported was obtained from published trials and US Food and Drug Administration drug inserts.

© 2024 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](#), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

cardiovascular AEs (CVAEs) have emerged as potentially important drivers of morbidity and mortality after T-cell–based therapy.^{14,15}

Although treatment-related cardiotoxicity is a recognized driver of morbidity and mortality in many cancer types,¹⁶⁻¹⁹ several considerations make T-cell–based therapies of special importance. First, these therapies are increasingly being administered to older patients who may have preexisting cardiovascular disease risk factors, such as hypertension, coronary artery disease, diabetes, or a combination thereof, increasing their susceptibility to cardiotoxicity.²⁰⁻²³ Second, they are often used in the relapsed/refractory setting, in which patients may have already received anthracyclines or other cardiotoxic chemotherapies that compound cardiac risk.²⁴ Finally, lymphodepletion regimens required before CAR-T therapy often incorporate agents that have known cardiotoxic potential.²⁵

To establish best practices regarding cardiovascular care for patients receiving novel cellular immunotherapies, we aim to comprehensively review the existing evidence on cardiotoxicity associated with CAR-T and BTE treatment(s). Our goal is to synthesize current knowledge and provide clinically relevant guidance surrounding the cardiovascular safety of these emerging therapies.

Possible mechanisms for cellular therapy-associated cardiotoxicities

Although the exact mechanisms underlying the cardiotoxic effects of T-cell therapies remain poorly defined, theories have coalesced around 3 main pathways (Figure 1).^{26,27} The first proposes that cardiotoxicity arises indirectly through on-target inflammatory effects on the tumor microenvironment. These effects trigger the systemic release of proinflammatory cytokines, leading to cytokine release syndrome (CRS) and subsequent cardiac dysfunction. The second theory suggests T cells directly engage antigens shared between tumor and cardiac tissue, eliciting targeted toxicity. Finally, some evidence indicates T cells may also attack the heart via off-target interactions with myocardial antigens unrelated to tumor tissue. It is important to note that most mechanistic data described so far focus specifically on CAR-T therapy. Comparatively, less is known about the mechanisms of cardiotoxicity associated with BTE.

Role of CRS in cardiotoxicity susceptibility

CRS has traditionally been considered the key driver of cardiovascular toxicity associated with cellular therapies. Characterized by fever and hypotension, it is a common side effect of CAR-T and BTE therapies.^{28,29} A substantial proportion of patients treated with CAR-T in the landmark clinical trials developed CRS, and the incidence varies based on the type of CAR-T construct (CD28 vs 4-1-BB costimulatory molecules), as well as the underlying disease and tumor burden. The rate of CRS ranges from 77% to 92% in acute leukemia trials,^{1,30} 39% to 94% in B-cell lymphoma trials^{3,31-38} and up to 95% in multiple myeloma trials^{4,5,39,40} (Table 1). Notably, recent trials have reported up to a 76% incidence of CRS even with non-CD28 constructs,^{5,41,42} suggesting insidious associated events such as arrhythmia may still be not well defined.

On the contrary, a lower percentage of BTE patients experienced CRS, with reported rates of up to 16% in B-cell acute lymphoblastic leukemia (B-ALL),⁶ ~70% in multiple myeloma⁷ and ranging from 44% to ~70% in B-cell lymphoma⁸⁻¹⁰; additionally, when CRS occurs with BTE therapy, it is less severe.⁶²

CVAEs have been reported across all grades of CRS, with higher grades of CRS being associated with tumor burden as well as other risk factors such as early-onset CRS, infusion dose, as well as the addition of fludarabine to the lymphodepletion regimen.^{63,64} Occasionally, CVAEs (eg, heart failure and arrhythmias) may occur in the absence of CRS.^{14,23} Although the grading of CRS may differ among trials, the 2 most widely used criteria are those provided by the American Society for Transplantation and Cellular Therapy⁶⁵ and the Common Terminology Criteria for Adverse Events.⁶⁶ Although they may agree on the major criteria for each CRS grade, the American Society for Transplantation and Cellular Therapy system provides more detailed specifications regarding the management of hypotension and hypoxia.

The mechanism underlying CRS involves the release of proinflammatory cytokines including interleukin-6 (IL-6), IL-1, tumor necrosis factor α , and nitric oxide.^{67,68} Although cytokine release was initially thought to arise from activated T cells, a murine model of CAR-T showed that the severity of CRS was mediated by cytokines produced by macrophage and monocyte activation, perpetuating a vicious cycle of inflammation.^{69,70} The downstream signals of these cytokines enable the trafficking of immune cells out of circulation, resulting in vascular leakage and hypotension.⁷¹ Meanwhile, tumor necrosis factor α signaling induces endothelium to express tissue factors, contributing to the prothrombotic events in severe CRS.⁶⁸ Additionally, high levels of von Willebrand factor and angiopoietin-2 have been linked to severe states of CRS in patients receiving CAR-T therapy.^{63,72} These molecules may lead to endothelial activation and the activation of coagulation cascades, contributing to the development of coagulopathy.⁷³

Preclinical studies have identified IL-6 and IL-1 as the major mediators of CRS,^{69,70} a finding further corroborated by clinical studies showing an association between CRS severity and levels of IL-6,⁷⁴ leading to the US Food and Drug Administration (FDA) approval of the IL-6 inhibitor tocilizumab for the treatment of severe CRS in 2017.⁷⁵ Early results from clinical studies with IL-1 inhibitor anakinra have demonstrated the promising activity of IL-1 inhibitors in treating CRS.⁷⁶ Currently, anakinra is being investigated for its potential in both preventing and managing CRS associated with T-cell–based therapies.⁷⁷⁻⁷⁹

From a cardiovascular perspective, the downstream effects of CRS may explain many of the clinically observed CVAEs, including hypotension, arrhythmias, left ventricular dysfunction, decompensated heart failure, and shock.⁸⁰ Across CAR-T trials, the median onset of CRS has preceded cardiovascular events, with cardiac toxicity commonly occurring within the CRS interval.⁸¹ This association is further supported by real-world studies,^{14,82-84} with 1 pharmacovigilance study reporting an almost twofold risk of CVAEs in patients with CRS.⁸⁴ Additionally, more severe cardiac events such as fluid-refractory hypotension, myocardial ischemia, as evidenced by troponin elevation, stress cardiomyopathy, pericarditis, and pericardial effusion are associated with higher-grade CRS.^{14,22,82,83,85-88}

It has been hypothesized that the mechanism of CRS-mediated cardiotoxicity mirrors that of systemic inflammatory response syndrome.^{28,64} Specifically, the aforementioned proinflammatory cytokine release leads to vascular and cellular disruption due to oxidative stress and altered calcium signaling, which ultimately leads to impaired endothelial integrity,⁶⁴ all of which contribute to acute hemodynamic changes which may trigger the CVAEs




Figure 1. Cardiovascular toxicity associated with cellular therapies. (A) Current literature has identified several cardiotoxic side effects of CAR-Ts, including arrhythmia, coagulopathy, heart failure, pericardial disease, and hypotension/shock. Less is known about the cardiotoxicity of BTEs, with only coagulopathy myocarditis, and hypotension/shock having been associated with therapy. Finally, there has been no definitive association between cardiotoxicity and TILs, given their recent introduction. (B) The mechanism of cardiotoxicity associated with cellular therapies can be divided into 3 categories: CRS-mediated cardiovascular dysfunction, direct cardiotoxicity, and indirect cardiotoxicity. Activated CAR-Ts or BTEs release cytokines, including granulocyte-macrophage colony-stimulating factor (GM-CSF), tumor necrosis factor (TNF), and interferon gamma (IFN- γ), which lead to macrophage activation, resulting in a proinflammatory cascade that causes oxidative stress, altered calcium cycling, and endothelial dysfunction, ultimately resulting in cardiovascular dysfunction. Indirect cardiotoxicity has been documented with CAR-Ts, in which engineered cells attack the sarcomeric protein titin as a crossantigen. Direct cardiotoxicity has been proposed as an alternative pathway in which CAR-Ts directly attack the myocardium. NO, nitric oxide.

Table 1. CVAEs reported in landmark clinical trials evaluating cellular therapies

Study	Trial name	Study phase	Drug class	Drug name	Indication	Intervention arm sample size	Median follow-up duration, mo*	Rate of CRS†	Rate of treatment-emergent CVAEs			
									Arrhythmia‡	Hypertension	Coagulopathy§	Others
Maude et al ^{1,43}	ELIANA	Phase 1/2	CAR-T	Tisagenlecleucel	R/R preB-ALL	79	13.1	61 (77%)	2 (4%)	145 (19%)	5 (6%)	Cardiac failure, 10 (9%)
Schuster et al ^{32,43}	JULIET	Phase 2	CAR-T	Tisagenlecleucel	R/R DLBCL	115	28.6	85 (74%)	12 (10%)	5 (4%)	5 (4%)	Cardiac failure, 1 (1%)
Fowler et al ^{35,43}	ELARA	Phase 2	CAR-T	Tisagenlecleucel	R/R FL	97	16.6	51 (53%)	4 (4%)	5 (5%)	2 (2%)	NA
Neelapu et al ^{31,44}	ZUMA-1	Phase 1/2	CAR-T	Axicabtagene ciloleucel	R/R LBCL	108	8.7	101 (94%)	25 (23%)	16 (15%)	2 (2%)	Cardiac failure, 6 (6%) Cardiac arrest, 4 (4%)
Jacobson et al ^{36,44}	ZUMA-5	Phase 2	CAR-T	Axicabtagene ciloleucel	R/R iNHL	146	17.5	123 (84%)	31 (21%)	19 (13%)	9 (6%)	Cardiac failure, 3 (2%)
Locke et al ^{44,45}	ZUMA-7	Phase 3	CAR-T	Axicabtagene ciloleucel	R/R LBCL	168	24.9	155 (92%)	24 (14%)	15 (9%)	15 (9%)	Cardiac failure, 2 (1%)
Wang et al ^{33,46}	ZUMA-2	Phase 2	CAR-T	Brexucabtagene autoleucel	R/R MCL	82	12.3	75 (91%)	8 (10%)¶	15 (18%)	8 (10%)	NA
Shah et al ^{30,46}	ZUMA-3	Phase 1/2	CAR-T	Brexucabtagene autoleucel	R/R preB-ALL	78	16.4	72 (92%)	12 (15%)	10 (13%)	13 (17%)	Cardiac failure, 3 (4%)
Kamdar et al ^{3,47}	TRANSFORM	Phase 2	CAR-T	Lisocabtagene maraleucel	R/R LBCL	89	6.2	44 (49%)	13 (15%)#	6 (7%)	NA	NA
Sehgal et al ^{38,47}	PILOT	Phase 2	CAR-T	Lisocabtagene maraleucel	R/R LBCL	61	12.3	24 (39%)	6 (10%)#	6 (10%)	NA	NA
Abramson et al ^{34,47}	TRANSCEND	Phase 1	CAR-T	Lisocabtagene maraleucel	R/R LBCL	268	18.8	123 (46%)	16 (6%)	38 (14%)	NA	Cardiomyopathy 4 (1.5%)
Siddiqi et al ⁴⁸	TRANSCEND CLL	Phase 1/2	CAR-T	Lisocabtagene maraleucel	R/RCLL	117	21.1	99 (85%)	NA	NA	NA	NA
Munshi et al ^{4,49}	KarMMa	Phase 2	CAR-T	Idecabtagene vicleucel	R/R MM	127	13.3	108 (85%)	6 (4.7%)	14 (11%)	11 (9%)	Cardiomyopathy 2 (1.6%)
Rodriguez-Otero et al ^{49,50}	KarMMa-3	Phase 3	CAR-T	Idecabtagene vicleucel	R/R MM	222	18.8	202 (91%)	16 (7%)	31 (14%)	31 (14%)	NA
Berdeja et al ^{39,51}	CARTITUDE-1	Phase 1/2	CAR-T	Ciltacabtagene autoleucel	R/R MM	97	12.4	92 (95%)	8 (8%)	18 (19%)	21 (22%)	NA
Cohen et al ⁴⁰	CARTITUDE-2	Phase 2	CAR-T	Ciltacabtagene autoleucel	R/R MM	20	11.3	12 (60%)	NA	NA	NA	NA
San-Miguel et al ^{5,51}	CARTITUDE-4	Phase 3	CAR-T	Ciltacabtagene autoleucel	R/R MM	188	15.9	147 (78%)	6 (3%)	13 (7%)	9 (5%)	NA

Reported AEs are treatment emergent, unless otherwise specified. Absolute numbers are derived from percentages and may not represent actual counts.

AMI, acute myocardial infarction; aPTT, activated partial thromboplastin time; iNHL, indolent non-Hodgkin lymphoma; INR, international normalized ratio; MRD, minimal residual disease; NA, not available; PT, prothrombin time; R/R CLL, relapse/refractory chronic lymphocytic lymphoma; R/R DLBCL, relapse/refractory diffuse large B-cell lymphoma; R/R FL, relapse/refractory follicular lymphoma; R/R LBCL, relapse/refractory large B-cell lymphoma; R/R MCL, relapse/refractory mantle cell lymphoma; R/R MM, relapse/refractory multiple myeloma; R/R preB-ALL, relapse/refractory precursor B-cell acute lymphoblastic leukemia.

*When applicable, median follow-up for overall survival is reported.

†Any-grade CRS.

‡When applicable, any of the following are reported as mention: atrial fibrillation, atrial flutter, atrial tachycardia, bradycardia, cardiac arrest, electrocardiogram QT prolonged, electrocardiogram T wave inversion, pulseless electrical activity, sinus bradycardia, supraventricular tachycardia, and ventricular tachycardia. Note, due to the lack of consistency of reporting among trials, the rate reported may coincide with any combination of the aforementioned arrhythmias.

§Coagulopathy refers to reporting of any of the following events: disseminated intravascular coagulation, elevated PT/aPTT, elevated INR, and elevated D-dimer.

¶When applicable, the reported AEs are from the FDA drug insert because it provides more comprehensive data on the reported CVAEs.

¶¶Reported rate for both bradyarrhythmia and nonventricular arrhythmia.

#Reported as tachycardia includes atrial fibrillation, atrial flutter, supraventricular tachycardia, sinus tachycardia, and tachycardia.

**Reported as treatment-related AEs.

Table 1 (continued)

Study	Trial name	Study phase	Drug class	Drug name	Indication	Intervention arm sample size	Median follow-up duration, mo*	Rate of CRS†	Rate of treatment-emergent CVAEs			
									Arrhythmia‡	Hypertension	Coagulopathy§	Others
Kantarjian et al ^{6,52}	TOWER	Phase 3	BTE	Blinatumomab	R/R preB-ALL	267	11.7	38 (14.2%)	NA	17 (6.4%)	NA	AMI, 1 (0.4%) Cardiac arrest, 1 (0.4%) Cardiac failure, 1 (0.4%) Pericardial effusion, 1 (0.4%)
Locatelli et al ^{52,53}	RIALTO	Phase 3	BTE	Blinatumomab	R/R preB-ALL	110	17.4	18 (16.4%)**	NA	NA	NA	NA
Martinelli et al ⁵⁴	ALCANTARA	Phase 2	BTE	Blinatumomab	R/R preB-ALL	45	16.1	4 (9%)	NA	NA	NA	NA
Gökbuget et al ⁵⁵	BLAST	Phase 2	BTE	Blinatumomab	MRD positive preB-ALL	116	30	4 (3%)	NA	NA	NA	NA
Moreau et al ^{7,56}	MajesTEC-1	Phase 2	BTE	Teclistamab	R/R MM	165	14.1	119 (72%)	26 (16%)	20 (12%)	NA	NA
Dickinson et al ^{8,57}	NP30179	Phase 2	BTE	Glofitamab	R/R DLBCL	154	12.6	102 (66%)	NA	NA	NA	NA
Budde et al ^{9,58}	GO29781	Phase 1	BTE	Mosunetuzumab	R/R FL	90	12.6	40 (44%)	NA	NA	NA	NA
Thieblemont et al ^{10,59}	EPCORE NHL-1	Phase 1/2	BTE	Epcoritamab	R/R LBCL	157	10.7	80 (51%)	16 (10%)	NA	NA	AMI, 1 (0.6%)
Chesney et al ^{60,61}	C-144-01	Phase 2	TIL	Lifileucel	Unresectable/metastatic melanoma	156	27.6	5 (3.2%)	74 (47.4%)#	21 (13.5%)	NA	Hypotension, 52 (33.3%)

Reported AEs are treatment emergent, unless otherwise specified. Absolute numbers are derived from percentages and may not represent actual counts.

AMI, acute myocardial infarction; aPTT, activated partial thromboplastin time; iNHL, indolent non-Hodgkin lymphoma; INR, international normalized ratio; MRD, minimal residual disease; NA, not available; PT, prothrombin time; R/R CLL, relapse/refractory chronic lymphocytic lymphoma; R/R DLBCL, relapse/refractory diffuse large B-cell lymphoma; R/R FL, relapse/refractory follicular lymphoma; R/R LBCL, relapse/refractory large B-cell lymphoma; R/R MCL, relapse/refractory mantle cell lymphoma; R/R MM, relapse/refractory multiple myeloma; R/R preB-ALL, relapse/refractory precursor B-cell acute lymphoblastic leukemia.

*When applicable, median follow-up for overall survival is reported.

†Any-grade CRS.

‡When applicable, any of the following are reported as mention: atrial fibrillation, atrial flutter, atrial tachycardia, bradycardia, cardiac arrest, electrocardiogram QT prolonged, electrocardiogram T wave inversion, pulseless electrical activity, sinus bradycardia, supraventricular tachycardia, and ventricular tachycardia. Note, due to the lack of consistency of reporting among trials, the rate reported may coincide with any combination of the aforementioned arrhythmias.

§Coagulopathy refers to reporting of any of the following events: disseminated intravascular coagulation, elevated PT/aPTT, elevated INR, and elevated D-dimer.

|| When applicable, the reported AEs are from the FDA drug insert because it provides more comprehensive data on the reported CVAEs.

¶Reported rate for both bradyarrhythmia and nonventricular arrhythmia.

#Reported as tachycardia includes atrial fibrillation, atrial flutter, supraventricular tachycardia, sinus tachycardia, and tachycardia.

**Reported as treatment-related AEs.

mentioned above.⁸⁹ Nonetheless, more research is still needed to fully explain the precise mechanisms connecting CRS to cardiotoxicity.

Non-CRS pathways of cardiotoxicity development

Other proposed hypotheses for CAR-T-mediated cardiotoxicity include direct myocardial damage from recognition of cardiac antigens, either due to antigen mimicry or alloreactivity. Fatal myocarditis in 2 patients has been reported, 1 with melanoma and the other with multiple myeloma, receiving CAR-T-cells targeting the MAGE-A3 tumor antigen, due to cross-reactivity with titin, a sarcomere protein in cardiomyocytes.⁶⁴ Analogous to graft-versus-host disease, alloreactivity has also been linked to cardiotoxicity, in which administered T cells respond to unrecognized major histocompatibility complex variants in host cardiac tissue, leading to cardiac damage.²⁶

Cardiovascular toxicities observed in clinical practice

Evidence from randomized controlled trials reporting the cardiovascular toxicity of cellular therapies is limited. These trials frequently excluded patients with preexisting cardiovascular conditions, a cohort that is at a higher risk of cardiotoxicity.^{10,11} Additionally, most trials were not powered to detect less common AEs, such as CVAEs, because their primary focus was on detecting differences in cancer-related outcomes. However, observational studies have highlighted the prevalence of CVAEs in patients receiving CAR-T and BTE therapies.^{14,15,22,23,82,83,85-88,90-94}

The most frequently reported cardiovascular side effects associated with CRS are hypotension and arrhythmia occurring with both CAR-T and BTE therapies.^{27,81} Trials on both report hypertension in up to 19% of patients (Table 1); however, many of these trials are single-armed studies, thus limiting the ability to determine whether it was directly related to therapy.

CAR-T therapy

In the past decade, 6 CAR-T therapies have received FDA approval for various hematologic malignancies in the relapsed or refractory setting, including precursor B-ALL, non-Hodgkin lymphomas, chronic lymphocytic leukemia, and multiple myeloma.^{1,3-5,30,33-40,80,95} In the major clinical trials, refractory shock, arrhythmias, cardiomyopathy, and heart failure were the reported cardiovascular side effects (Table 1).

Similarly, in practice, arrhythmias are the most commonly reported cardiovascular toxicities after CAR-T therapy, although the reported incidence in studies is inconsistent (Table 2). A comprehensive pharmacovigilance study of 2657 patients receiving CAR-T therapy found that CAR-T-related AEs were almost 3 times as likely to involve tachyarrhythmias.²² The most frequently reported arrhythmias included sinus tachycardia, atrial fibrillation, ventricular arrhythmia, and corrected QT interval prolongation, possibly corresponding to the inflammatory state and hemodynamic changes of CRS.²²

Other cardiovascular side effects reported with CAR-T therapy include cardiomyopathy, heart failure, pericardial disease, and coagulopathy. In a retrospective study, new or worsening cardiomyopathy was detected in 10.3% of 116 patients within months of

treatment.²³ In the aforementioned pharmacovigilance study, patients receiving CAR-T were 3.5 times as likely to report cardiomyopathy and twice as likely to report pericardial disease compared with non-CAR-T therapies.²² Moreover, coagulopathy, defined as an elevation in laboratory coagulation parameters and/or disseminated intravascular coagulopathy (DIC), was also reported in the major clinical trials,^{1,31,35-37} particularly with brexucabtagene autoleucel^{30,33} (Table 1); however, it is unclear whether it was directly treatment-related because these trials were single-armed and lacked a control group. Moreover, this finding is further supported by retrospective studies. In a study of patients with B-cell lymphomas, a coagulopathy rate of 13.3% was reported, with only 1 patient presenting with DIC.⁸⁶ Another study found an association between the occurrence of coagulation disorders and CRS, reporting a coagulopathy rate of 51%, with 7 patients developing DIC.⁹⁶

Hypotension, shock, and congestive heart failure have also been noted in major clinical trials and real-world studies (Tables 1 and 2). Hypotension and shock were generally attributed to severe CRS, whereas congestive heart failure was reported in only 1 patient at most in these trials (Table 1). Postmarketing studies have provided further insight into the incidence of heart failure with CAR-T therapy, reporting incidence rates that range from 1% to 15% (Table 2); however, due to the retrospective nature of these studies and heterogeneous patient populations, the exact incidence of these CVAEs remains uncertain. Similarly, despite considerations for potentially lower toxicity (eg, outpatient) in non-CD28-focused products, CRS and cardiac events have still been reported,^{5,41,42} and the long-term risk of more insidious or later CVAE events, such as arrhythmias and heart failure, remains undefined.

Additionally, studies reporting on major adverse cardiac event (MACE) rates after CAR-T therapy show inconsistent results. MACE is a composite outcome generally defined as the occurrence of ≥ 1 of the following: cardiovascular death, heart failure with reduced ejection fraction, acute coronary syndrome, ischemic stroke, or de novo (incident) symptomatic cardiac arrhythmia. Although the exact components included in MACE may vary between studies, it consistently includes at least 1 of these cardiovascular events or their variations.

A recent prospective study reported no MACE events in 137 patients treated with CAR-T cells.⁹⁴ Conversely, 1 retrospective study found a 21% cumulative incidence of MACE within 1 year after infusion,⁸³ whereas a more recent prospective study of 44 relatively healthier patients reported a MACE rate of $<5\%$.⁹³ Despite differing populations and study designs, the discrepancies reported emphasize the need for better characterization of the cardiotoxic profile of CAR-T therapy and the need to unify and standardize reporting outcomes.

A notable finding from the aforementioned pharmacovigilance study was the differing cardiotoxic profiles observed between axicabtagene ciloleucel and tisagenlecleucel, with tisagenlecleucel having a slightly more favorable cardiac safety profile and lower reporting rates of CRS, arrhythmias, and other severe CVAEs²²; this distinction was also reported in another pharmacovigilance study by Guha et al.⁸⁴ However, given the inherent limitations of retrospective studies, this observation serves merely as a signal that warrants further investigation into the potential reasons for this difference. In particular, this discrepancy raises questions regarding whether differences in CAR-T costimulatory domains,

Table 2. CVAEs reported in cohort and pharmacovigilance studies evaluating cellular therapies

Study	Study type	Drug class	Sample size	Median follow-up duration, mo	Rate of CRS*	Reported rates of CVAEs									
						Arrhythmia	Cardiac decompensation	Left ventricular dysfunction	Cardiac ischemia†	Cardiomyopathy	Cardiac death	Hypertension	Hypotension/shock	Coagulopathy	Others
Burstein et al ^{§1}	Retrospective	CAR-T	93	6‡	24 (25.8%)	NA	NA	10 (10%)	NA	NA	NA	NA	24 (25%)	NA	NA
Alvi et al ^{§2}	Retrospective	CAR-T	137	10	81 (59%)	5 (3.6%)	6 (4.4%)	NA	NA	NA	6 (4.4%)	NA	NA	NA	NA
Lefebvre et al ^{§3}	Retrospective	CAR-T	145	15	104 (72%)	11 (7.5%)	22 (15%)	NA	2 (1.4%)	NA	2 (1.4%)	NA	NA	NA	NA
Shalabi et al ^{§5}	Retrospective	CAR-T	52	NA	37 (71%)	NA	NA	6 (12%)	NA	NA	NA	NA	9 (17%)	NA	NA
Ganatra et al ^{§3}	Retrospective	CAR-T	187	5.5	155 (83%)	13 (7.0%)	NA	NA	NA	12 (10.3%)§	NA	NA	NA	NA	NA
Brammer et al ^{§7}	Retrospective	CAR-T	90	16.5	80 (88.9%)	11 (12.2%)	1 (1.1%)	NA	NA	NA	NA	NA	NA	NA	3 (3.3%); other myocarditis, 2 (2.2%)
Qi et al ^{§8}	Retrospective	CAR-T	126	1.6‡	103 (81.7%)	7 (6%)	15 (12%)	1 (1%)	9 (7%)	NA	NA	NA	NA	NA	New valve disease, 1 (1%)
Goldman et al ^{§2}	Pharmacovigilance	CAR-T	2657	NA	1457 (54.84%)	74 (2.79%)	NA	NA	12 (0.45%)	69 (2.6%)	NA	18 (0.68%)	286 (10.76%)	NA	Cardiogenic shock, 49 (1.84%); pericardial disease, 11 (0.41%); myocarditis, 2 (0.08%)
Korell et al ^{§4}	Prospective	CAR-T	137	8.8‡	75 (54.7%)	5 (4%)	17 (12%)	4 (3%)	NA	NA	NA	2 (1%)	25 (18%)	NA	NA
Wudhikarn et al ^{§5}	Retrospective	CAR-T	60	9	NA	8 (13.3%)	NA	3 (5%)	NA	NA	NA	5 (8.3%)	28 (46.6%)	8 (13.3%)	Pericardial effusion, 1 (1.67%)
Lefebvre et al ^{§3}	Prospective	CAR-T	44	16	23 (52%)	1 (1.67%)	1 (1.67%)	NA	NA	NA	NA	NA	NA	NA	NA
Jung et al ^{§2}	Retrospective	BTE	50	NA	10 (20%)	NA	NA	NA	NA	NA	NA	NA	NA	NA	7 (14%)¶
Sayed et al ^{§5}	Pharmacovigilance	BTE	3668	NA	NA	45 (1.22%)	52 (1.4%)	NA	NA	NA	NA	NA	193 (5.26%)	NA#	Myocarditis, 8 (0.22%); pericarditis, 4 (0.1%)
Fradley et al ^{§0}	Retrospective	TIL	43	NA	NA	6 (14%)	NA	NA	1 (2.3%)	NA	NA	NA	14 (32.6%)	NA	NA

ACS, acute coronary syndrome; MI, myocardial infarction; NA, not available.

*Includes any-grade CRS.

†Includes events reported as ACS or MI.

‡Reported as overall follow-up duration.

§Percentage out of 116 patients with available imaging.

||Reported rates from pharmacovigilance studies are the proportions representing the reporting of a particular AE of all drug-related AE reports.

¶Includes “cardiovascular disease/events” not otherwise specified.

#Increased reporting odds of DIC events was found to be statistically significant; however, the proportion of reported events was not stated.

patient characteristics, or dosing strategies could contribute to the onset of CVAEs; all of which need to be addressed in a prospective manner.

BTE therapy

To date, 7 BTE therapies have been FDA approved for the treatment of B-ALL, non-Hodgkin lymphomas, and multiple myeloma in the relapsed/refractory setting.^{6-10,54,97} However, the clinical trials for BTE therapies are mainly small-scale phase 2 studies, often enrolling <100 patients; consequently, they were underpowered to detect cardiotoxicity.

Arrhythmias were the most frequently reported CVAEs in major BTE trials.^{27,81} Trials for the relatively newer BTE therapies such as talquetamab,⁹⁸ elranatamab,⁹⁹ mosunetuzumab,⁹ epcoritamab,¹⁰ and glofitamab⁸ did not report specific CVAEs.

Despite the limitations of clinical trials, real-world data provide some insight into the cardiotoxicity associated with BTE therapy, although it remains somewhat inconsistent. In a large-scale pharmacovigilance study, CVAEs were reported in 1 of every 5 AEs related to BTE therapies.¹⁵ Of note, teclistamab, but not other BTE products, was associated with an excess of fatal cardiovascular events and myocarditis. Additionally, DIC was almost 3 times as likely to be reported with BTEs, a finding that was likely driven by blinatumomab. Interestingly, this study found no association between CVAEs and CRS; considering the lower incidence and severity of CRS observed with BTE therapy, alternative mechanisms may be contributing to cardiotoxicity. However, 1 limitation of this study was that it reported data only on blinatumomab and teclistamab, because few data were available on the more recently approved BTE therapies at the time.

Risk factors for cellular therapy-associated cardiotoxicities

Several key factors may increase the risk of cardiotoxicity, including advanced age, presence of baseline comorbidities such as uncontrolled diabetes, renal dysfunction, coronary artery disease, and heart failure, and prior receipt of cardiotoxic chemotherapy regimens such as anthracyclines or mediastinal radiotherapy.^{23,24} Thus, a thorough pretreatment assessment of cardiovascular health and risk stratification is essential for identifying patients requiring more intensive monitoring (Figure 2).

Prognostic implications of cellular therapy-associated cardiotoxicity

The long-term risks of sustained, cardiotoxicity-related morbidity and mortality associated with T-cell treatments remain unclear due to the lack of extended follow-up data. Current long-term studies have limited follow-up durations, ranging from 9 months¹⁴ to 16.5 months.⁸⁷ In studies with relatively longer follow-up (5-16.5 months),^{14,23,83,87} the majority of CVAEs were reported to occur within 3 months of CAR-T initiation, with a median time to cardiac event of ~1 to 2 weeks; however, the long-term impact of these events remains unclear. In a retrospective study of 202 CAR-T patients, 16% experienced severe cardiovascular events including myocardial infarction, cardiovascular shock, or heart failure.¹⁴ Patients with these severe CVAEs had a risk of mortality that was nearly 3 times that of those without severe CVAEs. Similarly, a pharmacovigilance study on BTE therapy shows

that when they do occur, CVAEs associated with BTE therapy are more fatal than other AEs reported with BTE therapy.¹⁵ Reassuringly, there is some evidence indicating that cardiotoxicity is often reversible without lasting impact^{23,91,100}; however, this evidence is inconclusive. This lack of long-term studies represents a gap in the literature that warrants investigation as we continue to develop and expand the use of cellular therapies.

Diagnostic algorithm for cellular therapy-associated cardiotoxicities

Pretreatment evaluation

Comprehensive baseline cardiac screening is recommended for all patients before initiating T-cell therapy (Figure 2). This includes cardiac biomarkers such as cardiac troponin (cTn) and natriuretic peptides (NPs); including brain natriuretic peptide and NT-pro-brain natriuretic peptide) to serve as a baseline for the early detection of infarction or heart failure. A 12-lead electrocardiography (ECG) is also recommended to allow for the evaluation of new-onset ischemia and arrhythmia. Additionally, transthoracic echocardiography with global longitudinal strain to assess baseline ventricular function and allow for risk stratification may be performed. In potentially high-risk patients, such as those with borderline reduced left ventricular ejection fraction (eg, 45%-50%), intracardiac malignant disease involvement, or even with inadequate transthoracic echocardiography image quality (Figure 2), more advanced imaging such as cardiac magnetic resonance imaging may be warranted to define and/or exclude underlying cardiac dysfunction; whereas cardiac stress testing may be most appropriate in those with potential coronary ischemia.^{101,102} Together, these may provide additional key insights into the ability to tolerate hemodynamic changes during therapy.^{81,100}

Posttreatment evaluation

Ongoing postinfusion surveillance may incorporate serial ECGs, repeat echocardiography for any new symptoms, and frequent assessment of cTn and NPs for myocardial injury/stress (Figure 2). Subject to their availability, tracking inflammatory markers, such as IL-6 and IL-1, may also provide early warning signals for cardiotoxicity.^{81,103} If echocardiography is inadequate or inconclusive, cardiac magnetic resonance imaging may be considered. Although evidence on predictive inflammatory/biomarker cutoffs remains limited, their elevations have been associated with increased cardiovascular events. For example, studies have identified associations between cardiovascular events and elevated IL-6, as well as ferritin and troponin.^{14,23,63,82,83,91} In a recent prospective study on patients receiving CAR-T, an association between high-sensitivity troponin and all-cause mortality was reported, suggesting that patients with elevated high-sensitivity troponin may benefit from more intensive postinfusion monitoring and early intervention.⁹⁴ However, further research is still needed to establish definitive biomarker thresholds for the prediction of cardiotoxicity.

Prevention and management of cellular therapy-associated cardiotoxicities

Management of CRS

Due to the close association between CRS and cardiotoxic symptoms, most recommendations aim at the prevention or early management of CRS (Figure 3). Supportive care and antipyretics




Figure 2. Identification and management of patients at high-risk of cardiotoxicity. Clinical considerations for the screening and management of patients at an increased risk of cardiotoxicity undergoing T-cell-directed therapy. CKD, chronic kidney disease; CMR, cardiac magnetic resonance imaging; ECG, electrocardiography; ECOG, Eastern Cooperative Oncology Group; ESRD, end stage renal disease; GLS, global longitudinal strain; INR, international normalized ratio; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association.

suffice for low-grade CRS, whereas severe CRS warrants tocilizumab, corticosteroids, and hemodynamic support.⁸⁰ Recently, a more potent IL-6 inhibitor, siltuximab, has shown promise for CRS in trials but requires further study.¹⁰⁴

Early intervention with preventive drugs in patients developing early signs of CRS has been suggested as a potential strategy to

mitigate severe CRS. In ZUMA-1, early intervention with tocilizumab and corticosteroids led to an absolute risk reduction of 11% in the rate of severe CRS (2% in the group with early intervention vs 13% in the group without early intervention).¹⁰⁵ Consistently, early preventive intervention has reduced CRS rates across various studies.^{106,107} However, despite these promising results, heterogeneity in management across centers and the lack of robust




Figure 3. Management of patients with CRS and/or cardiotoxicity. Proposed algorithm for the management of high-risk cardiac patients who develop CRS and/or cardiotoxicity during treatment with T-cell-directed therapy. ACLS, advanced cardiac life support; ECG, electrocardiography; GDMT, guideline-directed medical therapy.

evidence have limited the initiation of standard preventive recommendations on the use or timing of these drugs.

Moreover, the prophylactic use of tocilizumab on day 2 after infusion for the prevention of severe (grade ≥ 3) CRS was evaluated in subsequent analyses of the ZUMA-1 trial.^{2,31} Preliminary results showed an incidence of severe CRS of 13% in the cohort receiving no prophylaxis compared with 3% in the cohort receiving prophylactic tocilizumab. The same trial also evaluated the use of prophylactic corticosteroids (dexamethasone), demonstrating an incidence of severe CRS of 0% in the dexamethasone group compared with 13% in the group receiving no prophylaxis.¹⁰⁸ However, it is important to note that the cohort receiving corticosteroids had a lower tumor burden, which could have contributed to the overall lower incidence of CRS observed. Consequently, given the lack of robust data available, prophylactic management of CRS is not currently the standard of care and warrants further investigation in randomized controlled studies.

Management of cardiotoxicity

Beyond the early or prophylactic use of targeted therapies such as tocilizumab, prevention and management rely on retrospective data and traditional cardio-oncologic strategies. High-risk patients warrant monitoring for cardiotoxicity with telemetry, ECG, echocardiograms, cTn, and NP testing for the early detection of arrhythmias and circulatory collapse, as supported by the European Society of Cardiology cardio-oncology opinion guidelines.¹⁰³ Cardiovascular

complications should be managed by standard protocols. Additionally, permissive hypertension by withholding antihypertensive medication in high-risk patients has been proposed to address circulatory fluctuations,⁸¹ but limited evidence currently exists to support this recommendation.

Future directions

There is wider applicability of cellular therapies in hematologic and solid tumor cancers, and this will only continue to grow in the future. Although initial studies and pharmacovigilance data have revealed concerning safety signals, our current understanding of the cardiovascular risks and mechanisms of T-cell-based therapy is limited. Additionally, research on the cardiovascular effects of emerging T-cell therapies such as BTEs is scarce, with many gaps of knowledge in the available evidence (Table 3). With ongoing trials of BTEs in non-cancer-related indications, awareness and surveillance for potential cardiotoxicity in these wider patient populations is critical.

CAR-T therapy has the most well-documented evidence so far; yet, studies defining the underlying pathways, incidence rates, and risk factors for cardiovascular events are still insufficient. Current research on newer autologous CAR-T constructs, both in hematologic malignancies and solid tumors, as well as allogeneic and armored CAR-T approaches is promising, although it leaves us with the worry of potential cardiotoxicity. Although the promise of improved tumor specificity using armored CAR-Ts provides hope for possibly fewer CVAEs, the use of donor-derived cells in

Table 3. Key unanswered questions and future directions

Key unanswered questions	Future directions
Diagnosis and surveillance	
<ul style="list-style-type: none">• What is the optimal method of monitoring patients for the development of cardiotoxicity?• Can biomarkers be used for the early detection of cardiotoxicity?• What are the most effective biomarker cutoff values for the detection/monitoring of cardiotoxicity?	<ul style="list-style-type: none">• More support for large-scale post-marketing pharmacovigilance studies investigating the link between biomarker changes and the development of cardiovascular toxicity.• Large-scale RCTs comparing different surveillance strategies.
Management and prevention	
CRS <ul style="list-style-type: none">• Does prophylactic therapy against CRS improve overall survival?• Which drugs/drug combinations (tocilizumab, siltuximab, corticosteroids, and anakinra) are the most effective for the treatment/prevention of CRS?• When is the best timing to initiate prophylactic therapy against CRS? Non-CRS <ul style="list-style-type: none">• What is the role of contemporary cardioprotective therapies (eg, beta blockers, ACE inhibitor, and statins, etc) in the management and prevention of cardiotoxicity?• What is the ideal balance between maintaining adequate cancer therapy levels and cardiotoxicity, that is, permissive cardiotoxicity?	<ul style="list-style-type: none">• RCTs assessing the comparative efficacy of CRS prophylactic regimens.• Develop unified pretreatment risk assessment tools based on known risk factors and the integration of cardio-oncology services in high-risk patients.• RCTs assessing the comparative efficacy of cardioprotective therapies at preventing new-onset cardiac dysfunction.• RCTs comparing different levels of permissive cardiotoxicity with respect to eventual outcomes.
Risk factors	
<ul style="list-style-type: none">• What are the key risk factors for developing cardiotoxicity from cellular therapies?• What are the unique cardiotoxicity risks of individual cellular therapies?	<ul style="list-style-type: none">• Conduct studies to identify and validate risk factors for cardiotoxicity.
Long-term prognostic implications	
<ul style="list-style-type: none">• What are the long-term cardiovascular consequences of cellular therapies and how does this affect overall survival and patient quality of life?	<ul style="list-style-type: none">• Conduct long-term follow-up studies to assess the late-onset cardiovascular effects of cellular therapies.

RCT, randomized controlled trial.

allogeneic CAR-T regimens allowing for more widespread treatment is of concern. This broader reach, compounded with the unpredictable off-target risks that may occur, means the widespread use of allogeneic CAR-T therapy could magnify the CVAE signals that we are seeing now. Hence, there is a critical need to better characterize the cardiotoxicity profile of these emerging cellular therapies before large-scale adoption.

The expansion of adoptive cellular therapies to the treatment of solid tumor malignancies, as seen recently with the approval of tumor-infiltrating lymphocytes (TILs), brings forth the potential for cardiovascular toxicity. However, these risks are unclear, because several trials have failed to incorporate specific measures for detecting CVAEs in their design. Furthermore, recently reported studies with TILs for the treatment of melanoma^{60,109} have incidentally documented a severe risk of hypotension, possibly leading to cardiac ischemia. A retrospective study⁹⁰ has also identified the risk of CVAEs with TIL therapy.

Efforts to predict, identify, and prevent cardiovascular toxicity should remain a priority for studies on T-cell therapies. In line with this focus, future research should also explore the development of

targeted deterrence techniques, such as molecular safety switches and genetic modifications,¹¹⁰ aimed at preventing the onset of these AEs. However, it is important to note that these advancements are contingent upon establishing comprehensive cardiotoxic profiles for individual cellular therapies.

Multidisciplinary collaboration between cardiologists and oncologists and the strengthening of pharmacovigilance reporting are essential to fully evaluate cardiovascular outcomes. Additionally, standardized safety end points in trials along with registry data can help outline long-term cardiotoxicity profiles; thus, collaboration efforts among researchers, regulatory agencies, and physicians are paramount for maximizing the potential of these novel cellular therapies while ensuring patients safety.

Conclusion

Cellular therapies have revolutionized the treatment of hematologic malignancies and many solid tumors. Yet, cardiotoxicity remains a potential limitation to the effective application of these therapies. Understanding the effects of these rapidly proliferating therapies will facilitate the targeted prevention and control of these AEs. Multidisciplinary strategies inclusive of cardio-oncologic assessments and shared decision-making with patients, cardiologists, and oncologists are needed to facilitate increased applications of these impactful therapies. Despite gaps in current evidence, existing data may provide a basis for future investigations focused on defining the true incidence, effects, and burden of cardiotoxicity among cellular therapy-treated patients. Together, these provide compelling impetus for additional studies focused on defining the mechanisms and optimal preventive strategies for cardiac toxicities among cancers requiring these therapies.

Acknowledgments

This work was supported in part by National Institutes of Health (NIH) grant P50-CA140158. D.A. is supported by NIH grants K23-HL155890, R01HL168045, and R01HL170038, and an American Heart Association-Robert Wood Johnson Foundation (Harold Amos) grant.

The manuscript's content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Authorship

Contribution: N.E. and D.A. contributed to conceptualization and study design; M.M. and A.S. contributed to writing/editing the first draft of the manuscript; and N.E. and D.A. provided critical review and comments.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: M.M., [0000-0002-7306-8677](https://orcid.org/0000-0002-7306-8677); A.S., [0000-0003-0150-8917](https://orcid.org/0000-0003-0150-8917); D.A., [0000-0002-9113-8333](https://orcid.org/0000-0002-9113-8333); N.E., [0000-0002-8216-3457](https://orcid.org/0000-0002-8216-3457).

Correspondence: Narendranath Epperla, Division of Hematology and Hematologic Malignancies, Department of Medicine, Huntsman Cancer Institute, Salt Lake City, UT 84103; email: naren.epperla@hci.utah.edu.

References

1. Maude SL, Laetsch TW, Buechner J, et al. Tisagenlecleucel in children and young adults with B-cell lymphoblastic leukemia. *N Engl J Med*. 2018; 378(5):439-448.
2. Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1-2 trial. *Lancet Oncol*. 2019;20(1):31-42.
3. Kamdar M, Solomon SR, Arnason J, et al. Lisocabtagene maraleucel versus standard of care with salvage chemotherapy followed by autologous stem cell transplantation as second-line treatment in patients with relapsed or refractory large B-cell lymphoma (TRANSFORM): results from an interim analysis of an open-label, randomised, phase 3 trial. *Lancet*. 2022;399(10343):2294-2308.
4. Munshi NC, Anderson LD Jr, Shah N, et al. Idecabtagene vicleucel in relapsed and refractory multiple myeloma. *N Engl J Med*. 2021;384(8):705-716.
5. San-Miguel J, Dhakal B, Yong K, et al. Cilta-cel or standard care in lenalidomide-refractory multiple myeloma. *N Engl J Med*. 2023;389(4):335-347.
6. Kantarjian HM, Stein AS, Gökbuget N, et al. Blinatumomab versus chemotherapy for advanced acute lymphoblastic leukemia. *N Engl J Med*. 2017; 376(9):836-847.
7. Moreau P, Garfall AL, Van De Donk NWCJ, et al. Teclistamab in relapsed or refractory multiple myeloma. *N Engl J Med*. 2022;387(6):495-505.
8. Dickinson MJ, Carlo-Stella C, Morschhauser F, et al. Glofitamab for relapsed or refractory diffuse large B-cell lymphoma. *N Engl J Med*. 2022;387(24): 2220-2231.
9. Budde LE, Sehn LH, Matasar M, et al. Safety and efficacy of mosunetuzumab, a bispecific antibody, in patients with relapsed or refractory follicular lymphoma: a single-arm, multicentre, phase 2 study. *Lancet Oncol*. 2022;23(8):1055-1065.
10. Thieblemont C, Phillips T, Ghesquieres H, et al. Epcoritamab, a novel, subcutaneous CD3xCD20 bispecific T-cell-engaging antibody, in relapsed or refractory large B-cell lymphoma: dose expansion in a phase I/II trial. *J Clin Oncol*. 2023;41(12):2238-2247.
11. Sterner RC, Sterner RM. CAR-T cell therapy: current limitations and potential strategies. *Blood Cancer J*. 2021;11(4):69.
12. Goebeler M-E, Bargou RC. T cell-engaging therapies – BiTEs and beyond. *Nat Rev Clin Oncol*. 2020;17(7):418-434.
13. National Library of Medicine. Accessed 2 November 2024. <https://clinicaltrials.gov/>
14. Mahmood SS, Riedell PA, Feldman S, et al. Biomarkers and cardiovascular outcomes in chimeric antigen receptor T-cell therapy recipients. *Eur Heart J*. 2023;44(22):2029-2042.
15. Sayed A, Munir M, Ghazi SM, et al. Cardiovascular toxicities associated with bispecific T-cell engager therapy. *J Immunother Cancer*. 2024;12(2): e008518.
16. Johnson DB, Balko JM, Compton ML, et al. Fulminant myocarditis with combination immune checkpoint blockade. *N Engl J Med*. 2016;375(18): 1749-1755.
17. Moslehi JJ. Cardiovascular toxic effects of targeted cancer therapies. *N Engl J Med*. 2016;375(15):1457-1467.
18. Zaorsky NG, Churilla TM, Egleston BL, et al. Causes of death among cancer patients. *Ann Oncol*. 2017;28(2):400-407.
19. Strongman H, Gadd S, Matthews AA, et al. Does cardiovascular mortality overtake cancer mortality during cancer survivorship?: an English retrospective cohort study. *JACC CardioOncol*. 2022;4(1):113-123.
20. Armenian SH, Lacchetti C, Barac A, et al. Prevention and monitoring of cardiac dysfunction in survivors of adult cancers: American Society of Clinical Oncology Clinical Practice guideline. *J Clin Oncol*. 2017;35(8):893-911.
21. Accordino MK, Neugut AI, Hershman DL. Cardiac effects of anticancer therapy in the elderly. *J Clin Oncol*. 2014;32(24):2654-2661.
22. Goldman A, Maor E, Bomze D, et al. Adverse cardiovascular and pulmonary events associated with chimeric antigen receptor T-cell therapy. *J Am Coll Cardiol*. 2021;78(18):1800-1813.
23. Ganatra S, Redd R, Hayek SS, et al. Chimeric antigen receptor T-cell therapy-associated cardiomyopathy in patients with refractory or relapsed non-Hodgkin lymphoma. *Circulation*. 2020;142(17):1687-1690.
24. Herrmann J. Adverse cardiac effects of cancer therapies: cardiotoxicity and arrhythmia. *Nat Rev Cardiol*. 2020;17(8):474-502.
25. Nissani A, Lev-Ari S, Meirson T, et al. Comparison of non-myeloablative lymphodepleting preconditioning regimens in patients undergoing adoptive T cell therapy. *J Immunother Cancer*. 2021;9(5):e001743.
26. Baik AH, Oluwale OO, Johnson DB, et al. Mechanisms of cardiovascular toxicities associated with immunotherapies. *Circ Res*. 2021;128(11): 1780-1801.
27. Ganatra S, Dani SS, Yang EH, Zaha VG, Nohria A. Cardiotoxicity of T-cell antineoplastic therapies. *JACC CardioOncol*. 2022;4(5):616-623.
28. Lee DW, Gardner R, Porter DL, et al. Current concepts in the diagnosis and management of cytokine release syndrome. *Blood*. 2014;124(2):188-195.
29. Frey NV, Porter DL. Cytokine release syndrome with novel therapeutics for acute lymphoblastic leukemia. *Hematology Am Soc Hematol Educ Program*. 2016;2016(1):567-572.
30. Shah BD, Ghobadi A, Oluwale OO, et al. KTE-X19 for relapsed or refractory adult B-cell acute lymphoblastic leukaemia: phase 2 results of the single-arm, open-label, multicentre ZUMA-3 study. *Lancet*. 2021;398(10299):491-502.

31. Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-cell lymphoma. *N Engl J Med*. 2017;377(26):2531-2544.
32. Schuster SJ, Bishop MR, Tam CS, et al. Tisagenlecleucel in adult relapsed or refractory diffuse large B-cell lymphoma. *N Engl J Med*. 2019;380(1):45-56.
33. Wang M, Munoz J, Goy A, et al. KTE-X19 CAR T-cell therapy in relapsed or refractory mantle-cell lymphoma. *N Engl J Med*. 2020;382(14):1331-1342.
34. Abramson JS, Palomba ML, Gordon LI, et al. Lisocabtagene maraleucel for patients with relapsed or refractory large B-cell lymphomas (TRANSCEND NHL 001): a multicentre seamless design study. *Lancet*. 2020;396(10254):839-852.
35. Fowler NH, Dickinson M, Dreyling M, et al. Tisagenlecleucel in adult relapsed or refractory follicular lymphoma: the phase 2 ELARA trial. *Nat Med*. 2022;28(2):325-332.
36. Jacobson CA, Chavez JC, Sehgal AR, et al. Axicabtagene ciloleucel in relapsed or refractory indolent non-Hodgkin lymphoma (ZUMA-5): a single-arm, multicentre, phase 2 trial. *Lancet Oncol*. 2022;23(1):91-103.
37. Locke FL, Miklos DB, Jacobson CA, et al. Axicabtagene ciloleucel as second-line Therapy for large B-cell lymphoma. *N Eng J Med*. 2022;386(7):640-654.
38. Sehgal A, Hoda D, Riedell PA, et al. Lisocabtagene maraleucel as second-line therapy in adults with relapsed or refractory large B-cell lymphoma who were not intended for haematopoietic stem cell transplantation (PILOT): an open-label, phase 2 study. *Lancet Oncol*. 2022;23(8):1066-1077.
39. Berdeja JG, Madduri D, Usmani SZ, et al. Ciltacabtagene autoleucel, a B-cell maturation antigen-directed chimeric antigen receptor T-cell therapy in patients with relapsed or refractory multiple myeloma (CARTITUDE-1): a phase 1b/2 open-label study. *Lancet*. 2021;398(10297):314-324.
40. Cohen AD, Mateos M-V, Cohen YC, et al. Efficacy and safety of cilta-cel in patients with progressive multiple myeloma after exposure to other BCMA-targeting agents. *Blood*. 2023;141(3):219-230.
41. Del Bufalo F, De Angelis B, Caruana I, et al. GD2-CART01 for relapsed or refractory high-risk neuroblastoma. *N Engl J Med*. 2023;388(14):1284-1295.
42. Hu Y, Zhang M, Yang T, et al. Sequential CD7 CAR T-cell therapy and allogeneic HSCT without GVHD prophylaxis. *N Engl J Med*. 2024;390(16):1467-1480.
43. *KYMRIAH (tisagenlecleucel)*. US Food and Drug Administration; 2024.
44. *YESCARTA (axicabtagene ciloleucel) suspension for intravenous infusion*. US Food and Drug Administration; 2023.
45. Locke FL, Miklos DB, Jacobson CA, et al. Axicabtagene ciloleucel as second-line therapy for large B-cell lymphoma. *N Engl J Med*. 2022;386(7):640-654.
46. *TECARTUS (brexucabtagene autoleucel)*. US Food and Drug Administration; 2024.
47. *BREYANZI (lisocabtagene maraleucel)*. US Food and Drug Administration; 2024.
48. Siddiqi T, Maloney DG, Kenderian SS, et al. Lisocabtagene maraleucel in chronic lymphocytic leukaemia and small lymphocytic lymphoma (TRANSCEND CLL 004): a multicentre, open-label, single-arm, phase 1-2 study. *Lancet*. 2023;402(10402):641-654.
49. *ABECMA (idecabtagene vicleucel)*. US Food and Drug Administration; 2024.
50. Rodriguez-Otero P, Ailawadhi S, Arnulf B, et al. Ide-cel or standard regimens in relapsed and refractory multiple myeloma. *N Engl J Med*. 2023;388(11):1002-1014.
51. *CARVYKTI (ciltacabtagene autoleucel)*. US Food and Drug Administration; 2024.
52. *BLINCYTO (blinatumomab) for injection, for intravenous use*. US Food and Drug Administration; 2022.
53. Locatelli F, Zugmaier G, Mergen N, et al. Blinatumomab in pediatric relapsed/refractory B-cell acute lymphoblastic leukemia: RIALTO Expanded Access Study final analysis. *Blood Adv*. 2022;6(3):1004-1014.
54. Martinelli G, Boissel N, Chevallier P, et al. Long-term follow-up of blinatumomab in patients with relapsed/refractory Philadelphia chromosome-positive B-cell precursor acute lymphoblastic leukaemia: final analysis of ALCANTARA study. *Eur J Cancer*. 2021;146:107-114.
55. Gökbuget N, Dombret H, Bonifacio M, et al. Blinatumomab for minimal residual disease in adults with B-cell precursor acute lymphoblastic leukemia. *Blood*. 2018;131(14):1522-1531.
56. *TECVAYLI (teclistamab-cqyv) injection, for subcutaneous use*. US Food and Drug Administration; 2022.
57. *COLUMVI (glofitamab-gxbm) injection, for intravenous use*. US Food and Drug Administration; 2023.
58. *LUNSUMIO (mosunetuzumab-axgb) injection for intravenous use*. US Food and Drug Administration; 2022.
59. *EPKINLY (epcoritamab-bysp) injection, for subcutaneous use*. US Food and Drug Administration; 2023.
60. Chesney J, Lewis KD, Kluger H, et al. Efficacy and safety of lifileucel, a one-time autologous tumor-infiltrating lymphocyte (TIL) cell therapy, in patients with advanced melanoma after progression on immune checkpoint inhibitors and targeted therapies: pooled analysis of consecutive cohorts of the C-144-01 study. *J Immunother Cancer*. 2022;10(12):e005755.
61. *AMTAGVI (lifileucel)*. US Food and Drug Administration; 2024.
62. Cook J, Litzow M. Advances in supportive care for acute lymphoblastic leukemia. *Curr Hematol Malig Rep*. 2020;15(4):276-293.
63. Hay KA, Hanafi L-A, Li D, et al. Kinetics and biomarkers of severe cytokine release syndrome after CD19 chimeric antigen receptor-modified T-cell therapy. *Blood*. 2017;130(21):2295-2306.

64. Totzeck M, Michel L, Lin Y, Herrmann J, Rassaf T. Cardiotoxicity from chimeric antigen receptor-T cell therapy for advanced malignancies. *Eur Heart J*. 2022;43(20):1928-1940.
65. Lee DW, Santomaso BD, Locke FL, et al. ASTCT Consensus grading for cytokine release syndrome and neurologic toxicity associated with immune effector cells. *Biol Blood Marrow Transplant*. 2019;25(4):625-638.
66. Common Terminology Criteria for Adverse Events (CTCAE). US Department of Health and Human Services, National Institutes of Health; 2017.
67. Fitzgerald JC, Weiss SL, Maude SL, et al. Cytokine release syndrome after chimeric antigen receptor T cell therapy for acute lymphoblastic leukemia. *Crit Care Med*. 2017;45(2):e124-e131.
68. Morris EC, Neelapu SS, Giavridis T, Sadelain M. Cytokine release syndrome and associated neurotoxicity in cancer immunotherapy. *Nat Rev Immunol*. 2022;22(2):85-96.
69. Giavridis T, van der Stegen SJC, Eyquem J, Hamieh M, Piersigilli A, Sadelain M. CAR T cell-induced cytokine release syndrome is mediated by macrophages and abated by IL-1 blockade. *Nat Med*. 2018;24(6):731-738.
70. Norelli M, Camisa B, Barbiera G, et al. Monocyte-derived IL-1 and IL-6 are differentially required for cytokine-release syndrome and neurotoxicity due to CAR T cells. *Nat Med*. 2018;24(6):739-748.
71. Shimabukuro-Vornhagen A, Gödel P, Subklewe M, et al. Cytokine release syndrome. *J Immunother Cancer*. 2018;6(1):56.
72. Gust J, Hay KA, Hanafi L-A, et al. Endothelial activation and blood-brain barrier disruption in neurotoxicity after adoptive immunotherapy with CD19 CAR-T cells. *Cancer Discov*. 2017;7(12):1404-1419.
73. Chen J, Chung DW. Inflammation, von Willebrand factor, and ADAMTS13. *Blood*. 2018;132(2):141-147.
74. Teachey DT, Lacey SF, Shaw PA, et al. Identification of predictive biomarkers for cytokine release syndrome after chimeric antigen receptor T-cell therapy for acute lymphoblastic leukemia. *Cancer Discov*. 2016;6(6):664-679.
75. Le RQ, Li L, Yuan W, et al. FDA approval summary: tocilizumab for treatment of chimeric antigen receptor T cell-induced severe or life-threatening cytokine release syndrome. *Oncologist*. 2018;23(8):943-947.
76. Park JH, Nath K, Devlin SM, et al. CD19 CAR T-cell therapy and prophylactic anakinra in relapsed or refractory lymphoma: phase 2 trial interim results. *Nat Med*. 2023;29(7):1710-1717.
77. IL-1 receptor antagonist to prevent severe chimeric antigen receptor T-cell related encephalopathy syndrome ClinicalTrials.gov identifier: NCT04205838. Updated 5 July 2024. Accessed 1 January 2024. <https://clinicaltrials.gov/study/NCT04205838>
78. Phase 2 pilot study to evaluate efficacy and safety of anakinra to prevent CD19-targeted CAR-T cell-related cytokine release syndrome (CRS) and neurotoxicity in patients with B cell lymphoma. ClinicalTrials.gov identifier: NCT04359784. Updated 18 July 2024. Accessed 1 January 2024. <https://clinicaltrials.gov/study/NCT04359784>
79. Pilot study of anakinra to mitigate CAR-T toxicity in subjects with relapsed or refractory large B-cell lymphoma ClinicalTrials.gov identifier: NCT04432506. Updated 31 May 2024. Accessed 1 January 2024. <https://clinicaltrials.gov/study/NCT04432506>
80. Neelapu SS, Tummala S, Kebriaei P, et al. Chimeric antigen receptor T-cell therapy—assessment and management of toxicities. *Nat Rev Clin Oncol*. 2018;15(1):47-62.
81. Marar RI, Abbasi MA, Prathivadhi-Bhayankaram S, et al. Cardiotoxicities of novel therapies in hematologic malignancies: chimeric antigen receptor T-cell therapy and bispecific T-cell engager therapy. *JCO Oncol Pract*. 2023;19(6):331-342.
82. Alvi RM, Frigault MJ, Fradley MG, et al. Cardiovascular events among adults treated with chimeric antigen receptor T-cells (CAR-T). *J Am Coll Cardiol*. 2019;74(25):3099-3108.
83. Lefebvre B, Kang Y, Smith AM, Frey NV, Carver JR, Scherrer-Crosbie M. Cardiovascular effects of CAR T cell therapy: a retrospective study. *JACC CardioOncol*. 2020;2(2):193-203.
84. Guha A, Addison D, Jain P, et al. Cardiovascular events associated with chimeric antigen receptor T cell therapy: cross-sectional FDA Adverse Events Reporting System Analysis. *Biol Blood Marrow Transplant*. 2020;26(12):2211-2216.
85. Shalabi H, Sachdev V, Kulshreshtha A, et al. Impact of cytokine release syndrome on cardiac function following CD19 CAR-T cell therapy in children and young adults with hematological malignancies. *J Immunother Cancer*. 2020;8(2):e001159.
86. Wudhikarn K, Pennisi M, Garcia-Recio M, et al. DLBCL patients treated with CD19 CAR T cells experience a high burden of organ toxicities but low nonrelapse mortality. *Blood Adv*. 2020;4(13):3024-3033.
87. Brammer JE, Braunstein Z, Katapadi A, et al. Early toxicity and clinical outcomes after chimeric antigen receptor T-cell (CAR-T) therapy for lymphoma. *J Immunother Cancer*. 2021;9(8):e002303.
88. Qi K, Yan Z, Cheng H, et al. An analysis of cardiac disorders associated with chimeric antigen receptor T cell therapy in 126 patients: a single-centre retrospective study. *Front Oncol*. 2021;11:691064.
89. Hollenberg SM, Singer M. Pathophysiology of sepsis-induced cardiomyopathy. *Nat Rev Cardiol*. 2021;18(6):424-434.
90. Fradley MG, Damrongwatanasuk R, Chandrasekhar S, Alomar M, Kip KE, Sarnaik AA. Cardiovascular toxicity and mortality associated with adoptive cell therapy and tumor-infiltrating lymphocytes for advanced stage melanoma. *J Immunother*. 2020;44(2):86-89.
91. Burstein DS, Maude S, Grupp S, Griffis H, Rossano J, Lin K. Cardiac profile of chimeric antigen receptor T cell therapy in children: a single-institution experience. *Biol Blood Marrow Transplant*. 2018;24(8):1590-1595.

92. Jung SH, Lee SR, Yang DH, et al. Efficacy and safety of blinatumomab treatment in adult Korean patients with relapsed/refractory acute lymphoblastic leukemia on behalf of the Korean Society of Hematology ALL Working Party. *Ann Hematol*. 2019;98(1):151-158.
93. Lefebvre B, Kang Y, Vakilpour A, et al. Incidence of MACE in patients treated with CAR-T cell therapy. *JACC CardioOncol*. 2023;5(6):747-754.
94. Korell F, Entenmann L, Romann S, et al. Evaluation of all-cause mortality and cardiovascular safety in patients receiving chimeric antigen receptor T cell therapy: a prospective cohort study. *EClinicalMedicine*. 2024;69:102504.
95. Schuster SJ, Tam CS, Borchmann P, et al. Long-term clinical outcomes of tisagenlecleucel in patients with relapsed or refractory aggressive B-cell lymphomas (JULIET): a multicentre, open-label, single-arm, phase 2 study. *Lancet Oncol*. 2021;22(10):1403-1415.
96. Wang Y, Qi K, Cheng H, et al. Coagulation disorders after chimeric antigen receptor T cell therapy: analysis of 100 patients with relapsed and refractory hematologic malignancies. *Biol Blood Marrow Transplant*. 2020;26(5):865-875.
97. Goekbuget N, Dombret H, Bonifacio M, et al. BLAST: a confirmatory, single-arm, phase 2 study of blinatumomab, a Bispecific T-Cell Engager (BiTE®) antibody construct, in patients with minimal residual disease B-precursor acute lymphoblastic leukemia (ALL). *Blood*. 2014;124(21):379.
98. Chari A, Minnema MC, Berdeja JG, et al. Talquetamab, a T-cell-redirecting GPRC5D bispecific antibody for multiple myeloma. *N Engl J Med*. 2022;387(24):2232-2244.
99. Lesokhin AM, Tomasson MH, Arnulf B, et al. Elranatamab in relapsed or refractory multiple myeloma: phase 2 MagnetisMM-3 trial results. *Nat Med*. 2023;29(9):2259-2267.
100. Ghosh AK, Chen DH, Guha A, Mackenzie S, Walker JM, Roddie C. CAR T cell therapy-related cardiovascular outcomes and management: systemic disease or direct cardiotoxicity? *JACC CardioOncol*. 2020;2(1):97-109.
101. Addison D, Neilan TG, Barac A, et al. Cardiovascular imaging in contemporary cardio-oncology: a scientific statement from the American Heart Association. *Circulation*. 2023;148(16):1271-1286.
102. Lyon AR, López-Fernández T, Couch LS, et al. 2022 ESC guidelines on cardio-oncology developed in collaboration with the European Hematology Association (EHA), the European Society for Therapeutic Radiology and Oncology (ESTRO) and the International Cardio-Oncology Society (IC-OS). *Eur Heart J Cardiovasc Imaging*. 2022;23(10):e333-e465.
103. Lyon AR, López-Fernández T, Couch LS, et al. 2022 ESC Guidelines on cardio-oncology developed in collaboration with the European Hematology Association (EHA), the European Society for Therapeutic Radiology and Oncology (ESTRO) and the International Cardio-Oncology Society (IC-OS). *Eur Heart J*. 2022;43(41):4229-4361.
104. Riegler LL, Jones GP, Lee DW. Current approaches in the grading and management of cytokine release syndrome after chimeric antigen receptor T-cell therapy. *Ther Clin Risk Manag*. 2019;15:323-335.
105. Topp MS, van Meerten T, Houot R, et al. Earlier corticosteroid use for adverse event management in patients receiving axicabtagene ciloleucel for large B-cell lymphoma. *Br J Haematol*. 2021;195(3):388-398.
106. Gardner RA, Ceppi F, Rivers J, et al. Preemptive mitigation of CD19 CAR T-cell cytokine release syndrome without attenuation of antileukemic efficacy. *Blood*. 2019;134(24):2149-2158.
107. Kadauke S, Myers RM, Li Y, et al. Risk-adapted preemptive tocilizumab to prevent severe cytokine release syndrome after CTL019 for pediatric B-cell acute lymphoblastic leukemia: a prospective clinical trial. *J Clin Oncol*. 2021;39(8):920-930.
108. Oluwole OO, Bouabdallah K, Muñoz J, et al. Prophylactic corticosteroid use in patients receiving axicabtagene ciloleucel for large B-cell lymphoma. *Br J Haematol*. 2021;194(4):690-700.
109. Rohaan MW, van den Berg JH, Kvistborg P, Haanen JBAG. Adoptive transfer of tumor-infiltrating lymphocytes in melanoma: a viable treatment option. *J Immunother Cancer*. 2018;6(1):102.
110. Lin H, Cheng J, Zhu L, et al. Anti-CD5 CAR-T cells with a tEGFR safety switch exhibit potent toxicity control. *Blood Cancer J*. 2024;14(1):98.