

STATE-OF-THE-ART REVIEW

Navigating Hypoxemic Respiratory Failure in Critically Ill Cardiac Patients

Lee H. Sterling, MDCM,^{a,b} Shannon M. Fernando, MD, MSc,^{a,c,d,e} Patrick R. Lawler, MD, MPH,^{f,g} Susanna Price, MBBS, PhD,^h Eddy Fan, MD, PhD,^{f,i} Ewan Goligher, MD, PhD,^{f,i} Rebecca Mathew, MD^{a,j}

ABSTRACT

Acute respiratory failure is a common reason for admission to cardiac intensive care units and the prevalence of respiratory failure in this cohort is increasing over time. Hypoxemia can occur due to a variety of mechanisms; the most common cause in the cardiac intensive care units remains cardiogenic pulmonary edema, but other etiologies, such as pneumonia and acute respiratory distress syndrome, are also common. This article provides an update on mechanisms of hypoxemia among patients with cardiac critical illness, heart lung interactions during spontaneous and positive pressure ventilation, optimization of sedation and ventilation for cardiac patients including novel ventilation strategies, and management of refractory hypoxemia among patients with cardiac critical illness. (JACC Adv. 2025;■:101616)

© 2025 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Acute respiratory failure is a common reason for admission to cardiac intensive care units (CICUs)^{1,2} and the prevalence of respiratory failure is increasing. In their tertiary care CICUs, Jentzer and colleagues described a rising prevalence of respiratory failure, from 15% in 2007 to 38% in 2018.³ Acute respiratory failure is an umbrella term encompassing acute hypoxemic respiratory failure (AHRF) and hypercarbic failure. Hypoxemia, or a partial pressure of oxygen in arterial blood (PaO₂) below the normal range of 80–100 mm Hg, frequently leads to hypoxia, a failure of oxygenation at the tissue level. AHRF is often defined as a PaO₂ to fraction of inspiratory oxygen <300. This narrative review focuses on the pathophysiology, epidemiology, and

management of AHRF for patients with significant left ventricle (LV) or right ventricle (RV) dysfunction.

Physiologic mechanisms of hypoxemia in patients with cardiac critical illness are multifold and synergistic. They include ventilation/perfusion (V/Q) mismatch, or an imbalance between ventilation and perfusion in the lungs. An extreme V/Q mismatch occurs with right to left shunts, either due to anatomical intracardiac shunts, such as atrial and ventricular septal defects, or physiologic shunts from alveolar disorders (ie, atelectasis, cardiogenic or noncardiogenic pulmonary edema, or pneumonia). The other extreme is physiologic dead space due to decreased perfusion of gas exchange surfaces (ie, large pulmonary embolism). Other important

From the ^aCAPITAL Research Group, Division of Cardiology, University of Ottawa Heart Institute, Ottawa, Ontario, Canada;

^bDivision of Critical Care Medicine, University of British Columbia, Vancouver, British Columbia, Canada; ^cDepartment of Critical Care Medicine, Queen's University, Kingston, Ontario, Canada; ^dDepartment of Critical Care, Lakeridge Health, Oshawa, Ontario, Canada; ^eClinical Epidemiology Program, Ottawa Hospital Research Institute, Ottawa, Ontario, Canada; ^fInterdepartmental Division of Critical Care Medicine, University of Toronto, Toronto, Ontario, Canada; ^gDivision of Cardiology, McGill University Health Centre, Montreal, Quebec, Canada; ^hDepartments of Cardiology & Critical Care, Royal Brompton Hospital, National Heart & Lung Institute, Imperial College, London, United Kingdom; ⁱToronto General Hospital Research Institute, University Health Network, Toronto, Ontario, Canada; and the ^jDepartment of Critical Care, The Ottawa Hospital, Ottawa, Ontario, Canada.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

Manuscript received September 25, 2024; revised manuscript received December 2, 2024, accepted December 10, 2024.

**ABBREVIATIONS
AND ACRONYMS****AHRF** = acute hypoxemic respiratory failure**ARDS** = acute respiratory distress syndrome**CICU** = cardiac intensive care unit**CS** = cardiogenic shock**ECMO** = extracorporeal membrane oxygenation**FiO₂** = fraction of inspiratory oxygen**IL** = interleukin**IMV** = invasive mechanical ventilation**INO** = inhaled nitric oxide**LTVV** = low tidal volume ventilation**LV** = left ventricle**PaO₂** = partial pressure of oxygen in arterial blood**PEEP** = positive end expiratory pressure**PPV** = positive pressure ventilation**RV** = right ventricle**SBT** = spontaneous breathing trial**V/Q** = ventilation/perfusion

mechanisms of hypoxemia include diffusion impairment at the alveolar-pulmonary capillary interface (ie, interstitial lung disease); and hypoventilation secondary to reduced minute ventilation from central neurologic conditions (ie, sedation, central apnea in cardiac arrest survivors) or peripheral neuromuscular disorders (ie, cervical spine injuries, diaphragmatic weakness). Finally, of particular importance in a cardiac population is decreased mixed venous oxygen content (ie, low cardiac output due to pump failure). In low cardiac output states, low mixed venous hemoglobin saturation can cause severe hypoxemia if even a small degree of V/Q mismatch or shunt simultaneously coexists (**Central Illustration**).

AN OVERVIEW OF AHRF AND INVASIVE MECHANICAL VENTILATION IN THE CICUs

A familiar cause of AHRF in the CICUs is cardiogenic pulmonary edema.² This occurs when a rise in LV end-diastolic and left atrial pressures produce increased hydrostatic pressures in the pulmonary capillaries. Elevated hydrostatic pressures overwhelm the active lymphatic resorption of fluids leading to alveolar edema and impaired gas exchange.⁴ Importantly, multiple processes may coexist. For example, in cardiogenic shock (CS), there is often an accompanying systemic inflammatory cascade due to upregulation of inflammatory cytokines such as interleukin (IL)-1 β , IL-6, IL-8, tissue necrosis factor- α , and the complement system.⁵⁻⁷ In some patients, this inflammatory cascade may increase permeability of the pulmonary vascular membrane, leading to noncardiogenic pulmonary edema. Mechanical circulatory support strategies that increase end-diastolic pressure may also promote these mechanisms.⁸ The downstream effect of both high filling pressures and capillary leak in CS may result in worsening hypoxemia leading to further myocardial ischemia, tissue hypoxia, and progressive multi-organ failure.

A proportion of patients with AHRF admitted to CICUs are managed with either high flow nasal cannula or other forms of noninvasive positive pressure ventilation (PPV). High flow nasal cannula uses heated and humidified flow rates of up to 60 L/minute, providing a more reliable FiO₂ and minimal positive end expiratory pressure (PEEP). Other forms of noninvasive PPV include bilevel positive airway

HIGHLIGHTS

- Acute hypoxemic respiratory failure is common in contemporary cardiac intensive care units.
- When initiating invasive mechanical ventilation, it is crucial to consider the hemodynamic effects of both positive pressure ventilation and the impact of analgesic and sedation strategies.
- Adjunctive treatments for refractory hypoxemia in patients with cardiac critical illness remain understudied.


pressure which can support both hypoxia and hypercarbia. The most common indication for invasive mechanical ventilation (IMV) in CICUs remains AHRF. However, IMV may also be required for airway protection postcardiac arrest and to facilitate invasive procedures. When contemplating initiation of IMV, it is crucial to consider the hemodynamic effects of induction, intubation, and PPV, as discussed below and additionally reviewed elsewhere.⁹

HEART-LUNG INTERACTIONS AND THE IMPACTS OF SEDATION DURING IMV

On initiating IMV, the heart-lung unit undergoes several important changes related to the shift to PPV. The hemodynamic effects of IMV are mediated through a reduction in respiratory muscle effort, lung and chest wall inflation, and via the switch from negative to positive pleural pressures. Importantly, these hemodynamic changes have differential effects on the right and left sides of the heart (**Figure 1**).

RV preload is equal to cardiac venous return. Venous return is principally determined by the difference between mean systemic filling pressures and right atrial pressure.¹⁰ Initiation of IMV increases intrathoracic pressure and subsequently right atrial pressure, decreasing the gradient between mean systemic filling pressures and the right atrium, and impairing venous return.¹¹ During spontaneous respiration, pleural pressures become more negative during inspiration, and transmural cardiac pressures drop, increasing the gradient between the mean systemic filling pressures and the right atrium. In contrast, during IMV, each breath is accompanied by increased intrathoracic pressure without variation in right atrial pressures.^{12,13}

IMV has significant effects on RV afterload, classically described as a U-shaped curve mediated through changes in lung volume and pulmonary vascular


CENTRAL ILLUSTRATION Hypoxemic Respiratory Failure in Critically Ill Cardiac Patients

Sterling LH, et al. JACC Adv. 2025;■(■):101616.

The figure highlights pathophysiological mechanisms of hypoxemia. (A) A heart-lung unit with normal blood flows through a nondiseased lung. (B) An example of intraparenchymal shunt with zero ventilation to a perfused portion of lung, (C) V/Q mismatch, (D) an example of dead space ventilation, where a fully ventilated alveolus has zero perfusion, (E) diffusion impairment, (F) low mixed venous oxygen saturation seen in situations such as cardiogenic shock, and (G) hypoventilation. V/Q = ventilation/perfusion.

resistance. PEEP may reduce atelectasis and, for a subset of patients, decrease RV afterload. However, in most cases, IMV increases RV afterload. The combination of decreased RV preload and increased RV afterload culminates in decreased right-sided cardiac output. Ultimately, the cautious use of PEEP with RV dysfunction is more nuanced than a simplistic minimization or worse yet, avoidance. Common scenarios where PEEP titration must be undertaken cautiously include the patient struggling with refractory hypoxemia or acidosis as both produce synergistic

pulmonary vasoconstriction increasing RV afterload, and those with noncompliant lung parenchyma without a PPV-responsive condition, that is, interstitial lung disease. In this latter scenario, small changes in alveolar pressure produce marked rises in RV afterload. Finally, in patients with systemic hypotension, high PEEP raises central venous pressures, lowers RV perfusion pressure and may exacerbate RV ischemia. Conversely, patients with preserved RV function and adequate preload are less sensitive to the effects of PEEP and more liberal titration can be

FIGURE 1 Effects of Positive Pressure Ventilation on Preload and Afterload

Differential effects of positive pressure ventilation on left and right ventricular preload and afterload. LV = left ventricle; PEEP = positive end expiratory pressure; P_{Pl} = pleural pressures; PVR = pulmonary vascular resistance; RV = right ventricle; SVR = systemic vascular resistance.

undertaken to unload both the respiratory muscles and LV.⁹

In the absence of intracardiac shunts, LV preload is determined by cardiac output from the RV. Given the hemodynamic impact of IMV on the RV, there is often decreased LV preload during IMV. During spontaneous respiration, decreases in pleural pressures increase LV afterload.¹⁴ However, PEEP during IMV decreases LV afterload through multiple mechanisms. First, PEEP increases pressure on the LV and thoracic aorta causing a pressure gradient between intrathoracic structures and the systemic circulation, promoting forward flow.¹⁵ Second, increased intrathoracic pressures activate aortic baroreceptors, lowering systemic vascular resistance.¹¹ Finally, positive pleural pressures decrease transmural LV pressure and reduce LV wall tension. In patients with LV dysfunction, IMV may have numerous beneficial effects on LV performance by reducing preload and afterload, mimicking the effects of diuretics and afterload reducing agents. Conversely, in rare situations, IMV may not only induce neurohormonal activation and volume retention, but the necessity of sedation may necessitate vasopressors which could increase the risk of arrhythmias and myocardial ischemia.¹⁶ Ultimately, the effects of IMV on hemodynamics will depend on the

sum of the effects on RV and LV preload and afterload, as well as the systemic effects of PPV and will vary from patient to patient.

The optimal timing for endotracheal intubation and initiation of IMV in AHRF is unknown with no randomized controlled trial data to guide decision-making (Table 1).¹⁷ Even less evidence is available to guide this decision in the CICUs.⁹ Timing of intubation is thus highly individualized. However, the available data suggest that once the decision has been made to initiate IMV, delays are associated with increased mortality.^{18,19} There is a risk of peri-intubation hypotension and cardiac arrest with the attenuation in sympathetic tone inherent to all induction agents—a prominent concern in the patient with low cardiac output reliant on systemic vascular resistance to maintain systemic perfusion. This is compounded by the potential cardiodepressant effects of induction medications. Propofol was recently compared vs etomidate as an induction agent for patients requiring IMV in context of acute myocardial infarction: propofol was associated with reduced in-hospital mortality.²⁰ However, these data remain retrospective and at high risk for unmeasured confounding. In absence of prospective, randomized data, it may be most appropriate to use agents one is most

TABLE 1 Clinical Pearls for the Clinician Managing Hypoxemic Respiratory Failure in Critically Ill Cardiac Patients

| | |
|--|---|
| What is the optimal timing for intubation? | <ul style="list-style-type: none"> Optimal timing for endotracheal intubation and in AHRF is unknown and highly individualized. Once the decision has been made to initiate IMV, delays are associated with increased mortality |
| What are the hemodynamic considerations during induction and initiation of IMV | <ul style="list-style-type: none"> All induction agents attenuate sympathetic tone and may cause peri-intubation hypotension. Patients who are preload dependent, particularly those with RV dysfunction or pulmonary hypertension, should be placed on the lowest effective PEEP to maintain adequate oxygenation and prevent lung derecruitment LV preload and afterload will be reduced with positive pressure ventilation |
| How should my patient with AHRF be assessed for extubation? | <ul style="list-style-type: none"> Extubation readiness includes resolution of the condition requiring intubation, absence of significant tracheal secretions, adequate cough reflex and mentation, and stable cardiovascular and respiratory status. Standardized breathing trial reduce total mechanical ventilation days and identify objective and subjective indices of failure Weaning-induced pulmonary edema is a common cause of weaning failure and volume status should be optimized prior to extubation. |

AHRF = acute hypoxemic respiratory failure; IMV = invasive mechanical ventilation; LV = left ventricle; PEEP = positive end expiratory pressure; RV = right ventricle.

familiar with, with the smallest cardiodepressant effects and at lowest possible doses.²¹ Additionally, important strategies in those with RV preload dependent states include avoidance of intubation if feasible, and if not, to maintain spontaneous respiration during induction, maintain mean arterial pressure above pulmonary artery pressure, and optimize preload.

Most patients require sedation during IMV. Guidelines recommend intravenous opioid analgesics and sedatives at lowest possible effective doses, with titration based on established sedation scores, like the Richmond Agitation-Sedation Score.²² Opioids have minimal effect on cardiac output and may reduce myocardial oxygen demand.²³ Opioid-sparing strategies may also be considered, including regional blocks for rib fractures post cardiac arrest, or ketamine or lidocaine infusions. There is a paucity of evidence for the optimal sedative medication for intubated patients with cardiac critical illness. Recommended first-line sedatives in a general ICU population include propofol, short-acting benzodiazepines, and dexmedetomidine.²² A full overview of all available sedative medications and their relative utility in various clinical scenarios encountered in the CICUs is outside the scope of this article but has been the focus of prior expert reviews.²⁴ Ultimately the sedation strategy for patients requiring IMV in the CICUs should be individualized considering hemodynamics, baseline cardiac function, anticipated duration of IMV, and target level of sedation.

OPTIMIZING VENTILATORY SETTINGS IN THE CICUs

There is sparse evidence to inform optimal ventilator settings in the CICUs. Best practices are drawn from

the general ICU literature, and in cases of severe hypoxemia, from patients with acute respiratory distress syndrome (ARDS). In general, the various modes of IMV involve setting the fraction of inspired oxygen, level of PEEP, mandatory or patient-initiated ventilation (referred to as controlled or support modes) and choice of tidal volumes or peak pressures as a target for ventilation.²⁵

Mechanistically, the focus during initial resuscitation is to offload the respiratory muscles with controlled modes of ventilation with a transition to an assisted mode when patients can tolerate the workload of spontaneous respiration. Achieving lung protective settings with low tidal volume ventilation (LTVV) and noninjurious ventilator settings is also commonly emphasized. LTVV minimizes alveolar overdistension and ventilator-induced lung injury and reduces mortality in ARDS.²⁶⁻²⁸ The role of LTVV in AHRF not due to ARDS is less clear. A trial of 961 intubated patients without ARDS found that LTVV was not superior to intermediate tidal volume ventilation.²⁹ In contrast, a meta-analysis of multiple interventional and observational studies suggested benefit of LTVV in non-ARDS respiratory failure, though mortality benefit was only seen among observational studies.³⁰ The evidence for a LTVV strategy in the CICUs remains limited. In a cohort of 51 ventilated patients with heart failure or cardiac arrest in a tertiary care CICU, the median initial tidal volume was 9.3 mL/kg. All-cause mortality was 23% in patients who received a tidal volume below the median value and 36% in those who received a tidal volume above the median value.³¹ Similarly, in an analysis of the Extracorporeal Life Support Organization registry, intubated patients with CS supported on venoarterial extracorporeal membrane

oxygenation (ECMO) managed with a LTVV strategy had lower in-hospital mortality.³² Unfortunately, tidal volumes and driving pressure could only be inferred from the reported PEEP and peak inspiratory pressures, highlighting the need for high-quality observational data in a CICU population.

Pathophysiologically, volutrauma, atelectotrauma, and barotrauma all increase concentrations of inflammatory cytokines and may perpetuate ongoing multi-organ failure.²⁷ The inflammatory cascade of CS is profound and a growing area of exploration in CS management. Optimization of tidal volumes to minimize secondary sources of inflammation may provide an additional avenue for therapeutic benefit. Conversely, permissive hypercapnia from LTVV can cause acidosis, a myocardial depressant, so acid-base status should be monitored. Currently, LTVV is recommended for intubated patients with CS mostly based on expert opinion³³ with no randomized trial data yet showing benefit.

Lung protective ventilation aims to minimize the risk of ventilator-associated lung and diaphragm injury. Another major cause of ventilator-associated lung injury remains patient-ventilator dyssynchrony. Asynchrony can occur at each stage of breath triggering, delivery, and termination. Patient ventilation dyssynchrony is associated with worse clinical outcomes and in some cases contributes to ongoing lung injury.³⁴ Strategies to minimize dyssynchrony include changing ventilation parameters to match patient demand, optimizing sedation and occasionally, paralysis. A full overview of all patient-ventilator dyssynchrony has been well covered in prior reviews.³⁵

Another mainstay of lung protective ventilation involves administration of PEEP to optimize and sustain alveolar recruitment. Selection of PEEP should consider underlying cardiovascular pathology, hemodynamics, and time point in disease course. Patients who are preload dependent, particularly those with RV dysfunction or pulmonary hypertension, should be placed on the lowest effective PEEP to maintain adequate oxygenation and prevent lung derecruitment. In contrast, for patients who are predominantly afterload sensitive, most commonly those with LV dysfunction, higher PEEPs may have beneficial effects. In a mechanistic study, application of higher PEEPs (from 3 to 20 cm H₂O) improved mitral annular dimensions and reduced LV end systolic volumes. High PEEP strategies may thus be particularly beneficial in patients with LV dysfunction and significant mitral regurgitation.³⁶

Beyond the use of LTVV and higher PEEP, several alternative ventilation strategies are under

investigation for treatment of AHRF. The first involves targeting a low respiratory system driving pressure. Driving pressure is defined as the difference between plateau pressure and PEEP. In a post hoc analysis of several randomized trials of patients with ARDS, reduction in driving pressure was the best predictor of survival, even beyond achievement of LTVV and noninjurious PEEP.³⁷ Several ongoing randomized trials are examining the efficacy of lower driving pressures in ARDS and non-ARDS forms of AHRF.^{38,39} Second, several trials have examined an “open lung” ventilation strategy in patients with ARDS. No standardized definition exists for “open lung” ventilation, but it usually involves serial recruitment maneuvers or the use of inverse ratio ventilation. While there has been considerable interest in recruitment maneuvers, when compared against conventional LTVV in ARDS, there is no mortality benefit or improvement in ICU length of stay.³⁰

REFRACTORY HYPOXEMIA IN CARDIAC CRITICAL ILLNESS

Despite optimization, a subset of patients in the CICUs will develop refractory hypoxemia; defined as a PaO₂ ≤60 mm Hg or a PaO₂ to fraction of inspiratory oxygen ≤100 with FiO₂ ≥0.8 and either PEEP >15 cm H₂O or plateau pressures >30 cm H₂O for >12 hours despite LTVV.⁴⁰ Management of refractory hypoxemia may involve adjunctive treatment strategies including prone positioning, neuromuscular blockade, inhaled pulmonary vasodilators (eg, nitric oxide or epoprostenol), inverse I/E ventilation, and ECMO. There is a knowledge gap in the use of adjunctive treatments for refractory AHRF in CICU patients. Below is summarized the most common adjunctive treatments of refractory hypoxemia and the evidence for their use in patients with significant cardiac disease.

PRONE POSITIONING. Prone positioning for at least 16 hours daily reduces mortality in patients with moderate to severe ARDS.²⁷ Prone positioning can improve RV hemodynamics and may be helpful in context of acute cor pulmonale from AHRF. The hemodynamic effects on the LV are less clear. Improvement in RV hemodynamics increase LV preload and rises in pulmonary capillary wedge pressures have been reported with prone positioning.⁴¹ Additionally, ventricular arrhythmias remain a significant concern in patients admitted to CICUs. Even under optimal conditions, supination to facilitate chest compressions may take up to 3 minutes.⁴² The use of prone positioning in the CICUs or patients with LV predominant CS should therefore be undertaken

on a case-by-case basis, accounting for the potentially deleterious hemodynamic effects of higher LV filling pressures and arrhythmic risk.

NEUROMUSCULAR BLOCKADE. Neuromuscular blockade in patients with severe ARDS is theorized to improve outcomes by relaxation of expiratory muscle effort, reducing patient-ventilator dyssynchrony, lowering work of breathing and potential anti-inflammatory mechanisms.⁴³ However, paralysis is associated with several important adverse effects including myopathy, polyneuropathy, pressure ulcers, deep venous thrombosis, and corneal abrasions.⁴⁴ Guidelines conditionally recommend neuromuscular blockade in early severe ARDS. While ACURASYS (the Neuromuscular blockers in early acute respiratory distress syndrome) trial demonstrated improved 90-day survival, ROSE (the Early Neuromuscular Blockade in the Acute Respiratory Distress Syndrome) trial failed to show benefit when using a lighter sedation strategy, which is more reflective of contemporary practice.⁴⁵⁻⁴⁷ Practically, use of neuromuscular blockade is reserved for patients who remain dyssynchronous with the ventilator despite adequate sedation or with severe hypoxemia. Neuromuscular blocking agents are unlikely to cause direct hemodynamic effects but may reduce myocardial oxygen demand. In the comatose patient following cardiac arrest, prior exposure to neuromuscular blockade may represent an important confounder. Residual paralysis should be excluded prior to formal neuroprognostication.

INHALED NITRIC OXIDE AND PULMONARY VASODILATORS. Inhaled nitric oxide (iNO) may improve V/Q mismatch via local pulmonary vasodilation in well-ventilated alveoli. While iNO often improves oxygenation, there is no established reduction in mortality and concerning associations with renal impairment when used for longer durations.⁴⁸ In patients with CS with predominantly RV failure, iNO may have a role to temporarily reduce pulmonary vascular resistance and augment cardiac index.⁴⁹ In patients with severe LV dysfunction, augmentation of right-sided cardiac output may increase LV preload.⁵⁰ While a potential tool in managing patients with RV dysfunction or severe AHRF, the decision to use iNO in the CICUs should be individualized.

EXTRACORPOREAL MEMBRANE OXYGENATION. V-V ECMO reduces mortality in severe ARDS^{51,52} and may indirectly improve RV hemodynamics by allowing for ultra-lung protective ventilation and improvement in hypoxic vasoconstriction.⁵³ A subset of patients will have marked hemodynamic improvement with V-V

ECMO if their RV dysfunction improves with optimization of oxygenation and ventilation. If V-V ECMO fails to improve RV hemodynamics other options include switching to venoarterial ECMO or addition of right-sided circulatory support. However, if ECMO is being considered in context of LV predominant CS and refractory hypoxemia, V-A configurations are more appropriate. Notably, the routine use of V-A ECMO in acute myocardial infarction-associated CS does not improve outcomes and is associated with vascular complications.⁵⁴

WEANING FROM INVASIVE MECHANICAL VENTILATION

Ideally, patients improve to the point of being ready for liberation from IMV.⁵⁵ An assessment of extubation readiness includes resolution of the condition requiring intubation, absence of significant tracheal secretions, adequate cough reflex and mentation, and stable cardiovascular and respiratory status. Thereafter, a spontaneous breathing trial (SBT) should be performed⁵⁶ to assess for objective and subjective indices of failure.

The weaning process constitutes up to 50% of ventilator days and 26% to 42% of patients fail their first wean attempt.⁵⁷ Importantly, weaning-induced alveolar edema is a common cause of failed SBTs.⁵⁸ Ensuring optimization of filling pressures, commonly through diuresis or cautious fluid loading in preload dependent states, is therefore crucial. Other important causes of failed SBTs include excessive respiratory load, neuromuscular dysfunction, critical illness neuromuscular abnormalities, neuropsychological factors, and metabolic/endocrine disorders.⁵⁹ Standardized ventilator liberation protocols reduce mechanical ventilation days, ICU length of stay, and the risk of prolonged IMV.⁶⁰

CONCLUSIONS

This review provides an overview of the epidemiology and management of AHRF and refractory hypoxemia in the CICUs. Several research gaps remain. First, a better understanding of optimal induction and sedation strategies in the decompensated cardiac patient is required. Second, the current use of lung protective ventilation in the CICUs and the ideal ventilatory strategy for AHRF in the CICUs remains unknown. Finally, the approach to refractory hypoxemia in the CICU patient who is failing despite lung protective strategies remains understudied.

FUNDING SUPPORT AND AUTHOR DISCLOSURES

Dr Fan has received personal fees from ALung Technologies, Aerogen, Baxter, GE Healthcare, Inspira, and Vasomune, outside of the submitted work. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

ADDRESS FOR CORRESPONDENCE: Dr Rebecca Mathew, CAPITAL Research Group, Division of Cardiology, University of Ottawa Heart Institute, 40 Ruskin Street, Ottawa, Ontario K1Y 4W7, Canada. E-mail: rmathew@ottawaheart.ca.

REFERENCES

- Bohula EA, Katz JN, van Diepen S, et al. Demographics, care patterns, and outcomes of patients admitted to cardiac intensive care units: the critical care cardiology trials network prospective North American multicenter registry of cardiac critical illness. *JAMA Cardiol*. 2019;4:928-935.
- Metkus TS, Miller PE, Alviar CL, et al. Advanced respiratory support in the contemporary cardiac ICU. *Crit Care Explor*. 2020;2:e0182.
- Jentzer JC, Alviar CL, Miller PE, et al. Trends in therapy and outcomes associated with respiratory failure in patients admitted to the cardiac intensive care unit. *J Intensive Care Med*. 2022;37:543-554.
- Starling EH. On the absorption of fluids from the connective tissue spaces. *J Physiol*. 1986;19:312-326.
- Kohsaka S, Menon V, Lowe AM, et al. Systemic inflammatory response syndrome after acute myocardial infarction complicated by cardiogenic shock. *Arch Intern Med*. 2005;165:1643-1650.
- Orrem HL, Nilsson PH, Pischke SE, et al. Acute heart failure following myocardial infarction: complement activation correlates with the severity of heart failure in patients developing cardiogenic shock. *ESC Heart Fail*. 2018;5:292-301.
- Shpektor A. Cardiogenic shock: the role of inflammation. *Acute Card Care*. 2010;12:115-118.
- Burkhoff D, Sayer G, Doshi D, Uriel N. Hemodynamics of mechanical circulatory support. *J Am Coll Cardiol*. 2015;66:2663-2674.
- Alviar CL, Miller PE, McAreavey D, et al. Positive pressure ventilation in the cardiac intensive care unit. *J Am Coll Cardiol*. 2018;72:1532-1553.
- Beard DA, Feigl EO. Understanding Guyton's venous return curves. *Am J Physiol Heart Circ Physiol*. 2011;301:H629-H633.
- Pinsky MR. Determinants of pulmonary arterial flow variation during respiration. *J Appl Physiol Respir Environ Exerc Physiol*. 1984;56:1237-1245.
- Kirby RR, Perry JC, Calderwood HW, Ruiz BC, Lederman DS. Cardiorespiratory effects of high positive end-expiratory pressure. *Anesthesiology*. 1975;43:533-539.
- Lansdorp B, Hofhuizen C, van Lavieren M, van Swieten H, et al. Mechanical ventilation-induced intrathoracic pressure distribution and heart-lung interactions. *Crit Care Med*. 2014;42:1983-1990.
- Magder SA, Lichtenstein S, Adelman AG. Effect of negative pleural pressure on left ventricular hemodynamics. *Am J Cardiol*. 1983;52:588-593.
- Klinger JR. Hemodynamics and positive end-expiratory pressure in critically ill patients. *Crit Care Clin*. 1996;12:841-864.
- Frazier SK. Neurohormonal responses during positive pressure mechanical ventilation. *Heart Lung*. 1999;28:149-167.
- Yarnell CJ, Angriman F, Ferreyro BL, et al. Oxygenation thresholds for invasive ventilation in hypoxemic respiratory failure: a target trial emulation in two cohorts. *Crit Care*. 2023;27:67.
- Kangelaris KN, Ware LB, Wang CY, et al. Timing of intubation and clinical outcomes in adults with acute respiratory distress syndrome. *Crit Care Med*. 2016;44:120-129.
- Bauer PR, Gajic O, Nanchal R, et al. Association between timing of intubation and outcome in critically ill patients: a secondary analysis of the ICON audit. *J Crit Care*. 2017;42:1-5.
- Thomas A, Banna S, Shahu A, et al. Propofol vs etomidate for induction prior to invasive mechanical ventilation in patients with acute myocardial infarction. *Am Heart J*. 2024;272:116-125.
- Acquisto NM, Mosier JM, Bittner EA, et al. Society of critical care medicine clinical practice guidelines for rapid sequence intubation in the critically ill adult patient. *Crit Care Med*. 2023;51:1411-1430.
- Devlin JW, Skrobik Y, Gélinas C, et al. Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med*. 2018;46:e825-e873.
- Alderman EL, Barry WH, Graham AF, Harrison DC. Hemodynamic effects of morphine and pentazocine differ in cardiac patients. *N Engl J Med*. 1972;287:623-627.
- Schenone AL, Chen K, Andress K, Militello M, Cho L. Editor's Choice- Sedation in the coronary intensive care unit: an adapted algorithm for critically ill cardiovascular patient. *Eur Heart J Acute Cardiovasc Care*. 2019;8:167-175.
- Pham T, Brochard LJ, Slutsky AS. Mechanical ventilation: state of the art. *Mayo Clin Proc*. 2017;92:1382-1400.
- Putensen C, Theuerkauf N, Zinserling J, Wrigge H, Pelosi P. Meta-analysis: ventilation strategies and outcomes of the acute respiratory distress syndrome and acute lung injury. *Ann Intern Med*. 2009;151:566-576.
- Grasselli G, Calfee CS, Camporota L, et al. ESICM guidelines on acute respiratory distress syndrome: definition, phenotyping and respiratory support strategies. *Intensive Care Med*. 2023;49:727-759.
- Acute Respiratory Distress Syndrome Network, Brower RG, Matthay MA. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med*. 2000;342:1301-1308.
- Simonis FD, Serpa Neto A, Binnekade JM, et al. Effect of a low vs intermediate tidal volume strategy on ventilator-free days in intensive care unit patients without ARDS: a randomized clinical trial. *JAMA*. 2018;320:1872-1880.
- Ball L, Serpa Neto A, Trifiletti V, et al. Effects of higher PEEP and recruitment manoeuvres on mortality in patients with ARDS: a systematic review, meta-analysis, meta-regression and trial sequential analysis of randomized controlled trials. *Intensive Care Med Exp*. 2020;8:39.
- Shorofsky M, D J, F L, R H, Lipes J. Mechanical ventilation with high tidal volume and associated mortality in the cardiac intensive care unit. *Acute Card Care*. 2014;16:9-14.
- Rali AS, Tran LE, Auvil B, et al. Modifiable mechanical ventilation targets are associated with improved survival in ventilated VA-ECLS patients. *JACC Heart Fail*. 2023;11:961-968.
- Wiesen J, Ornstein M, Tonelli AR, Menon V, Ashton RW. State of the evidence: mechanical ventilation with PEEP in patients with cardiogenic shock. *Heart*. 2013;99:1812-1817.
- de Wit M, Miller KB, Green DA, Ostman HE, Gennings C, Epstein SK. Ineffective triggering predicts increased duration of mechanical ventilation. *Crit Care Med*. 2009;37:2740-2745.
- Sottile PD, Albers D, Smith BJ, Moss MM. Ventilator dyssynchrony - detection, pathophysiology, and clinical relevance: a Narrative review. *Ann Thorac Med*. 2020;15:190-198.
- Patzelt J, Zhang Y, Seizer P, et al. Effects of mechanical ventilation on heart geometry and mitral valve leaflet coaptation during percutaneous edge-to-edge mitral valve repair. *JACC Cardiovasc Interv*. 2016;25:151-159.
- Amato MB, Meade MO, Slutsky AS, et al. Driving pressure and survival in the acute respiratory distress syndrome. *N Engl J Med*. 2015;372:747-755.
- Aoyama H, Yamada Y, Fan E. The future of driving pressure: a primary goal for mechanical ventilation? *J Intensive Care*. 2018;6:64.
- Goligher E. Platform of randomized adaptive clinical trials in critical illness (PRACTICAL) randomized control trial, Invasive Mechanical Ventilation (IMV) strategies domain. 2024. Accessed June 11, 2024. <https://practical->

platform.org/assets/images/uploads/doc/IMV_Strategies_Domain_Protocol.v.1.0_22Dec2022_.pdf

40. Alhurani RE, Oeckler RA, Franco PM, Jenkins SM, Gajic O, Pannu SR. Refractory hypoxemia and use of rescue strategies. A U.S. National survey of adult intensivists. *Ann Am Thorac Soc*. 2016;13:1105-1114.
41. Jozwiak M, Teboul J, Anguel N, et al. Beneficial hemodynamic effects of prone positioning in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2013;188:1428-1433.
42. Mullen L, Byrd D. Using simulation training to improve perioperative patient safety. *AORN J*. 2013;97:419-427.
43. Plens G, Droghe M, Alcala G, et al. Expiratory muscle activity counteracts positive end-expiratory pressure and is associated with fentanyl dose in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2024;209:563-572.
44. Slutsky AS. Neuromuscular blocking agents in ARDS. *N Engl J Med*. 2010;363:1176-1180.
45. Papazian L, Forel J, Gacouin A, et al. Neuromuscular blockers in early acute respiratory distress syndrome. *N Engl J Med*. 2010;363:1107-1116.
46. Moss M, Huang DT, Brower RG, et al. Early neuromuscular blockade in the acute respiratory distress syndrome. *N Engl J Med*. 2019;380:1997-2008.
47. Qadir N, Sahetya S, Munshi L, et al. An update on management of adult patients with acute respiratory distress syndrome: an official American Thoracic Society clinical practice guideline. *Am J Respir Crit Care Med*. 2024;209:24-36.
48. Adhikari NK, Burns KE, Friedrich JO, Granton JT, Cook DJ, Meade MO. Effect of nitric oxide on oxygenation and mortality in acute lung injury: systematic review and meta-analysis. *BMJ*. 2007;334:779.
49. Inglessis I, Shin J, Lepore J, et al. Hemodynamic effects of inhaled nitric oxide in right ventricular myocardial infarction and cardiogenic shock. *J Am Coll Cardiol*. 2004;44:793-798.
50. Semigran MJ, Cockrill BA, Kacmarek R, et al. Hemodynamic effects of inhaled nitric oxide in heart failure. *J Am Coll Cardiol*. 1994;24:982-988.
51. Combes A, Hajage D, Capellier G, et al. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. *N Engl J Med*. 2018;378:1965-1975.
52. Goligher EC, Tomlinson G, Hajage D, et al. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome and posterior probability of mortality benefit in a post hoc Bayesian analysis of a randomized clinical trial. *JAMA*. 2018;320:2251-2259.
53. Levy D, Desnos C, Lebreton G, et al. Early reversal of right ventricular dysfunction after venovenous extracorporeal membrane oxygenation in patients with COVID-19 pneumonia. *Am J Respir Crit Care Med*. 2023;207:784-787.
54. Thiele H, Zeymer U, Akin I, et al. Extracorporeal life support in infarct-related cardiogenic shock. *N Engl J Med*. 2023;389:1286-1297.
55. Shahu A, Banna S, Applefeld W, et al. Liberation from mechanical ventilation in the cardiac intensive care unit. *JACC Adv*. 2023;2:100173.
56. Schmidt G, Girard T, Kress J, et al. Official executive summary of an American Thoracic Society/American College of Chest Physicians clinical practice guideline: liberation from mechanical ventilation in critically ill adults. *Am J Respir Crit Care Med*. 2017;195:115-119.
57. Pham T, Heunks L, Bellani G, et al. Weaning from mechanical ventilation in intensive care units across 50 countries (WEAN SAFE): a multicentre, prospective, observational cohort study. *Lancet Respir Med*. 2023;11:465-476.
58. Liu J, Shen F, Teboul JL, et al. Cardiac dysfunction induced by weaning from mechanical ventilation: incidence, risk factors, and effects of fluid removal. *Crit Care*. 2016;20:369.
59. Boles J, Bion J, Connors A, et al. Weaning from mechanical ventilation. *Eur Respir J*. 2007;29:1033-1056.
60. Blackwood B, Alderdice F, Burns K, Cardwell C, Lavery G, O'Halloran P. Protocolized versus non-protocolized weaning for reducing the duration of mechanical ventilation in critically ill adult patients. *Cochrane Database Syst Rev*. 2010:CD006904.

KEY WORDS cardiac intensive care, heart-lung interactions, invasive mechanical ventilation