

## ORIGINAL RESEARCH

# Effects of Sacubitril/Valsartan According to Natriuretic Peptide Levels in Patients Enrolled in PARADIGM-HF and PARAGON-HF

Toru Kondo, MD, PhD,<sup>a,b</sup> Pardeep S. Jhund, MB ChB, MSc, PhD,<sup>a</sup> Inder S. Anand, MD, DPHIL,<sup>c</sup> Brian L. Claggett, PhD,<sup>d</sup> Akshay S. Desai, MD, MPH,<sup>d</sup> Kieran F. Docherty, MB ChB, PhD,<sup>a</sup> Carolyn S.P. Lam, MBBS, PhD,<sup>e</sup> Martin P. Lefkowitz, MD,<sup>f</sup> Aldo P. Maggioni, MD,<sup>g</sup> Felipe A. Martinez, MD,<sup>h</sup> Margaret M. Redfield, MD,<sup>i</sup> Jean L. Rouleau, MD,<sup>j</sup> Dirk J. Van Veldhuisen, MD, PhD,<sup>k</sup> Faiez Zannad, MD, PhD,<sup>l</sup> Michael R. Zile, MD,<sup>m</sup> Milton Packer, MD,<sup>n</sup> Scott D. Solomon, MD,<sup>d</sup> John J.V. McMurray, MD<sup>a</sup>

## ABSTRACT

**BACKGROUND** Recent trials of new heart failure (HF) treatments suggest the effect of therapy may vary by N-terminal pro-B type natriuretic peptide (NT-proBNP) level.

**OBJECTIVES** The authors examined the efficacy of sacubitril/valsartan according to NT-proBNP levels in patients with reduced, mildly reduced, and preserved left ventricular ejection fraction (LVEF) enrolled in PARADIGM-HF (Prospective Comparison of Angiotensin Receptor–Neprilysin Inhibitor with Angiotensin–Converting–Enzyme Inhibitor to Determine Impact on Global Mortality and Morbidity in Heart Failure Trial) and PARAGON-HF (Prospective Comparison of Angiotensin Receptor–Neprilysin Inhibitor with Angiotensin–Receptor Blockers Global Outcomes in HF with Preserved Ejection Fraction).

**METHODS** Individual patient data from PARADIGM-HF and PARAGON-HF were pooled and participants were divided into categories defined by quintiles of NT-proBNP level. The primary outcome examined was the composite of HF hospitalization or cardiovascular death.

**RESULTS** Among the 13,195 patients enrolled in both trials, 13,142 (99.6%) had a baseline NT-proBNP level measured. The rate of the primary outcome (per 100 person-years) increased with NT-proBNP levels: quintile 1, 5.9 (95% CI: 5.3–6.5); quintile 2, 7.5 (95% CI: 6.9–8.2); quintile 3, 9.0 (95% CI: 8.2–9.7); quintile 4, 12.0 (95% CI: 11.1–12.9); and quintile 5, 20.8 (95% CI: 19.6–22.2). The relative risk reduction in the primary outcome with sacubitril/valsartan was consistent across NT-proBNP levels: the HR in quintile 1 was 0.79 (95% CI: 0.65–0.96); quintile 2, 0.87 (95% CI: 0.72–1.04); quintile 3, 0.79 (95% CI: 0.66–0.93); quintile 4, 0.85 (95% CI: 0.73–0.99); and quintile 5, 0.86 (95% CI: 0.76–0.97; *P* for interaction = 0.86). The absolute risk reduction was greatest in NT-proBNP quintile 5; the number needed to treat for the primary outcome over the median follow-up of 31 months was 16 in quintile 5 vs 37 in quintile 1.

**CONCLUSIONS** The relative risk reductions with sacubitril/valsartan were consistent irrespective of NT-proBNP level in HF patients across the range of LVEF. Consequently, the absolute risk reductions were greatest in patients with higher NT-proBNP levels. (PARADIGM-HF; [NCT01035255](https://doi.org/10.1016/j.jchf.2019.03.001); and PARAGON-HF; [NCT01920711](https://doi.org/10.1016/j.jchf.2020.03.001)). (JACC Heart Fail. 2025;■:■–■)  
© 2025 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

ABBREVIATIONS  
AND ACRONYMS**BNP** = B-type natriuretic peptide**HF** = heart failure**LVEF** = left ventricular ejection fraction**NP** = natriuretic peptide**NT-proBNP** = N-terminal pro-B-type natriuretic peptide

Natriuretic peptides (NPs) are commonly used in clinical practice to help diagnose heart failure (HF), assess prognosis, and evaluate response to treatment.<sup>1,2</sup> It has recently been reported that the benefits of the soluble guanylate cyclase stimulator vericiguat were attenuated in patients with a higher N-terminal pro-B-type natriuretic peptide (NT-proBNP) level.<sup>3,4</sup> The latter observation raises the question as to whether high NP levels identify patients with HF so advanced that they might not respond to vericiguat and perhaps other treatments. This hypothesis is potentially supported by older findings from I-PRESERVE (Irbesartan in Heart Failure with Preserved Ejection Fraction Study) and TOPCAT (Treatment of Preserved Cardiac Function Heart Failure with an Aldosterone Antagonist), although not by newer findings with cardiac-specific myosin activator omecamtiv mecarbil which appeared to have greater benefits in patients with higher NT-proBNP levels.<sup>5,6</sup> Alternatively, the attenuated effect of vericiguat in patients with high NP levels could result from crosstalk between the particulate and soluble guanylate cyclase pathways, which has been described in experimental animals and humans.<sup>7,8</sup> This finding with vericiguat also raises a question about whether the effect of sacubitril/valsartan varies according to NP level as neprilysin inhibition increases NPs (acting through particulate guanylate cyclase) and other vasoactive peptides some of which (eg, bradykinin) act through the nitric oxide-soluble guanylate cyclase pathway. To investigate this further, we examined the effect of sacubitril/valsartan according to baseline NT-proBNP level using pooled patient-level data from the PARADIGM-HF (Prospective Comparison of Angiotensin Receptor-Neprilysin Inhibitor with Angiotensin-Converting-Enzyme Inhibitor to Determine

Impact on Global Mortality and Morbidity in Heart Failure Trial; [NCT01035255](#)) trial and the PARAGON-HF (Efficacy and Safety of LCZ696 Compared to Valsartan, on Morbidity and Mortality in Heart Failure Patients With Preserved Ejection Fraction; [NCT01920711](#)).<sup>9,10</sup> In this large data set, we describe patient characteristics according to NT-proBNP level across the range of left ventricular ejection fraction (LVEF), outcomes related to NT-proBNP level adjusted for baseline characteristics, and the relative and absolute risk reductions with a neprilysin inhibitor plus a renin-angiotensin blocker (ie, sacubitril/valsartan) compared to a renin-angiotensin blocker alone (ie, enalapril or valsartan).

## METHODS

The design and primary results of PARADIGM-HF and PARAGON-HF are already published.<sup>9-11</sup> Each trial was approved by local ethics committees and written informed consent was obtained from each patient.

**STUDY PATIENTS.** Briefly, PARADIGM-HF included 8,399 patients with an LVEF  $\leq 40\%$  (later changed to  $\leq 35\%$  by amendment). Patients were required to have NYHA functional class II-IV, and an NT-proBNP  $\geq 600$  pg/mL (or B-type natriuretic peptide [BNP]  $\geq 150$  pg/mL) or NT-proBNP  $\geq 400$  pg/mL (or BNP  $\geq 100$  pg/mL) if hospitalized for HF within the last 12 months. Patients were also required to be treated with a stable dose of a beta-blocker and an angiotensin-converting enzyme inhibitor (or angiotensin receptor blocker) equivalent to at least 10 mg of enalapril. Patients entered a single-blind run-in phase and received enalapril 10 mg twice daily for 2 weeks. If no unacceptable side effects occurred, they were switched to single-blind treatment with sacubitril/valsartan for an additional 4 to 6 weeks (49/51 mg twice daily, increasing to 97/103 mg twice daily). Thereafter, patients were randomized in a 1:1

From the <sup>a</sup>British Heart Foundation Cardiovascular Research Centre, University of Glasgow, Glasgow, United Kingdom;

<sup>b</sup>Department of Cardiology, Nagoya University Graduate School of Medicine, Nagoya, Japan; <sup>c</sup>VA Medical Center and University of Minnesota, Minneapolis, Minnesota; <sup>d</sup>Cardiovascular Division, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA; <sup>e</sup>National Heart Centre Singapore and Duke-National University of Singapore, Singapore; <sup>f</sup>Novartis, East Hanover, New Jersey, USA; <sup>g</sup>ANMCO Research Center, Florence, Italy; <sup>h</sup>University of Cordoba, Cordoba, Argentina; <sup>i</sup>Department of Cardiovascular Medicine, Mayo Clinic, Rochester, Minnesota, USA; <sup>j</sup>Institut de Cardiologie de Montréal, Université de Montréal, Montréal, Quebec, Canada; <sup>k</sup>Department of Cardiology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands; <sup>l</sup>Investigation Network Initiative-Cardiovascular and Renal Clinical Trialists, University Hospital, University of Lorraine, Nancy, France; <sup>m</sup>Department of Medicine, Medical University of South Carolina, Charleston, South Carolina, USA; and the <sup>n</sup>Baylor Heart and Vascular Institute, Baylor University Medical Center, Dallas, Texas, USA.

Biyyem Bozkurt, MD, PhD, served as acting Editor-in-Chief and main adjudicator for this paper.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

Manuscript received June 28, 2024; revised manuscript received November 19, 2024, accepted December 13, 2024.

ratio to double-blind treatment with either sacubitril/valsartan 97/103 mg twice daily or enalapril 10 mg twice daily. The primary outcome was the composite of HF hospitalization or cardiovascular death. The median follow-up was 27 months.

PARAGON-HF included 4,796 patients who were  $\geq 50$  years of age and who had an LVEF of 45% or higher. Patients were required to have signs and symptoms of HF, NYHA functional class II-IV, evidence of structural heart disease, and to be treated with a diuretic. For patients hospitalized for HF within 9 months and those not in atrial fibrillation (AF) on screening electrocardiography (ECG) were required to have an NT-proBNP  $\geq 200$  pg/mL, and those in AF were required to have an NT-proBNP level  $\geq 600$  pg/mL. For patients without hospitalization for HF within 9 months, those not in AF on screening ECG were required to have an NT-proBNP  $\geq 300$  pg/mL, and those in AF were required to have an NT-proBNP  $\geq 900$  pg/mL. Patients entered a single-blind run-in phase and received valsartan 40 mg or 80 mg twice daily for 1 to 2 weeks. If no unacceptable side effects occurred, they were switched to single-blind treatment with sacubitril/valsartan 49/51 mg twice daily for 2 to 4 weeks. Thereafter, patients were randomized in a 1:1 ratio to double-blind treatment with either sacubitril/valsartan 97/103 mg twice daily or valsartan 160 mg twice daily. The primary outcome was the composite of total (first and recurrent) hospitalizations for HF and cardiovascular death. The median follow-up was 35 months.

**NT-proBNP EVALUATION.** NT-proBNP was measured in both trials using samples collected at screening (before the run-in period), using the same assay (Roche Elecsys) in a central laboratory.<sup>12,13</sup> Patients without a baseline NT-proBNP measurement were excluded from this analysis.

**STUDY OUTCOMES.** For this analysis, the primary outcome used was the composite of time to the first hospitalization for HF or cardiovascular death. HF hospitalization, cardiovascular death, all-cause death, total (first and recurrent) HF hospitalizations, and the composite of total HF hospitalizations and cardiovascular death were also evaluated as secondary outcomes. All these events were centrally adjudicated by the same clinical events committee in both trials.

**STATISTICAL ANALYSES.** Baseline characteristics were summarized as mean  $\pm$  SD or median (Q1-Q3) for continuous variables and counts and percentages for categorical variables according to the quintiles of NT-proBNP levels. Differences in baseline characteristics were tested using the Cochran-Armitage trend

test for binary variables and the Jonckheere Terpstra test for continuous variables, respectively.

To evaluate the association between NT-proBNP levels and outcomes, time-to-event data were evaluated using the Kaplan-Meier estimator, and Cox proportional-hazards models, stratified by region and trial, and adjusted for randomized treatment, and HRs with 95% CIs were computed for the quintiles of NT-proBNP. Total (first and recurrent) events were evaluated with semiparametric proportional-rates models, stratified by region and trial, and adjusted for randomized treatment, and rate ratios with 95% CIs were computed. In addition, HRs and rate ratios stratified by region and trial and adjusted for randomized treatment, age, sex, race, body mass index (BMI), systolic blood pressure, heart rate, NYHA functional class III or IV, time from diagnosis of HF, prior HF hospitalization, LVEF, hypertension, history of myocardial infarction, history of stroke, AF rhythm on ECG at screening, and estimated glomerular filtration rate (eGFR) were reported.

To examine the effects of sacubitril/valsartan compared to enalapril or valsartan according to NT-proBNP levels, time-to-first event was evaluated with Cox proportional-hazards models and total events were evaluated with semiparametric proportional-rates models, respectively, and these models were stratified according to trial and region. In addition to the quintile categories of NT-proBNP, these estimates were obtained for patients with NT-proBNP  $< 1,000$  pg/mL or  $\geq 1,000$  pg/mL. The effect of randomized treatment across the range of NT-proBNP (log-transformed) levels as a continuous variable was modelled flexibly using restricted cubic splines with 3 knots (10th, 50th, and 90th percentile of log-transformed NT-proBNP). The effects of sacubitril/valsartan were also examined for subgroups of interest (males and females; nonobese and obese; no AF and AF on screening electrocardiography; eGFR  $\geq 60$  mL/min/1.73 m<sup>2</sup> and eGFR  $< 60$  mL/min/1.73 m<sup>2</sup>). Regarding BMI, analysis for treatment effect was explored in more detailed subgroups (ie,  $< 25.0$  kg/m<sup>2</sup>, 25.0-29.9 kg/m<sup>2</sup>, 30-34.9 kg/m<sup>2</sup>, and  $\geq 35.0$  kg/m<sup>2</sup>). By applying a consistent relative risk reduction with sacubitril/valsartan (observed in the overall population) to event rates seen in the enalapril/valsartan treated population, differences in the incidence rate of the primary outcome were calculated continuously across the spectrum of log-transformed NT-proBNP. All analyses were conducted based on complete case analysis.

All analyses were conducted using STATA version 17.0.

**TABLE 1** Baseline Characteristics According to Categories Defined by Quintiles of NT-proBNP Level at Baseline

	Quintile 1 ≤620 pg/mL (n = 2,634)	Quintile 2 621-1,039 pg/mL (n = 2,627)	Quintile 3 1,040-1,654 pg/mL (n = 2,625)	Quintile 4 1,655-2,995 pg/mL (n = 2,628)	Quintile 5 ≥2,996 pg/mL (n = 2,628)	P Value
Age, y	67.8 ± 10.6	65.9 ± 11.3	67.6 ± 10.9	67.7 ± 11.5	66.2 ± 11.8	0.14
Region						
North America	246 (9.3)	266 (10.1)	224 (8.5)	221 (8.4)	193 (7.3)	0.001
Latin America	281 (10.7)	354 (13.5)	334 (12.7)	366 (13.9)	466 (17.7)	<0.001
Western Europe	670 (25.4)	706 (26.9)	772 (29.4)	688 (26.2)	586 (22.3)	0.010
Central Europe	1,002 (38.0)	875 (33.3)	881 (33.6)	902 (34.3)	868 (33.0)	0.002
Asia-Pacific	435 (16.5)	426 (16.2)	414 (15.8)	451 (17.2)	515 (19.6)	0.002
Male	1,484 (56.3)	1,815 (69.1)	1,778 (67.7)	1,808 (68.8)	1,975 (75.2)	<0.001
Race						
White	2,016 (76.5)	1,905 (72.5)	1,950 (74.3)	1,871 (71.2)	1,666 (63.4)	<0.001
Black	83 (3.2)	119 (4.5)	102 (3.9)	90 (3.4)	132 (5.0)	0.029
Asian	378 (14.4)	402 (15.3)	383 (14.6)	437 (16.6)	511 (19.4)	<0.001
Other race or missing data	157 (6.0)	201 (7.7)	190 (7.2)	230 (8.8)	319 (12.1)	<0.001
Body-mass index, kg/m <sup>2</sup>	30.5 ± 5.3	29.5 ± 5.5	29.3 ± 5.3	28.4 ± 5.2	26.8 ± 5.1	<0.001
Weight category						
Underweight	11 (0.4)	25 (1.0)	21 (0.8)	37 (1.4)	72 (2.7)	<0.001
Normal	380 (14.4)	524 (20.0)	515 (19.6)	636 (24.2)	947 (36.1)	<0.001
Overweight	915 (34.8)	947 (36.1)	1,016 (38.7)	1,044 (39.8)	991 (37.8)	0.001
Obese	1,327 (50.4)	1,130 (43.0)	1,071 (40.8)	907 (34.6)	614 (23.4)	<0.001
Vital signs						
Systolic blood pressure, mm Hg	128.6 ± 15.9	124.6 ± 15.7	124.2 ± 15.6	124.2 ± 16.1	121.8 ± 15.8	<0.001
Diastolic blood pressure, mm Hg	74.4 ± 9.9	73.7 ± 10.2	74.0 ± 10.3	73.9 ± 10.4	73.2 ± 10.4	<0.001
Heart rate, beats/min	69.2 ± 11.2	70.7 ± 11.8	71.8 ± 12.3	72.6 ± 12.4	74.1 ± 12.5	<0.001
HF characteristics						
NYHA functional class III or IV	508 (19.3)	506 (19.3)	558 (21.3)	658 (25.1)	782 (29.8)	<0.001
Time from diagnosis of HF, y						
≤1	1,068 (40.7)	910 (34.7)	856 (32.6)	817 (31.1)	831 (31.6)	<0.001
1-5	908 (34.6)	921 (35.1)	971 (37.0)	1,019 (38.8)	1,064 (40.5)	<0.001
≥5	651 (24.8)	793 (30.2)	797 (30.4)	792 (30.1)	732 (27.9)	0.029
Prior HF hospitalization	1,531 (58.1)	1,383 (52.6)	1,406 (53.6)	1,531 (58.3)	1,702 (64.8)	<0.001
Ischemic cause	1,263 (48.0)	1,375 (52.3)	1,326 (50.5)	1,392 (53.0)	1,385 (52.7)	0.001
LVEF	49.0 ± 15.1	40.2 ± 14.7	39.7 ± 14.6	37.6 ± 14.1	31.7 ± 11.6	<0.001
Clinical history						
Atrial fibrillation	752 (28.6)	938 (35.7)	1,336 (50.9)	1,386 (52.7)	1,211 (46.1)	<0.001
Myocardial infarction	869 (33.0)	1,042 (39.7)	908 (34.6)	933 (35.5)	953 (36.3)	0.42
PCI or CABG	830 (31.5)	858 (32.7)	819 (31.2)	799 (30.4)	636 (24.2)	<0.001
Hypertension	2,305 (87.5)	2,075 (79.0)	2,084 (79.4)	2,110 (80.3)	1,903 (72.4)	<0.001
Diabetes	1,095 (41.6)	1,006 (38.3)	1,006 (38.3)	944 (35.9)	898 (34.2)	<0.001
Stroke	213 (8.1)	218 (8.3)	266 (10.1)	268 (10.2)	264 (10.1)	0.001
COPD	353 (13.4)	328 (12.5)	327 (12.5)	357 (13.6)	377 (14.3)	0.16
ECG findings						
AF in screening ECG	162 (6.2)	521 (20.0)	1,006 (38.5)	1,091 (41.8)	928 (35.5)	<0.001
Heart rate in ECG, beats/min	67.8 ± 12.6	70.4 ± 14.0	72.9 ± 14.8	74.2 ± 15.0	77.3 ± 16.5	<0.001
Heart rate without AF on screening ECG, beats/min	66.9 ± 11.7	67.9 ± 12.3	69.6 ± 13.2	71.4 ± 13.6	75.4 ± 14.7	<0.001
Heart rate with AF on screening ECG, beats/min	81.8 ± 16.7	80.5 ± 15.9	78.1 ± 15.7	78.2 ± 16.0	80.9 ± 18.7	0.95
Laboratory values <sup>a</sup>						
NT-proBNP, pg/mL	424 (327-516)	807 (703-918)	1,307 (1,163-1,472)	2,140 (1,877-2,513)	5,017 (3,735-7,691)	<0.001
In patients without AF on screening ECG	420 (327-513)	788 (698-900)	1,293 (1,150-1,462)	2,153 (1,894-2,523)	5,238 (3,799-7,999)	<0.001
In patients with AF on screening ECG	479 (291-554)	875 (738-959)	1,328 (1,181-1,484)	2,121 (1,852-2,486)	4,718 (3,647-6,780)	<0.001
eGFR, mL/min/1.73 m <sup>2</sup>	69.9 ± 20.4	68.5 ± 19.4	66.8 ± 19.0	65.1 ± 18.9	62.6 ± 18.8	<0.001
eGFR <60	865 (32.9)	919 (35.0)	1,001 (38.3)	1,100 (41.9)	1,256 (47.9)	<0.001
Serum creatinine, μmol/L	90.3 ± 24.4	95.2 ± 24.5	96.3 ± 24.9	98.8 ± 25.4	105.0 ± 27.3	<0.001
Blood urea nitrogen, mmol/L	7.2 ± 2.7	7.2 ± 2.8	7.5 ± 2.8	7.7 ± 2.9	8.2 ± 3.4	<0.001
Serum sodium, mmol/L	140.9 ± 3.1	140.9 ± 3.0	141.1 ± 3.3	141.1 ± 3.2	141.2 ± 3.4	<0.001
Hemoglobin, g/L	136.0 ± 15.0	139.2 ± 15.4	138.6 ± 15.6	137.6 ± 16.4	135.6 ± 16.5	0.12
Anemia <sup>b</sup>	544 (21.0)	481 (18.6)	514 (19.9)	581 (22.5)	772 (29.9)	<0.001

Continued on the next page

TABLE 1 Continued

	Quintile 1 ≤620 pg/mL (n = 2,634)	Quintile 2 621-1,039 pg/mL (n = 2,627)	Quintile 3 1,040-1,654 pg/mL (n = 2,625)	Quintile 4 1,655-2,995 pg/mL (n = 2,628)	Quintile 5 ≥2,996 pg/mL (n = 2,628)	P Value
Treatment						
Pretrial use of ACEI or ARB	2,443 (92.7)	2,472 (94.1)	2,488 (94.8)	2,499 (95.1)	2,568 (97.7)	<0.001
Beta-blocker	2,182 (82.8)	2,321 (88.4)	2,345 (89.3)	2,350 (89.4)	2,394 (91.1)	<0.001
Mineralocorticoid antagonist	925 (35.1)	1,149 (43.7)	1,179 (44.9)	1,229 (46.8)	1,411 (53.7)	<0.001
Loop diuretic agent	662 (25.1)	1,160 (44.2)	1,248 (47.5)	1,414 (53.8)	1,847 (70.3)	<0.001
Implantable cardioverter-defibrillator	116 (4.4)	285 (10.8)	283 (10.8)	281 (10.7)	294 (11.2)	<0.001

Values are mean ± SD, n (%), or median (Q1-Q3), unless otherwise indicated. Body mass index was missing in 12 cases, blood pressure and heart rate in 1 case, NYHA functional class in 15 cases, time from diagnosis of HF in 12 cases, ischemic cause in 1 case, LVEF in 1 case, history of atrial fibrillation in 4 cases, history of stroke in 10 cases, COPD in 4 cases, electrocardiography data in 66 cases, eGFR and serum creatinine in 27 cases, blood urea nitrogen in 95 cases, serum sodium in 111 cases, and hemoglobin in 209 cases. <sup>a</sup>Laboratory data at screening are shown. <sup>b</sup>Hemoglobin <130 g/L in males, <120 g/L in females.

ACEI = angiotensin-converting enzyme inhibitor; AF = atrial fibrillation; ARB = angiotensin receptor blocker; CABG = coronary artery bypass graft; COPD = chronic obstructive pulmonary disease; ECG = electrocardiography; eGFR = estimated glomerular filtration rate; HF = heart failure; LVEF = left ventricular ejection fraction; NT-proBNP = N-terminal pro-B-type natriuretic peptide; PCI = percutaneous catheter intervention.

## RESULTS

Among the 13,195 patients enrolled in PARADIGM-HF and PARAGON-HF, 14 in PARADIGM-HF and 39 in PARAGON-HF did not have a baseline NT-proBNP level measured, leaving 13,142 (99.6%) for analysis in this study. Of these, 6,570 patients were randomized to sacubitril/valsartan, and 6,572 patients were randomized to enalapril or valsartan. The median level of NT-proBNP at baseline was 1,307 pg/mL (Q1-Q3: 702-2,513 pg/mL); 1,615 pg/mL (Q1-Q3: 888-3,231 pg/mL) in PARADIGM-HF and 911 pg/mL (Q1-Q3: 464-1,613 pg/mL) in PARAGON-HF, respectively (Supplemental Figure 1). Patients were categorized by the quintile of NT-proBNP level; patients in quintile 1 had NT-proBNP level ≤620 pg/mL, those in quintile 2 621-1,039 pg/mL, those in quintile 3 1,040-1,654 pg/mL, those in quintile 4 1,655-2,995 pg/mL, and those in quintile 5 ≥2,996 pg/mL.

**PATIENT CHARACTERISTICS.** Many baseline characteristics including demographics, comorbidities, and functional status differed according to NT-proBNP level (Table 1). Patients with higher NT-proBNP levels were more often male, less frequently White (and more frequently Asian), and had a lower average BMI. Age was similar across quintiles of NT-proBNP level. Regarding clinical history, patients with higher NT-proBNP levels were more likely to have a history of AF (28.6% in quintile 1 vs 46.1% in quintile 5) and were more likely to have a history of stroke. Conversely, patients with higher NT-proBNP levels less often had a history of hypertension and diabetes mellitus.

Regarding HF characteristics, higher NT-proBNP levels were associated with higher NYHA functional class, longer duration of HF, and lower LVEF (49.0% in quintile 1 vs 31.7% in quintile 5). The median

NT-proBNP level was higher in patients with AF on their ECG at screening (1,774 pg/mL [Q1-Q3: 1,178-2,998 pg/mL] vs compared 1,481 pg/mL [Q1-Q3: 747-2,586 pg/mL] in patients without AF). Only 6.2% of patients in quintile 1 had AF on their ECG compared to 20.0% in quintile 2, 38.5% in quintile 3, 41.8% in quintile 4, and 35.5% in quintile 5.

As NT-proBNP levels increased, eGFR values decreased. Patients with higher NT-proBNP also had lower systolic blood pressure and higher heart rates.

Pretrial use of an angiotensin-converting-enzyme inhibitor or angiotensin receptor blocker, beta-blocker, mineralocorticoid receptor antagonist, loop diuretic, and implantable cardioverter-defibrillator was more common among patients with higher NT-proBNP.

**PRIMARY AND SECONDARY OUTCOMES ACCORDING TO NT-proBNP LEVELS.** The rate (per 100 patient-years) of the primary composite outcome of HF hospitalization or cardiovascular death increased with increasing NT-proBNP levels was 5.9 (95% CI: 5.3-6.5) in quintile 1; 7.5 (95% CI: 6.9-8.2) in quintile 2; 9.0 (95% CI: 8.2-9.7) in quintile 3; 12.0 (95% CI: 11.1-12.9) in quintile 4; and 20.8 (95% CI: 19.6-22.2) in quintile 5 (Table 2, Figure 1). The adjusted HR, with quintile 1 as the reference group, was 1.30 (95% CI: 1.13-1.49) for quintile 2, 1.56 (95% CI: 1.36-1.79) for quintile 3, 2.03 (95% CI: 1.77-2.32) for quintile 4, and 3.33 (95% CI: 2.91-3.81) for quintile 5. Similar trends were observed for the other outcomes.

**EFFECTS OF SACUBITRIL/VALSARTAN ON OUTCOMES ACCORDING TO BASELINE NT-proBNP LEVEL.** The benefit of sacubitril/valsartan, compared to enalapril or valsartan, was consistent across the quintile categories of NT-proBNP levels for all outcomes examined (Table 3). The HR for the primary composite outcome

**TABLE 2 Outcomes According to Quintile Categories of NT-proBNP Levels at Baseline**

	Quintile 1 ≤620 pg/mL (n = 2,634)	Quintile 2 621-1,039 pg/mL (n = 2,627)	Quintile 3 1,040-1,654 pg/mL (n = 2,625)	Quintile 4 1,655-2,995 pg/mL (n = 2,628)	Quintile 5 ≥2,996 pg/mL (n = 2,628)
<b>HF hospitalization or CV death</b>					
Number of events (%)	403 (15.3)	468 (17.8)	545 (20.8)	685 (26.1)	1,002 (38.1)
Rate (95% CI)	5.9 (5.3-6.5)	7.5 (6.9-8.2)	9.0 (8.2-9.7)	12.0 (11.1-12.9)	20.8 (19.6-22.2)
Unadjusted HR (95% CI) <sup>a</sup>	Ref.	1.27 (1.11-1.46)	1.52 (1.33-1.73)	2.02 (1.78-2.30)	3.47 (3.07-3.93)
Adjusted HR (95% CI) <sup>b</sup>	Ref.	1.30 (1.13-1.49)	1.56 (1.36-1.79)	2.03 (1.77-2.32)	3.33 (2.91-3.81)
<b>HF hospitalization</b>					
Number of events (%)	304 (11.5)	306 (11.7)	360 (13.7)	448 (17.1)	607 (23.1)
Rate (95% CI)	4.4 (3.9-4.9)	4.9 (4.4-5.5)	5.9 (5.3-6.6)	7.9 (7.2-8.6)	12.6 (11.7-13.7)
Unadjusted HR (95% CI) <sup>a</sup>	Ref.	1.17 (1.00-1.38)	1.41 (1.21-1.65)	1.90 (1.64-2.21)	3.20 (2.76-3.71)
Adjusted HR (95% CI) <sup>b</sup>	Ref.	1.21 (1.02-1.42)	1.46 (1.24-1.72)	1.93 (1.64-2.27)	3.12 (2.65-3.68)
<b>CV death</b>					
Number of events (%)	154 (5.9)	224 (8.5)	264 (10.1)	352 (13.4)	669 (25.5)
Rate (95% CI)	2.1 (1.8-2.5)	3.4 (3.0-3.9)	4.0 (3.6-4.5)	5.6 (5.0-6.2)	12.3 (11.4-13.2)
Unadjusted HR (95% CI) <sup>a</sup>	Ref.	1.45 (1.18-1.78)	1.73 (1.41-2.11)	2.32 (1.91-2.82)	4.69 (3.90-5.64)
Adjusted HR (95% CI) <sup>b</sup>	Ref.	1.45 (1.17-1.79)	1.72 (1.39-2.11)	2.24 (1.83-2.74)	4.16 (3.41-5.07)
<b>All-cause death</b>					
Number of events (%)	252 (9.6)	315 (12.0)	374 (14.3)	493 (18.8)	795 (30.3)
Rate (95% CI)	3.4 (3.0-3.9)	4.8 (4.3-5.3)	5.7 (5.1-6.3)	7.8 (7.2-8.6)	14.6 (13.6-15.6)
Unadjusted HR (95% CI) <sup>a</sup>	Ref.	1.34 (1.13-1.58)	1.59 (1.36-1.88)	2.17 (1.86-2.53)	3.90 (3.36-4.54)
Adjusted HR (95% CI) <sup>b</sup>	Ref.	1.31 (1.11-1.56)	1.54 (1.30-1.83)	2.03 (1.72-2.40)	3.37 (2.86-3.97)
<b>Total HF hospitalizations</b>					
Number of events	512	488	581	737	1,085
Rate (95% CI)	7.0 (6.1-8.1)	7.4 (6.4-8.6)	8.8 (7.8-10.1)	11.7 (10.5-13.1)	19.9 (18.1-21.9)
Unadjusted rate ratio (95% CI) <sup>a</sup>	Ref.	1.15 (0.94-1.40)	1.37 (1.14-1.66)	1.85 (1.55-2.22)	3.40 (2.85-4.06)
Adjusted rate ratio (95% CI) <sup>b</sup>	Ref.	1.19 (0.98-1.46)	1.43 (1.17-1.73)	1.91 (1.58-2.31)	3.37 (2.78-4.09)
<b>Total HF hospitalizations and CV death</b>					
Number of events	666	712	845	1,089	1,754
Rate (95% CI)	9.1 (8.1-10.3)	10.8 (9.7-12.1)	12.9 (11.7-14.2)	17.4 (15.9-19.0)	32.1 (29.9-34.6)
Unadjusted rate ratio (95% CI) <sup>a</sup>	Ref.	1.22 (1.03-1.44)	1.46 (1.24-1.70)	1.96 (1.69-2.28)	3.73 (3.21-4.33)
Additional rate ratio (95% CI) <sup>b</sup>	Ref.	1.25 (1.06-1.48)	1.50 (1.27-1.76)	1.99 (1.69-2.33)	3.58 (3.05-4.21)

Rates are given per 100 patient-years. <sup>a</sup>Baseline model adjusted for randomized treatment and stratified by region and trial. <sup>b</sup>Adjusted for randomized treatment, age, sex, race, body mass index, systolic blood pressure, heart rate, NYHA class III or IV, time from diagnosis of HF, prior HF hospitalization, left ventricular ejection fraction, hypertension, history of myocardial infarction, history of stroke, AF on electrocardiography at screening, and eGFR, and stratified by region and trial.

CV = cardiovascular; Ref. = Reference; other abbreviations as in Table 1.


in the overall pooled population was 0.84 (95% CI: 0.78-0.90), and it was 0.79 (95% CI: 0.65-0.96) in quintile 1; 0.87 (95% CI: 0.72-1.04) in quintile 2; 0.79 (95% CI: 0.66-0.93) in quintile 3; 0.85 (95% CI: 0.73-0.99) in quintile 4; and 0.86 (95% CI: 0.76-0.97) in quintile 5, with *P* for interaction = 0.86. The consistent benefit was similarly observed if patients were divided according to NT-proBNP <1,000 pg/mL or ≥1,000 pg/mL (Supplemental Table 1). Examined as a continuous variable, there was no interaction between NT-proBNP and the effect of randomized treatment on the primary outcome (*P* = 0.49) (Figure 2). The absolute rate difference between sacubitril/valsartan and enalapril or valsartan was greater in patients with higher NT-proBNP levels as a result of the higher event rate among people with a higher NT-proBNP level (Central Illustration). The

number of patients needed to be treated (NNT) over the median trial duration of 31 months (Q1-Q3: 22-38 months) to prevent 1 patient from experiencing the primary endpoint was 37 (95% CI: 27-59) in quintile 1 and 16 (95% CI: 12-25) in quintile 5 of NT-proBNP.

The effects of sacubitril/valsartan on the other outcomes of interest were also consistent across quintiles of NT-proBNP (*P* for interaction for all outcomes ≥0.5) (Table 3). The results were also consistent when NT-proBNP was modelled as a continuous variable (*P* for interaction for all outcomes > 0.1) (Figure 2).

**EFFECTS OF SACUBITRIL/VALSARTAN ON HF HOSPITALIZATION OR CARDIOVASCULAR DEATH ACCORDING TO NT-proBNP LEVELS IN SUBGROUPS OF INTEREST.** The effect of sacubitril/valsartan, compared to enalapril



**FIGURE 1** Kaplan-Meier Curves for Each Outcome According to Quintiles of NT-proBNP

Kaplan-Meier curves according to quintiles of N-terminal pro-B-type natriuretic peptide (NT-proBNP) for the composite of hospitalization or cardiovascular (CV) death (A), heart failure (HF) hospitalization (B), CV death (C), and all-cause death (D).

or valsartan, on the primary outcome, according to NT-proBNP level in the interest subgroups (male vs female; no AF vs AF; obese vs nonobese; and  $\text{eGFR} \geq 60 \text{ mL/min/1.73 m}^2$  vs  $\text{eGFR} < 60 \text{ mL/min/1.73 m}^2$ ) are shown in [Supplemental Table 2](#). There was no significant interaction between quintiles of NT-proBNP and the effect of randomized treatment on the primary outcome in each subgroup of interest ( $P$  for interaction for all outcomes  $> 0.5$ ). Even when analyzed in more detailed subgroups of BMI, the consistent benefit was similarly observed ([Supplemental Table 3](#)). The results were also consistent when NT-proBNP was modelled as a continuous variable ( $P$  for interaction for all outcomes  $> 0.2$ ) ([Figure 3](#)).


## DISCUSSION

As expected, in this pooled analysis of PARADIGM-HF and PARAGON-HF, demographics, comorbidities, and HF status varied according to NT-proBNP level at baseline. Also as expected, higher NT-proBNP levels were associated with a greater risk of clinical outcomes in the overall HF population, although this has rarely been demonstrated across such a wide range of LVEFs. NT-proBNP appears useful for risk stratification in patients HF, irrespective of LVEF phenotype, and those in the highest NT-proBNP quintile had a 3-times higher risk of HF hospitalization or cardiovascular death compared to those in the lowest quintile, even after adjustment for other prognostic

**TABLE 3** Effect of Sacubitril/Valsartan on Outcomes According to Quintile Categories of NT-proBNP Levels at Baseline

	Quintile 1 ≤620 pg/mL		Quintile 2 621-1,039 pg/mL		Quintile 3 1,040-1,654 pg/mL	
	Sacubitril/Valsartan (n = 1,313)	Enalapril or Valsartan (n = 1,321)	Sacubitril/Valsartan (n = 1,317)	Enalapril or Valsartan (n = 1,310)	Sacubitril/Valsartan (n = 1,308)	Enalapril or Valsartan (n = 1,317)
HF hospitalization or CV death						
Number of events (%)	178 (13.6)	225 (17.0)	224 (17.0)	244 (18.6)	244 (18.7)	301 (22.9)
Rate (95% CI)	5.1 (4.4-5.9)	6.6 (5.8-7.5)	7.1 (6.2-8.1)	8.0 (7.0-9.0)	7.9 (7.0-9.0)	10.0 (8.9-11.2)
HR (95% CI) <sup>a</sup>	0.79 (0.65-0.96)		0.87 (0.72-1.04)		0.79 (0.66-0.93)	
HF hospitalization						
Number of events (%)	127 (9.7)	177 (13.4)	146 (11.1)	160 (12.2)	163 (12.5)	197 (15.0)
Rate (95% CI)	3.7 (3.1-4.4)	5.2 (4.5-6.0)	4.6 (3.9-5.4)	5.2 (4.5-6.1)	5.3 (4.5-6.2)	6.5 (5.7-7.5)
HR (95% CI) <sup>a</sup>	0.72 (0.57-0.90)		0.84 (0.67-1.06)		0.80 (0.65-0.98)	
CV death						
Number of events (%)	68 (5.2)	86 (6.5)	108 (8.2)	116 (8.9)	114 (8.7)	150 (11.4)
Rate (95% CI)	1.9 (1.5-2.4)	2.3 (1.9-2.9)	3.2 (2.7-3.9)	3.6 (3.0-4.3)	3.5 (2.9-4.2)	4.6 (3.9-5.4)
HR (95% CI) <sup>a</sup>	0.78 (0.57-1.08)		0.90 (0.69-1.17)		0.76 (0.59-0.97)	
All-cause death						
Number of events (%)	114 (8.7)	138 (10.5)	154 (11.7)	161 (12.3)	171 (13.1)	203 (15.4)
Rate (95% CI)	3.1 (2.6-3.7)	3.8 (3.2-4.4)	4.6 (3.9-5.4)	5.0 (4.2-5.8)	5.2 (4.5-6.0)	6.2 (5.4-7.1)
HR (95% CI) <sup>a</sup>	0.82 (0.64-1.05)		0.92 (0.74-1.15)		0.84 (0.68-1.03)	
Total HF hospitalizations						
Number of events	212	300	235	253	248	333
Rate (95% CI)	5.8 (4.6-7.4)	8.2 (6.9-9.8)	7.0 (5.8-8.6)	7.8 (6.3-9.7)	7.5 (6.3-9.1)	10.2 (8.6-12.1)
Rate ratio (95% CI) <sup>a</sup>	0.73 (0.55-0.98)		0.85 (0.63-1.14)		0.73 (0.57-0.94)	
Total HF hospitalizations and CV death						
Number of events	280	386	343	369	362	483
Rate (95% CI)	7.7 (6.4-9.3)	10.5 (9.0-12.3)	10.2 (8.8-12.0)	11.4 (9.7-13.4)	11.0 (9.5-12.8)	14.8 (12.9-17.0)
Rate ratio (95% CI) <sup>a</sup>	0.75 (0.58-0.95)		0.86 (0.69-1.09)		0.74 (0.60-0.91)	
TABLE 3 Continued						
	Quintile 4 1,655-2,995 pg/mL		Quintile 5 ≥2,996 pg/mL		P Value for Interaction	
	Sacubitril/Valsartan (n = 1,326)	Enalapril or Valsartan (n = 1,302)	Sacubitril/Valsartan (n = 1,306)	Enalapril or Valsartan (n = 1,322)		
HF hospitalization or CV death						
Number of events (%)	321 (24.2)	364 (28.0)	468 (35.8)	534 (40.4)	0.86	
Rate (95% CI)	11.1 (9.9-12.3)	13.0 (11.7-14.4)	19.2 (17.6-21.1)	22.5 (20.6-24.4)		
HR (95% CI) <sup>a</sup>	0.85 (0.73-0.99)		0.86 (0.76-0.97)			
HF hospitalization						
Number of events (%)	209 (15.8)	239 (18.4)	293 (22.4)	314 (23.8)	0.54	
Rate (95% CI)	7.2 (6.3-8.2)	8.5 (7.5-9.7)	12.1 (10.7-13.5)	13.2 (11.8-14.7)		
HR (95% CI) <sup>a</sup>	0.85 (0.71-1.02)		0.91 (0.77-1.06)			
CV death						
Number of events (%)	164 (12.4)	188 (14.4)	307 (23.5)	362 (27.4)	0.86	
Rate (95% CI)	5.2 (4.5-6.1)	6.0 (5.2-6.9)	11.2 (10.0-12.6)	13.3 (12.0-14.7)		
HR (95% CI) <sup>a</sup>	0.85 (0.69-1.05)		0.85 (0.73-1.00)			
All-cause death						
Number of events (%)	237 (17.9)	256 (19.7)	373 (28.6)	422 (31.9)	0.93	
Rate (95% CI)	7.5 (6.6-8.5)	8.2 (7.2-9.2)	13.6 (12.3-15.1)	15.5 (14.1-17.0)		
HR (95% CI) <sup>a</sup>	0.92 (0.77-1.10)		0.89 (0.77-1.02)			
Total HF hospitalizations						
Number of events	328	409	511	574	0.69	
Rate (95% CI)	10.4 (8.9-12.2)	13.1 (11.2-15.3)	18.7 (16.2-21.7)	21.1 (18.6-24.0)		
Rate ratio (95% CI) <sup>a</sup>	0.81 (0.65-1.01)		0.88 (0.73-1.07)			
Total HF hospitalizations and CV death						
Number of events	492	597	818	936	0.59	
Rate (95% CI)	15.6 (13.8-17.7)	19.1 (16.9-21.7)	29.9 (26.8-33.5)	34.4 (31.2-37.9)		
Rate ratio (95% CI) <sup>a</sup>	0.82 (0.69-0.98)		0.87 (0.75-1.01)			
Rates are given per 100 patient years. <sup>a</sup> Stratified by region and trial. Abbreviations as in Tables 1 and 2.						




**FIGURE 2** Treatment Effect of Sacubitril/Valsartan on Outcomes Across the Range of NT-proBNP Levels at Baseline

Effect of sacubitril/valsartan on the composite of HF hospitalization or CV death (A), HF hospitalization (B), CV death (C), all-cause death (D), total (first and recurrent) HF hospitalizations (E), composite of total HF hospitalizations and CV death (F), across the range of NT-proBNP levels at baseline. The range of NT-proBNP levels shown is between the 2nd and 98th percentile. Abbreviations as in [Figure 1](#).

variables, including LVEF. Of more interest, the relative risk reductions of all outcomes with sacubitril/valsartan were consistent across NT-proBNP levels at baseline. Consequently, there was a greater absolute risk reduction in patients with higher NT-proBNP levels. Finally, the benefits of sacubitril/valsartan were similar irrespective of NT-proBNP concentration in the subgroups defined by baseline

characteristics known to influence NT-proBNP level.<sup>1,2,14-19</sup>

As described in the introduction, an interaction has been described between NT-proBNP and the effect of 2 recently developed therapies for HF. Among patients with HFrEF and LVEF  $\leq 35\%$  enrolled in GALACTIC-HF (Global Approach to Lowering Adverse Cardiac Outcomes through Improving Contractility


**CENTRAL ILLUSTRATION** Incidence Rate and Rate Differences With Sacubitril/Valsartan Treatment for the Composite of HF Hospitalization or CV Death Across the Range of NT-proBNP Levels at Baseline

Kondo T, et al. JACC Heart Fail. 2025;■(■):■-■.

Incidence rates for the composite of heart failure (HF) hospitalization or cardiovascular (CV) death were calculated in patients with HF with reduced ejection fraction/HF with mildly reduced ejection fraction/HF with preserved ejection fraction across the range of N-terminal pro-B-type natriuretic peptide (NT-proBNP) levels at baseline. Rate differences for the incidence rate of the composite of HF hospitalization or CV death are calculated by applying a consistent relative risk reduction with sacubitril/valsartan (observed in the overall population) to participants in the control group across the range of NT-proBNP levels at baseline. The range of NT-proBNP levels shown is between the 2nd and 98th percentile.

in Heart Failure), the median NT-proBNP was 1,998 pg/mL (Q1-Q3: 993-4,079 pg/mL). There was an interaction between baseline NT-proBNP level and the relative risk reduction in the primary endpoint (a worsening HF event or cardiovascular death) due to treatment with omecamtiv mecarbil, with a greater effect in patients with a higher NT-proBNP ( $P$  for interaction = 0.005 for NT-proBNP analyzed as a continuous variable).<sup>4</sup> Conversely, among patients with LVEF <45% participating in VICTORIA, the median NT-proBNP was 2,816 pg/mL (Q1-Q3: 1,556-5,314 pg/mL) and the relative risk reduction in the primary endpoint (hospitalization for HF or cardiovascular death) with vericiguat was less in patients with a higher NT-proBNP ( $P$  for interaction = 0.002 for NT-proBNP analyzed as a continuous variable).<sup>3</sup> Some support for the hypothesis that higher baseline NT-proBNP levels might identify “treatment non-responders” comes from analyses of TOPCAT (median NT pro-BNP 900 pg/mL [Q1-Q3: 557-1,920 pg/mL]) with spironolactone and I-PRESERVE (median NT pro-BNP 339 pg/mL [Q1-Q3: 133-964 pg/mL]) with irbesartan although these trials did not show a significant effect of treatment overall meaning interpretation of subgroup findings and interactions is even more difficult than usual.<sup>5,6</sup> No such interaction

between NT-proBNP level and treatment effect was seen in the present study (where the median NT pro-BNP was 1,307 pg/mL [Q1-Q3: 702-2,513 pg/mL]) and the relative risk reduction with sacubitril/valsartan was consistent across the full range of LVEF in patients with HF in PARADIGM-HF and PARAGON-HF. Because of this, the absolute risk reduction was larger in patients with a higher NT-proBNP level who were at greater risk and the resulting NNT over the trial duration in the highest NT-proBNP quintile was only 16 for the primary outcome compared to an NNT of 37 for patients with the lowest NT-proBNP quintile. The only other similar finding we know of is with sodium-glucose cotransporter 2 inhibitors, specifically in analyses of the DAPA-HF (Dapagliflozin And Prevention of Adverse outcomes in Heart Failure), DELIVER (Dapagliflozin Evaluation to Improve the Lives of Patients with Preserved Ejection Fraction Heart Failure), and the EMPEROR-Reduced (Empagliflozin Outcome Trial in Patients With Chronic Heart Failure With Reduced Ejection Fraction) and EMPEROR-Preserved (Empagliflozin Outcome Trial in Patients With Preserved Ejection Fraction) trials.<sup>20-23</sup> These observations are important for clinical practice as more severely ill patients (with higher NT-proBNP levels) tend to receive less guideline-

**FIGURE 3** Effect of Sacubitril/Valsartan on the Primary Outcome (HF Hospitalization or CV Death) Across the Range of NT-ProBNP Level in Subgroups of Interest

Effects of sacubitril/valsartan on the composite of HF hospitalization or CV death in males (A), females (B), nonobese (C), obese (D), no-atrial fibrillation (no-AF) (E), in AF (F), in  $eGFR \geq 60 \text{ mL/min/1.73 m}^2$  (G), and in  $eGFR < 60 \text{ mL/min/1.73 m}^2$  (H) across the range of NT-proBNP levels at baseline. The range of NT-proBNP levels shown is between the 2nd and 98th percentile.  $eGFR$  = estimated glomerular filtration rate; other abbreviations as in Figure 1.

recommended therapy yet have, potentially, the most to gain from treatment.

**STUDY LIMITATIONS.** Patients enrolled in these clinical trials were selected according to specific inclusion and exclusion criteria; therefore, our results may not be generalizable to all patients with HF in the general population. The analysis of NT-proBNP and outcomes by quintiles was determined post hoc. There were different NT-proBNP enrollment thresholds for patients with AF or prior HF hospitalization which may have affected the relationship between NT-proBNP level and outcome. As all patients had to have an entry NT-proBNP level above at least 200 pg/mL, we cannot be sure about the effect of sacubitril/valsartan in patients with an NT-proBNP concentration below this cutoff. NT-proBNP levels were higher in VICTORIA than in other trials although overlapped substantially with GALACTIC-HF. Moreover, a qualitatively similar interaction between NT-proBNP level and the effect of omecamtiv mecarbil was seen for participants enrolled in the inpatient setting (where NT-proBNP levels were much higher) and the outpatient setting. In addition, although PIONEER-HF was small and not a morbidity/mortality trial, sacubitril/valsartan improved clinical outcomes in patients with a median baseline NT-proBNP of 4,821 pg/mL (Q1-Q3: 3,109-8,767 pg/mL).<sup>24</sup>

## CONCLUSIONS

In patients with HF across the range of LVEF, higher NT-proBNP levels were associated with many adverse prognostic features, including lower BMI, worse symptoms, AF, and renal impairment. Among these patients, higher NT-proBNP levels were also associated with higher risks of HF hospitalization and cardiovascular death. The benefits of sacubitril/valsartan on these outcomes were consistent regardless of NT-proBNP levels, with greater absolute benefits in patients with higher NT-proBNP levels.

## FUNDING SUPPORT AND AUTHOR DISCLOSURES

PARADIGM-HF and PARAGON-HF were supported by Novartis. Dr Kondo is supported by the Great Britain Sasakawa Foundation and grant 20K17112 from Grant-in-Aid for Scientific Research; and he has received lecture fees from AstraZeneca KK, Abbott Japan LLC, Boehringer Ingelheim, Ono Pharmaceutical Co Ltd, Kowa Company Ltd, Kyowa Kirin Co Ltd, and Novartis Pharma KK. Dr Jhund has received grants from the British Heart Foundation Centre of Research Excellence Grant RE/18/6/34217 and the Vera Melrose Heart Failure Research Fund; speakers fees from AstraZeneca, Novartis, Alkem Metabolics, ProAdWise Communications, Sun Pharmaceuticals, and Intas Pharmaceuticals; advisory board fees from AstraZeneca, Boehringer Ingelheim, and Novartis; and research funding from AstraZeneca, Boehringer Ingelheim, and Analog Devices Inc. Dr Jhund is also the Director, Global Clinical Trial Partners (GCTP). Dr Jhund's institution, the University of Glasgow, has been remunerated for clinical

trial work from AstraZeneca, Bayer AG, Novartis, and NovoNordisk. Dr Anand has received consulting fees from Amgen, ARCA Pharma, AstraZeneca, Boehringer Ingelheim, Boston Scientific, LivaNova, and Novartis. Dr Claggett has received consulting fees from Boehringer Ingelheim. Dr Desai has received grants and personal fees from AstraZeneca during the conduct of the study; personal fees from Abbott, Biofourmis, Boston Scientific, Boehringer Ingelheim, Corvidia, DalCor Pharma, Relysa, Regeneron, and Merck; grants and personal fees from Alnylam and Novartis; and personal fees from Amgen, outside of the submitted work. Dr Docherty's institution, the University of Glasgow, has been remunerated by AstraZeneca for his work on clinical trials; he has received speaker fees from AstraZeneca, Boehringer Ingelheim, Pharmacosmos, Translational Medical Academy, and Radcliffe Cardiology; has served on Advisory Boards for Us2.ai and holds equity in the company; has served on an advisory board and served on a clinical endpoint committee for Bayer AG; and has received research grant support (paid to his institution) from AstraZeneca, Roche, Novartis, and Boehringer Ingelheim. Dr Lam has received a Clinician Scientist Award from the National Medical Research Council of Singapore; has received research support from NovoNordisk and Roche Diagnostics; has served as consultant or on the advisory board/steering committee/executive committee for Alleviant Medical, Allysta Pharma, Amgen, AnaCardio AB, Applied Therapeutics, AstraZeneca, Bayer, Boehringer Ingelheim, Boston Scientific, CardioRenal, Cytokinetics, Dharma Inc, EchoNous Inc, Eli Lilly, Impulse Dynamics, Intellia Therapeutics, Ionis Pharmaceutical, Janssen Research and Development LLC, Medscape/WebMD Global LLC, Merck, Novartis, Novo Nordisk, Prosciento Inc, Radcliffe Group Ltd, Recardio Inc, ReCor Medical, Roche Diagnostics, Sanofi, Siemens Healthcare Diagnostics, and Us2.ai; and has served as co-founder and nonexecutive director of Us2.ai. Dr Lefkowitz is an employee and shareholder of Novartis. Dr Maggioni has received fees for serving on a study committee from Bayer and Fresenius. Dr Martinez has received personal fees from AstraZeneca. Dr Redfield has served as an unpaid consultant for Novartis. Dr Rouleau has received grants and consulting fees from Novartis; and consulting fees from Abbott, AstraZeneca, MyoKardia, and Sanofi. Dr van Veldhuisen has received fees for serving on a steering committee and travel support from ARCA biopharma and Corvia Medical. Dr Zannad has received consulting fees from Amgen, Applied Therapeutics, AstraZeneca, Bayer, Boehringer, Cardior, Cereno Scientific, Cellprothera, CEVA, CVRx, Merck, Novartis, Servier, and Vifor-Pharma; and has been a founder of CardioRenal and CVCT. Dr Zile has received research funding from Novartis and has been a consultant for Novartis, Abbott, Boston Scientific, CVRx, EBR, Endotronics, Ironwood, Merck, Medtronic, and Myokardia V Wave. Dr Packer has received consulting fees from AbbVie, Akcea, Actavis, Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Cardior, Daiichi Sankyo, Gilead, Johnson and Johnson, Novo Nordisk, Pfizer, Relysa, Sanofi, Synthetic Biologics, and Theravance. Dr Solomon has received research grants from Actelion, Alnylam, Amgen, AstraZeneca, Bellerophon, Bayer, Bristol Myers Squibb, Celladon, Cytokinetics, Eidos, Gilead, GlaxoSmithKline, Ionis, Lilly, Mesoblast, MyoKardia, National Institutes of Health/NHLBI, Neurotronik, Novartis, NovoNordisk, Respicardia, Sanofi Pasteur, Theracos, US2.AI; and has received consulting fees from Abbott, Action, Akros, Alnylam, Amgen, Arena, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol Myers Squibb, Cardior, Cardurion, Corvia, Cytokinetics, Daiichi-Sankyo, GlaxoSmithKline, Lilly, Merck, Myokardia, Novartis, Roche, Theracos, Quantum Genomics, Cardurion, Janssen, Cardiac Dimensions, Tenaya, Sanofi-Pasteur, Nanaqor, Trembeau, CellPro-Thera, Moderna, American Regent, and Sarepta. Dr McMurray has received payments through Glasgow University from work on clinical trials, consulting, and other activities from Amgen, AstraZeneca, Bayer, Cardurion, Cytokinetics, GlaxoSmithKline, KBP Biosciences, and Novartis; consulting fees from Alnylam Pharma, Bayer, BMS, George Clinical PTY Ltd, Ionis Pharma, Novartis, Regeneron Pharma, and River 2 Renal Corporation; personal lecture fees from Abbott, Alkem Metabolics, AstraZeneca, Blue Ocean

Scientific Solutions Ltd, Boehringer Ingelheim, Canadian Medical and Surgical Knowledge, Emcure Pharma Ltd, Eris Lifesciences, European Academy of CME, Hikma Pharmaceuticals, Imagica health, Intas Pharma, J.B. Chemicals and Pharma Ltd, Lupin Pharma, Medscape/Heart.Org, ProAdWise Communications, Radcliffe Cardiology, Sun Pharma, The Corpus, Translation Research Group, and Translational Medicine Academy; has been a director of Global Clinical Trial Partners Ltd; and is supported by a British Heart Foundation Centre of Research Excellence Grant RE/18/6/34217 and the Vera Melrose Heart Failure Research Fund.

**ADDRESS FOR CORRESPONDENCE:** Dr John J.V. McMurray, British Heart Foundation Cardiovascular Research Centre, University of Glasgow, 126 University Place, Glasgow G12 8TA, United Kingdom. E-mail: [john.mcmurray@glasgow.ac.uk](mailto:john.mcmurray@glasgow.ac.uk). X handle: [@UoGHeartFailure](https://twitter.com/UoGHeartFailure).

## PERSPECTIVES

### COMPETENCY IN PATIENT CARE AND PROCEDURAL

**SKILLS:** The effects of sacubitril/valsartan were consistent irrespective of NT-proBNP level in patients with HF across the range of LVEF, with greater absolute risk reduction in patients with higher NT-proBNP levels who were at greater risk.

**TRANSLATIONAL OUTLOOK:** The effect of some HF drugs varies according to NP level; but this is not the case for sacubitril/valsartan. Future drugs added for HF treatment should be evaluated for whether their efficacy differs according to NP levels.

## REFERENCES

- McDonagh TA, Metra M, Adamo M, et al. 2021 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J*. 2021;42:3599-3726.
- Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association joint committee on clinical practice guidelines. *J Am Coll Cardiol*. 2022;79(17):e263-e421.
- Ezekowitz JA, O'Connor CM, Troughton RW, et al. N-terminal pro-B-type natriuretic peptide and clinical outcomes: vericiguat heart failure with reduced ejection fraction study. *JACC Heart Fail*. 2020;8:931-939.
- Docherty KF, McMurray JJV, Claggett BL, et al. Efficacy of omecamtiv mecarbil in heart failure with reduced ejection fraction according to N-terminal pro-B-type natriuretic peptide level: insights from the GALACTIC-HF trial. *Eur J Heart Fail*. 2023;25:248-259.
- Anand IS, Rector TS, Cleland JG, et al. Prognostic value of baseline plasma amino-terminal pro-brain natriuretic peptide and its interactions with irbesartan treatment effects in patients with heart failure and preserved ejection fraction: findings from the I-PRESERVE trial. *Circ Heart Fail*. 2011;4:569-577.
- Anand IS, Claggett B, Liu J, et al. Interaction between spironolactone and natriuretic peptides in patients with heart failure and preserved ejection fraction: from the TOPCAT trial. *JACC Heart Fail*. 2017;5:241-252.
- Madhani M, Okorie M, Hobbs AJ, MacAllister RJ. Reciprocal regulation of human soluble and particulate guanylate cyclases in vivo. *Br J Pharmacol*. 2006;149:797-801.
- Madhani M, Scotland RS, MacAllister RJ, Hobbs AJ. Vascular natriuretic peptide receptor-linked particulate guanylate cyclases are modulated by nitric oxide-cyclic GMP signalling. *Br J Pharmacol*. 2003;139:1289-1296.
- McMurray JJ, Packer M, Desai AS, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med*. 2014;371:993-1004.
- Solomon SD, McMurray JJV, Anand IS, et al. Angiotensin-neprilysin inhibition in heart failure with preserved ejection fraction. *N Engl J Med*. 2019;381:1609-1620.
- McMurray JJ, Packer M, Desai AS, et al. Dual angiotensin receptor and neprilysin inhibition as an alternative to angiotensin-converting enzyme inhibition in patients with chronic systolic heart failure: rationale for and design of the Prospective Comparison of ARNI With ACEI to Determine Impact on Global Mortality and Morbidity in Heart Failure trial (PARADIGM-HF). *Eur J Heart Fail*. 2013;15:1062-1073.
- Khan MS, Kristensen SL, Vaduganathan M, et al. Natriuretic peptide plasma concentrations and risk of cardiovascular versus non-cardiovascular events in heart failure with reduced ejection fraction: insights from the PARADIGM-HF and ATMOSPHERE trials. *Am Heart J*. 2021;237:45-53.
- Pabon MA, Cunningham JW, Claggett BL, et al. Natriuretic peptide-based inclusion criteria in heart failure with preserved ejection fraction clinical trials: insights from PARAGON-HF. *Eur J Heart Fail*. 2022;24:672-677.
- Mueller C, McDonald K, de Boer RA, et al. Heart Failure Association of the European Society of Cardiology practical guidance on the use of natriuretic peptide concentrations. *Eur J Heart Fail*. 2019;21:715-731.
- Lam CS, Cheng S, Choong K, et al. Influence of sex and hormone status on circulating natriuretic peptides. *J Am Coll Cardiol*. 2011;58:618-626.
- Mogensen UM, Jhund PS, Abraham WT, et al. Type of atrial fibrillation and outcomes in patients with heart failure and reduced ejection fraction. *J Am Coll Cardiol*. 2017;70:2490-2500.
- Butt JH, Kondo T, Jhund PS, et al. Atrial fibrillation and dapagliflozin efficacy in patients with preserved or mildly reduced ejection fraction. *J Am Coll Cardiol*. 2022;80:1705-1717.
- Mehra MR, Uber PA, Park MH, et al. Obesity and suppressed B-type natriuretic peptide levels in heart failure. *J Am Coll Cardiol*. 2004;43:1590-1595.
- Madamanchi C, Alhosaini H, Sumida A, Runge MS. Obesity and natriuretic peptides, BNP and NT-proBNP: mechanisms and diagnostic implications for heart failure. *Int J Cardiol*. 2014;176:611-617.
- Jhund PS, Kondo T, Butt JH, et al. Dapagliflozin across the range of ejection fraction in patients with heart failure: a patient-level, pooled meta-analysis of DAPA-HF and DELIVER. *Nat Med*. 2022;28:1956-1964.
- Butt JH, Adamson C, Docherty KF, et al. Efficacy and safety of dapagliflozin in heart failure with reduced ejection fraction according to N-terminal pro-B-type natriuretic peptide: insights from the DAPA-HF trial. *Circ Heart Fail*. 2021;14:e008837.
- Myhre PL, Vaduganathan M, Claggett BL, et al. Influence of NT-proBNP on efficacy of dapagliflozin in heart failure with mildly reduced or preserved ejection fraction. *JACC: Heart Failure*. 2022;10(12):902-913.
- Januzzi JL Jr, Zannad F, Anker SD, et al. Prognostic importance of NT-proBNP and effect of empagliflozin in the EMPEROR-Reduced trial. *J Am Coll Cardiol*. 2021;78:1321-1332.
- Velazquez EJ, Morrow DA, DeVore AD, et al. Angiotensin-neprilysin inhibition in acute decompensated heart failure. *N Engl J Med*. 2019;380:539-548.

**KEY WORDS** clinical trial, heart failure, natriuretic peptide, outcomes, sacubitril/valsartan

**APPENDIX** For supplemental tables and a figure, please see the online version of this paper.