

# Emerging EGFR-Targeted Therapy in Head and Neck Cancer

## A Review

Woochan Hwang, MBBS; Jong Seok Ahn, MBBS; Hyun Ae Jung, MD, PhD; Lori J. Wirth, MD; Jong Chul Park, MD

**IMPORTANCE** Head and neck squamous cell carcinoma (HNSCC) represents a significant global health burden with limited therapeutic options for patients with recurrent or metastatic disease. Epidermal growth factor receptor (EGFR) is overexpressed in most patients with HNSCC and represents a biologically compelling target, yet current EGFR-directed therapies have demonstrated only modest clinical benefit.

**OBSERVATIONS** This review describes the evolving landscape of EGFR-targeted therapeutics in HNSCC, including cetuximab-based combination regimens as well as novel agents, such as bispecific antibodies, antibody-drug conjugates, immune cell engagers, and adaptive cell therapies. The biological rationale behind these approaches, and the results of early-phase trials, are presented in this review. Notably, cetuximab and other EGFR-targeted therapies have demonstrated inferior efficacy in patients with human papillomavirus (HPV)-positive disease.

**CONCLUSIONS AND RELEVANCE** The emerging data of combinatorial approaches and novel EGFR-targeting therapeutic agents offer renewed optimism for patients with advanced HNSCC who have limited treatment options. Future progress will depend on novel agents leveraging a deeper understanding of tumor biology, innovative approaches to reduce on-target off-tumor toxic effects of targeting EGFR, and improving efficacy in the growing population of patients with HPV-positive HNSCC.

*JAMA Oncol.* 2025;11(11):1373-1384. doi:10.1001/jamaoncol.2025.3136  
Published online September 25, 2025.

 CME at [jamacmelookup.com](https://jamacmelookup.com)

**Author Affiliations:** Division of Hematology-Oncology, Department of Medicine, Massachusetts General Hospital, Boston (Hwang, Ahn, Jung, Wirth, Park); Lunit Inc, Seoul, Republic of Korea (Hwang, Ahn); Division of Hematology and Oncology, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Republic of Korea (Jung); Harvard Medical School, Boston, Massachusetts (Wirth, Park).

**Corresponding Author:** Jong Chul Park, MD, Massachusetts General Hospital, 55 Fruit St, Boston, MA 02114 ([jpark73@mgh.harvard.edu](mailto:jpark73@mgh.harvard.edu)).

Head and neck cancer is the seventh most common cancer worldwide, responsible for more than 660 000 new diagnoses and 325 000 deaths each year.<sup>1-3</sup> Approximately 90% of these cancers are classified as head and neck squamous cell carcinoma (HNSCC), which originates from the epithelial lining of the oral cavity, pharynx, and larynx. The incidence of HNSCC varies by region, with higher prevalence in areas of high tobacco, alcohol, and areca nut use. Recent epidemiological trends indicate a rising incidence of human papillomavirus (HPV)-related oropharyngeal cancers, while tobacco-related HNSCC has been declining consistently. HPV-positive (or p16-positive) HNSCC has a distinct molecular profile and a better prognosis compared to HPV-negative HNSCC.

Despite significant advances in the management of locally advanced HNSCC, treatment options for recurrent or metastatic (R/M) HNSCC are limited, and the prognosis remains poor. The lack of dominant driver molecular pathways and modest immune responses poses a great challenge in new drug development for HNSCC. The current standard therapeutic options for R/M HNSCC include cytotoxic chemotherapy, programmed cell death 1 (PD-1) immune checkpoint blockade therapy, and epidermal growth factor receptor (EGFR) blockade therapy. EGFR overexpression is observed in the majority of patients with HNSCC, particularly when the disease is not associated with HPV, and plays a crucial role in promoting tumor cell proliferation, survival, and metastasis, making it an ideal

target for therapeutic intervention.<sup>4</sup> While traditional EGFR-targeted therapies have shown only modest efficacy and are typically used after standard chemotherapy and immune checkpoint inhibitors (ICIs), EGFR-targeted strategies have garnered renewed interest due to the emergence of novel therapeutic modalities. This review provides a comprehensive overview of the evolving landscape of EGFR-targeted therapies in HNSCC, ranging from established traditional treatments, novel combination strategies, and emerging EGFR-targeting agents under clinical investigation.

## Discussion

### EGFR Pathways in Head and Neck Cancer

The EGFR, also referred to as ERBB1, serves as the prototype of the EGFR family, which also includes ERBB2, ERBB3, and ERBB4. Structurally, EGFR consists of the extracellular domain, the transmembrane domain, the intracellular juxtamembrane region, the intracellular tyrosine kinase domain, and the C-terminal regulatory region.

EGFR-ligand binding triggers a conformational change exposing dimerization domains. The resulting homodimerization or heterodimerization (with other ERBB receptors) activates intracellular tyrosine kinase domains through transautophosphorylation, where each receptor phosphorylates tyrosine residues on the other. These phosphorylated residues act as docking sites for

downstream signaling molecules that activate multiple signaling cascades, including the RAS/RAF/MEK/ERK (MAPK), PI3K/AKT/mTOR, PLC $\gamma$ /PKC, and JAK/STAT pathways.

EGFR overexpression occurs in 80% to 100% of patients with HNSCC, based on elevated messenger RNA, resulting from dysregulated p53, polymorphisms in intron 1 of *EGFR*, and *EGFR* amplification.<sup>5</sup> EGFR expression level is an independent negative prognostic factor in HNSCC.<sup>6,7</sup> Other pathogenic alterations include point variants, found in less than 3% of patients, with the majority occurring in exon 19, and rarely EGFRvIII, which is a variant that removes exons 2 to 7 of the extracellular domain, causing constitutive activation.<sup>8,9</sup>

### Traditional EGFR Monotherapy

Conventional therapeutics targeting EGFR, such as monoclonal antibodies and small-molecule tyrosine kinase inhibitors (TKIs), have been evaluated in patients with HNSCC. Cetuximab is a chimeric IgG1 monoclonal antibody against the ligand-binding domain of EGFR, inhibiting ligand-induced downstream signaling activation and promoting internalization and degradation of the receptor. It also induces immune-mediated cytotoxic pathways through antibody-dependent cellular cytotoxicity (ADCC) and antibody-dependent cellular phagocytosis (ADCP) via Fc $\gamma$  engagement on natural killer (NK) cells and macrophages.<sup>10</sup>

The study by Bonner et al<sup>11</sup> led to cetuximab's 2006 approval for treating patients with locally advanced HNSCC as a radiosensitizing agent, demonstrating improved locoregional control (hazard ratio [HR], 0.68 [95% CI, 0.52-0.89];  $P = .005$ ), overall survival (OS) (HR, 0.74 [95% CI, 0.57-0.97];  $P = .03$ ), and progression-free survival (PFS) (HR, 0.70 [95% CI, 0.54-0.90];  $P = .006$ ) compared to radiation therapy alone. Cetuximab monotherapy later gained approval as a second-line option after platinum-based chemotherapy, based on a phase 2 study that demonstrated a modest objective response rate (ORR) of 13%.<sup>12</sup> In 2008, the EXTREME regimen, a combination of cetuximab with platinum-based doublet chemotherapy, received approval for R/M HNSCC based on OS (HR, 0.80 [95% CI, 0.64-0.99];  $P = .04$ ) and PFS (HR, 0.54 [95% CI, 0.43-0.67];  $P < .001$ ) improvement over chemotherapy alone.<sup>13</sup> Grade 3 skin reactions were observed in 20 of 219 patients (9%) who received the EXTREME regimen compared with 1 of 215 of those who received chemotherapy alone ( $P < .001$ ).

Panitumumab is another EGFR-targeting monoclonal antibody but lacks ADCC activity owing to its IgG2-based structure. It demonstrated a modest efficacy as second-line monotherapy with an ORR of 4% (2 of 51) and a median (IQR) PFS of 1.4 (95% CI, 1.3-2.4) months.<sup>14</sup> The addition of panitumumab to platinum-based chemotherapy in the first-line setting failed to improve OS for patients with R/M HNSCC.<sup>15</sup>

Small-molecule TKIs competitively inhibit adenosine triphosphate binding at the kinase's active site in the intracellular domain, preventing the phosphorylation of downstream proteins. EGFR-targeting TKIs have been extensively evaluated in HNSCC. First-generation reversible EGFR-targeting TKIs, such as gefitinib and erlotinib, showed limited efficacy with a low ORR and no survival benefit.<sup>16,17</sup> Afatinib, a second-generation irreversible inhibitor targeting multiple ERBB family receptors (EGFR, ERBB2, ERBB4), achieved a modest ORR of 10.2% but failed to improve OS over standard chemotherapy in patients with disease refractory to platinum-

based chemotherapy.<sup>18</sup> Notably, efficacy was restricted to HPV-negative tumors. Despite lacking formal regulatory approval for treating patients with HNSCC, afatinib is endorsed in the National Comprehensive Cancer Network guideline (category B).

### HPV Status and Anti-EGFR Therapy

While anti-EGFR therapies are indicated regardless of HPV status, multiple studies have reported reduced efficacy in patients with HPV-positive HNSCC.<sup>15,19,20</sup> This disparity may be partly attributed to differences in the underlying molecular characteristics. HPV-positive tumors harbor higher rates of genetic alterations in EGFR downstream pathways, including *PIK3CA* variants and *KRAS* variants.<sup>21-23</sup> Additionally, *EGFR* amplification was exclusively observed in HPV-negative tumors within the Cancer Genome Atlas dataset. Emerging EGFR-targeted treatments are exploring innovative strategies to enhance the efficacy and overcome resistance in patients with both HPV-positive and HPV-negative HNSCC.

### Emerging Approaches in EGFR-Targeted Treatments

This review highlights recent advances in EGFR-directed therapies, focusing on (1) combination regimens with existing EGFR agents (Figure 1; Table 1) and (2) novel EGFR-targeting therapeutics with innovative mechanisms (Figure 2 and Figure 3; Table 2).


#### Cetuximab-Based Combination Regimens

**PD-1/Programmed Cell Death Ligand 1 Inhibitors** | PD-1/programmed cell death 1 ligand 1 (PD-L1) inhibitors are recommended as first-line therapy for patients with R/M HNSCC, either alone or in combination with platinum-based chemotherapy, which function by blocking T-cell exhaustion. These agents may have a synergistic effect with cetuximab through promoting antigen presentation and NK cell engagement. This interaction provides a solid rationale for combining PD-1/PD-L1 inhibitors with cetuximab to potentiate antigen-specific T-lymphocyte responses and improve therapeutic efficacy.

Cetuximab plus nivolumab showed promising results with an ORR of 22% and a median (IQR) PFS of 3.4 months in the second-line cohort and an ORR of 37% and a median (IQR) PFS of 6.1 months in the first-line cohort.<sup>24</sup> Biomarker analysis showed that p16/HPV negativity correlated with higher response rates (19 [41%] vs 7 [18%];  $P = .02$ ). Similarly, promising results were seen when cetuximab was combined with pembrolizumab, resulting in an ORR of 45%, median PFS of 6.5 (95% CI, 2.1 to NR) months, and median OS of 18.4 (95% CI, 11.0 to NR) months from mainly treatment-naïve patients.<sup>25</sup> Again, most responses were seen in patients with HPV-negative disease, with an ORR of 57% (12 of 21).

The combination of cetuximab and other PD-1/PD-L1 ICIs, such as durvalumab and toripalimab, demonstrated similar efficacy, further supporting the hypothesis of an immune-based anticancer mechanism of cetuximab.<sup>26,27</sup> Consistent with other observational reports, a meta-analysis of multiple anti-PD-1-based HNSCC trials demonstrated that adding cetuximab to PD-1 inhibitors significantly improved ORR (15% [95% CI, 12%-18%] vs 46% [95% CI, 34%-58%];  $P < .001$ ) and 1-year OS rate (36% [95% CI, 32%-41%] vs 59% [95% CI, 47%-71%];  $P < .001$ ) in HPV-negative HNSCC but offered no advantage in HPV-positive disease.<sup>46</sup> A phase 3 randomized trial (NCT06589804) of pembrolizumab with or without

Figure 1. Cetuximab-Based Combination Regimens and Their Molecular Targets



cetuximab has recently started enrollment and may provide data supporting the findings.

**NK Cell Activators** | NK cells play a crucial role in tumor immunosurveillance, with their activity regulated by a balance of activating and inhibitory receptors. Many solid tumors, including HNSCC, upregulate nonclassical major histocompatibility (MHC) molecules, such as HLA-E and HLA-G, which interact with inhibitory receptors NKG2A and ILT2/ILT4 on immune cells, respectively, allowing them to escape immune surveillance. Targeting these inhibitory axes can restore NK cell function and improve the antitumor immunity of cetuximab through enhanced ADCC.

The potential synergy between cetuximab and NK cell activators has been explored clinically through monalizumab, an anti-NKG2A antibody, which blocks the interaction with HLA-E. Despite the encouraging results from early phase trials, the phase 3 INTERLINK-1 study, which evaluated cetuximab with or without monalizumab, failed to meet the primary end point of OS in patients with R/M HNSCC after platinum chemotherapy and ICI therapy.<sup>20,47</sup> Interestingly, the response rate of 19% reported for cetuximab monotherapy was higher than the previously reported ORR of 13% in the pre-ICI era, which may be due to the effect of prior ICI exposure.<sup>12</sup>

Another strategy to restore NK cell function in HNSCC is targeting the ILT-HLA-G axis. TTX-080, a monoclonal antibody that neutralizes HLA-G, in combination with cetuximab, showed a preliminary efficacy signal in HPV-negative HNSCC, with an ORR of 57%, while patients with HPV-positive disease experienced an ORR of 0%.<sup>28</sup>

Interleukin 15 (IL-15) has a role in supporting NK cell proliferation and tumor killing through JAK/STAT and MAPK signaling, which provides the rationale for combining with monoclonal antibodies with ADCC effect. An ongoing study is investigating engineered NK cells, PD-L1 tumor-targeted high-affinity NK, combined with an IL-15 superagonist (N-803), along with cetuximab, in patients with advanced HNSCC (NCT06239220). Similarly, NKTR-255, a recombinant human IL-15 agonist, is being explored in combination with cetuximab.<sup>29</sup>

NK cell activators enhance cetuximab efficacy primarily by improving NK cell cytotoxicity, but they can also indirectly boost ADCP through the secretion of interferon- $\gamma$  and tumor necrosis factor- $\alpha$ , which increases the phagocytic capacity of macrophages. However, tumors frequently evade ADCP through the CD47-SIRP $\alpha$  axis, where CD47 on tumor cells functions as a "do not eat me" signal, protecting them from macrophage-mediated phagocytosis. This makes the combination of cetuximab with an anti-CD47 agent an appealing approach.

Table 1. Key Clinical Trials of Cetuximab-Based Combination Regimens for Recurrent or Metastatic Head and Neck Squamous Cell Carcinoma (R/M HNSCC)

Agent	Description	Study design (ClinicalTrials.gov identifier)	Intervention	Evaluate patients with R/M HNSCC, No. and characteristics	Efficacy	Participants with grade ≥3 TRAEs (discontinuation rate)
<b>PD-1/PD-L1 inhibitors</b>						
Nivolumab <sup>24</sup>	Anti-PD-1 IgG4 mAb	Phase 2 single-arm study (NCT03370276)	Cetuximab plus nivolumab	Overall cohort (N = 88) grouped into prior-therapy cohort (n = 45) (22 with HPV-positive disease) and treatment-naïve cohort (n = 43) (18 with HPV-positive disease)	Prior-therapy cohort: median OS, <sup>a</sup> 11.4 mo; ORR, 22%; median PFS, 3.4 mo; treatment-naïve cohort: median OS, <sup>a</sup> 20.2 mo; ORR, 37%; PFS, 6.1 mo	Prior-therapy cohort: NA; treatment-naïve cohort: 23.3% (NA)
Pembrolizumab <sup>25</sup>	Anti-PD-1 IgG4 mAb	Phase 2 multiarm nonrandomized trial (NCT03082534) Phase 3 randomized clinical trial (NCT06589804)	Cetuximab plus pembrolizumab Pembrolizumab with or without cetuximab	Patients resistant to platinum-based chemotherapy (n = 33); other arms not reported to date NA	ORR, <sup>a</sup> 45%; PFS, 6.5 mo; DOR, 13.1 mo; OS, 18.4 mo NA	42.4% (15% discontinued) NA
Durvalumab <sup>26</sup>	Anti-PD-L1 IgG1 mAb	Phase 2 single-arm study (NCT03691714)	Cetuximab plus durvalumab	All patients resistant to platinum-based chemotherapy (N = 33) (10 received prior ICI, 1 prior cetuximab)	ORR, <sup>a</sup> 39%; median PFS, 5.8 mo; median OS, 9.6 mo	48.6% (NA)
Toripalimab <sup>27</sup>	Anti-PD-1 IgG4 mAb	Phase 1b/2 single-arm study (NCT04856631)	Cetuximab plus toripalimab	All patients resistant to platinum-based chemotherapy (N = 12) (9 with PD-L1 CPS ≥1)	ORR, 50%	0% (0% discontinued)
<b>NK cell activators</b>						
Monalizumab <sup>20</sup>	Anti-NKG2A IgG1 mAb	Phase 3 INTERLINK-1 randomized clinical trial (NCT04590963)	Cetuximab with or without monalizumab	All patients received prior platinum-based chemotherapy and ICI and no prior cetuximab (N = 264) (216 with HPV-unrelated disease)	Overall study population OS: HR, 1.03; P = .89; HPV-unrelated cohort OS: HR, 1.00; P = .99	Monalizumab: 38.6% (5.7% discontinued); placebo: 30.3% (1.6% discontinued)
TTX-080 <sup>28</sup>	Anti-HLA-G IgG1 mAb	Phase 1a/1b study (NCT04485013)	Cetuximab plus TTX-080	All patients received prior ICI (N = 17) (7 with HPV-negative and 10 with HPV-positive disease)	HPV-negative cohort: ORR, <sup>a</sup> 57% (1 CR, 3 PR); median PFS, 23.9 mo; HPV-positive cohort: ORR, <sup>a</sup> 0%; median PFS, 9.1 mo	NA
N-803	IL-15 superagonist antibody cytokine fusion protein	Phase 1 study (NCT06239220)	Cetuximab plus PD-L1 tumor-targeted high-affinity NK plus N-803	NA	NA	NA
NKTR-255 <sup>29</sup>	Recombinant IL-15 agonist	Phase 1b/2 study (NCT04616196)	Cetuximab plus NKTR-255	All patients received prior therapy (N = 3) (2 prior ICI, 1 prior cetuximab)	ORR, 0%; DCR, 66%	NA
<b>Multireceptor TKIs</b>						
Lenvatinib <sup>30</sup>	TKI against VEGF/FGF/PDGFRα	Phase 1/1b study (NCT03524326)	Cetuximab plus lenvatinib	Patients with R/M HNSCC or skin cancer (N = 9)	ORR, 67%; median PFS, 3.6 mo	NA
Cabozantinib <sup>31</sup>	TKI against AXL/c-MET/VEGFR	Phase 1 study (NCT03667482)	Cetuximab plus cabozantinib	Patients may have received prior therapy (N = 20) (19 received prior ICI, 19 prior platinum-based chemotherapy, 16 prior cetuximab)	Overall study population: ORR, 20%; DCR, 75%; median PFS, 3.4 mo; median OS, 8.1 mo; cetuximab-naïve cohort: ORR, 50%	65% (NA)
Sorafenib <sup>32</sup>	TKI against VEGF/PDGFR/RAF	Phase 2 randomized clinical trial (NCT00939627)	Cetuximab with or without sorafenib	Patients naïve to sorafenib or cetuximab (N = 43)	Both arms: ORR, 8%; OS, NS; PFS, NS	NA

(continued)

Table 1. Key Clinical Trials of Cetuximab-Based Combination Regimens for Recurrent or Metastatic Head and Neck Squamous Cell Carcinoma (R/M HNSCC) (continued)

Agent	Description	Study design (ClinicalTrials.gov identifier)	Intervention	Evaluate patients with R/M HNSCC, No. and characteristics	Efficacy	Participants with grade ≥ 3 TRAEs (discontinuation rate)
<b>c-MET/HGF inhibitors</b>						
Ficlatuzumab <sup>33</sup>	Anti-HGF IgG1 mAb	Phase 2 randomized clinical trial (NCT03422536)	Ficlatuzumab with or without cetuximab	Patients resistant to cetuximab (N = 58); randomized to monotherapy (n = 26) or combination therapy (n = 32) (in combination therapy cohort, 16 had HPV-positive disease, 16 had HPV-negative disease)	Monotherapy closed early for futility; combination therapy cohort: ORR, 19%; median PFS, 3.7 mo; HPV positive: ORR, 0%; median PFS, 2.3 mo; HPV negative: ORR, 38% (2 CR, 4 PR), median PFS, 4.1 mo	NA
		Phase 3 FIERCE-HN randomized clinical trial (NCT06064877)	Cetuximab with or without ficlatuzumab	NA	NA	NA
Tivantinib <sup>34</sup>	TKI against c-MET	Phase 2 randomized clinical trial (NCT01696955)	Cetuximab with or without tivantinib	All patients received prior platinum-based chemotherapy and were cetuximab-naïve (N = 78); randomized to combination therapy (n = 40) or monotherapy (n = 38)	Combination therapy: ORR, 7.5% (1 CR); monotherapy: ORR, 7.9%; overall study population: median PFS, 4 mo; NS; median OS, 8 mo; NS; HPV-positive cohort ORR, 0%	NA
Capmatinib (INC280)	TKI against c-MET	Phase 1b study; terminated (NCT02205398)	Cetuximab plus capmatinib	NA	NA	NA
<b>Other targeting agents</b>						
ERAS-601	SHP2 inhibitor	FLAGSHIP-1 phase 1 study (NCT04670679)	Cetuximab plus ERAS-601	NA	NA	NA
Palbociclib <sup>35-37</sup>	CDK 4/6 inhibitor	Phase 1 study (NCT03498378)	Cetuximab plus palbociclib plus avelumab	Patients with no prior therapy using EGFR inhibitor or PD-1 or PD-L1 inhibitor in the recurrent or metastatic setting (N = 12) (5 with HPV-positive disease)	ORR, 41.7% (3 CR, 2 PR); DCR, 75%; median PFS, 6.5 mo; median OS, NA	75% (NA)
		Phase 2 nonrandomized clinical trial (NCT02101034)	Cetuximab plus palbociclib	Patients may or may not have received prior therapy (N = 55); grouped into patients who are cetuximab-naïve (n = 28) and cetuximab-resistant (n = 27)	HPV-unrelated disease: ORR, <sup>a</sup> 39%; median PFS, 5.4 mo, median OS, 9.5 mo; cetuximab-naïve: Median ORR, <sup>a</sup> 19%; median PFS, 3.7 mo; median OS, 6.3 mo	HPV-unrelated disease: 60% (NA); cetuximab-naïve: 44% (NA)
		PALATINUS phase 2 randomized clinical trial (NCT02499120)	Cetuximab with or without palbociclib	Patients received prior platinum-based chemotherapy and were cetuximab-naïve with HPV-unrelated disease (N = 125); grouped into monotherapy (n = 65) and combination therapy (n = 60)	Combination therapy: median OS, <sup>a</sup> 9.7 mo; median PFS, 3.9 mo; monotherapy: median OS, <sup>a</sup> 7.8 mo, median PFS, 4.6 mo	Combination therapy: 54.7% (NA); monotherapy: 15% (NA)
		Phase 3 randomized clinical trial (NCT04966481)	Cetuximab with or without palbociclib	All patients had CDK inhibitor 2A-altered, HPV-unrelated disease and received prior ICI (N = NA)	NA	NA

Abbreviations: CDK, cyclin-dependent kinase; CPS, combined positive score; CR, complete response; DCR, disease control rate; DOR, duration of response; HPV, human papillomavirus; HR, hazard ratio; ICI, immune checkpoint inhibitor; IL-15, interleukin 15; NA, not available; NS, not significant; ORR, objective response rate; OS, overall survival; PD-1, programmed cell death 1; PD-L1, programmed cell death ligand 1; PFS, progression-free survival; PR, partial response; TKI, tyrosine kinase inhibitor; TRAE, treatment-related adverse events.

<sup>a</sup> Primary end point.



Figure 2. Bispecific Antibodies and Antibody-Drug Conjugates Targeting EGFR





Figure 3. Immune Cell Engagers and Adaptive Cell Therapies Targeting EGFR



**Multireceptor TKIs** | The rationale for combining multireceptor TKIs with cetuximab to treat patients with R/M HNSCC is based on the recognized role of alternative receptor tyrosine kinase activation in both intrinsic and acquired resistance to cetuximab.

Lenvatinib, a multireceptor TKI with activity against the VEGFR 1, 2, and 3; FGFR 1, 2, 3, and 4; PDGFR $\alpha$ ; KIT; and RET pathways, was

evaluated in combination with cetuximab. The unique activity of lenvatinib against FGFR 1, 2, 3, and 4, a recognized resistance mechanism for EGFR inhibition, seemed promising based on the initial responses from 9 evaluable patients. However, extensive thromboembolic events and atherosclerosis were observed in some patients, which are well-established adverse events of VEGF TKIs.<sup>30</sup>

Table 2. Key Clinical Trials of Novel Therapeutics Targeting Epidermal Growth Factor Receptor (EGFR) in Recurrent or Metastatic Head and Neck Squamous Cell Carcinoma (R/M HNSCC)

Agent	Description	Study design (ClinicalTrials.gov identifier)	Intervention	Evaluable patients with R/M HNSCC, No. and characteristics	Efficacy	Participants with grade ≥3 TRAEs (discontinuation rate)
<b>Bispecific antibodies</b>						
Ficerafusp alfa (BCA-101) <sup>38</sup>	Bifunctional anti-EGFR mAb with TGF-β trap	Phase 1/1b study (NCT04429542)	Ficerafusp alfa plus pembrolizumab	All patients received first-line therapy for R/M HNSCC and had PD-L1 CPS ≥1 (N = 39)	ORR, 46%; HPV-negative disease: ORR, 57%	40% (14% discontinued)
Petosemtamab <sup>39</sup>	Targets EGFR, LGR5	Phase 2/3 randomized clinical trial (NCT06788990)	Pembrolizumab with or without ficerafusp alfa	All patients received first-line therapy for R/M HNSCC, had PD-L1 CPS ≥1, and had HPV-negative disease (N = NA)	NA	NA
		Single-arm phase 2 study (NCT03526835)	Petosemtamab plus pembrolizumab	All patients received first-line therapy for R/M HNSCC and had PD-L1 CPS ≥1 (N = 24) (4 had HPV-positive disease, 20 had HPV-negative disease)	ORR, <sup>a</sup> 67%; HPV-positive disease: ORR, 75%; HPV-negative disease: ORR, 65%	24% (4% discontinued)
		LiGeR-HN1 phase 3 randomized clinical trial (NCT06525220)	Pembrolizumab with or without petosemtamab	All patients received first-line therapy for R/M HNSCC and had PD-L1 CPS ≥1	NA	NA
		LiGeR-HN2 phase 3 randomized clinical trial (NCT06496178)	Petosemtamab vs investigator's choice of cetuximab, methotrexate, or docetaxel	All patients received prior ICI and platinum-based chemotherapy (N = NA)	NA	NA
Amivantamab	Targets EGFR, c-MET	OrigAMI-4 phase 1b/2 study (NCT06385080)	NA	NA	NA	NA
MCLA-129 <sup>40</sup>	Targets EGFR, c-MET	Phase 1/2 study (NCT04868877)	MCLA-129	Patients whose prior treatment failed (N = 12)	ORR, <sup>a</sup> 17% (unconfirmed); DCR, 67%	NA
SI-B001 <sup>41</sup>	Targets EGFR, ERBB3	S209 single-arm phase 2 study (NCT05044897)	SI-B001	All patients received prior ICI plus platinum-based chemotherapy (N = 9)	ORR, <sup>a</sup> 22.2%; DCR, 77.8%; median PFS, 2.7 mo	NA
		S206 single-arm phase 2 study (NCT05054439)	SI-B001 (plus paclitaxel if patients were paclitaxel naive)	All patients received prior ICI with or without platinum-based chemotherapy (N = 22)	ORR, <sup>a</sup> 45.5%; DCR, 81.8%; median PFS, 5.1 mo	NA
<b>Antibody-drug conjugates</b>						
MRG003 <sup>42</sup>	EGFR targeting ADC with MMAE payload via valine-citrulline linker	Phase 2 study (NCT04868162)	MRG003, 2.0 mg/kg (DL1), or MRG003, 2.3 mg/kg (DL2)	Patients with EGFR-positive disease (N = 62) grouped into DL1 dosage (n = NA) and DL2 dosage (n = 14) (95.6% received prior platinum-based chemotherapy, 76.1% prior ICI, 47.8% prior cetuximab)	DL1 dosage: data NA; DL2 dosage: ORR, <sup>a</sup> 43% (1 CR); DCR, 86%; median PFS, 4.2 mo; median OS, 11.3 mo	NA
		Phase 3 randomized clinical trial (NCT05751512)	MRG003 vs cetuximab/methotrexate	NA	NA	NA
Cetuximab sarotalocan (RM-1929) <sup>43</sup>	Cetuximab with IRDye700DX payload	Phase 1/2a study (NCT02422979)	Cetuximab sarotalocan	Patients received prior platinum-based chemotherapy unless contraindicated (N = 30) (21 received prior chemotherapy, 10 received prior ICI, 7 received prior cetuximab)	ORR, 26.7%; DCR, 80%	63.3% (NA)
		ECLIPSE phase 3 randomized clinical trial (NCT06699212)	Cetuximab sarotalocan plus pembrolizumab	First-line setting	NA	NA

(continued)

Table 2. Key Clinical Trials of Novel Therapeutics Targeting Epidermal Growth Factor Receptor (EGFR) in Recurrent or Metastatic Head and Neck Squamous Cell Carcinoma (R/M HNSCC) (continued)

Agent	Description	Study design (ClinicalTrials.gov identifier)	Intervention	Evaluable patients with R/M HNSCC, No. and characteristics	Efficacy	Participants with grade ≥3 TRAEs (discontinuation rate)
Tilatamab samrotectan (AZD9592)	Bispecific ADC targeting EGFR and c-MET with TOP1i inhibitor payload	EGRET phase 1 study (NCT05647122)	Tilatamab samrotectan	NA	NA	NA
GEN1286 (PRO1286)	Bispecific ADC targeting EGFR and c-MET with TOP1i inhibitor payload	Phase 1/2 study (NCT06685068)	GEN1286	NA	NA	NA
Izalontamab brengitecan (BL-B01D1) <sup>44</sup>	Bispecific ADC targeting EGFR and ERBB3 with TOP1i inhibitor payload via cathepsin B cleavable linker	Phase 1 study (NCT05194982)	Izalontamab brengitecan	Patients received prior platinum-based chemotherapy (N = 20) (1 with disease progression on standard of care; 1 received prior TKI, 18 received prior ICI)	ORR, 15%; DCR 80%	36% (3% discontinued)
M1231	Bispecific ADC targeting EGFR and MUC1 with microtubule inhibitor payload	Phase 1 study (NCT04695847)	M1231	NA	NA	NA
<b>Immune cell engagers</b>						
JANX008	T-cell engager targeting EGFR and CD3	Phase 1 study (NCT05783622)	JANX008	NA	NA	NA
CX-904	T-cell engager targeting EGFR and CD3	Phase 1 study (NCT05387265)	CX-904	NA	NA	NA
REGN7075	T-cell engager targeting EGFR and CD28	COMBINE-EGFR-1 phase 1/2 study (NCT04626635)	REGN7075 plus cemiplimab	NA	NA	NA
DF9001	NK cell engager targeting EGFR and NKG2D	Phase 1 study (NCT05597839)	DF9001 with or without pembrolizumab	NA	NA	NA
PF-08046052 (SGN-EGFRd2)	T-cell engager targeting EGFR and T-cell receptor δ2	Phase 1 study (NCT05983133)	PF-08046052	NA	NA	NA
<b>Adaptive cell therapies</b>						
T4 immunotherapy <sup>45</sup>	Autologous panERBB-specific CAR T cells	Phase 1 study (NCT01818323)	T4 immunotherapy	Patients for whom no standard therapy remains or is suitable (N = 15)	ORR, 0%; DCR, 60%	0% (NA)
A2B395	Allogenic logic-gated EGFR CAR T cell	DENALI-1 phase 1/2 study (NCT06682793)	A2B395	NA	NA	NA
ACE2016	Allogenic EGFR targeting γδ2 T cells	Phase 1 study (NCT06415487)	ACE2016	NA	NA	NA

Abbreviations: ADC, antibody-drug conjugate; CAR, chimeric antigen receptor; CPS, combined positive score; CR, complete response; DCR, disease-control rate; HPV, human papillomavirus; ICI, immune checkpoint inhibitor; mAB, monoclonal antibody; MMAE, monomethyl auristatin E; MUC1, mucin 1; NA, not available; NK, natural killer; ORR, objective response rate; PD-1, programmed cell death 1; PD-L1, programmed cell death ligand 1; PFS, progression-free survival; OS, overall survival; TGF-β, transforming growth factor β; TKI, tyrosine kinase inhibitor; TOP1i, topoisomerase 1 inhibitor; TRAE, treatment-related adverse events.

<sup>a</sup> Primary end point.



Cabozantinib, a TKI blocking the VEGFR 1, 2, and 3; cMET; AXL; and RET pathways, and sorafenib, a TKI blocking the VEGFR 1, 2, and 3; PDGFR $\beta$ ; KIT; and RET pathways, were also evaluated in combination with cetuximab. But further development was hampered by the lack of robust efficacy or the high rate of treatment-related adverse events, including skin rash, fatigue, and liver toxic effects.<sup>31,32</sup>

**c-MET/HGF Inhibitors** | The c-MET receptor tyrosine kinase has been implicated in HNSCC progression, with c-MET overexpression being linked to advanced disease stage. Given the role of abnormal c-MET/HGF pathway activation in EGFR TKI resistance and the success of combined c-MET/HGF inhibition in lung cancers, similar combination strategies are being explored in HNSCC.

Ficlatuzumab, a monoclonal antibody inhibiting the HGF pathway, showed a potential signal of synergy with cetuximab with an ORR of 19% in patients with platinum and ICI-refractory R/M HNSCC.<sup>33</sup> All responses were seen in the HPV-negative subgroup with an ORR of 38%. A randomized phase 3 trial (FIERCE-HN) is ongoing to further evaluate the efficacy of this combination compared to cetuximab monotherapy in HPV-negative disease refractory to ICI and platinum chemotherapy.

Despite the appealing mechanism, not all combinations of cetuximab and HGF/c-MET targeting agents have been successful. A phase 2 randomized clinical trial of cetuximab with or without tivantinib, a MET TKI, reported an increase in toxic effects, especially myelosuppression, without improvement in response rate or survival.<sup>34</sup> Notably, all responses were seen in patients with HPV-negative disease in both treatment arms.

**Other Targeting Agents** | SHP2, an oncogenic nonreceptor tyrosine phosphatase, links receptor tyrosine kinase activation to RAS-MAPK signaling. This pathway activates through EGFR and alternative receptor tyrosine kinases involved in cetuximab resistance. The combination of ERAS-601, an allosteric SHP2 inhibitor, with cetuximab is under investigation in the FLAGSH-1 trial.<sup>48</sup>

Cell-cycle dysregulation in HNSCC occurs through cyclin D1 (*CCND1*), cyclin-dependent kinase (CDK) inhibitor 2A (*CDKN2A*), or RB transcriptional corepressor 1 (*RB1*) alterations and may cause EGFR inhibition resistance. Palbociclib, a CDK 4/6 inhibitor, combined with cetuximab showed promising response rates in HPV-negative disease but failed to show OS improvement over cetuximab monotherapy in biomarker-unselected patients.<sup>35,36</sup> An ongoing phase 3 study ([NCT04966481](#)) is investigating cetuximab with or without palbociclib in patients with *CDKN2A*-altered, HPV-negative HNSCC.

## Novel Agents Targeting EGFR

**Bispecific Antibodies** | Bispecific antibodies offer the advantage of engaging multiple targets, potentially enhancing antitumor activity and overcoming resistance mechanisms. Ficerafusp alfa (BCA101) is a bifunctional antibody targeting the EGFR and TGF- $\beta$ 1 pathways through the extracellular domain of transforming growth factor  $\beta$  (TGF- $\beta$ ) receptor 2. TGF- $\beta$  promotes cancer progression via epithelial-mesenchymal transition and immune suppression, with epithelial-mesenchymal transition being implicated as a mechanism of cetuximab resistance in patients with HNSCC.<sup>49</sup> In an ongoing phase 1/1b study, ficerafusp alfa plus pembrolizumab demonstrated a response rate of 46% in treatment-naïve R/M HNSCC. Notably, the

response rate was higher in HPV-negative compared to HPV-positive disease (57% vs 18%).<sup>38</sup> A randomized phase 2/3 clinical trial ([NCT06788990](#)) comparing the efficacy of ficerafusp alfa plus pembrolizumab to pembrolizumab alone as first-line therapy in HPV-negative R/M HNSCC is ongoing.

Petosemtamab is a bispecific antibody targeting the EGFR and leucine-rich repeat 16 containing G protein-coupled receptor 5 (LGR5) pathways. LGR5 is a cancer stem cell marker with expression in 52% to 89% of patients with HNSCC and association with tumor self-renewal, metastasis, and resistance to chemotherapy.<sup>50</sup> Petosemtamab is thought to inhibit tumor growth by blocking EGFR-dependent signaling, promoting LGR5-mediated cointernalization and degradation of EGFR, and enhancing ADCC and ADCP through its low-fucose Fc domain. Petosemtamab demonstrated a robust monotherapy activity with an ORR of 36%, median (IQR) PFS of 5.0 (3.2-6.8) months, and median (IQR) OS of 11.5 (7.2-20.6) months in patients with R/M HNSCC whose disease progressed following treatment with platinum-based chemotherapy and ICI therapy.<sup>51</sup> The combination of petosemtamab with pembrolizumab in the first-line setting resulted in a response rate of 67%.<sup>39</sup> Notably, responses were seen in patients with both HPV-positive and HPV-negative disease. Two phase 3 randomized clinical trials that include patients with HPV-positive and HPV-negative disease are underway: petosemtamab monotherapy vs standard single-agent in the second-line or third-line setting (LiGeR-HN2) and pembrolizumab with or without petosemtamab as first-line therapy in R/M HNSCC (LiGeR-HN1).

Amivantamab is a bispecific antibody targeting the EGFR and c-MET pathways. An ongoing phase 1b/2 clinical trial ([NCT06385080](#)) is investigating amivantamab alone and in combination with standard-of-care treatments. MCLA-129 is another EGFR and c-MET bispecific antibody, which reported a preliminary efficacy but a high infusion-related reaction rate, leading to discontinuation of further development.<sup>40</sup>

SI-BOO1 is a bispecific antibody that targets the EGFR and ERBB3 pathways. While ERBB3 lacks intrinsic kinase activity, it forms heterodimers with other ERBB family members, particularly ERBB2, and activates downstream signaling involved in cell proliferation and survival. Given its suggested role in EGFR-directed therapy resistance, ERBB3 has emerged as an important target, and SI-BOO1 combined with taxane therapy has demonstrated an encouraging efficacy in HNSCC refractory to platinum chemotherapy and ICI therapy.<sup>41</sup>

**Antibody-Drug Conjugates** | Antibody-drug conjugates (ADCs) combine an antibody, linker, and payload to deliver cytotoxic drug payloads with high specificity to target cells. ADCs have seen immense success in solid tumors with multiple new approvals in recent years, yet none have been approved to treat patients with R/M HNSCC. While various antigens are being explored for ADCs in HNSCC, we will focus on a few key examples that target EGFR.

MRGO03 is an ADC with an IgG1 monoclonal antibody with an affinity to EGFR higher than that of cetuximab, a valine-citrulline cleavable linker, and monomethyl auristatin E (MMAE), a highly toxic antimicrotubule payload. In a phase 2 study, MRGO03 demonstrated a single-agent response rate of 30.6% in heavily pretreated patients with EGFR-positive HNSCC.<sup>42</sup> A randomized phase 3 study ([NCT05751512](#)) evaluating MRGO03 in comparison with standard cetuximab or methotrexate as second-line and third-line therapy is ongoing.

The development of EGFR-targeting ADCs to treat patients with HNSCC faces the challenge of overcoming on-target off-tumor toxic effects. While many ADCs in clinical development use tubulin or topoisomerase 1 inhibitors (TOP1i) as their payloads, some use novel approaches using noncytotoxic payloads to tackle this issue. For example, cetuximab sarotalocan combines cetuximab with IRDye700DX, a unique payload activated by near-infrared light illumination at the tumor site.<sup>43</sup> A phase 3 trial (ECLIPSE, [NCT06699212](#)) for cetuximab sarotalocan plus pembrolizumab as first-line treatment for patients with locoregional recurrent HNSCC is ongoing.

To enhance the efficacy of conventional single antigen-targeting ADCs, a multiantigen targeting strategy is being explored. Tilatamig samrotescan (AZD9592) and GEN1286 (PRO1286) are bispecific ADCs targeting EGFR and c-MET, designed to deliver TOP1i payloads. Tilatamig samrotescan was engineered with a higher affinity for c-MET compared to EGFR (greater than 15-fold) to reduce EGFR-driven toxic effects.<sup>52</sup> It is under investigation in a phase 1 study (EGRET).<sup>53</sup> GEN1286 is also in early-phase clinical investigation ([NCT06685068](#)) and comprises a monovalent IgG1 fine-tuned affinity for EGFR and c-MET, conjugated with a TOP1i by a hydrophilic linker.

Other targets are also being explored for bispecific ADCs. Izalontamab brengitecan (BL-B01D1) is a bispecific tetravalent ADC targeting EGFR with a high affinity and ERBB3 with a low affinity, delivering a TOP1i payload.<sup>44</sup> M1231 is a bispecific ADC targeting the EGFR and mucin 1 (MUC1), a transmembrane glycoprotein coexpressed with EGFR in HNSCC.

**Immune Cell Engagers** | Immune cell engagers are bispecific molecules that redirect T cells or NK cells to kill cancer cells by binding both a tumor-associated antigen and an immune receptor (eg, CD3, CD28, or CD16). The dual-targeting ensures specificity, limits off-target effects, and can trigger a bystander effect, allowing immune cells to eliminate nearby tumor cells lacking the tumor-associated antigen.<sup>54</sup>

T-cell engagers (TCEs) are a subclass of immune cell engagers that redirect cytotoxic T cells by binding both the tumor-associated antigen and CD3, enabling MHC-independent activation. Early-generation TCEs often caused cytokine release syndrome, necessitating step-up dosing. JANX008 and CX-904 are conditionally activated bispecific TCEs targeting EGFR and CD3 with protease-cleavable masking domains, remaining inactive in circulation until activated by tumor-associated proteases. This enables localized T-cell engagement and reduces the risk of cytokine release syndrome.<sup>55,56</sup> Both are being investigated in phase 1 clinical trials ([NCT05783622](#), [NCT05387265](#)) with results yet to be reported.

While CD3-directed TCEs rely primarily on T-cell receptor-mediated T-cell redirection, REGN7075 targets CD28, which provides a costimulatory signal crucial for full T-cell activation, expansion, and sustained antitumor activity. An ongoing phase 1/2 study is evaluating REGN7075 in combination with cemiplimab, a PD-1 inhibitor ([NCT04626635](#)).

NK cells and  $\gamma\delta$  T cells are alternative targets for immune cell engagers that offer the advantage of allogeneic compatibility without the risk of graft-vs-host disease. DF9001 targets EGFR and NKG2D, a receptor expressed on NK cells, CD8-positive T cells, and  $\gamma\delta$  T cells, and is being evaluated as monotherapy and in combination with pembrolizumab ([NCT05597839](#)).<sup>57</sup> PF-08046052 (SGN-EGFRd2) is a bispecific antibody that targets EGFR and the

$\gamma\delta$  T-cell receptor on  $\gamma\delta$  T cells, directing their intrinsic antitumor activity to EGFR-expressing cancer cells while requiring a secondary phosphoantigen stress signal to minimize off-target effects. It also binds to EGFR extracellular domain epitopes distinct from cetuximab's to address cetuximab-induced escape variants. A phase 1 clinical trial ([NCT05983133](#)) is evaluating PF-08046052 in selected EGFR-expressing tumors.

**Adaptive Cell Therapies** | Adaptive cell therapy is a type of immunotherapy using immune cells engineered to target specific cells of interest, typically involving isolation of immune cells from patients (autologous) or donors (allogeneic), ex vivo expansion with or without genetic manipulation, and reinfusion of manufactured cells. While chimeric antigen receptor (CAR) T-cell therapy has transformed treatment for hematologic cancers, applications in solid tumors remain limited due to manufacturing complexity, lack of tumor-specific antigens, inefficient cell trafficking and infiltration, and an immunosuppressive tumor microenvironment.<sup>58</sup> Next-generation CAR T cells use a variety of strategies to overcome these challenges.

The T4 immunotherapy study ([NCT01818323](#)) evaluates autologous CAR T cells targeting ERBB receptors in HNSCC. Patient-derived T cells are engineered to express 2 synthetic peptides: T1E28Z (pan-ERBB chimeric peptide derived from the epidermal growth factor [EGF] protein) and 4 $\alpha$ B (IL-2/IL-15 signaling on IL-4 binding), enabling selective expansion in the presence of IL-4 during manufacturing. To enhance local activity and reduce systemic toxic effects, T4 CAR T cells are delivered intratumorally. Preliminary phase 1 data demonstrated feasibility, safety, and undetectable peripheral T4 CAR T cells.<sup>45</sup>

Logic-gated CAR T cells require 2 or more signals for activation, improving specificity and reducing off-tumor toxic effects. A2B395 is an allogeneic, logic-gated CAR T-cell therapy targeting EGFR with an HLA-binding inhibitory receptor creating an "AND-NOT" logic gate. EGFR-expressing cells lacking HLA-A02, through loss of heterozygosity in approximately 30% of solid tumors, are targeted while normal cells expressing HLA-A02 are protected. The DENALI-1 trial ([NCT06682793](#)) is enrolling patients with EGFR-expressing solid tumors and HLA-A02 loss of heterozygosity.

While  $\alpha\beta$  T-cell therapies often require MHC knockout and T-cell receptor removal to prevent graft-vs-host disease or rejection for allogeneic use,  $\gamma\delta$  T cells recognize antigens independently of HLA, potentially lowering these risks.<sup>59</sup> ACE2016, an allogeneic  $\gamma\delta$  T-cell therapy with conjugated anti-EGFR antibodies, is under investigation in the first human trial ([NCT06415487](#)).<sup>60</sup>

## Conclusions

Recent advances have revitalized EGFR as a therapeutic target in HNSCC, with encouraging results from cetuximab-immunotherapy combinations and novel bispecific antibodies now progressing to registration trials. Despite this promise, significant challenges remain, including on-target off-tumor toxic effects from EGFR expression in normal tissues and limited efficacy in the growing population of patients with HPV-positive disease. The path forward will require a deeper understanding of tumor biology, resistance mechanisms, and tailored strategies for distinct tumor subtypes to meaningfully improve outcomes in this challenging cancer.

## ARTICLE INFORMATION

**Accepted for Publication:** June 13, 2025.

**Published Online:** September 25, 2025.  
doi:10.1001/jamaoncol.2025.3136

**Conflict of Interest Disclosures:** Dr Hwang reported personal fees from Lunit and being an employee of Lunit Inc outside the submitted work. Dr Jung reported receiving consulting and advisory board fees from Yuhan, Astrazeneca, Aimedbio, Merck, Sanofi, and Novartis. Dr Wirth reported receiving advisory board fees from Ayala Pharmaceuticals, Blueprint Medicines, Cue Biopharma, Cullinan Oncology, Genentech USA, Loxo Oncology, Merck, NewLink Genetics, Novartis, and Rakuten Medical, consulting fees and advisory board fees from Bayer HealthCare Pharmaceuticals and Eisai, advisory board fees and fees for serving on a steering committee from Eli Lilly, and fees for serving on a data and safety monitoring board from Iovance Biotherapeutics. Dr Park reports receiving research funding from Inhibrx, CUE Biopharma, ALX Oncology, Ankyra Therapeutics, A2 Biotherapeutics, and ISA Pharmaceuticals. No other disclosures were reported.

## REFERENCES

- Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2021;71(3):209-249. doi:10.3322/caac.21660
- Johnson DE, Burtneis B, Leemans CR, Lui VWY, Bauman JE, Grandis JR. Head and neck squamous cell carcinoma. *Nat Rev Dis Primers*. 2020;6(1):92. doi:10.1038/s41572-020-00224-3
- Barsouk A, Aluru JS, Rawla P, Saginala K, Barsouk A. Epidemiology, risk factors, and prevention of head and neck squamous cell carcinoma. *Med Sci (Basel)*. 2023;11(2):42. doi:10.3390/medsci11020042
- Hynes NE, Lane HA. ERBB receptors and cancer: the complexity of targeted inhibitors. *Nat Rev Cancer*. 2005;5(5):341-354. doi:10.1038/nrc1609
- Kalyankrishna S, Grandis JR. Epidermal growth factor receptor biology in head and neck cancer. *J Clin Oncol*. 2006;24(17):2666-2672. doi:10.1200/JCO.2005.04.8306
- Pivot X, Magné N, Guardiola E, et al. Prognostic impact of the epidermal growth factor receptor levels for patients with larynx and hypopharynx cancer. *Oral Oncol*. 2005;41(3):320-327. doi:10.1016/j.oraloncology.2004.09.011
- Ang KK, Berkey BA, Tu X, et al. Impact of epidermal growth factor receptor expression on survival and pattern of relapse in patients with advanced head and neck carcinoma. *Cancer Res*. 2002;62(24):7350-7356.
- Perisanidis C. Prevalence of EGFR tyrosine kinase domain mutations in head and neck squamous cell carcinoma: cohort study and systematic review. *In Vivo*. 2017;31(1):23-34. doi:10.21873/invivo.11020
- Sok JC, Coppelli FM, Thomas SM, et al. Mutant epidermal growth factor receptor (EGFRvIII) contributes to head and neck cancer growth and resistance to EGFR targeting. *Clin Cancer Res*. 2006;12(17):5064-5073. doi:10.1158/1078-0432.CCR-06-0913
- Li B, Hao Y, He H, et al. CD47-SIRPα blockade sensitizes head and neck squamous cell carcinoma to cetuximab by enhancing macrophage adhesion to cancer cells. *Cancer Res*. 2024;84(19):3189-3206. doi:10.1158/0008-5472.CAN-24-0176
- Bonner JA, Harari PM, Giral J, et al. Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck. *N Engl J Med*. 2006;354(6):567-578. doi:10.1056/NEJMoa053422
- Vermorken JB, Trigo J, Hitt R, et al. Open-label, uncontrolled, multicenter phase II study to evaluate the efficacy and toxicity of cetuximab as a single agent in patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck who failed to respond to platinum-based therapy. *J Clin Oncol*. 2007;25(16):2171-2177. doi:10.1200/JCO.2006.06.7447
- Vermorken JB, Mesia R, Rivera F, et al. Platinum-based chemotherapy plus cetuximab in head and neck cancer. *N Engl J Med*. 2008;359(11):1116-1127. doi:10.1056/NEJMoa0802656
- Rischin D, Spigel DR, Adkins D, et al. PRISM: phase 2 trial with panitumumab monotherapy as second-line treatment in patients with recurrent or metastatic squamous cell carcinoma of the head and neck. *Head Neck*. 2016;38(S1)(suppl 1):E1756-E1761. doi:10.1002/hed.24311
- Vermorken JB, Stöhlmacher-Williams J, Davidenko I, et al; SPECTRUM investigators. Cisplatin and fluorouracil with or without panitumumab in patients with recurrent or metastatic squamous-cell carcinoma of the head and neck (SPECTRUM): an open-label phase 3 randomised trial. *Lancet Oncol*. 2013;14(8):697-710. doi:10.1016/S1470-2045(13)70181-5
- Soulieres D, Senzer NN, Vokes EE, Hidalgo M, Agarwala SS, Siu LL. Multicenter phase II study of erlotinib, an oral epidermal growth factor receptor tyrosine kinase inhibitor, in patients with recurrent or metastatic squamous cell cancer of the head and neck. *J Clin Oncol*. 2004;22(1):77-85. doi:10.1200/JCO.2004.06.075
- Stewart JSW, Cohen EEW, Licitra L, et al. Phase III study of gefitinib compared with intravenous methotrexate for recurrent squamous cell carcinoma of the head and neck [corrected]. *J Clin Oncol*. 2009;27(11):1864-1871. doi:10.1200/JCO.2008.17.0530
- Machiels JPH, Haddad RI, Fayette J, et al; LUX-H&N 1 investigators. Afatinib versus methotrexate as second-line treatment in patients with recurrent or metastatic squamous-cell carcinoma of the head and neck progressing on or after platinum-based therapy (LUX-Head & Neck 1): an open-label, randomised phase 3 trial. *Lancet Oncol*. 2015;16(5):583-594. doi:10.1016/S1470-2045(15)70124-5
- Vermorken JB, Psyrri A, Mesia R, et al. Impact of tumor HPV status on outcome in patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck receiving chemotherapy with or without cetuximab: retrospective analysis of the phase III EXTREME trial. *Ann Oncol*. 2014;25(4):801-807. doi:10.1093/annonc/mdt574
- Fayette J, Licitra LFL, Harrington KJ, et al. 8540 INTERLINK-1: phase III study of cetuximab (CTX) & monalizumab (M) in participants (pts) with recurrent/metastatic head and neck squamous cell carcinoma (R/M HNSCC) with disease progression on/after platinum chemotherapy (CT) and previously treated with an immune checkpoint inhibitor (ICI). *Ann Oncol*. 2023;34:S554-S555. doi:10.1016/j.annonc.2023.09.2000
- Lui VWY, Hedberg ML, Li H, et al. Frequent mutation of the PI3K pathway in head and neck cancer defines predictive biomarkers. *Cancer Discov*. 2013;3(7):761-769. doi:10.1158/2159-8290.CD-13-0103
- Seiwert TY, Zuo Z, Keck MK, et al. Integrative and comparative genomic analysis of HPV-positive and HPV-negative head and neck squamous cell carcinomas. *Clin Cancer Res*. 2015;21(3):632-641. doi:10.1158/1078-0432.CCR-13-3310
- Lawrence MS, Sougnez C, Lichtenstein L, et al; Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*. 2015;517(7536):576-582. doi:10.1038/nature14129
- Chung CH, Li J, Steuer CE, et al. Phase II multi-institutional clinical trial result of concurrent cetuximab and nivolumab in recurrent and/or metastatic head and neck squamous cell carcinoma. *Clin Cancer Res*. 2022;28(11):2329-2338. doi:10.1158/1078-0432.CCR-21-3849
- Sacco AG, Chen R, Worden FP, et al. Pembrolizumab plus cetuximab in patients with recurrent or metastatic head and neck squamous cell carcinoma: an open-label, multi-arm, non-randomised, multicentre, phase 2 trial. *Lancet Oncol*. 2021;22(6):883-892. doi:10.1016/S1470-2045(21)00136-4
- Gulati S, Crist M, Riaz MK, et al. Durvalumab plus cetuximab in patients with recurrent or metastatic head and neck squamous cell carcinoma: an open-label, nonrandomized, phase II clinical trial. *Clin Cancer Res*. 2023;29(10):1906-1915. doi:10.1158/1078-0432.CCR-22-3886
- Guo Y, Li Z, Chen W, Fang M, Liu Z. Preliminary safety and efficacy of toripalimab combined with cetuximab in platinum-refractory recurrent or metastatic head and neck squamous cell carcinoma (R/M-HNSCC): a phase Ib/II clinical trial. 2022. *J Clin Oncol*. doi:10.1200/JCO.2022.40.16\_suppl.602
- Ulahannan SV, Marron TU, Park H, et al. Results from phase 1a/1b analyses of TTX-080, a first in class HLA-G antagonist, in combination with cetuximab in patients (pts) with metastatic colorectal cancer and head and neck squamous cell carcinoma. *J Clin Oncol*. 2024;42(16)(suppl):2524. doi:10.1200/JCO.2024.42.16\_suppl.2524
- Altan M, Patnaik A, Barve M, et al. 957 NKTR-255+cetuximab in patients with solid tumors: interim safety and efficacy results from the phase 1b dose-escalation study. *J Immunother Cancer*. 2021;9(suppl 2):A1007. doi:10.1136/jitc-2021-SITC2021.957
- Dunn L, Ho AL, Eng J, et al. A phase I/Ib study of lenvatinib and cetuximab in patients with recurrent/metastatic (R/M) head and neck squamous cell carcinoma (HNSCC). *J Clin Oncol*. 2020;38(15)(suppl):6541. doi:10.1200/JCO.2020.38.15\_suppl.6541



31. Desilets A, Pfister DG, Stein S, et al. A phase 1 study of concurrent cabozantinib and cetuximab in recurrent or metastatic head and neck squamous cell cancer. *Oral Oncol.* 2024;154:106861. doi:10.1016/j.oraloncology.2024.106861
32. Gilbert J, Schell MJ, Zhao X, et al. A randomized phase II efficacy and correlative studies of cetuximab with or without sorafenib in recurrent and/or metastatic head and neck squamous cell carcinoma. *Oral Oncol.* 2015;51(4):376-382. doi:10.1016/j.oraloncology.2014.12.011
33. Bauman JE, Saba NF, Roe D, et al. Randomized phase II trial of ficlatuzumab with or without cetuximab in pan-refractory, recurrent/metastatic head and neck cancer. *J Clin Oncol.* 2023;41(22):3851-3862. doi:10.1200/JCO.22.01994
34. Kochanny SE, Worden FP, Adkins DR, et al. A randomized phase 2 network trial of tivantinib plus cetuximab versus cetuximab in patients with recurrent/metastatic head and neck squamous cell carcinoma. *Cancer.* 2020;126(10):2146-2152. doi:10.1002/cncr.32762
35. Adkins D, Ley J, Neupane P, et al. Palbociclib and cetuximab in platinum-resistant and in cetuximab-resistant human papillomavirus-unrelated head and neck cancer: a multicentre, multigroup, phase 2 trial. *Lancet Oncol.* 2019;20(9):1295-1305. doi:10.1016/S1470-2045(19)30405-X
36. Adkins D, Lin JC, Sacco AG, et al. Palbociclib plus cetuximab versus placebo plus cetuximab in platinum-resistant, cetuximab-naïve, HPV-unrelated head and neck cancer: A double-blind randomized phase II trial (PALATINUS). *J Clin Oncol.* 2021;37(15)(suppl):6013. doi:10.1200/JCO.2019.37.15\_suppl.6013
37. Dennis MJ, Sacco AG, Qi Y, et al. A phase I study of avelumab, palbociclib, and cetuximab in patients with recurrent or metastatic head and neck squamous cell carcinoma. *Oral Oncol.* 2022;135:106219. doi:10.1016/j.oraloncology.2022.106219
38. Hanna GJ, Kaczmar J, Zandberg DP, et al. Updated dose expansion results of a phase 1/1b study of the bifunctional EGFR/TGFB inhibitor BCA101 with pembrolizumab in patients with recurrent, metastatic head and neck squamous cell carcinoma. *Int J Radiat Oncol Biol Phys.* 2024;118(5):e88. doi:10.1016/j.ijrobp.2024.01.194
39. Fayette J, Clatot F, Brana I, et al. Petosemtamab (MCLA-158) with pembrolizumab as first-line (1L) treatment of recurrent/metastatic (r/m) head and neck squamous cell carcinoma (HNSCC): phase 2 study. *J Clin Oncol.* 2024;42(16)(suppl):6014. doi:10.1200/JCO.2024.42.16\_suppl.6014
40. Bossi P, Minuti G, Braña I, et al. 362P efficacy and safety of MCLA-129, an anti-EGFR/c-MET bispecific antibody, in head and neck squamous cell cancer (HNSCC). *Ann Oncol.* 2023;34:S1610-S1611. doi:10.1016/j.annonc.2023.10.470
41. Xue L, Yang K, Fang M, et al. Results from two phase II studies of SI-BO01, an EGFR×HER3 bispecific antibody, with/without chemotherapy in patients (pts) with recurrent and metastatic head and neck squamous cell carcinoma (HNSCC). *J Clin Oncol.* 2023;41(16)(suppl):6037. doi:10.1200/JCO.2023.41.16\_suppl.6037
42. Xue L, Han Y, Zhang Q, et al. 939P Efficacy and safety of a novel anti-EGFR ADC MRG003 in recurrent or metastatic squamous cell carcinoma of the head and neck patients. *Ann Oncol.* 2023;34:S590. doi:10.1016/j.annonc.2023.09.2083
43. Cognetti DM, Johnson JM, Curry JM, et al. Phase 1/2a, open-label, multicenter study of RM-1929 photoimmunotherapy in patients with locoregional, recurrent head and neck squamous cell carcinoma. *Head Neck.* 2021;43(12):3875-3887. doi:10.1002/hed.26885
44. Ma Y, Huang Y, Zhao Y, et al. BL-B01D1, a first-in-class EGFR-HER3 bispecific antibody-drug conjugate, in patients with locally advanced or metastatic solid tumours: a first-in-human, open-label, multicentre, phase 1 study. *Lancet Oncol.* 2024;25(7):901-911. doi:10.1016/S1470-2045(24)00159-1
45. Papa S, Adami A, Metoudi M, et al. Intratumoral pan-ErbB targeted CAR-T for head and neck squamous cell carcinoma: interim analysis of the T4 immunotherapy study. *J Immunother Cancer.* 2023;11(6):e007162. doi:10.1136/jitc-2023-007162
46. Zhang S, Zheng M, Nie D, et al. Efficacy of cetuximab plus PD-1 inhibitor differs by HPV status in head and neck squamous cell carcinoma: a systematic review and meta-analysis. *J Immunother Cancer.* 2022;10(10):e005158. doi:10.1136/jitc-2022-005158
47. Cohen RB, Bauman JR, Salas S, et al. Combination of monalizumab and cetuximab in recurrent or metastatic head and neck cancer patients previously treated with platinum-based chemotherapy and PD-(L)1 inhibitors. *J Clin Oncol.* 2020;38(15)(suppl):6516. doi:10.1200/JCO.2020.38.15\_suppl.6516
48. McKean M, Rosen E, Barve M, et al. Abstract CT184: preliminary dose escalation results of ERAS-601 in combination with cetuximab in FLAGSHIP-1: a phase I study of ERAS-601, a potent and selective SHP2 inhibitor, in patients with previously treated advanced or metastatic solid tumors. *Cancer Res.* 2023;83(8)(suppl):CT184-CT184. doi:10.1158/1538-7445.AM2023-CT184
49. Boreddy SR, Nair R, Pandey PK, et al. BCA101 is a tumor-targeted bifunctional fusion antibody that simultaneously inhibits EGFR and TGFβ signaling to durably suppress tumor growth. *Cancer Res.* 2023;83(11):1883-1904. doi:10.1158/0008-5472.CAN-21-4425
50. Morgan RG, Mortenson E, Williams AC. Targeting LGR5 in colorectal cancer: therapeutic gold or too plastic? *Br J Cancer.* 2018;118(11):1410-1418. doi:10.1038/s41416-018-0118-6
51. Cohen EE, Fayette J, Daste A, et al. Abstract CT012: clinical activity of MCLA-158 (petosemtamab), an IgG1 bispecific antibody targeting EGFR and LGR5, in advanced head and neck squamous cell cancer (HNSCC). *Cancer Res.* 2023;83(8)(suppl):CT012. doi:10.1158/1538-7445.AM2023-CT012
52. Comer F, Mazor Y, Hurt E, et al. Abstract 5736: AZD9592: an EGFR-cMET bispecific antibody-drug conjugate (ADC) targeting key oncogenic drivers in non-small-cell lung cancer (NSCLC) and beyond. *Cancer Res.* 2023;83(7)(suppl):5736. doi:10.1158/1538-7445.AM2023-5736
53. Aggarwal C, Azzoli CG, Spira AI, et al. EGRET: a first-in-human study of the novel antibody-drug conjugate (ADC) AZD9592 as monotherapy or combined with other anticancer agents in patients (pts) with advanced solid tumors. *J Clin Oncol.* 2023;41(16)(suppl):TPS3156. doi:10.1200/JCO.2023.41.16\_suppl.TPS3156
54. Fucà G, Spagnoletti A, Ambrosini M, de Braud F, Di Nicola M. Immune cell engagers in solid tumors: promises and challenges of the next generation immunotherapy. *ESMO Open.* 2021;6(1):100046. doi:10.1016/j.esmoop.2020.100046
55. Salek-Ardakani S, DiRaimondo T, Budimir N, et al. 1123 Preclinical activity and safety profile of JANX008, a novel EGFR-targeting tumor-activated T cell engager for treatment of solid tumors. *J Immunother Cancer.* 2022;10(suppl 2):A1167. doi:10.1136/jitc-2022-SITC2022.1123
56. Autio KA, Boni V, Humphrey RW, Naing A. Probody therapeutics: an emerging class of therapies designed to enhance on-target effects with reduced off-tumor toxicity for use in immuno-oncology. *Clin Cancer Res.* 2020;26(5):984-989. doi:10.1158/1078-0432.CCR-19-1457
57. Bigelow M, LaCroix C, Chawla MK, et al. 1060 DF9001, an EGFR targeted immune engager, stimulates innate and adaptive anti-tumor immunity with a distinctive safety profile. *J Immunother Cancer.* 2024;12(suppl 2):A1180. doi:10.1136/jitc-2024-SITC2024.1060
58. Sterner RC, Sterner RM. CAR-T cell therapy: current limitations and potential strategies. *Blood Cancer J.* 2021;11(4):69. doi:10.1038/s41408-021-00459-7
59. Xu Y, Xiang Z, Alnaggar M, et al. Allogeneic Vy9Vδ2 T-cell immunotherapy exhibits promising clinical safety and prolongs the survival of patients with late-stage lung or liver cancer. *Cell Mol Immunol.* 2021;18(2):427-439. doi:10.1038/s41423-020-0515-7
60. Li HK, Wu TS, Leng PR, et al. Abstract LBO89: ACE2016: an off-the-shelf EGFR-targeting yδ2 T cell therapy against EGFR-expressing solid tumors. *Cancer Res.* 2023;83(8)(suppl):LBO89. doi:10.1158/1538-7445.AM2023-LBO89