

# Pectus Arcuatum Definitions, Diagnostics, and Surgical Guidelines

## A Systematic Review

Rawan M. Zeineddine, MD; Ahmed E. Khedr, MBBS; Juan M. Farina, MD; Nour B. Odeh, MBBS; Abdulrahman Senjab, MBBS; J. Asher Jenkins, MD; Dawn E. Jaroszewski, MD, MBA

**IMPORTANCE** Pectus arcuatum is a rare anterior chest wall deformity that is often confused with excavatum, carinatum, or a mixed deformity. A complete understanding of its clinical and physiological basis will improve diagnostic accuracy and allow for standardized management.

**OBJECTIVES** To gain a comprehensive understanding of pectus arcuatum, focusing on its nomenclatures, epidemiology, classification, pathophysiology, preoperative presentation and assessments, management options, and surgical outcomes.

**EVIDENCE REVIEW** PubMed, Google Scholar, and subsequent reference searches were used to extract original studies addressing pectus arcuatum fully or partially. The searches were performed on February 2, 2025. Two independent reviewers screened articles based on preestablished inclusion and exclusion criteria. Quality assessment was performed, but given the rarity of arcuatum and the limited, heterogeneous nature of available literature, none of the studies were excluded based on the level of evidence or quality assessments. Using a standardized data extraction form, results were synthesized and presented as a review with recommendations to guide clinical decision-making.

**FINDINGS** A total of 65 studies were included in the final review. Of those, 37 (56.9%) were cohort studies, 18 (27.7%) were case reports, 8 (12.3%) were case series, and 2 (3.1%) were technical reports. A total of 455 cases of pectus arcuatum were represented. Heterogeneity was found in nomenclature, epidemiology, associated conditions, assessments, management, and outcomes. An open surgical approach and, more recently, hybrid open and minimally invasive technique are reported for surgical repair. Postoperative outcomes were good in most surgical series, with no significant difference when compared with the repair of other pectus deformities.

**CONCLUSIONS AND RELEVANCE** Results of this systematic review reveal that pectus arcuatum is a rare variant of pectus deformities that is distinct from carinatum and excavatum. Unified terminology and standardized diagnostic criteria are important to facilitate proper management. Minimally invasive hybrid approaches continue to refine outcomes, balancing invasiveness with efficacy.

JAMA Surg. doi:10.1001/jamasurg.2025.2365  
Published online July 23, 2025.

 [Supplemental content](#)

 [CME at jamacmelookup.com](#)

**Author Affiliations:** Department of Cardiovascular and Thoracic Surgery, Mayo Clinic, Phoenix, Arizona.

**Corresponding Author:** Dawn E. Jaroszewski, MD, MBA, Department of Cardiovascular and Thoracic Surgery, Mayo Clinic Arizona, 5881 E Mayo Boulevard, Phoenix, AZ 85054 ([jaroszewski.dawn@mayo.edu](mailto:jaroszewski.dawn@mayo.edu)).

**P**ectus arcuatum is a unique anterior chest wall deformity involving premature fusion of the sternal ossification centers and obliteration of the manubriosternal joint. It may be misdiagnosed as excavatum, carinatum, or mixed deformity. The literature shows significant variation in clinical aspects and terminology. Difficulties in obtaining clear information and misunderstanding of the deformity have hindered accurate assessment and standardized management. This systematic review aims to provide a comprehensive summary of the current literature to guide diagnostic and treatment recommendations.

### Methods

#### Study Design

This systematic review provides a comprehensive summary of arcuatum, focusing on its nomenclature, epidemiology, classification,

pathophysiology, preoperative presentation and assessments, management, and outcomes. It followed the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guidelines; the protocol is available on request.

#### Search Strategy

A structured search was conducted in PubMed and Google Scholar databases using MeSH terms and free-text keywords. PubMed used the following search terms: ("Pectus Carinatum"[Mesh]) OR (Currarino-Silverman syndrome) OR (Type II pectus carinatum) OR (Pouter pigeon chest) OR ("Horns of Steer") OR (Chondromanubrial deformity) OR (arcuatum). Google Scholar used the following search terms: ("Pectus Carinatum") OR (Currarino-Silverman syndrome) OR (Type II pectus carinatum) OR (Pouter pigeon chest) OR ("Horns of Steer") OR (Chondromanubrial deformity) OR (arcuatum). No time limits were set, and the search was performed on February 2, 2025.

## Screening and Selection

Initial search results were deduplicated and screened by title and abstract using an artificial intelligence–powered tool. Ambiguous or relevant papers underwent full-text screening. Additional articles were identified through citation tracking. Two reviewers (R.M.Z., A.E.K.) independently screened studies. Disagreements were resolved by consensus with an expert (D.E.J.).

### Inclusion Criteria

- Studies specifically addressing arcuatum (fully or as part of a mixed cohort).
- English-language publications.
- Technical papers and original studies (randomized clinical trials, cohort, case-control, series, reports).

### Exclusion Criteria

- Nonhuman studies.
- Full article text not available.
- Reports of superior sternal protrusion or mixed deformities without clear evidence of arcuatum.
- Studies on various pectus deformities without stratified results by type.

## Data Extraction

A single expert reviewer (R.M.Z.) extracted data using a standardized form (eTable 1 in the [Supplement](#)) to ensure accuracy and capture of nuances specific to arcuatum.

## Quality Assessment and Data Synthesis

Study quality was assessed using ROBINS-I for observational studies and JBI tools for case series/reports (eTable 2 in the [Supplement](#)). Given the rarity and heterogeneity of arcuatum studies, none were excluded based on level of evidence or quality. All relevant studies were included for comprehensive coverage. No meta-analysis was conducted. Outcomes were synthesized descriptively where available.

## Results

From the database screening, 971 articles were identified. After removing 46 duplicates, 925 papers underwent title and abstract screening. Of these, 260 studies were selected for full-text review. Applying eligibility criteria and adding 8 papers from reference screening, 65 papers were included. Of those, 37 (56.9%)<sup>1-37</sup> were cohort studies, 18 (27.7%)<sup>38-55</sup> were case reports, 8 (12.3%)<sup>56-63</sup> were case series, and 2 (3.1%)<sup>64,65</sup> were technical reports. Cohen  $\kappa$  of 0.7909 indicates substantial reviewer agreement. A total of 455 arcuatum cases were represented. Study characteristics are summarized in eTable 3 in the [Supplement](#); the PRISMA flow diagram appears in the eFigure in the [Supplement](#).

## Nomenclatures

Arcuatum ([Figure 1](#)) has been referred to by various names as follows: (1) Currarino-Silverman syndrome,<sup>56</sup> (2) angulated synostosis,<sup>1</sup> (3) arcuate deformity,<sup>2</sup> (4) chondromanubrial deformity,<sup>66</sup> (5) manubriosternal prominence,<sup>3</sup> (6) type II pectus carinatum,<sup>4,5</sup> (7) horns of steer deformity,<sup>57</sup> (8) pouter pigeon breast deformity,<sup>6,38</sup> (9) pectus carinatum superior,<sup>7</sup> and (10) Wenlin chest.<sup>39</sup>

## Key Points

**Question** What are the standardized definitions, diagnostics, and surgical guidelines for pectus arcuatum?

**Findings** In this systematic review including 65 studies, preoperative computed tomography/magnetic resonance imaging and echocardiography were recommended to assess cardiopulmonary function and screen for congenital heart disease. Symptomatic patients should be considered for surgery, with expected improvement.

**Meaning** Results suggest that surgical correction relied on wedge osteotomy, with titanium plating and bar placement being key for stabilization and lower cartilage remodeling. Hybrid techniques continue to evolve, aiming to balance invasiveness and efficacy.

## Epidemiology

The prevalence and demographics of arcuatum remain unclear in part due to an underreporting and misdiagnosis of these deformities. Arcuatum is commonly considered a rare pectus variant, accounting for 1% or fewer cases.<sup>8,58</sup> It is likely more prevalent. In a 10-year study, Kelly et al<sup>9</sup> found 7% of patients (31 patients) had arcuatum. Haje et al<sup>10</sup> reported 5% among 4012 patients over 32 years.

Our review identified 455 cases: 15% female, 22% male, 63% unreported (eTable 3 in the [Supplement](#)). Lacquet et al<sup>11</sup> reported equal sex distribution. Zeineddine et al<sup>12</sup> noted slight male predominance (11 males vs 9 females).

A recent study (2017-2021)<sup>13</sup> of 1520 children with anterior chest wall deformities found 17 arcuatum cases: 6 Hispanic, 10 non-Hispanic White, and 1 unspecified race and ethnicity.


## Classification

Despite the description by Currarino-Silverman of the hallmarks of arcuatum in 1958,<sup>56</sup> its unique combination of features, some overlapping with other deformities, has led to the deformity being misdiagnosed as other pectus variants.

Some viewed arcuatum as a type of protrusion deformity or carinatum variant. In 1952, Lester divided protrusion deformities into midline (sternum/cartilage) and lateral anterior chest wall types, further separating midline into upper arcuate and lower oblique forms.<sup>2,14</sup> In 1957, Lester categorized pectus into excavatum, carinatum, or mixed types, proposing a shared etiologic basis.<sup>14</sup> In 1958, Brodtkin identified 2 protrusion types: upper chondromanubrial and lower chondrogladiolar prominence, the former involving manubrium and adjacent first and second cartilages with a depressed gladiolus.<sup>59</sup> Robiscek et al<sup>5</sup> divided carinatum into 3 types, with type II (pouter pigeon breast) described as a variant featuring chondromanubrial prominence, protrusion of the first 2 cartilages, a posteriorly arched sternal body, and an anteriorly pointing xiphoid tip. In 1999, Haje et al<sup>7</sup> categorized pectus into 5 types based on sternal protrusion/depression localization, referring to arcuatum as *pectus carinatum superior*.

Given the lower degree of depression seen in arcuatum, some consider it an excavatum variant.<sup>8,57</sup> Others suggest arcuatum as a mixed deformity.<sup>15</sup> In 1991, Saxena and Willital<sup>15</sup> classified chest wall deformities into 11 types with arcuatum (type 9) being a transitional form: funnel chest (4 types), pigeon chest (4 types), funnel-pigeon combinations (type 9), thoracic wall aplasia (type 10), and cleft sternum (type 11).

Figure 1. Physical Features of Pectus Arcuatum



A-C, Typical physical features of pectus arcuatum: manubriosternal prominence, lower sternal depression, and abnormally inserted upper ribs forming the characteristic "horns of a steer" appearance.

Acastello<sup>67</sup> classified chest wall deformities by defect origin: cartilaginous (type 1), costal (type 2), chondrocostal (type 3), sternal (type 4), and clavicle-scapular (type 5), attributing arcuatum to cartilaginous type 2 or superior carinatum.<sup>16</sup>

Recently, surgeons have stressed on distinguishing arcuatum from other pectus deformities, emphasizing that it is not a mixture of these conditions.<sup>12,17,18</sup>

### Pathophysiology

Arcuatum was recognized as a congenital deformity but often unnoticed until adolescence.<sup>14,59</sup> Brodtkin proposed his diaphragmatic theory in 1949<sup>66</sup> with reinforcement in 1957,<sup>59</sup> suggesting unbalanced contractions of a malformed diaphragm (V-shaped fibrotendinous structure with thin anterior and hypertrophied lateral segments) drive the deformity. These forces push the sternum and cartilage forward, causing upper protrusion and lower retraction during inspiration. As the thorax elongates, upper protrusion persists while the lower gladiolus flattens. Limited evidence without control groups hinders confirmation of observed patterns as unique to arcuatum.<sup>59</sup>

In 1952 and 1957,<sup>2,14</sup> Lester combined published observations<sup>66</sup> with intraoperative findings, suggesting disproportionate second to fourth rib overgrowth and diaphragm shape drives the deformity. Rib overgrowth creates force and buckles at weak points, typically the costochondral junction, pushing the sternum forward. If the diaphragm fails to adjust, the xiphoid restrains it, causing arcuate sternal protrusion.

Premature sternal ossification centers fusion and manubriosternal joint obliteration were coined by Currarino-Silverman in 1958<sup>56</sup> and supported by others.<sup>1,7</sup> Normally, ossification begins at 5 to 6 months in fetal development, with segments fusing gradually. Complete sternal fusion is rare, and manubriosternal fusion occurs in only 10% of adults. However, arcuatum patients uniformly exhibit early manubriosternal fusion, confirmed radiologically.<sup>7,12,60</sup> Fokin<sup>1</sup> compared premature sternal fusion radiograms with earlier ones, found abnormally narrowed sternal sutures, supporting premature obliteration over failed segmentation. Causes include undeveloped fibrous lamina or pathological

joint degeneration.<sup>56</sup> Ravitch supported Currarino-Silverman, emphasizing rib overgrowth as primary drivers of arcuatum rather than sternal abnormalities. He dismissed the diaphragmatic theory, arguing that the diaphragm only accentuates the deformity. He suggested rib lateral incurvature makes the sternum appear prominent and compensates for the latent depressed area,<sup>6</sup> but in arcuatum, the ribs are bulky and override the sternum rather than curve inward.<sup>12</sup> Fokin also identified second to fifth cartilage overgrowth with hyaline cartilage degeneration, atypical fibrils, reduced chondrocytes, and thin periosteum.<sup>1</sup>

Abnormal differentiation of anterior mesenchymal cells during cardiogenesis may disrupt endocardial cushions, sternum, and aortic derivatives, explaining arcuatum's link to congenital heart diseases (CHDs) and supporting a potential genetic component.<sup>68</sup>


### Preoperative Presentation and Assessment

#### History

- **Congenital:** typically noticeable before age 3 years, often within the first year of life.<sup>1,18,38,56,59</sup>
- **Progression:** Fokin observed progression in 50% of patients (n = 18), with a 2° per year nonlinear progression rate uncorrelated with lower sternal depression.<sup>1</sup> Currarino-Silverman reported 1 case with straight sternum at 1 year, featuring narrowed, irregular sutures, became angulated by age 4 years, and another with arcuatum evident at 8 months, worsening over time.<sup>56</sup> Multiple reports support progression (Figure 2A and B).<sup>40-42</sup>
- **Psychosocial implications:** the psychosocial effects are significant. Patients often avoid chest-revealing activities/clothing, show abnormal posture, low self-esteem, anxiety, or depression.<sup>1,12,16,59,64</sup> Fokin noted psychological effects begin around age 7 years and peak by 10 years,<sup>1</sup> although cosmetic concerns persist into adulthood.<sup>12</sup> Many prefer surgical scars over the deformity, sometimes viewing them as symbols of victimhood, reflecting the profound psychological impact.<sup>1</sup>
- **Cardiovascular symptoms:** chest pain and exercise intolerance are most common.<sup>1,12,19,61</sup> Cardiac compression, displacement, and pulmonary artery compression can result from inward lower sternum and cartilages angulation (Figure 3D).<sup>12,38,43</sup>




Figure 2. Patient With Progressive Features of Pectus Arcuatum: Before and After Hybrid Repair



Pectus arcuatum evident from a young age (A), which progressed over time (B). The patient subsequently underwent hybrid repair (C), during which her sternum was stabilized using an H plate after a single wedge osteotomy, her abnormal ribs were reapproximated and rigidly fixed bilaterally, and 2 bars were

placed to correct her lower sternal depression. One bar was positioned at a low point below the sternal end to support lower costal cartilage remodeling and relieve cardiac compression (D).

Figure 3. Preoperative Assessments



A 3-dimensional (3-D) reconstruction (A) and computed tomography (CT) views of a short, unsegmented (B), broad sternum with bulky and calcified rib insertions (C) and crowded lower costal cartilages compressing the heart (D). Positional transthoracic echocardiogram demonstrates right ventricular

compression at the lower costal cartilages, with the right ventricular diameter decreasing from 25.93 mm (supine) (E) to 14.58 mm (bending over) (F) and the right ventricular outflow tract velocity-time integral decreasing from 13.01 cm (supine) (G) to 9.50 cm (bending over) (H).

- Pulmonary issues: pulmonary symptoms are debated. Some link them to decreased respiratory excursion and increased anterior-posterior diameter, leading to infections.<sup>12,19</sup> Others consider them uncommon and milder than in excavatum.<sup>1</sup> Postoperative improvement and reduced asthma have been reported.<sup>20</sup> Pulmonary evaluation may help, as cases of diaphragmatic

dysfunction, restrictive/obstructive diseases, and recurrent infections exist.<sup>1,20,62</sup>

- Family history: many patients with pectus arcuatum have relatives with the condition. Chidambaram reported arcuatum in 2 siblings and their mother.<sup>60</sup> Fokin found 4 of 11 patients had close relatives with chest deformities, 2 with confirmed parental

sternal synostosis.<sup>1</sup> Abdellaoui et al<sup>18</sup> reported 32% (n = 34) with skeletal malformation family history. Zeineddine et al<sup>12</sup> noted 7 of 16 patients had a family history of pectus, none being arcuatum. Currarino-Silverman included 3 cases, 1 with an affected maternal aunt, and frequent manubriosternal synchondrosis in parents.<sup>56</sup> Inheritance appears polygenic or multifactorial. Some reported shared genetic links across anterior chest deformities.<sup>14,62,69</sup>

- CHD: arcuatum has been associated with various CHD (Table).<sup>1,11,12,16-18,21,40,42,44-48,56,58,60,62,64</sup> Chidambaram et al<sup>60</sup> noted mitral valve prolapse with mild regurgitation in 1 sibling and their mother.
- Syndromic associations: arcuatum may be isolated or associated with connective tissue diseases, spinal deformities, and syndromes. Lacquet et al<sup>11</sup> reported 3 of 37 hemiarctuatum cases over 25 years, associated with contralateral Poland syndrome. Kuzmichev et al<sup>64</sup> noted occasional Poland syndrome. Zeineddine et al<sup>12</sup> reported 1 of 20 patients with Klinefelter syndrome. Noonan syndrome has also been reported.<sup>18,47,48</sup> The Table summarizes all anomalies.<sup>1,11,12,16-18,21,40,42,44-48,56,58,60,62,64</sup>

### Physical Examination

- Chest morphology: upper sternal convexity and lower concavity with prominent rib insertions create the classic horn of steer appearance (Figure 1A-C).
- Functional murmurs: Fokin reported functional murmurs in 7 of 18 patients, likely caused by right ventricular outflow tract obstruction.<sup>1</sup>
- Associated anomalies: evaluation for evidence of connective tissue disorders or other musculoskeletal abnormalities.

### Imaging

**Radiographs** | Chest radiograph and computed tomography (CT) are essential for preoperative assessment; 3-dimensional CT or magnetic resonance imaging are considered best.<sup>16,18</sup> Findings include upper sternal protrusion, lower depression, and abnormal costosternal junctions. The sternum is short, broad, and nonsegmented (1-2 segments) (Figure 3A-C).<sup>6,7,12,18,22,56</sup> Manubriosternal fusion has been reported,<sup>7</sup> with a reduced angle of Louis (normal: 175°-145°), increasing sagittal chest diameter.<sup>1,6,18,56</sup> Acute proximal third sternal angulation is common in arcuatum, unlike the gradual curvature in carinatum.<sup>22</sup> The manubrial orientation also differs from excavatum, where it is more vertical.<sup>23</sup> A Z-shaped sternum, or angulated synostosis, appears on lateral views.<sup>1,6</sup>

The xiphoid-vertebral body distance is often normal, yielding a normal Haller Index.<sup>12,14,24</sup> Lower sternum depression occurs in at least one-third of cases.<sup>1</sup> The lower sternal end tends to be vertical, unlike excavatum (posterior) and carinatum (anterior).<sup>22</sup> Less commonly, the xiphoid may be asymmetric, bifid, or absent.<sup>6,16,63</sup>

The deformity is usually symmetric.<sup>1</sup> Costosternal insertions are often bulky and heavily calcified (hyperostosis), primarily affecting second to fourth/fifth ribs.<sup>1,12,25,26,63</sup> Lower cartilages are often longitudinally clustered at their sternal insertion,<sup>12,58</sup> possibly causing right ventricular compression.<sup>12,43</sup> The inward buckling and compression from these lower cartilages can be position-dependent and may be missed on supine imaging (Figure 3D-H).<sup>12</sup>

**Table. Details of the Articles Reporting Associated Congenital Heart Diseases Among Other Anomalies**

Source	Cases with pectus arcuatum, No.	Associated anomalies
<b>Associated congenital heart diseases</b>		
Currarino and Silverman, <sup>56</sup> 1958	3	(1/3) Large PDA, large VSD, pulmonary artery and arteriolar stenosis
Briinner, <sup>44</sup> 1961	1	ToF and PDA
Kaplan et al, <sup>45</sup> 1968	1	Pulmonic stenosis
Welch and Vos, <sup>62</sup> 1973	2	(1/2) Double aortic arch and PDA
Chidambaram and Mehta, <sup>60</sup> 1992	5	(1/5) Parachute mitral valve with mild mitral stenosis and regurgitation, (1/5) endocardial cushion type of atrioventricular valve with mild to moderate MR, (2/5) MVP with mild MR, (1/5) aortic coarctation, (1/5) ostium secundum type of ASD that closed spontaneously
Mehta, <sup>46</sup> 1996	1	Aortic coarctation and parachute mitral valve
Fokin, <sup>1</sup> 2000	18	(2/18) VSD, (1/18) ToF
Kuzmichev et al, <sup>64</sup> 2016	NA	Unspecified CHD
Hysi et al, <sup>21</sup> 2015	3	(1/3) ASD, (1/3) ToF, recurrent VSD and pulmonary and aortic stenosis
Gritsiuta et al, <sup>16</sup> 2021	4	(1/4) MR and tricuspid regurgitation, (1/4) MVP (3 mm) and VSD (3 mm), (1/4) MVP (5 mm)
Abdellaoui et al, <sup>18</sup> 2023	34	(2/34) Unspecified CHD
St-Louis et al, <sup>17</sup> 2024	5	(1/5) Mild anterior mitral valve leaflet thickening
<b>Other associated anomalies</b>		
Currarino and Silverman, <sup>56</sup> 1958	3	2 Kyphosis, 1 kyphoscoliosis, case 1: mild congenital micrognathia, bilateral clubfoot, inguinal hernia, and cryptorchidism. Case 2: underdeveloped physical growth, craniofacial asymmetry, occipital and frontal flattening, microphthalmia, cataract, incontinentia pigmenti, aberrant subclavian artery causing dysphagia
Kaplan et al, <sup>45</sup> 1968	1	NS
Shamberger and Welch, <sup>58</sup> 1988	5	1 Bilateral clubfeet
Lacquet et al, <sup>11</sup> 1998	37	3 Poland syndrome
Fokin, <sup>1</sup> 2000	18	1 Ossification of T3-T4 disk
Kuzmichev et al, <sup>64</sup> 2016	NA	Poland syndrome
Alkhunaizi et al, <sup>47</sup> 2018	1	NS
Sahu et al, <sup>48</sup> 2020	1	NS
Gritsiuta et al, <sup>16</sup> 2021	4	2 Kyphosis, 1 kyphoscoliosis
Wang et al, <sup>40</sup> 2022	1	Poland syndrome
Wang et al, <sup>42</sup> 2022	1	Scoliosis
Abdellaoui et al, <sup>18</sup> 2023	34	4 NS, 3 scoliosis, 1 cervical rib, 1 cervical-dorsal vertebral malformation, 1 Baraitser-Winter syndrome, diaphragmatic hernia
Zeineddine et al, <sup>12</sup> 2024	20	1 Klinefelter syndrome, 6 scoliosis

Abbreviations: ASD, atrial septal defect; CHD, congenital heart diseases; MR, mitral regurgitation; MVP, mitral valve prolapse; NA, not available; NS, Noonan syndrome; PDA, patent ductus arteriosus; ToF, tetralogy of Fallot; VSD, ventricular septal defect.

**Echocardiography** | Echocardiography is recommended preoperatively due to the high association between arcuatum and CHD.<sup>1</sup> Positional echocardiography is particularly effective in detecting transient cardiac compression during positional changes, often missed in standard examinations (Figure 3E-H).<sup>12</sup>

**Other Tests** | Cardiopulmonary exercise testing (CPET) has been used in patients with exercise limitation. Zeineddine et al<sup>12</sup> reported decreased maximal oxygen consumption (VO<sub>2</sub>; median, 67.0%; IQR, 57.3%-69.0% of predicted) and mild restrictive patterns in 38.5% on pulmonary testing. González-Barba et al<sup>24</sup> found no CPET abnormalities in 3 patients.

## Management

### Noninvasive Approach

As a rule, bracing is largely ineffective in arcuatum due to early sternal rigidity and is not recommended.<sup>9,18,27-30</sup> Haje et al,<sup>10</sup> however, reported outcomes for his bracing protocol, which included individualized orthosis, prolonged treatment, exercise to balance internal and external pressures, close follow-up, and expert pressure adjustments. In his 32-year study of 69 children and 4 adults with arcuatum, 7.3% of children had good-to-excellent improvement and 21.7% moderate improvement.<sup>10</sup> In adults, only 1 patient noted some moderate improvement.<sup>10</sup>

### Suggested Surgical Indications

Cosmetic dissatisfaction is the most reported surgical indication.<sup>16,17,24,49,59,64</sup> Others emphasized symptoms and cardiopulmonary compromise as indication for surgery.<sup>9,12,20</sup> Fokin provided recommendations based on disease progression, syndromic associations, cardiopulmonary findings, and angle of Louis:

- Mild deformities (angle >130°): annual follow-up.
- Moderate deformities (angle 130°-115°): surgery, especially with lower depression.
- Severe deformities (angle <115°): surgery.

### Surgical Approach

The evolution of arcuatum surgical techniques reflects ongoing refinement. Ravitch introduced the first operative approach in 1952, involving extensive cartilage resection and sternal osteotomies.<sup>6</sup> Over decades, surgeons modified and supported this open technique.<sup>9</sup> In 1957, Chin proposed xiphosternopexy, reattaching the xiphoid at a higher level to reposition the sternum over time, though it was not applied to arcuatum.<sup>3</sup> Brodtkin later referenced it for 2 cases of unilateral chondromanubrial deformities.<sup>15,59</sup> Although Chin's xiphosternopexy supported Brodtkin's diaphragmatic theory, concerns remain. Brodtkin described chondromanubrial deformity as an upper protrusion with a flat or depressed lower part featuring the Harrison groove, sometimes unilateral. This describes a carinatum variant, not arcuatum. He later noted surgery was ineffective for non-unilateral cases, which likely represent true arcuatum.<sup>59</sup> Howard was successful with Chin's method only after modifications with cartilage and sternal resections.<sup>30</sup>

Advancements in surgical expertise have led to a shift toward less invasive procedures.<sup>12,17,29</sup> Since Onen introduced the hybrid technique in 2016,<sup>31</sup> it has shown good outcomes (Figure 2 and Figure 4) with low complications.<sup>12,17,18,24,29,31,32,50</sup> eTable 4 in the Supplement outlines the chronological progression of techniques.

Age recommendations for surgery remain controversial. Some advocate waiting for rib growth completion,<sup>64</sup> whereas others prefer early adolescence when cartilage is most flexible and surgery less demanding.<sup>11,20</sup> Challenges with younger patients include postoperative noncompliance and thoracic dystrophy,<sup>11,16</sup> whereas older patients face greater surgical difficulty due to sternum rigidity.<sup>11</sup> For patients with connective tissue disorders, surgery is advised after growth completion due to high regression risk.<sup>11</sup>

For patients with concurrent cardiac abnormalities, whether to perform surgery concurrently or in stages remains debated. Hysi et al<sup>21</sup> recommend Ravitch for optimal exposure and postoperative management. The hybrid approach is an option but may risk reduced cardiac compression efficacy and increased midsternotomy dehiscence. If intrathoracic bars are placed, pericardial closure or a barrier is required to prevent cardiac adhesions that may cause injury during bar removal.<sup>21,70</sup>

### Postoperative Complications and Outcomes

Few complications are reported in the literature,<sup>1,9,17,24</sup> likely due to the lack of stratified, dedicated studies. For the open approach, Ravitch<sup>38</sup> reported in his first case of right pleural effusion requiring thoracentesis. Kelly et al<sup>9</sup> reported in their 10-year experience 1 wound infection managed with antibiotics. For the hybrid approach, Zeineddine et al<sup>12</sup> reported few complications, the most significant being 1 hemothorax requiring pigtail insertion and transfusion and the other complications being 3 pleural effusions requiring thoracentesis.

Recovery was satisfactory and comparable to other pectus deformities.<sup>1,12,25,64</sup> Most had good cosmetic outcomes<sup>1,11,12,24,38</sup>; however, residual prominence or asymmetry due to inadequate resection was reported.<sup>11</sup> Long-term cardiopulmonary improvement was subjectively validated by patients.<sup>1,12</sup>

No special pain management strategies are needed beyond standard pectus protocols. Intercostal cryoablation<sup>24</sup> reduced epidural anesthesia use<sup>64</sup> and shorten hospital stays.<sup>12,17</sup>


## Discussion

The one consistency in arcuatum literature is the variability in nomenclature and classification, leading to inconsistency in identification and treatment recommendations. Recognizing arcuatum as a separate entity is essential due to its origin, features, and unique management, a concept acknowledged by some.<sup>11,12,18,71</sup> Misclassification has been extensive: Ravitch misidentifying arcuatum,<sup>38</sup> Lester's initial diagnosis of arcuatum was excavatum,<sup>14</sup> later corrected,<sup>2</sup> but still considered a carinatum variant,<sup>14</sup> and more recent papers classifying arcuatum as an excavatum variant<sup>1,6,8,16,40,57</sup> or a mixed deformity.<sup>8,72</sup>

Arcuatum and superior carinatum are used interchangeably, highlighting confusion in the literature. Superior pectus carinatum features high sternal protrusion but lacks arcuatum's hallmark: fused, short, broad sternum supporting its sternal origin. Although once grouped with type II carinatum and considered a cartilaginous anomaly, Torre et al<sup>73</sup> modified Acastello's classification, distinguishing type II carinatum (cartilaginous) from arcuatum (sternal). Unlike true type II carinatum, arcuatum is often associated with lower sternal depression and frequently overlooked.<sup>74</sup> Unlike excavatum,



**Figure 4. Radiographic Studies and Intraoperative Transesophageal Echocardiogram (TEE) Demonstrating Cardiac Compression Before Surgery and Its Complete Resolution After Hybrid Repair**



Computed tomography (CT) (A) and TEE (B) showing right ventricular compression (pink arrowhead and dotted line) by the lower costal cartilages. This resolved after the hybrid approach (C), as demonstrated by intraoperative TEE (D) (blue dotted line).

arcuatum typically has normal Haller Index despite some depression, which can cause cardiac compression and symptoms.<sup>3,12,14,24</sup> Arcuatum is not merely a coexistence of excavatum and carinatum but a unique deformity with varying protrusions and depressions.<sup>11,12,18,71</sup>

History taking and a workup, including cardiopulmonary testing in symptomatic patients, is recommended. Arcuatum's prominent upper sternum is harder to conceal than excavatum, contributing to its significant psychosocial impact and possibly explaining the near-equal sex distribution, as breast tissue masking is less effective. Labeling arcuatum as purely cosmetic overlooks its physiologic potential.<sup>16,61</sup> The incidence of symptomatic patients remains

unclear for multiple reasons. The deformity is likely underdiagnosed, and even when recognized, many believe it to be purely cosmetic due to its relatively normal Haller Index,<sup>17,59,64</sup> a factor affecting physicians' clinical assessment and decisions for ordering testing. Symptoms can be nonspecific, with the most common being exercise-associated dyspnea, chest pain, and tachycardia. Experienced surgeons noted symptom resolution after surgical repair.<sup>6,12,19</sup>

Though early literature highlights cardiopulmonary effects,<sup>6,38</sup> strong evidence remains limited. A recent study by Zeineddine et al<sup>12</sup> underscored dynamic cardiac compression from crowded lower costal cartilages during positional changes (eg, leaning

forward), offering insight into physiologic symptoms. Thorough history-taking and follow-up after management can help link symptoms to their true etiology and better documents surgical outcomes.

An important association in literature is CHD in patients with abnormal sternal ossification, particularly arcuatum.<sup>1,60,68,75,76</sup> Although Currarino-Silverman first described this syndrome by presenting cases with CHD<sup>56</sup> and patterns exist (Table), including CHD as diagnostic criteria is inaccurate, as many confirmed cases lack this association.<sup>24,51</sup> Preoperative echocardiogram is recommended, as CT alone may miss functional compromise and does not screen for CHD.<sup>16,59,64</sup>

Connective tissue disease associations exist (Table) but remain poorly supported.<sup>16</sup> Studies often exclude such patients,<sup>57</sup> report them as mixed deformities without clarification,<sup>77</sup> or fail to establish such association. Still, targeted evaluation is recommended when history or physical findings suggest such conditions.<sup>12,17</sup>

Surgical management has traditionally included modified Ravitch techniques,<sup>16</sup> but the hybrid approach, combining minimally invasive repair of pectus excavatum (MIRPE) with a midline incision to correct sternal protrusions and abnormal costochondral insertions, is gaining traction.<sup>12,17,24,72</sup> Recent hybrid techniques emphasize using bars to correct lower depression and enable superior, less invasive, correction.<sup>12,17,24</sup> Main differences lie in surgical sequence. St-Louis et al<sup>17</sup> reported inserting dissectors first, to delineate incisions, whereas Zeineddine et al<sup>12</sup> began with the open portion to minimize resisting forces and optimize reapproximation. St-Louis et al<sup>17</sup> performed cartilage resections before sternal wedge osteotomies. Others start with sternal osteotomy and fixation before cartilage reapproximation to minimize excision.<sup>12</sup> Many surgeons recommend rigid sternal fixation to prevent malunion and minimize pain.<sup>12,24,64,78</sup> St-Louis et al<sup>17</sup> felt Nuss bars eliminated the need for sternal plating and avoided vertical incisions for better aesthetics.<sup>78,79</sup>

Unlike excavatum, arcuatum may not require multiple bars, even in adults.<sup>12,17</sup> Due to the shortened sternum, bars may need to be placed below the sternal end, especially if lower costal cartilages are compressing the heart. Combined fixation methods of these lower bars are necessary to minimize migration risks.<sup>12,19</sup>

Regardless of an open or minimally invasive approach, the critical step is the wedge sternal osteotomy.<sup>12,17,33</sup> The unique sternal angulation and rigidity render MIRPE alone insufficient and may exaggerate angulations.<sup>40,71</sup> Multilevel osteotomies are usually not necessary to complete repair; a single wedge often suffices, preserving sternal integrity.<sup>12,17,33,58,64</sup> Determining the optimal wedge angle and width for realignment can be challenging and patient specific. Starting with a minimal cut width and expanding as needed is recommended.<sup>12</sup> Others suggest autologous bone/cartilage grafts to fill residual gaps.<sup>24,43,51</sup> Recent computer-aided design modeling innovations offer patient-specific cutting templates and may reduce surgical time.<sup>32,65,80</sup>

Extensive subperiosteal cartilage resection is unnecessary. In adults, heavily calcified cartilages may preclude subperiosteal dissection, making minimal cartilage resection essential. Rigid rib-to-sternal fixation is recommended in some cases.<sup>12,19</sup>

### Limitations

This study has some limitations. Despite an extensive search, some articles on arcuatum may have been missed. Stratification was difficult, as many mixed-cohort studies did not differentiate arcuatum-specific findings, preventing data inclusion. Single-reviewer data extraction may introduce bias despite adherence to a standardized protocol to minimize this risk. Among included studies, 56.9% were small cohort studies, 27.7% case reports, 12.3% case series, and 3.1% technical reports, limiting generalizability. About 6.3% contained low-accuracy data but were included to ensure comprehensive coverage given the limited, heterogeneous literature.

### Conclusions

This systematic review reveals that pectus arcuatum is an under-recognized and often misdiagnosed pectus deformity that is an entity distinct from carinatum and excavatum. Unified terminology and standardized diagnostic criteria are vital to improve management strategies and document surgical outcomes. Minimally invasive hybrid approaches continue to refine outcomes, balancing invasiveness with efficacy.

#### ARTICLE INFORMATION

**Accepted for Publication:** April 29, 2025.

**Published Online:** July 23, 2025.

doi:10.1001/jamasurg.2025.2365

**Author Contributions:** Dr Jaroszewski had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** Zeineddine, Khedr, Farina, Odeh, Jenkins, Jaroszewski.

**Acquisition, analysis, or interpretation of data:** Zeineddine, Farina, Odeh, Senjab, Jenkins, Jaroszewski.

**Drafting of the manuscript:** Zeineddine, Khedr, Farina, Odeh, Senjab, Jaroszewski.

**Critical review of the manuscript for important intellectual content:** Zeineddine, Farina, Jenkins, Jaroszewski.

**Statistical analysis:** Zeineddine, Khedr, Farina, Odeh, Senjab.

**Administrative, technical, or material support:**

Zeineddine, Jaroszewski.

**Supervision:** Zeineddine, Farina, Jaroszewski.

**Conflict of Interest Disclosures:** Dr Jaroszewski reported receiving consulting and IP/royalties from Zimmer Biomet Inc through Mayo Clinic Ventures and speaker fees from Atricure Inc outside the submitted work. No other disclosures were reported.

#### REFERENCES

1. Fokin AA. Pouter pigeon breast. *Chest Surg Clin N Am*. 2000;10(2):377-391. doi:10.1016/S1052-3359(25)00502-2
2. Lester CW. Pigeon breast (pectus carinatum) and other protrusion deformities of the chest of developmental origin. *Ann Surg*. 1953;137(4):482-489. doi:10.1097/00000658-195304000-00008
3. Chin EF. Surgery of funnel chest and congenital sternal prominence. *Br J Surg*. 1957;44(186):360-376. doi:10.1002/bjs.18004418607
4. Robicsek F, Daugherty HK, Mullen DC, et al. Technical considerations in the surgical management of pectus excavatum and carinatum. *Ann Thorac Surg*. 1974;18(6):549-564. doi:10.1016/S0003-4975(10)64401-3
5. Robicsek F, Cook JW, Daugherty HK, Selle JG. Pectus carinatum. *Coll Works Cardipulm Dis*. 1979;22:65-78.
6. Ravitch MM. The operative correction of pectus carinatum. *Bull Soc Int Chir*. 1975;34(2):117-120.
7. Haje SA, Harcke HT, Bowen JR. Growth disturbance of the sternum and pectus deformities: imaging studies and clinical correlation. *Pediatr Radiol*. 1999;29(5):334-341. doi:10.1007/s002470050602
8. Kelly RE Jr, Quinn A, Varela P, Redlinger RE Jr, Nuss D. Dysmorphology of chest wall deformities: frequency distribution of subtypes of typical pectus



- excavatum and rare subtypes. *Arch Bronconeumol*. 2013;49(5):196-200. doi:10.1016/j.arbres.2012.09.013
9. Edward KR, John OR, Jay GM, et al. Ten-year experience with staged management of pectus carinatum: Results and lessons learned. *J Pediatr Surg*. 2021;56(10):1835-1840. doi:10.1016/j.jpedsurg.2021.01.027
  10. Haje SA, de Podestá Haje D. Orthopedic approach to pectus deformities: 32 years of studies. *Rev Bras Ortop*. 2015;44(3):191-198.
  11. Lacquet LK, Morshuis WJ, Folgering HT. Long-term results after correction of anterior chest wall deformities. *J Cardiovasc Surg (Torino)*. 1998;39(5):683-688.
  12. Zeineddine RM, Farina JM, Shawwaf KA, et al. Surgical treatment and outcomes of pectus arcuatum. *JTCVS Tech*. 2024;28:194-202. doi:10.1016/j.jtc.2024.09.015
  13. Pitt JB, Zeineddin S, Carter M, et al. Demographics of anterior chest wall deformity patients: a tertiary children's hospital experience. *J Surg Res*. 2024;293:451-457. doi:10.1016/j.jss.2023.09.040
  14. Lester CW. The etiology and pathogenesis of funnel chest, pigeon breast, and related deformities of the anterior chest wall. *J Thorac Surg*. 1957;34(1):1-10. doi:10.1016/S0096-5588(20)30366-4
  15. Saxena AK, Willital GH. Surgical repair of pectus carinatum. *Int Surg*. 1999;84(4):326-330.
  16. Gritsiuta AI, Bracken A, Beebe K, Pechetov AA. Currarino-Silverman syndrome: diagnosis and treatment of rare chest wall deformity, a case series. *J Thorac Dis*. 2021;13(5):2968-2978. doi:10.21037/jtd-20-3472
  17. St-Louis E, Labege JM, Wissanji H, Emil S. A minimally invasive hybrid procedure to correct pectus arcuatum. *J Plast Reconstr Aesthet Surg*. 2024;93:127-132. doi:10.1016/j.bjps.2024.04.043
  18. Abdellaoui S, Scalabre A, Piolat C, et al. Pectus arcuatum: a pectus unlike any other. *J Pediatr Surg*. 2023;58(9):1679-1685. doi:10.1016/j.jpedsurg.2023.03.013
  19. Fonkalsrud EW. Surgical correction of pectus carinatum: lessons learned from 260 patients. *J Pediatr Surg*. 2008;43(7):1235-1243. doi:10.1016/j.jpedsurg.2008.02.007
  20. Fonkalsrud EW, Beanes S. Surgical management of pectus carinatum: 30 years' experience. *World J Surg*. 2001;25(7):898-903. doi:10.1007/s00268-001-0048-x
  21. Hysi I, Vincentelli A, Juthier F, et al. Cardiac surgery and repair of pectus deformities: when and how? *Int J Cardiol*. 2015;194:83-86. doi:10.1016/j.ijcard.2015.05.100
  22. Haje DP, Teixeira KO, Silva M, Volpon JB, Mendlovitz PS, Dolabela P. Analysis of sternal curvature patterns in patients with pectus and control. *Acta Ortop Bras*. 2021;29(5):258-262. doi:10.1590/1413-785220212905243733
  23. Haje DP, Teixeira KO, Silva Neto M, Volpon JB, Mendlovitz PS, Vasconcelos PD. Sternal tomographic evaluation in pectus patients: analysis of sagittal parameters. *Acta Ortop Bras*. 2022;30(5):e250612. doi:10.1590/1413-785220223005e250612
  24. González-Barba I, Laín A, García L, Giné C, López M. Pectus arcuatum: description of a hybrid surgical technique. *JPS Open*. 2023;3:100030. doi:10.1016/j.jypso.2023.100030
  25. Wurtz A, Rousse N, Benhamed L, et al. Simplified open repair for anterior chest wall deformities: analysis of results in 205 patients. *Orthop Traumatol Surg Res*. 2012;98(3):319-326. doi:10.1016/j.otsr.2011.11.005
  26. Haje SA, Haje DdeP, Silva Neto M, et al. Pectus deformities: tomographic analysis and clinical correlation. *Skeletal Radiol*. 2010;39(8):773-782. doi:10.1007/s00256-010-0874-8
  27. Cohee AS, Lin JR, Frantz FW, Kelly RE Jr, AS C. Staged management of pectus carinatum. *J Pediatr Surg*. 2013;48(2):315-320. doi:10.1016/j.jpedsurg.2012.11.008
  28. van Braak H, de Beer SA, Zwaveling S, Oomen MWN, de Jong JR. Ravitch surgery or dynamic compression bracing for pectus carinatum: a retrospective cohort study. *Ann Thorac Surg*. 2024;117(1):144-150. doi:10.1016/j.athoracsur.2022.10.041
  29. Shang Z, Hong C, Duan X, Li X, Si Y. Orthotic bracing or minimally invasive surgery? a summary of 767 pectus carinatum cases for 9 years. *Biomed Res Int*. 2021;2021:6942329. doi:10.1155/2021/6942329
  30. Howard R. Pigeon chest (protrusion deformity of the sternum). *Med J Aust*. 1958;45(20):664-666.
  31. Önen A. A new surgical approach for mixed-type pectus carinatum: the Onen procedure. *Türk Gogus Kalp Dama*. 2014;22:878-883. doi:10.5606/tgkdc.dergisi.2014.7940
  32. Leng S, Bici K, Facchini F, et al. Customized cutting template to assist sternotomy in pectus arcuatum. *Ann Thorac Surg*. 2019;107(4):1253-1258. doi:10.1016/j.athoracsur.2018.10.057
  33. Kara M, Gundogdu AG, Kadioglu SZ, Cayirci EC, Taskin N, M K. The use of sternal wedge osteotomy in pectus surgery: when is it necessary? *Asian Cardiovasc Thorac Ann*. 2016;24(7):658-662. doi:10.1177/0218492316657243
  34. Robicsek F, Sanger PW, Taylore FH, Thomas MJ. The surgical treatment of chondrosternal prominence (pectus carinatum). *J Thorac Cardiovasc Surg*. 1963;45:691-701.
  35. Nowacki W, Kruczyński J, Futyma J, Stencel P. Results of operative treatment of the pectus carinatum. *Chir Narzadow Ruchu Ortop Pol*. 2008;73(3):203-206, 192-195.
  36. Del Frari B, Schwabegger AH. Diced autologous rib cartilage for primary treatment or refinement of minor chest wall deformities. *Plast Reconstr Surg*. 2011;128(1):154-162. doi:10.1097/PRS.0b013e318217425f
  37. Servi M, Buonamici F, Carfagni M, et al. CAD-based automatic modeling of customized cutting templates for pectus arcuatum surgical correction. *Annu Int Conf IEEE Eng Med Biol Soc*. 2020;2020:6044-6048. doi:10.1109/EMBC44109.2020.9175824
  38. Ravitch MM. Unusual sternal deformity with cardiac symptoms operative correction. *J Thorac Surg*. 1952;23(2):138-144. doi:10.1016/S0096-5588(20)31203-4
  39. Wang W, Long W, Liu Y, Cai B, Luo J. Morphological characteristics of Wenlin chest. *Int J Surg Case Rep*. 2022;4:22-24.
  40. Wang W, Long W, Liu Y, Cai B, Luo J. Reoperation after failure of Nuss procedure on Wenlin chest with Poland syndrome. *Int J Orthop Sci*. 2022;8(3):396-399. doi:10.22271/ortho.2022.v8.i3f.3224
  41. Wang W, Long W, Liu Y, Cai B, Luo J. Surgical treatment of Wenlin chest. *Int J Surg Case Rep*. 2022;4:62-64. doi:10.22271/27081494.2022.v4.i2b.52
  42. Wang W, Long W, Liu Y, Cai B, Luo J. Surgical treatment of Wenlin chest after spinal orthopedics. *Int J Surg Case Rep*. 2022;4:65-67. doi:10.22271/27081494.2022.v4.i2b.53
  43. Kim SY, Park S, Kim ER, Park IK, Kim YT, Kang CH. A case of successful surgical repair for pectus arcuatum using chondrosteronoplasty. *Korean J Thorac Cardiovasc Surg*. 2016;49(3):214-217. doi:10.5090/kjtc.2016.49.3.214
  44. Brünner S. Premature synostosis in the sternum (Silverman disease). *Acta Paediatr (Stockh)*. 1961;50(3):288-290. doi:10.1111/j.1651-2227.1961.tb08174.x
  45. Kaplan MS, Opitz JM, Gosset FR. Noonan syndrome—a case with elevated serum alkaline phosphatase levels and malignant schwannoma of the left forearm. *AJDC*. 1968;116(4):359-366. doi:10.1001/archpedi.1968.02100020363003
  46. Mehta AV. Currarino-Silverman syndrome and congenital heart disease. *Tex Heart Inst J*. 1996;23(1):69.
  47. Alkhunaizi E, Walkiewicz MA, Chitayat D. Mutation in the ADNP gene associated with Noonan syndrome features. *Clin Dysmorphol*. 2018;27(2):53-57. doi:10.1097/MCD.0000000000000215
  48. Sahu S, Chaudhury S, Saldanha D. Noonan syndrome with somnambulism: a rare case report. *Ind Psychiatry J*. 2020;29(2):339-341. doi:10.4103/ipj.ipj.84\_19
  49. Brichon PY, Wihlm JM. Correction of a severe pouter pigeon breast by triple sternal osteotomy with a novel titanium rib bridge fixation. *Ann Thorac Surg*. 2010;90(6):e97-e99. doi:10.1016/j.athoracsur.2010.08.068
  50. Akkas Y, Kocer B, Peri NG. A new technique for mixed-type pectus carinatum: modified Onen method. *Asian Cardiovasc Thorac Ann*. 2016;24(2):227-230. doi:10.1177/0218492315627943
  51. Soria-Gondek A, Oviedo-Gutiérrez M, Martín-Lluís A, López de Castro P, de Diego M. Modified Ravitch procedure and autologous cartilage graft for pectus arcuatum. *Ann Thorac Surg*. 2022;114(2):e105-e107. doi:10.1016/j.athoracsur.2021.10.053
  52. Mannatrizio D, Domenico M, Fascia G, Giacomo F, Guglielmi G, Giuseppe G. superior pectus carinatum (currarino-silverman syndrome) in a 66-year-old woman: a case report. *Digit Diagn*. 2022;3(2):141-148.
  53. Wang W, Long W, Liu Y, Cai B, Luo J. Minimally invasive surgery of Wenlin chest. *Int J Surg Case Rep*. 2022;4:59-61.
  54. Wang W, Long W, Liu Y, Cai B, Luo J. Reoperation after failure of Nuss procedure on severe Wenlin chest. *Int J Surg Sci*. 2022;6:14-17.
  55. Wang W. Minimally invasive surgical treatment for severe Wenlin chest. Accessed June 26, 2025.

<https://www.casereportsofsurgery.com/article/62/4-2-21-220.pdf>

56. Currarino G, Silverman FN. Premature obliteration of the sternal sutures and pigeon-breast deformity. *Radiology*. 1958;70(4):532-540. doi:10.1148/70.4.532
57. Cartoski MJ, Nuss D, Goretsky MJ, et al. Classification of the dysmorphology of pectus excavatum. *J Pediatr Surg*. 2006;41(9):1573-1581. doi:10.1016/j.jpedsurg.2006.05.055
58. Shamberger RC, Welch KJ. Surgical correction of chondromanubrial deformity (Currarino Silverman syndrome). *J Pediatr Surg*. 1988;23(4):319-322. doi:10.1016/S0022-3468(88)80197-0
59. Brodtkin HA. Pigeon breast; congenital chondrosternal prominence; etiology and surgical treatment by xiphosternopexy. *AMA Arch Surg*. 1958;77(2):261-270. doi:10.1001/archsurg.1958.01290020111022
60. Chidambaram B, Mehta AV. Currarino-Silverman syndrome (pectus carinatum type 2 deformity) and mitral valve disease. *Chest*. 1992;102(3):780-782. doi:10.1378/chest.102.3.780
61. Lam CR, Taber RE. Surgical treatment of pectus carinatum. *Arch Surg*. 1971;103(2):191-194. doi:10.1001/archsurg.1971.01350080107016
62. Welch KJ, Vos A. Surgical correction of pectus carinatum (pigeon breast). *J Pediatr Surg*. 1973;8(5):659-667. doi:10.1016/0022-3468(73)90404-1
63. Allwyn Joshua S, Shetty L, Pare VS, Sebastian R. Chondromanubrial deformity and bifid rib, rare variations seen in pectus carinatum: a radiological finding. *Surg Radiol Anat*. 2013;35(5):443-447. doi:10.1007/s00276-012-1046-4
64. Kuzmichev V, Ershova K, Adamyan R. Surgical correction of pectus arcuatum. *J Vis Surg*. 2016;2:55. doi:10.21037/jovs.2016.02.28
65. Carfagni M, Facchini F, Furferi R, et al. Towards a CAD-based automatic procedure for patient specific cutting guides to assist sternal osteotomies in pectus arcuatum surgical correction. *J Comput Des Eng*. 2019;6(1):118-127. doi:10.1016/j.jcde.2018.01.001
66. Brodtkin HA. Congenital chondrosternal prominence (pigeon breast) a new interpretation. *Pediatrics*. 1949;3(3):286-295.
67. Acastello E. *Patologías de la Pared Torácica en Pediatría*. Editorial El Ateneo; 2006.
68. Gabrielsen TO, Ladyman GH. Early closure of the sternal sutures and congenital heart disease. *Am J Roentgenol Radium Ther Nucl Med*. 1963;89:975-983.
69. Farina JM, Olson RJ, Dhamija R, et al. The complexity of familial inheritance in pectus excavatum: a 10-family exome sequencing analysis. *Genes (Basel)*. 2024;15(11):1429. doi:10.3390/genes15111429
70. Stephens EH, Dearani JA, Jaroszewski DE. Pectus excavatum in cardiac surgery patients. *Ann Thorac Surg*. 2023;115(5):1312-1321. doi:10.1016/j.athoracsur.2023.01.040
71. Fokin AA, Steuerwald NM, Ahrens WA, Allen KE. Anatomical, histologic, and genetic characteristics of congenital chest wall deformities. *Semin Thorac Cardiovasc Surg*. 2009;21(1):44-57. doi:10.1053/j.semtcvs.2009.03.001
72. Emil S. Current options for the treatment of pectus carinatum: when to brace and when to operate? *Eur J Pediatr Surg*. 2018;28(4):347-354. doi:10.1055/s-0038-1667297
73. Torre M, Rapuzzi G, Jasonni V, Varela PJ. Chest wall deformities: an overview on classification and surgical options. In: Cardoso PFG. *Topics in Thoracic Surgery*. IntechOpen; 2012:117-136.
74. Pickard LR, Tepas JJ, Shermeta DW, Haller JA Jr; LR P. Pectus carinatum: results of surgical therapy. *J Pediatr Surg*. 1979;14(3):228-230. doi:10.1016/S0022-3468(79)80476-5
75. Andren L, Hall P. Diminished segmentation or premature ossification of the sternum in congenital heart disease. *Br Heart J*. 1961;23(2):140-142. doi:10.1136/hrt.23.2.140
76. Gritsiuta A, Elmore LR, Petrov RV. Surgical correction of Currarino-Silverman syndrome. *J Pulm Med*. 2021;5(4):1000117.
77. Behr CA, Denning NL, Kallis MP, et al. The incidence of Marfan syndrome and cardiac anomalies in patients presenting with pectus deformities. *J Pediatr Surg*. 2019;54(9):1926-1928. doi:10.1016/j.jpedsurg.2018.11.017
78. Tam DY, Nedadur R, Yu M, Yanagawa B, Fremes SE, Friedrich JO. Rigid plate fixation vs wire cerclage for sternotomy after cardiac surgery: a meta-analysis. *Ann Thorac Surg*. 2018;106(1):298-304. doi:10.1016/j.athoracsur.2018.02.043
79. Lazar HL, Salm TV, Engelman R, Orgill D, Gordon S. Prevention and management of sternal wound infections. *J Thorac Cardiovasc Surg*. 2016;152(4):962-972. doi:10.1016/j.jtcvs.2016.01.060
80. Servi M, Buonamici F, Carfagni M, et al. CAD-based automatic modeling of customized cutting templates for pectus arcuatum surgical correction. *Annu Int Conf IEEE Eng Med Biol Soc*. 2020;2020:6044-6048. doi:10.1109/EMBC44109.2020.9175824