

# Risk of Poststroke Epilepsy Among Young Adults With Ischemic Stroke or Intracerebral Hemorrhage

Esmée Verburgt, MSc; Lina Fellah, MSc; Merel S. Ekker, MD, PhD; Mijntje M. I. Schellekens, MD; Esther M. Boot, MD; Maikel H. M. Immens, MD; Mayte E. van Alebeek, MD, PhD; Paul J. A. M. Brouwers, MD, PhD; Renate M. Arntz, MD, PhD; Gert W. van Dijk, MD, PhD; Rob A. R. Gons, MD, PhD; Inge W. M. van Uden, MD, PhD; Tom den Heijer, MD, PhD; Julia H. van Tuijl, MD, PhD; Karlijn F. de Laat, MD, PhD; Anouk G.W. van Norden, MD, PhD; Sarah E. Vermeer, MD, PhD; Marian S. G. van Zagten, MD, PhD; Robert J. van Oostenbrugge, MD, PhD; Marieke J. H. Wermer, MD, PhD; Paul J. Nederkoorn, MD, PhD; Henk Kerkhoff, MD, PhD; Fergus A. Rooyer, MD; Frank G. van Rooij, MD, PhD; Ido R. van den Wijngaard, MD, PhD; Anil M. Tuladhar, MD, PhD; Jamie I. Verhoeven, MD; Nina A. Hilken, MD, PhD; Frank-Erik de Leeuw, MD, PhD

 Supplemental content

**IMPORTANCE** Poststroke epilepsy (PSE) is a major complication among young adults and is associated with problems with functional recovery and daily life. Although scores have been developed to predict risk of PSE, they have not been validated among patients with stroke at a young age.

**OBJECTIVES** To investigate both the risk of and risk factors for PSE at a young age and validate current PSE risk scores among a cohort of young adults.

**DESIGN, SETTING, AND PARTICIPANTS** This cohort study used data from ODYSSEY (Observational Dutch Young Symptomatic Stroke Study), a prospective cohort study conducted among 17 hospitals in the Netherlands between May 27, 2013, and March 3, 2021, with follow-up until February 28, 2024. Participants included 1388 consecutive patients aged 18 to 49 years with neuroimaging-proven ischemic stroke or intracerebral hemorrhage (ICH) and without a history of epilepsy. Statistical analysis took place between June and August 2024.

**EXPOSURE** First-ever neuroimaging-proven ischemic stroke or ICH.

**MAIN OUTCOMES AND MEASURES** Poststroke epilepsy was defined as at least 1 remote symptomatic seizure (>7 days). Cumulative incidence functions were used to calculate the 5-year risk of PSE. Fine-Gray regression models were used to identify risk factors associated with PSE (age, sex, clinical stroke, and neuroimaging variables). The performances of the SeLECT (severity of stroke, large-artery atherosclerosis, early seizure, cortical involvement, and territory of middle cerebral artery) 2.0 risk score (for ischemic stroke) and the CAVE (cortical involvement, age, bleeding volume, and early seizure) risk score (for ICH) were assessed with C statistics and calibration bar plots.

**RESULTS** This study included 1388 patients (ischemic stroke, 1231 [88.7%]; ICH, 157 [11.3%]; median age, 44.1 years [IQR, 38.0-47.4 years]; 736 men [53.0%]; median follow-up, 5.3 years [IQR, 3.4-7.4 years]), of whom 57 (4.1%) developed PSE. The 5-year cumulative risk of PSE was 3.7% (95% CI, 0.2%-4.8%) after ischemic stroke and 7.6% (95% CI, 3.5%-11.8%) after ICH. Factors associated with PSE after ischemic stroke were an acute symptomatic seizure (<7 days) (hazard ratio [HR], 10.83 [95% CI, 2.05-57.07];  $P = .005$ ) and cortical involvement (HR, 5.35 [95% CI, 1.85-15.49];  $P = .002$ ). The only factor associated with PSE after ICH was cortical involvement (HR, 8.20 [95% CI, 2.22-30.25];  $P = .002$ ). The C statistic was 0.78 (95% CI, 0.71-0.84) for the SeLECT 2.0 risk score and 0.83 (95% CI, 0.76-0.90) for the CAVE risk score, and calibration was good for both scores.

**CONCLUSION** This study suggests that the risk of PSE among young adults is relatively low and that the factors that were associated with PSE were similar to variables included in the existing risk scores, which can therefore also be applied for young adults after stroke. Future clinical trials should investigate the optimal primary and secondary prophylaxis for patients at high risk.

JAMA Neurol. doi:10.1001/jamaneurol.2025.0465  
Published online April 14, 2025.

**Author Affiliations:** Author affiliations are listed at the end of this article.

**Corresponding Author:** Frank-Erik de Leeuw, MD, PhD, Department of Neurology, Radboud Institute of Medical Innovation, Donders Institute for Brain, Cognition and Behavior, PO Box 9101, 6500 HB Nijmegen, the Netherlands (frank.erik.deleeuw@radboudumc.nl).

Stroke at a young age occurs during a demanding phase of life, which may result in disability, and thus poses a risk of major complications. One of these complications is poststroke epilepsy (PSE). Patients with PSE have a higher risk of mortality, worse functional outcomes, and lower quality of life.<sup>1-6</sup> They often need long-term treatment with antiepileptic drugs, which may cause adverse events such as fatigue, dizziness, behavioral changes, and depression.<sup>7-9</sup>

Only a few studies have investigated the incidence and cumulative risk of PSE specifically among young adults.<sup>10-13</sup> These studies reported a 5-year cumulative risk between 5.5% and 15% for patients with ischemic stroke and 21% for patients with intracerebral hemorrhage (ICH). However, these estimates are based on studies in which neuroimaging-based proof of ischemic stroke was not required, and these studies were conducted decades ago. Recently, several risk scores that predict the individual risk of PSE have been developed.<sup>14</sup> However, they have all been derived from the general stroke population, including mainly patients older than 65 years, with untested external validity in young adults. To accurately inform patients about the long-term risks of PSE, it is important to investigate the risk of PSE among young adults specifically and to identify which patients are at highest risk.

This study investigated the incidence of PSE among young adults aged 18 to 49 years in a large prospective cohort of patients with neuroimaging-proven strokes. In addition, we aimed to identify clinical and neuroimaging characteristics that are associated with PSE. Last, we assessed the performance of existing risk scores of PSE in a cohort of young adults.

## Methods

### Patients and Study Design

In this cohort study, we included patients from the Observational Dutch Young Symptomatic Stroke Study (ODYSEY), a multicenter observational study with 17 participating hospitals in the Netherlands, which has been described in more detail previously.<sup>15</sup> It includes patients aged 18 to 49 years with a first-ever neuroimaging-proven ischemic stroke or ICH between May 27, 2013, and March 3, 2021. Patients with transient symptoms lasting less than 24 hours were included as having (minor) strokes according to the tissue-based definition.<sup>16</sup> Hemorrhagic transformation of an ischemic stroke was classified as an ischemic stroke. Exclusion criteria were transient monocular blindness or retinal infarction, a history of transient ischemic attack or stroke, traumatic ICH, subarachnoid hemorrhage, ICH due to a known ruptured aneurysm, ICH in a known cerebral malignant neoplasm, cerebral venous sinus thrombosis, and not being a permanent resident of the Netherlands. For the present study, we also excluded patients with a history of epilepsy ( $n = 30$ ). The Medical Review Ethics Committee Region East-Netherlands approved this cohort study, and all participants provided written informed consent. The study followed the Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline for cohort studies.

### Key Points

**Question** What is the risk of poststroke epilepsy (PSE) among young adults (<50 years) with ischemic stroke or intracerebral hemorrhage?

**Findings** In this cohort study including 1388 patients, the 5-year cumulative risk of PSE was 3.7% after ischemic stroke and 7.6% after intracerebral hemorrhage. Factors associated with PSE were similar to variables in current risk scores for PSE.

**Meaning** This study suggests that the risk of PSE among young adults is relatively low and that current risk scores for PSE can be applied to identify young patients at high risk.

### Outcome

The outcome was assessed using a 2-stage approach. First, information about acute symptomatic seizures was retrieved from medical files during baseline data collection by physicians. Information about the occurrence of remote symptomatic seizures was collected through contacting patients by telephone or online survey until February 28, 2024, using a standardized structured questionnaire in Dutch (translated for this article). Second, all first-reported seizures for which the patient sought medical attention, as well as the reported use of antiseizure medication, were cross-checked with medical records from the participating hospital or general practitioner. We additionally validated the self-reported method by cross-checking the medical records of 64 patients from 6 hospitals who did not report any PSE and found 100% agreement. The questionnaires are shown in the eAppendix in Supplement 1.

An acute symptomatic seizure was defined as a seizure within 7 days after the initial stroke, and a remote symptomatic seizure was defined as a seizure within 7 days or later after the initial stroke. We defined PSE as 1 or more remote acute symptomatic seizures, according to the guidelines of the International League Against Epilepsy (ILAE).<sup>17</sup> Seizure type was defined as focal (with or without impaired awareness) or generalized.

### Risk Factors

The following cardiovascular risk factors were systematically assessed through medical files at the time of the index event: hypertension, dyslipidemia, diabetes, smoking, excessive alcohol use, illicit drug use, and obesity, and were described previously.<sup>18</sup> Cause of ischemic stroke was defined according to the modified Trial of Org 10172 in Acute Stroke Treatment (TOAST) classification.<sup>19</sup> Cause of ICH was defined in 4 categories: macrovascular causes, deep perforating vasculopathy, other cause determined, and undetermined. Severity of stroke was defined by the National Institutes of Health Stroke Scale (NIHSS) score at admission.<sup>20</sup> We stratified severity of stroke as minor stroke (NIHSS score  $\leq 3$ ), and major stroke (NIHSS score  $> 3$ ).<sup>21</sup>

### External Validation Risk Scores

We calculated the SeLECT (severity of stroke, large-artery atherosclerosis, early seizure, cortical involvement, and ter-

territory of middle cerebral artery) 2.0 risk score for each patient with ischemic stroke: severity of stroke (1 point for an NIHSS score of 4-10 and 2 points for an NIHSS score of  $\geq 11$ ), large-artery atherosclerosis (1 point), acute symptomatic seizure (3 points), acute symptomatic status epilepticus (7 points), cortical involvement (2 points), and territory of the middle cerebral artery (1 point).<sup>22</sup> For patients with ICH, we calculated the CAVE (cortical involvement, age, bleeding volume, and early seizure) risk score: cortical involvement (1 point), younger than 65 years of age (1 point), bleeding volume of 10 mL or more (1 point), and an early seizure (1 point).<sup>23</sup> Cortical involvement of the stroke was assessed on computed tomography or magnetic resonance imaging scans or based on clinical symptoms (eg, neglect, aphasia, or hemianopsia). Bleeding volume was determined by the ABC/2 formula.<sup>24</sup>

We assessed the performance of the SeLECT 2.0 and CAVE risk scores through discrimination and calibration. Discrimination determines how well a model can distinguish between patients with and patients without PSE and was assessed with the C statistic. Calibration shows the agreement between the predicted risk and the observed risk, which is illustrated with bar plots. The observed risk was calculated only for the risk scores with at least 30 patients.

### Statistical Analysis

Statistical analysis took place between June and August 2024. Differences in baseline characteristics between patients with and patients without PSE were examined with a Mann-Whitney test for nonnormally distributed continuous variables or a  $\chi^2$  test for categorical variables. The number of patient-years was calculated until the date of first remote symp-

tomatic seizure for patients with PSE, death, or last available follow-up. Cumulative incidence functions were used to investigate the 5-year cumulative incidence of PSE stratified per type of stroke (ischemic stroke or ICH), to account for the competing risk of death. Second, we calculated the incidence rate per 100 person-years per type of stroke, per year, by dividing the number of PSE cases by the number of person-years at risk.

Age- and sex-adjusted Fine-Gray proportional hazard models were used to investigate the association between predefined risk factors (age, sex, NIHSS score at admission, occurrence of an acute symptomatic seizure, cortical involvement, infarction in the middle cerebral artery territory, and modified TOAST classification) and PSE among patients with ischemic stroke. Second, we investigated the association between predefined risk factors (age, sex, NIHSS score at admission, occurrence of an acute symptomatic seizure, cortical involvement, and bleeding volume of  $\geq 10$  mL) and PSE among patients with ICH. Multivariable analyses were done including variables that were significant in the age- and sex-adjusted models. All *P* values were from 2-sided tests and results were deemed statistically significant at *P* < .05. Analyses were performed with RStudio, version 4.1.3 (2022-03-10; The R Project for Statistical Computing).

### Results

This study included 1388 patients (median age, 44.1 years [IQR, 38.0-47.4 years]; 736 men [53.0%] and 652 women [47.0%]), of whom 1231 (88.7%) had an ischemic stroke and 157 (11.3%) had an ICH (Table 1 and Figure 1). The median follow-up was 5.3 years (IQR, 3.4-7.4 years), with a total of

Table 1. Baseline Characteristics

Characteristic	All patients, No./total No. (%)	Patients without PSE, No./total No. (%)	Patients with PSE, No./total No. (%)	<i>P</i> value
Type of stroke				
Ischemic stroke	1231/1388 (88.7)	1187/1331 (89.2)	44/57 (77.2)	.01
Intracerebral hemorrhage	157/1388 (11.3)	144/1331 (10.8)	13/57 (22.8)	
Age, median (IQR), y	44.1 (38.0-47.4)	44.1 (38.2-47.4)	44.3 (35.3-47.4)	.68
Men	736/1388 (53.0)	707/1331 (53.1)	29/57 (50.9)	.84
Women	652/1388 (47.0)	624/1331 (46.9)	28/57 (49.1)	
Follow-up, median (IQR), y	5.3 (3.4-7.4)	5.3 (3.4-7.4)	5.4 (3.9-7.8)	.41
NIHSS score at admission, median (IQR) <sup>a</sup>	3 (1-7)	3 (1-7)	6 (2-14)	.002
Minor stroke ( $\leq 3$ )	776/1384 (56.1)	753/1328 (56.7)	23/56 (41.1)	.03
Major stroke ( $> 3$ )	608/1384 (43.9)	575/1328 (43.3)	33/56 (58.9)	
Diabetes	130/1388 (9.4)	125/1331 (9.4)	5/57 (8.8)	< .99
Hypertension	522/1388 (37.6)	505/1331 (37.9)	17/57 (29.8)	.27
Smoking	630/1388 (45.4)	608/1331 (45.7)	2/57 (3.6)	.36
Excess alcohol use	86/1388 (6.2)	84/1331 (6.3)	2/57 (3.5)	.56
Dyslipidemia	837/1388 (60.3)	807/1331 (60.6)	30/57 (52.6)	.28
Illicit drug use <sup>b</sup>	90/1079 (6.5)	89/1031 (6.7)	1/48 (1.8)	.12
Acute symptomatic seizure (<7 d)	19/1388 (1.4)	13/1331 (1.0)	6/57 (10.5)	< .001
Acute status epilepticus	1/1388 (0.1)	0	1/57 (1.8)	.68
Cortical involvement <sup>c</sup>	659/1364 (47.5)	610/1308 (45.8)	49/56 (86.0)	< .001

(continued)

Table 1. Baseline Characteristics (continued)

Characteristic	All patients, No./total No. (%)	Patients without PSE, No./total No. (%)	Patients with PSE, No./total No. (%)	P value
Ischemic stroke				
TOAST classification				
Atherothrombotic	55/1231 (4.5)	52/1187 (4.4)	3/44 (6.8)	.08
Likely atherothrombotic	174/1231 (14.1)	167/1187 (14.1)	7/44 (15.9)	
Small vessel disease	161/1231 (13.1)	161/1187 (13.6)	0	
Cardioembolic	202/1231 (16.4)	194/1187 (16.3)	8/44 (18.2)	
Rare causes	280/1231 (22.7)	264/1187 (22.2)	16/44 (36.4)	
Cryptogenic	304/1231 (24.7)	295/1187 (24.9)	9/44 (20.5)	
Multiple causes	55/1231 (4.5)	54/1187 (4.5)	1/44 (2.3)	
SeLECT 2.0 score, median (IQR) <sup>d</sup>	2 (1-3)	2 (1-3)	4/44 (3-5)	<.001
0	215/1227 (17.5)	213/1184 (18.0)	2/43 (4.7)	
1	235/1227 (19.2)	234/1184 (19.8)	1/43 (2.3)	
2	211/1227 (17.2)	209/1184 (17.7)	2/43 (4.7)	
3	291/1227 (23.7)	277/1184 (23.4)	14/43 (32.6)	
4	147/1227 (12.0)	142/1184 (12.0)	5/43 (11.6)	
5	110/1227 (9.0)	94/1184 (7.9)	16/43 (37.2)	
6	14/1227 (1.1)	13/1184 (1.1)	1/43 (2.3)	
7	1/1227 (0.1)	0	1/43 (2.3)	
8	2/1227 (0.2)	2/1184 (0.2)	0	
9	0	0	0	
10	0	0	0	
11	0	0	0	
12	0	0	0	
13	1/1227 (0.1)	0	1/43 (2.3)	
Intracerebral hemorrhage				
Cause of stroke				
Macrovascular cause	37/157 (23.6)	31/144 (21.5)	6/13 (46.3)	.19
Deep perforating vasculopathy	57/157 (36.3)	55/144 (38.2)	2/13 (15.4)	
Other determined cause	12/157 (7.6)	11/144 (7.6)	1/13 (7.7)	
Undetermined	51/157 (32.5)	47/144 (32.6)	4/13 (30.8)	
CAVE score, median (IQR) <sup>e</sup>	2 (1-2)	1 (1-2)	3 (2-3)	<.001
1	70/148 (47.3)	70/135 (51.9)	0	
2	46/148 (31.1)	40/135 (29.6)	6/13 (46.2)	
3	30/148 (20.3)	24/135 (17.8)	6/13 (46.2)	
4	2/148 (1.4)	1/135 (0.7)	1/13 (7.7)	

Abbreviations: CAVE, cortical involvement, age, bleeding volume, and early seizure; NIHSS, National Institutes of Health Stroke Scale; PSE, poststroke epilepsy; SeLECT, severity of stroke, large-artery atherosclerosis, early seizure, cortical involvement, and territory of middle cerebral artery; TOAST, Trial of Org 10172 in Acute Stroke Treatment.

<sup>a</sup> NIHSS score was missing for 4 patients.

<sup>b</sup> Illicit drug use was missing for 309 patients.

<sup>c</sup> Cortical involvement was missing for 24 patients.

<sup>d</sup> SeLECT 2.0 score was missing for 4 patients.

<sup>e</sup> CAVE score was missing for 9 patients.


7002 patient-years of follow-up. Overall, 57 patients (4.1%) developed PSE (median age, 44.3 [IQR, 35.3-47.4 years]; 29 men [50.9%] and 28 women [49.1%]). The median time to first late seizure was 7 months (IQR, 2-14 months). Table 1 shows the baseline characteristics of the included patients. The median NIHSS score at admission was 3 (IQR, 1-7) for patients without PSE and 6 (IQR, 2-14) for patient with PSE. Of 13 patients with acute symptomatic seizures who did not develop PSE, only 2 (15.4%; 1 with ischemic stroke and 1 with ICH) did not receive any antiseizure medication after the

first seizure. The eTable in [Supplement 1](#) shows the type of seizures in patients with PSE.

### Ischemic Stroke and the Risk of PSE

In total, 44 of 1231 patients with an ischemic stroke (3.6%) developed PSE. The 5-year cumulative risk of PSE after an ischemic stroke was 3.7% (95% CI, 0.2%-4.8%) (**Figure 2**). The incidence rate per 100 person-years was 2.2 (95% CI, 1.8-2.6) in the first year after the stroke, which decreased to 0.8 (95% CI, 0.5-1.1) in the second year. Variables that were associated with

Figure 1. Flowchart of Included Patients



ODYSSEY indicates Observational Dutch Young Symptomatic Stroke Study.

PSE in the multivariable analysis were an acute symptomatic seizure (hazard ratio [HR], 10.83 [95% CI, 2.05-57.07];  $P = .005$ ) and cortical involvement (HR, 5.35 [95% CI, 1.85-15.49];  $P = .002$ ) (Table 2). None of the patients with PSE had ischemic stroke caused by small vessel disease.

### ICH and the Risk of PSE

A total of 13 of 157 patients with an ICH (8.3%) developed PSE during follow-up. The 5-year cumulative risk of PSE after ICH was 7.6% (95% CI, 3.5%-11.8%) (Figure 2). The incidence rate per 100 person-years was 10.8 (95% CI, 7.6-13.9) in the first year and 1.1 (95% CI, 0.0-2.1) in the second year. The only variable that was associated with PSE in multivariable analyses was cortical involvement (HR, 8.20 [95% CI, 2.22-30.25];  $P = .002$ ) (Table 2).


### External Validation Risk Scores

The C statistic of the SeLECT 2.0 risk score was 0.78 (95% CI, 0.71-0.84). The predicted and observed risks of PSE are depicted in Figure 3. The observed risk of PSE according to the SeLECT 2.0 risk score in our study was lower compared with the predicted risk of the original study for all scores, however, with mostly overlapping 95% CIs between the predicted and observed risk. The C statistic of the CAVE risk score was 0.83 (95% CI, 0.76-0.90). The observed risk of PSE according to the CAVE risk score varied in overestimation and underestimation across the scores, but with wide 95% CIs.


## Discussion

We showed a 5-year cumulative risk of PSE of 3.7% among patients with ischemic stroke and 7.6% among patients with ICH. For both types of stroke, the risk was highest within the first year after stroke. Variables that were associated with PSE were an acute symptomatic seizure and cortical involve-

Figure 2. Five-Year Cumulative Risk and Incidence Per 100 Person-Years of Poststroke Epilepsy (PSE) After an Ischemic Stroke or Intracerebral Hemorrhage



B Incidence rate



The incidence rate per 100 person-years was calculated by dividing the number of events by the total person-years at risk per 1-year intervals. The middle of those intervals was chosen as the point estimator in the graph. The shaded areas indicate 95% CIs.

ment. These variables were also included in the SeLECT 2.0 and CAVE risk scores. Both risk scores showed good discrimination and reasonable calibration in this young cohort of patients with stroke.

The risk of PSE in our study was lower compared with previous studies.<sup>10-13</sup> First, a previous Dutch cohort study reported a 5-year risk of 15% for patients with ischemic stroke and 21% for patients with ICH.<sup>10</sup> However, that study defined PSE by including both acute symptomatic seizures (between 1 day and 7 days) and remote symptomatic seizures based on the ILAE definitions at that time, whereas we included late seizures only due to the suggested different pathophysiological mechanisms. Although a remote symptomatic seizure is associated with chronic changes in the brain, an acute symptomatic seizure reflects acute and possibly reversible changes.<sup>25</sup> This difference likely resulted in an




Table 2. Age- and Sex-Adjusted and Multivariable Fine-Gray Proportional Hazard Models

Characteristic	Age and sex adjusted		Multivariable	
	HR (95% CI)	P value	HR (95% CI)	P value
<b>Ischemic stroke</b>				
Age	1.01 (0.96-1.05)	.63	NA	NA
Women	1.02 (0.56-1.85)	.94	NA	NA
NIHSS score at admission				
Minor stroke ( $\leq 3$ )	1 [Reference]	NA	1 [Reference]	NA
Major stroke ( $> 3$ )	2.26 (1.23-4.13)	.009	1.59 (0.83-3.02)	.16
Acute symptomatic seizure	12.97 (3.32-50.66)	<.001	10.83 (2.05-57.07)	.005
Cortical involvement	9.95 (3.45-28.70)	<.001	5.35 (1.85-15.49)	.002
Infarction in MCA territory	2.61 (1.21-5.67)	.01	1.61 (0.66-3.92)	.29
Cause of stroke				
Atherothrombotic	1.76 (0.45-6.84)	.17	1.13 (0.25-5.19)	.99
Likely atherothrombotic	1.33 (0.48-3.65)	.59	1.65 (0.59-4.56)	.34
Small vessel disease	NA	NA	NA	NA
Cardioembolic	1.38 (0.54-3.55)	.50	1.46 (0.55-3.90)	.45
Rare causes	2.00 (0.88-4.54)	.10	1.92 (0.77-4.78)	.16
Multiple causes	0.59 (0.08-4.61)	.61	0.75 (0.10-5.85)	.79
Cryptogenic	1 [Reference]	NA	1 [Reference]	NA
<b>Intracerebral hemorrhage</b>				
Age	0.94 (0.89-0.98)	.02	0.94 (0.89-1.00)	.65
Women	1.83 (0.63-5.28)	.26	NA	NA
NIHSS score at admission				
Minor stroke ( $\leq 3$ )	1 [Reference]	NA	NA	NA
Major stroke ( $> 3$ )	0.60 (0.20-1.79)	.36	NA	NA
Acute symptomatic seizure	7.10 (1.40-36.31)	.02	4.02 (0.87-18.70)	.08
Cortical involvement	9.30 (2.60-33.34)	<.001	8.20 (2.22-30.25)	.002
Bleeding volume $\geq 10$ mL	2.1 (0.7-6.9)	.21	NA	NA

Abbreviations: NIHSS, National Institutes of Health Stroke Scale; MCA, middle cerebral artery; NA, not applicable.

Figure 3. Bar Plots Showing Predicted and Observed Risk of Poststroke Epilepsy (PSE)



A, Risk after ischemic stroke according to the SeLECT (severity of stroke, large-artery atherosclerosis, early seizure, cortical involvement, and territory of middle cerebral artery) 2.0 score. B, Risk after intracerebral hemorrhage according to the CAVE (cortical involvement, age, bleeding volume, and early

seizure) score. The observed risk was calculated only for risk scores with at least 30 patients. Error bars indicate 95% CIs, which were not reported for the predicted risk per CAVE score.

overestimation of the PSE risk in the previous study. Second, the previous study included ischemic strokes without neuroimaging-based proof of cerebral ischemia,<sup>10</sup> as was the case in another retrospective study from Helsinki that

reported a cumulative 5-year risk of 9.5% (95% CI, 7.6%-11.4%) after an ischemic stroke.<sup>12</sup> The risk of including stroke mimics is higher without neuroimaging-based confirmation of ischemia.<sup>25</sup> As a consequence, previous studies might

have misclassified some seizures as transient ischemic attacks as the index event, leading to an overestimation of PSE risk. Last, a large Taiwanese retrospective cohort study showed a 5-year cumulative risk of 5.5% after ischemic stroke, which is more in line with our findings.<sup>11</sup> In that study, stroke and seizures were ascertained based on *International Classification of Diseases, Ninth Revision, Clinical Modification* codes, without verification of codes, which increases the risk of misclassification. We were able to confirm the relatively low risk of PSE among young adults through a well-defined cohort of patients with ischemic stroke.

We found characteristics that are associated with PSE to be similar to the variables in the SeLECT 2.0 and CAVE risk scores. These risk scores were derived from cohorts including mostly patients older than 65 years. Our study shows that these risk scores are also applicable to young adults, as is evident from the C statistics in our cohort that are similar to the original studies. We found a C statistic of 0.78 (95% CI, 0.71-0.84) for the SeLECT 2.0 score in our young cohort of patients with stroke, compared with the a C statistic of 0.77 (95% CI, 0.71-0.82) in their external validation.<sup>22</sup> The C statistic of the CAVE score in our study was 0.83 (95% CI, 0.76-0.90) compared with 0.81 (95% CI, 0.76-0.86) in the original derivation cohort and 0.69 (95% CI, 0.59-0.78) in an external validation cohort.<sup>23</sup> The CAVE score slightly underestimated the risk of PSE after ICH in our cohort, whereas the SeLECT 2.0 score slightly overestimated the observed risk. There is potential to improve discrimination of the risk scores among young adults. First, electroencephalographic (EEG) asymmetric background activity measured within the first 72 hours after stroke was a factor associated with PSE after ischemic stroke.<sup>26</sup> However, the clinical relevance and added value of EEG outcomes in the SeLECT 2.0 score have yet to be determined.<sup>27</sup> Second, cortical superficial siderosis improved the prediction of the SeLECT and CAVE scores; however, it has not been validated in a young cohort of patients with stroke.<sup>28</sup> Finally, it has been found that the addition of the inflammatory cytokine interleukin-1 $\beta$  improved the predictive value of the SeLECT score in a cohort of patients of all ages with ischemic stroke.<sup>29</sup> This finding could be explained by the release of an inflammatory cascade after neuronal damage.<sup>30</sup> Because trigger factors such as fever and flulike disease are highly prevalent among young patients with ischemic stroke, the addition of this cytokine might be of value, especially in the younger population.<sup>31</sup>

The present study aids neurologists in informing young patients with a stroke about the risk of PSE. In line with previous studies, we found the risk of PSE after an ICH to be higher compared with the risk after an ischemic stroke and that risk was highest in the first years after stroke. This is probably due to an increased risk of (cortical) damage in the brain after a large amount of bleeding and early processes after an ICH, such as edema and inflammation.<sup>32</sup> Although the risk was lower than previously described, the effect of PSE on daily life remains major, especially for young adults who have to deal with that uncertainty for decades to come (eg, not being

able to drive a car, difficulties in returning to work, or starting a family). Therefore, it is important to identify which patients are at the highest risk. Our study found that the SeLECT 2.0 and CAVE risk scores can be accurately applied in this relatively young patient group as well.

Apart from informing neurologists and patients, these risk scores may also be important for stratification in future clinical trials about optimal primary and secondary prophylactic treatment of PSE. Antiseizure medication as primary prophylactic treatment for patients with ischemic stroke and ICH is currently not recommended.<sup>25</sup> The PEACH (Prevention of Epileptic Seizures at the Acute Phase of Intracerebral Hemorrhage) trial investigated whether levetiracetam in the acute phase of an ICH reduced the risk of acute symptomatic seizures among patients with a mean age of 77 years.<sup>33</sup> Because an acute symptomatic seizure is a risk factor for remote symptomatic seizures, reducing the risk of acute symptomatic seizures might reduce the risk of PSE. The PEACH trial showed that levetiracetam as the primary prophylactic drug was safe but did not have any association with the occurrence of clinical seizures and did not show differences in functional outcome. In addition, several studies have suggested a reduced risk of PSE among statin users with ischemic stroke or ICH compared with patients who do not take statins, although the evidence is conflicting.<sup>34</sup> Much remains uncertain regarding the prevention of PSE after stroke. Future studies should recruit patients at high risk for which risk scores can support identifying these patients.

### Strengths and Limitations

Our study has some strengths. ODYSSEY is a large prospective multicenter study, with only patients with neuroimaging-proven strokes, ensuring a very small risk of misclassifying the index stroke. Second, all the variables to validate existing risk scores were available within the ODYSSEY cohort. Third, we included both patients with ischemic stroke and patients with ICH, whereas most studies included only patients with ischemic strokes.

Nevertheless, there are also some limitations. First, all seizures were self-reported by the patients, which could have resulted in an underestimation of the occurrence of PSE. However, we cross-checked this method and found perfect agreement. Second, the number of PSE cases in our study is relatively low, especially for patients with ICH. This could have hampered the validity of the association with variables. Third, the median NIHSS score in ODYSSEY was relatively low but comparable with the NIHSS score in previous studies of young patients with stroke.<sup>10-12</sup> This indicates that although limited generalization cannot be excluded, this risk seems limited. Our results are not generalizable to a young stroke population that is more severely affected. Fourth, data on the use and type of antiseizure medication were collected only during the first seizure. We therefore lack data to report on how many patients developed drug-resistant seizures according to the current definition of the ILAE.<sup>35</sup>

## Conclusions

This cohort study found that the risk of PSE among young adults with ischemic stroke or ICH was relatively low and was highest

in the first year after stroke. Existing risk scores may be implemented for patients experiencing a stroke at a young age and can be used to reliably inform young patients about the risk of PSE. Future clinical trials should investigate the optimal primary and secondary prophylaxis for patients at high risk.

### ARTICLE INFORMATION

**Accepted for Publication:** December 19, 2024.

**Published Online:** April 14, 2025.

doi:10.1001/jamaneurol.2025.0465

**Author Affiliations:** Department of Neurology, Radboud Institute of Medical Innovation, Donders Institute for Brain, Cognition and Behavior, Nijmegen, the Netherlands (Verburt, Fella, Ekker, Schellekens, Boot, Immens, Tuladhar, Verhoeven, Hilken, de Leeuw); Department of Neurology, Gelre Hospital, Apeldoorn, the Netherlands (van Alebeek); Department of Neurology, Medisch Spectrum Twente, Enschede, the Netherlands (Brouwers, Arntz); Department of Neurology, Canisius-Wilhelmina Hospital, Nijmegen, the Netherlands (van Dijk); Department of Neurology, Catharina Hospital, Eindhoven, the Netherlands (Gons, van Uden); Department of Neurology, Franciscus Gasthuis & Vlietland Hospital, Rotterdam, the Netherlands (den Heijer); Department of Neurology, Elisabeth-TweeSteden Hospital, Tilburg, the Netherlands (van Tuijl); Department of Neurology, Haga Hospital, The Hague, the Netherlands (de Laat); Department of Neurology, Amphia Hospital, Breda, the Netherlands (van Norden); Department of Neurology, Rijnstate Hospital, Arnhem, the Netherlands (Vermeer); Department of Neurology, Jeroen Bosch Hospital, 's-Hertogenbosch, the Netherlands (van Zagten); Department of Neurology, Maastricht University Medical Centre, Maastricht, the Netherlands (van Oostenbrugge); Department of Neurology, Leiden University Medical Centre, Leiden, the Netherlands (Wermer); Department of Neurology, University Medical Centre Groningen, Groningen, the Netherlands (Wermer); Department of Neurology, Amsterdam University Medical Centre, Amsterdam, the Netherlands (Nederkoorn); Department of Neurology, Albert Schweitzer Hospital, Dordrecht, the Netherlands (Kerkhoff); Department of Neurology, Zuyderland Hospital, Sittard-Geleen, the Netherlands (Rooyer); Department of Neurology, Medical Centre Leeuwarden, Leeuwarden, the Netherlands (van Rooij); Department of Neurology, Haaglanden Medical Centre, The Hague, the Netherlands (van den Wijngaard).

**Author Contributions:** Ms Verburt and Dr de Leeuw had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** Verburt, Brouwers, Arntz, van Uden, den Heijer, Vermeer, Tuladhar, Verhoeven, Hilken, de Leeuw.

**Acquisition, analysis, or interpretation of data:** Verburt, Fella, Ekker, Schellekens, Boot, Immens, van Alebeek, Brouwers, van Dijk, Gons, van Tuijl, de Laat, van Norden, van Zagten, van Oostenbrugge, Wermer, Nederkoorn, Kerkhoff, Rooyer, van Rooij, van den Wijngaard, Tuladhar, Verhoeven, Hilken, de Leeuw.

**Drafting of the manuscript:** Verburt, Fella, Nederkoorn.  
**Critical review of the manuscript for important**

**intellectual content:** All authors.

**Statistical analysis:** Verburt, Fella, Verhoeven, Hilken.

**Obtained funding:** Tuladhar, de Leeuw.

**Administrative, technical, or material support:** Verburt, Immens, van Alebeek, van Norden, van Oostenbrugge, Rooyer.

**Supervision:** Ekker, Brouwers, van Zagten, Kerkhoff, van den Wijngaard, Tuladhar, Hilken, de Leeuw.

**Conflict of Interest Disclosures:** Dr van den Wijngaard reported receiving personal fees from Philips and Medtronic outside the submitted work. Dr de Leeuw reported serving as associate editor of the *International Journal of Stroke*. No other disclosures were reported.

**Funding/Support:** This study was supported by the Radboud University Medical Centre (Dr Verhoeven), Dutch Heart Foundation (grant 2016T86 [Dr Wermer], grant 03-005-2022-0031 [Dr Hilken], grant 2016T044 [Dr Tuladhar], and grant 2014 T060 [Dr de Leeuw]), and Vidi (grant 9171337 of the ZorgOnderzoek Nederland and the Medical Sciences/Nederlandse Organisatie voor Wetenschappelijk Onderzoek [Dr Wermer] and grant 016-126-351 [Dr de Leeuw]).

**Role of the Funder/Sponsor:** The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Data Sharing Statement:** See Supplement 2.

### REFERENCES

- Zelano J, Redfors P, Åsberg S, Kumlien E. Association between poststroke epilepsy and death: a nationwide cohort study. *Eur Stroke J*. 2016;1(4):272-278. doi:10.1177/2396987316669000
- Arntz RM, Rutten-Jacobs LC, Maaijwee NA, et al. Poststroke epilepsy is associated with a high mortality after a stroke at young age: follow-up of Transient Ischemic Attack and Stroke Patients and Unelucidated Risk Factor Evaluation study. *Stroke*. 2015;46(8):2309-2311. doi:10.1161/STROKEAHA.115.010115
- Misra S, Kasner SE, Dawson J, et al. Outcomes in patients with poststroke seizures: a systematic review and meta-analysis. *JAMA Neurol*. 2023;80(11):1155-1165. doi:10.1001/jamaneurol.2023.3240
- Arntz RM, Maaijwee NA, Rutten-Jacobs LC, et al. Epilepsy after TIA or stroke in young patients impairs long-term functional outcome: the FUTURE Study. *Neurology*. 2013;81(22):1907-1913. doi:10.1212/01.wnl.0000436619.25532.f3
- Baranowski CJ. The quality of life of older adults with epilepsy: a systematic review. *Seizure*. 2018;60:190-197. doi:10.1016/j.seizure.2018.06.002
- de Boer HM, Mula M, Sander JW. The global burden and stigma of epilepsy. *Epilepsy Behav*. 2008;12(4):540-546. doi:10.1016/j.yebeh.2007.12.019
- Tanaka T, Fukuma K, Abe S, et al; PROPOSE Study Investigators. Antiseizure medications for post-stroke epilepsy: a real-world prospective cohort study. *Brain Behav*. 2021;11(9):e2330. doi:10.1002/brb3.2330
- Mula M, von Oertzen TJ, Cock HR, Yogarajah M, Lozsadi DA, Agrawal N. Fatigue during treatment with antiepileptic drugs: a levetiracetam-specific adverse event? *Epilepsy Behav*. 2017;72:17-21. doi:10.1016/j.yebeh.2017.04.030
- Weintraub D, Buchsbaum R, Resor SR Jr, Hirsch LJ. Psychiatric and behavioral side effects of the newer antiepileptic drugs in adults with epilepsy. *Epilepsy Behav*. 2007;10(1):105-110. doi:10.1016/j.yebeh.2006.08.008
- Arntz R, Rutten-Jacobs L, Maaijwee N, et al. Post-stroke epilepsy in young adults: a long-term follow-up study. *PLoS One*. 2013;8(2):e55498. doi:10.1371/journal.pone.0055498
- Do PT, Chen LY, Chan L, Hu CJ, Chien LN. Risk factors for postischemic stroke epilepsy in young adults: a nationwide population-based study in Taiwan. *Front Neurol*. 2022;13:880661. doi:10.3389/fneur.2022.880661
- Roivainen R, Haapaniemi E, Putaala J, Kaste M, Tatlisumak T. Young adult ischaemic stroke related acute symptomatic and late seizures: risk factors. *Eur J Neurol*. 2013;20(9):1247-1255. doi:10.1111/ene.12157
- Naess H, Nyland HI, Thomassen L, Aarseth J, Myhr KM. Long-term outcome of cerebral infarction in young adults. *Acta Neurol Scand*. 2004;110(2):107-112. doi:10.1111/j.1600-0404.2004.00273.x
- Zou S, Chen Y. Research progress on the prediction of post-stroke epilepsy. *Acta Epileptol*. 2020;2:24. doi:10.1186/s42494-020-00031-z
- Arntz RM, van Alebeek ME, Synhaeve NE, et al. Observational Dutch Young Symptomatic Stroke study (ODYSSEY): study rationale and protocol of a multicenter prospective cohort study. *BMC Neurol*. 2014;14:55. doi:10.1186/1471-2377-14-55
- Easton JD, Saver JL, Albers GW, et al; American Heart Association; American Stroke Association Stroke Council; Council on Cardiovascular Surgery and Anesthesia; Council on Cardiovascular Radiology and Intervention; Council on Cardiovascular Nursing; Interdisciplinary Council on Peripheral Vascular Disease. Definition and evaluation of transient ischemic attack: a scientific statement for healthcare professionals from the American Heart Association/American Stroke Association Stroke Council; Council on Cardiovascular Surgery and Anesthesia; Council on Cardiovascular Radiology and Intervention; Council on Cardiovascular Nursing; and the Interdisciplinary Council on Peripheral Vascular Disease. *Stroke*. 2009;40(6):2276-2293. doi:10.1161/STROKEAHA.108.192218
- Fisher RS, Acevedo C, Arzamanoglou A, et al. ILAE official report: a practical clinical definition of epilepsy. *Epilepsia*. 2014;55(4):475-482. doi:10.1111/epi.12550



18. Verburgt E, Hillkens NA, Ekker MS, et al. Short-term and long-term risk of recurrent vascular event by cause after ischemic stroke in young adults. *JAMA Netw Open*. 2024;7(2):e240054. doi:10.1001/jamanetworkopen.2024.0054
19. Boussier MG, Amarenco P, Chamorro A, et al; PERFORM Study Investigators. Rationale and design of a randomized, double-blind, parallel-group study of terutroban 30 mg/day versus aspirin 100 mg/day in stroke patients: the prevention of cerebrovascular and cardiovascular events of ischemic origin with terutroban in patients with a history of ischemic stroke or transient ischemic attack (PERFORM) study. *Cerebrovasc Dis*. 2009;27(5):509-518. doi:10.1159/000212671
20. Brott T, Adams HP Jr, Olinger CP, et al. Measurements of acute cerebral infarction: a clinical examination scale. *Stroke*. 1989;20(7):864-870. doi:10.1161/01.STR.20.7.864
21. Fischer U, Baumgartner A, Arnold M, et al. What is a minor stroke? *Stroke*. 2010;41(4):661-666. doi:10.1161/STROKEAHA.109.572883
22. Sinka L, Abaira L, Imbach LL, et al. Association of mortality and risk of epilepsy with type of acute symptomatic seizure after ischemic stroke and an updated prognostic model. *JAMA Neurol*. 2023;80(6):605-613. doi:10.1001/jamaneurol.2023.0611
23. Haapaniemi E, Strbian D, Rossi C, et al. The CAVE score for predicting late seizures after intracerebral hemorrhage. *Stroke*. 2014;45(7):1971-1976. doi:10.1161/STROKEAHA.114.004686
24. Kothari RU, Brott T, Broderick JP, et al. The ABCs of measuring intracerebral hemorrhage volumes. *Stroke*. 1996;27(8):1304-1305. doi:10.1161/01.STR.27.8.1304
25. Zelano J, Holtkamp M, Agarwal N, Lattanzi S, Trinka E, Brigo F. How to diagnose and treat post-stroke seizures and epilepsy. *Epileptic Disord*. 2020;22(3):252-263. doi:10.1684/epd.2020.1159
26. Bentes C, Martins H, Peralta AR, et al. Early EEG predicts poststroke epilepsy. *Epilepsia Open*. 2018;3(2):203-212. doi:10.1002/epi4.12103
27. Doerrfuss JI, Kilic T, Ahmadi M, Holtkamp M, Weber JE. Quantitative and qualitative EEG as a prediction tool for outcome and complications in acute stroke patients. *Clin EEG Neurosci*. 2020;51(2):121-129. doi:10.1177/1550059419875916
28. Tanaka T, Fukuma K, Abe S, et al. Association of cortical superficial siderosis with post-stroke epilepsy. *Ann Neurol*. 2023;93(2):357-370. doi:10.1002/ana.26497
29. Shen L, Yang J, Tang Y. Predictive values of the SeLECT Score and IL-1 $\beta$  for post-stroke epilepsy. *Neuropsychiatr Dis Treat*. 2021;17:2465-2472. doi:10.2147/NDT.S324271
30. Esenwa CC, Elkind MS. Inflammatory risk factors, biomarkers and associated therapy in ischaemic stroke. *Nat Rev Neurol*. 2016;12(10):594-604. doi:10.1038/nrneurol.2016.125
31. Ekker MS, Verhoeven JI, Rensink KML, et al. Trigger factors for stroke in young adults: a case-crossover study. *Neurology*. 2023;100(1):e49-e61. doi:10.1212/WNL.000000000000201341
32. Ravikummar A, Bohoopathy RM, Laha A, Mani R. Post-stroke seizures: a comprehensive review of epidemiology, pathophysiology, risk factors, clinical spectrum and outcomes. *Med Res Arch*. 2024;12(10):1-16. doi:10.18103/mra.v12i10.5768
33. Peter-Derex L, Philippeau F, Garnier P, et al. Safety and efficacy of prophylactic levetiracetam for prevention of epileptic seizures in the acute phase of intracerebral haemorrhage (PEACH): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Neurol*. 2022;21(9):781-791. doi:10.1016/S1474-4422(22)00235-6
34. Nucera B, Rinaldi F, Nardone R, Lattanzi S, Brigo F. Statins in primary prevention of poststroke seizures and epilepsy: a systematic review. *Epilepsy Behav*. 2020;112:107400. doi:10.1016/j.yebeh.2020.107400
35. Kwan P, Arzimanoglou A, Berg AT, et al. Definition of drug resistant epilepsy: consensus proposal by the ad hoc Task Force of the ILAE Commission on Therapeutic Strategies. *Epilepsia*. 2010;51(6):1069-1077. doi:10.1111/j.1528-1167.2009.02397.x