

RESEARCH LETTER

Oral Anticoagulation and Factor VIII Replacement Therapy in Patients With Hemophilia Undergoing Pulsed-Field or Radiofrequency Catheter Ablation for Atrial Fibrillation

Sanghamitra Mohanty¹, MD, MS; Michela Casella², MD; Paolo Compagnucci³, MD; Prem Geeta Torlapati⁴, MD, MPH; Vincenzo Mirco La Fazio⁵, MD; Carola Gianni⁶, MD, PhD; Bryan MacDonald⁷, MD; Angel Quintero Mayedo⁸, MD; Domenico Giovanni Della Rocca⁹, MD, PhD; John Allison¹⁰, MD; Mohamed Bassiouny¹¹, MD; G. Joseph Gallinghouse¹², MD; John D. Burkhardt¹³, MD; Rodney Horton¹⁴, MD; Amin Al-Ahmad¹⁵, MD; Luigi Di Biase¹⁶, MD, PhD; Antonio Dello Russo¹⁷, MD; Andrea Natale¹⁸, MD

Hemophilia is characterized by reduced or unmeasurable levels of coagulation factor VIII (hemophilia A) or factor IX (hemophilia B).¹ Since the introduction of clotting factor concentrates, the life expectancy of patients with hemophilia has increased to over 70 years making them more likely to be confronted with age-related comorbidities like atrial fibrillation (AF). Anticoagulation is one of the main components of AF management to reduce the risk of AF-related strokes. However, hemophiliacs have a high bleeding tendency making it challenging to find a balance between coagulation and anticoagulation in patients with coexistent hemophilia and AF.¹ According to guidelines and expert opinions, coagulation factor concentrate is given prophylactically to maintain trough factor levels >80% during the periprocedural period and ≥30% during the next 4 to 6 weeks in patients undergoing AF ablation.¹

Pulsed-field ablation (PFA) has demonstrated tissue selectivity via application of ultra-rapid electrical pulses causing irreversible electroporation and subsequent cellular death. Notably, cardiomyocytes exhibit a lower threshold for irreversible electroporation than other cell types including the endothelial cells that remain unaffected during the procedure.² Thus, given the fact that endothelial injury promotes thrombus formation, it is

prudent to speculate that in its absence anticoagulation management would be less challenging in patients with AF with hemophilia undergoing PFA.

In this case series, we report our experiences with patients with hemophilia undergoing radiofrequency ablation (RFA) or PFA.

This multicenter study included consecutive patients with hemophilia undergoing AF ablation.

Patients' data were collected from IRB-approved institutional databases and retrospective medical record review. The data that support the findings of this study are available from the corresponding author upon reasonable request.

ABLATION PROCEDURE

All patients received pulmonary vein isolation plus isolation of left atrial posterior wall using either RFA or PFA. The ablation procedures have been described in detail in earlier publications.³

COAGULATION MANAGEMENT

Patients received clotting factor replacement to bring the preprocedure level to 60% to 150%. During the

Key Words: atrial fibrillation ■ hemophilia ■ interquartile range ■ life expectancy ■ oral anticoagulation ■ pulsed-field ablation

Correspondence to: Andrea Natale, MD, Texas Cardiac Arrhythmia Institute, St. David's Medical Center, Austin, TX. Email dr.natale@gmail.com

For Sources of Funding and Disclosures, see page 747.

© 2024 American Heart Association, Inc.

Circulation: Arrhythmia and Electrophysiology is available at www.ahajournals.org/journal/circep

Nonstandard Abbreviations and Acronyms

| | |
|------------|-------------------------|
| AF | atrial fibrillation |
| IQR | interquartile range |
| PFA | pulsed-field ablation |
| RFA | radiofrequency ablation |

ablation procedure, heparin boluses of 6000 to 12000 units were given to maintain an ACT of 300 to 500 seconds.

Anticoagulation was continued for at least 45 days following the RFA and 1 week in the PFA group. In addition, patients received factor VIII replacement therapy for 1 week in the PFA group to up to 5 weeks in the RFA group. All patients were monitored for thromboembolic events for 1-year.

This series included 8 patients (no. 6: hemophilia B, others: hemophilia A; age: 61.8 ± 6.978 (median, 62.5; interquartile range [IQR], 68–57) years of which 3 (37.5%) received PFA and 5 subjects underwent RFA for drug-refractory AF. Mean $\text{CHA}_2\text{DS}_2\text{-VASc}$ score were 2.67 ± 1.15 (median, 2; IQR, 4–2) and 2.80 ± 1.48

(median, 3; IQR, 4–1.5) and the HAS-BLED score were 3.33 ± 1.53 (median, 3; IQR, 5–2) and 2.80 ± 1.10 (median, 2; IQR, 4–2) in the PFA and the RFA cohorts, respectively.

Details of pre- and postprocedural factor VIII replacement therapy and anticoagulation are provided in the Table. Preprocedural clotting factor level (PFA, $93.43 \pm 47.99\%$ [median, 61; IQR, 161.3–58] versus RFA, $66.44 \pm 4.61\%$ [median, 59; IQR, 68.6–56.5]) and the amount of clotting factor administered (PFA, 3323.33 ± 365.56 IU [median, 3250; IQR, 3720–3000] versus RFA, 2096.00 ± 839.48 [median, 1670; IQR, 3000–1405]) were similar in the patients receiving PFA or RFA. Following the procedure, patients with PFA received less clotting factor replacement therapy compared with the RFA group (PFA, 78866.67 ± 1761.63 ; median, 79 100; IQR, 80 500–77 000) versus RFA (152260.00 ± 29704.51 IU; median, 140 000; IQR, 182950–127700). The duration of the replacement therapy was also lower in the PFA group (6.65 ± 0.229 ; median, 6; IQR, 7–5) versus (31 \pm 3.316; median, 30; IQR, 32–30) days.

No bleeding or thromboembolic events were reported at 1-year follow-up.

Table. Pre- and Postprocedure Oral anticoagulation and Clotting Factor Replacement Therapy

| Pre-procedure | | | | | Post-procedure | | | | | |
|-----------------------------|-----------------------------------|---|---------------------------|--|-----------------------------|-----------------------------------|--|---------------------------|------|-------|
| Patient (type of procedure) | Clotting factor administered (IU) | Administration strategy (dosage) | Clotting factor level (%) | Pre-procedure OAC | Post-procedure OAC | Clotting factor administered (IU) | Administration strategy (dosage) | Clotting factor level (%) | | |
| | | | | | | | | 24 h | 72 h | 1 wk |
| 1 (PFA-PVI) | 3720 | 2 doses of 1860 IU–12 h apart | 161.30 | Apixaban, 5 mg BID | Apixaban, 5 mg BID for 7 d | 79 100 | 5 doses of 1130 units 12 hourly over a week | 84 | 78.9 | 137 |
| 2 (PFA-PVI) | 3000 | 2 doses of 1500 IU–12 h apart | 58.00 | Apixaban, 5 mg BID | Apixaban, 5 mg BID for 7 d | 77 000 | 5 doses of 1100 units 12 hourly over a week | 67 | 81 | 128.9 |
| 3 (PFA-PVI) | 3250 | 2 doses of 1575 IU–12 h apart | 61.00 | Apixaban, 5 mg BID | Apixaban, 5 mg BID for 7 d | 80 500 | 5 doses of 1150 units 12 hourly over a week | 72 | 79 | 130 |
| 4 (RFA-PVI) | 1250 | 1 dose of 1250 IU–24 h before the procedure | 66.00 | Initially warfarin then Apixaban, 5 mg BID | Apixaban, 5 mg BID for 45 d | 120 400 | 2150 units, 12 hourly for 28 d | 54.3 | 156 | 131.6 |
| 5 (RFA-PVI) | 1560 | 1 dose of 1560 IU–24 h before the procedure | 71.20 | Warfarin, 5 mg daily | Apixaban, 5 mg BID for 45 d | 192 500 | 2750 units, 12 hourly for 35 d | 56.9 | 82.7 | 100.8 |
| 6 (RFA-PVI) | 3000 | 1 dose of 3000 IU–24 h before the procedure | 59.00 | Enoxaparin, 90 mg BID | No OAC | 140 000 | One dose of 5000 units followed by 2500 units 12 hourly for 28 d | 38.7 | 54 | 62.9 |
| 7 (RFA-PVI) | 1670 | 1 dose of 1670 IU–24 h before the procedure | 56.00 | Apixaban, 5 mg BID | Apixaban, 5 mg BID for 45 d | 135 000 | 2250 units, 12 hourly for 30 d | 58.5 | 67 | 89 |
| 8 (RFA-PVI) | 3000 | 1 dose of 3000 IU–24 h before the procedure | 57.00 | Apixaban, 5 mg BID | Apixaban, 5 mg BID for 45 d | 173 400 | 2550 units, 12 hourly for 34 d | 59.1 | 69 | 88 |

BID indicates twice a day; OAC, oral anticoagulation; PFA, pulsed-field ablation; PVI, pulmonary vein isolation; and RFA, radiofrequency ablation.

To the best of our knowledge, this is the first series to report the difference in anticoagulation and clotting factor replacement therapy in patients with AF with hemophilia undergoing PFA or RFA. The current gold-standard treatment for hemophilia is prophylactic factor VIII replacement therapy that requires intravenous (IV) infusion. With a half-life of 10 hours, the IV infusion is essential twice daily to maintain the concentrations of the clotting factor high enough to provide adequate support for coagulation to prevent spontaneous bleeding.⁴ The inconvenience of such frequent IV infusions and maintenance of a central venous line for an extended period cannot be overemphasized. Moreover, the burden of frequent IV infusions compromises the quality of life by making active lifestyles challenging.⁴ In patients with AF with hemophilia, the type, intensity, and duration of antithrombotic therapy are determined, by the risk of AF-associated ischemic stroke, the bleeding risk, and the need for replacement therapy ensuring a level of coagulation factor that allows for safe antithrombotic treatment.⁵ Endothelial injury induced by the thermal effect of RF ablation is known to enhance the risk of thromboembolism whereas PFA is documented to not cause any injury to the vascular endothelium. In this series, following the PFA procedure, the factor VIII replacement therapy and oral anticoagulation could be safely discontinued after 1 week without any bleeding or thromboembolic events during the 1-year follow-up period.

The findings from this case series suggest that in hemophilia patients undergoing AF ablation, it might be possible to safely discontinue factor VIII replacement therapy and oral anticoagulation after a relatively shorter period following PFA compared with radiofrequency ablation. However, larger studies are required to replicate these findings before any major conclusions can be reached.

ARTICLE INFORMATION

Affiliations

Texas Cardiac Arrhythmia Institute, St. David's Medical Center, Austin, TX (S.M., P.G.T., V.M.L.F., C.G., B.M., A.Q.M., J.A., M.B., G.J.G., J.D.B., R.H., A.A.-A., A.N.). Department of Electrophysiology, Ospedale Riuniti Hospital, Ancona, Italy (M.C., P.C., A.D.R.). Department of Cardiology, University Hospital (UZ) Brussels, Belgium (D.G.D.R.). Department of Electrophysiology, Albert Einstein College of Medicine at Montefiore Hospital, Bronx, NY (L.D.B.). Interventional Electrophysiology, Scripps Clinic, San Diego, CA (A.N.). Department of Internal Medicine, Metro Health Medical Center, Case Western Reserve University School of Medicine, Cleveland, OH (A.N.). Department of Biomedicine and Prevention, Division of Cardiology, University of Tor Vergata, Rome, Italy (A.N.).

Sources of Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Disclosures

Dr Natale is a consultant for Abbott, Biosense Webster, Biotronik, Boston Scientific and iRhythm. Dr Dello Russo is a consultant for Abbott Medical. Dr Burkhardt is a consultant for Biosense Webster and Stereotaxis. Dr Di Biase is a consultant for Biosense Webster, Boston Scientific, Stereotaxis, and St. Jude Medical and has received speaker honoraria from Medtronic, Atricure, EPIEP, and Biotronik. The other authors report no conflicts.

REFERENCES

- Franchini M, Focosi D, Mannucci PM. How we manage cardiovascular disease in patients with hemophilia. *Haematologica*. 2023;108:1748–1757. doi: 10.3324/haematol.2022.282407
- Inoue K. Pulsed-Field ablation for atrial fibrillation - Achieving excellence with a simplified technique. *Circ J*. 2023;87:1727–1729. doi: 10.1253/circj.CJ-23-0596
- Verma A, Haines DE, Boersma LV, Sood N, Natale A, Marchlinski FE, Calkins H, Sanders P, Packer DL, Kuck KH, et al; PULSED AF Investigators. Pulsed field ablation for the treatment of atrial fibrillation: PULSED AF Pivotal Trial. *Circulation*. 2023;147:1422–1432. doi: 10.1161/CIRCULATIONAHA.123.063988
- Lambert T, Benson G, Dolan G, Hermans C, Jiménez-Yuste V, Ljung R, Morfini M, Zupančič-Salek S, Santagostino E. Practical aspects of extended half-life products for the treatment of haemophilia. *Ther Adv Hematol*. 2018;9:295–308. doi: 10.1177/2040620718796429
- Badescu MC, Badulescu OV, Butnariu LI, Floria M, Ciocoiu M, Costache II, Popescu D, Bratoiu I, Buliga-Finis ON, Rezus C. Current therapeutic approach to atrial fibrillation in patients with congenital hemophilia. *J Pers Med*. 2022;12:519. doi: 10.3390/jpm12040519