


Journal Pre-proof

A decision algorithm for assessing indirectness in Core GRADE

Gordon Guyatt, distinguished professor, Prashanti Eachempati, adjunct professor, Monica Hultcrantz, head of HTA Region Stockholm, Victor Montori, professor, Per Olav Vandvik, professor, Alfonso Iorio, professor, Thomas Agoritsas, professor



PII: S0895-4356(25)00432-9

DOI: <https://doi.org/10.1016/j.jclinepi.2025.112099>

Reference: JCE 112099

To appear in: *Journal of Clinical Epidemiology*

Received Date: 18 November 2025

Accepted Date: 3 December 2025

Please cite this article as: Guyatt G, Eachempati P, Hultcrantz M, Montori V, Vandvik PO, Iorio A, Agoritsas T, A decision algorithm for assessing indirectness in Core GRADE, *Journal of Clinical Epidemiology* (2026), doi: <https://doi.org/10.1016/j.jclinepi.2025.112099>.

This is a PDF of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability. This version will undergo additional copyediting, typesetting and review before it is published in its final form. As such, this version is no longer the Accepted Manuscript, but it is not yet the definitive Version of Record; we are providing this early version to give early visibility of the article. Please note that Elsevier's sharing policy for the Published Journal Article applies to this version, see: <https://www.elsevier.com/about/policies-and-standards/sharing#4-published-journal-article>. Please also note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 Elsevier Inc. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

Title: A decision algorithm for assessing indirectness in Core GRADE

Authors:

Gordon Guyatt, distinguished professor^{1 2 3*}, **Prashanti Eachempati, adjunct professor**,^{3 4 5*}, Monica Hultcrantz, head of HTA Region Stockholm^{6 7}, Victor Montori, professor^{8 9}, Per Olav Vandvik, professor^{3 10 11}, Alfonso Iorio, professor^{1 2}, Thomas Agoritsas, professor^{1 3 12}

***Both authors contributed equally (joint first authors)**

1. Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada.
2. Department of Medicine, McMaster University, Hamilton, ON, L8S 4L8, Canada
3. MAGIC Evidence Ecosystem Foundation, Oslo, Norway
4. Peninsula Dental School, University of Plymouth, United Kingdom
5. Faculty of Dentistry, Manipal University College Malaysia, Melaka, Malaysia
6. HTA Region Stockholm, Centre for Health Economics, Informatics and Health Care Research (CHIS), Stockholm Health Care Services, Sweden
7. Department of Learning, Informatics, Management and Ethics, Karolinska Institutet, Stockholm, Sweden
8. Division of Endocrinology, Department of Medicine, Mayo Clinic, Rochester, MN, USA
9. Knowledge and Evaluation Research Unit, Mayo Clinic, Rochester, MN, USA
10. Institute of Health and Society, University of Oslo Faculty of Medicine, Oslo, Norway
11. Department of Medicine, Lovisenberg Diakonale Hospital, Oslo, Norway
12. Division General Internal Medicine, University Hospitals of Geneva, University of Geneva, Geneva, Switzerland

Corresponding author: Prashanti Eachempati

Email: prashanti@magicevidence.org

Phone: +447917977090

Gordon Guyatt, distinguished professor ^{1 2 3*}, Prashanti Eachempati, adjunct professor, ^{3 4 5*}, Monica Hultcrantz, head of HTA Region Stockholm ^{6 7}, Victor Montori, professor ^{8 9}, Per Olav Vandvik, professor ^{3 10 11}, Alfonso Iorio, professor ^{1 2}, Thomas Agoritsas, professor ^{1 3 12}

1. Department of Health Research Methods, Evidence, and Impact, McMaster University, Hamilton, Ontario, Canada
2. Department of Medicine, McMaster University, Hamilton, ON, L8S 4L8, Canada
3. MAGIC Evidence Ecosystem Foundation, Oslo, Norway
4. Peninsula Dental School, University of Plymouth, United Kingdom
5. Faculty of Dentistry, Manipal University College Malaysia, Melaka, Malaysia
6. HTA Region Stockholm, Centre for Health Economics, Informatics and Health Care Research (CHIS), Stockholm Health Care Services, Sweden
7. Department of Learning, Informatics, Management and Ethics, Karolinska Institutet, Stockholm, Sweden
8. Division of Endocrinology, Department of Medicine, Mayo Clinic, Rochester, MN, USA
9. Knowledge and Evaluation Research Unit, Mayo Clinic, Rochester, MN, USA
10. Institute of Health and Society, University of Oslo Faculty of Medicine, Oslo, Norway
11. Department of Medicine, Lovisenberg Diakonale Hospital, Oslo, Norway
12. Division General Internal Medicine, University Hospitals of Geneva, University of Geneva, Geneva, Switzerland

***Both authors contributed equally (joint first authors).**

Letter to Editor:

In the recent *Core GRADE 5: Rating certainty of evidence—assessing indirectness* [1], published in The BMJ, and in the complementary Journal of Clinical Epidemiology article, *Core GRADE Unpacked: A Summary of Recent Innovations in Complementary GRADE Methodology* [2], we outlined the Core GRADE principles and explained how Core GRADE users can judge whether or not to rate down for indirectness when Population, Intervention, Comparator, or Outcome (PICO) elements diverge from the target PICO. Subsequent to these publications, we have reflected on the desirability of an algorithm to assist users in making this decision and decided such an algorithm would be helpful. *Figure 1* presents the algorithm as a logical, stepwise complement to the existing text.

The process begins by determining whether the evidence is direct or indirect. Although indirectness is rarely a concern in systematic reviews of direct evidence, it is frequently an issue in guidelines or HTAs in which the PICO in the relevant studies differs from the target PICO. Thus, when recognition of the limitations of direct evidence (low or very low certainty

for a particular outcome) prompts an explicit search for indirect evidence, serious concerns about indirectness are very likely to arise.

For Population-Intervention-Comparator, the algorithm first prompts assessors to judge whether there are differences – in, for example, age, disease severity, intervention dose, clinician expertise or comparators - between the target and study PICOs. If there are differences, it asks whether biological or social factors are sufficiently similar to expect the same impact on the target outcome. The degree of dissimilarity then dictates whether to rate down once, twice, or not at all.

For Outcome, the algorithm focuses on two common sources of indirectness: surrogate outcomes and mismatches in outcome definition or timing. It directs assessors to judge how closely a surrogate links to the patient-important endpoint and whether the measurement aligns with what matters to patients. [3]

We anticipate this visual decision-support tool will improve consistency and transparency in indirectness assessments, enhance GRADE training, and serve as a reference for systematic reviewers, health technology assessors, and guideline developers.

Competing interests: None declared.


References

1. Guyatt G, Iorio A, De Beer H, Owen A, Agoritsas T, Murad MH, et al. Core GRADE 5: rating certainty of evidence—assessing indirectness. *BMJ*. 2025;389: e083865. doi:10.1136/bmj-2024-083865.
2. Colunga-Lozano LE, Wang Y, Agoritsas T, Hultcrantz M, Iorio A, Montori VM, Guyatt G, Core GRADE unpacked: A summary of recent innovations in complementary GRADE Methodology, *Journal of Clinical Epidemiology* (2025), doi: <https://doi.org/10.1016/j.jclinepi.2025.112047>.
3. Christensen R, Ciani O, Manyara AM, Taylor RS. Surrogate endpoints: a key concept in clinical epidemiology. *J Clin Epidemiol*. 2024; 167:111242. doi: 10.1016/j.jclinepi.2023.111242.


Figure legend:

Figure 1. Decision-algorithm for assessing indirectness in Core GRADE.

Journal Pre-proof



*Although some outcomes may be considered surrogates, they are so closely and directly linked to the disease process that they always represent a patient-important outcome. For example, a hip fracture diagnosed on radiograph inevitably leads to pain, immobility, and the need for treatment, and in malaria, treatment failure (recurrence or recrudescence) confirmed by uncorrected or corrected PCR is a patient-important outcome.



Conflict of Interest Declaration

The authors declare that they have no conflicts of interest in relation to this submission. No financial, personal, or institutional relationships have influenced the development of this algorithm or the preparation of the accompanying letter to the editor