Check for updates

Circulation: Genomic and Precision Medicine

REVIEW

High-Throughput Methods for Variant Functional Assessment in Cardiac Disease

Richard E. Dolder III, BS*; Clayton E. Friedman, PhD*; Alexander M. Loiben, PhD*; Kai-Chun Yang, MD**; Andrew M. Glazer, PhD**

ABSTRACT: In vitro functional modeling of genetic variants has revolutionized our understanding of which variants can cause cardiac disorders, providing insights into their molecular underpinnings. This review provides an overview of high-throughput methods used for the functional assessment of variants implicated in inherited cardiac diseases. Advances in gene-editing technology now enable the efficient generation of cells expressing individual genetic variants or libraries of variants for robust functional studies. We discuss innovative assays that can evaluate dozens or hundreds of variants sequentially. For example, the electrophysiological properties of numerous cardiac ion channel variants in genes linked to inherited arrhythmias can be characterized using automated patch clamping. The mechanical properties of cardiomyocytes expressing candidate cardiomyopathy variants can be assessed using techniques such as atomic force microscopy, traction force microscopy, and impedance-based methods. Multiplexed assays of variant effect are an emerging family of techniques that use gene-specific or general assays, combined with next-generation sequencing, to characterize hundreds or thousands of pooled genetic variants. We examine the key advantages and limitations of each method and outline future goals for the field. Innovative in vitro studies of cardiac genetic variants will enhance our understanding of variant-disease relationships and improve diagnosis, screening, and treatment options for these disorders.

Key Words: arrhythmias, cardiac ■ cardiomyopathies ■ CRISPR-Cas systems ■ high-throughput screening assays ■ induced pluripotent stem cells ■ mechanics ■ patch-clamp techniques

Downloaded from http://ahajournals.org by on February 27, 2026

Cardiovascular diseases (CVDs) are responsible for approximately one-third of global deaths.[1] This group of disorders involves the heart, brain, and blood vessels, contributing to the burden of heart failure, sudden cardiac death, and stroke. While behavioral factors such as tobacco use, physical inactivity, and environmental factors such as pollution modify the risk of CVDs, many of these diseases have a strong genetic component. CVDs such as hypertrophic cardiomyopathy (HCM), dilated cardiomyopathy, and inherited arrhythmia syndromes have been attributed to disease-causing variants, primarily in genes encoding sarcomeric proteins and voltage-gated ion channels. The American College of Medical Genetics and Genomics variant classification framework categorizes variants as pathogenic, likely pathogenic, variant of uncertain significance (VUS), likely benign, or benign.[2] The discovery of a pathogenic variant in specific genes can often be clinically actionable, informing diagnosis, medical management, and family screening. Current guidelines recommend the reporting of incidentally detected pathogenic variants in 81 clinically actionable genes, including 41 genes associated with CVD.[3] In recent years, the cost of clinical genetic testing has decreased dramatically. However, many genetic testing results return VUS, which are not clinically actionable; this is known as the VUS problem.[2] High-throughput assays can provide functional evidence for the reclassification of VUS and improve the diagnosis and treatment of cardiac diseases. In addition, functional assays play a critical role in establishing and validating gene–disease relationships. The Clinical Genome Resource is spearheading systematic efforts to both confirm novel

Correspondence to: Kai-Chun Yang, MD, University of Washington, 850 Republican St, Brotman Bldg, Room 442, Seattle, WA 98109, Email kcyang@uw.edu; or Andrew M. Glazer, PhD, Vanderbilt University Medical Center, 2215 Garland Ave, 1235 Medical Research Bldg IV, Nashville, TN 27232, Email andrew.m.glazer@vumc.org
*R.E. Dolder III, C.E. Friedman, and A.M. Loiben contributed equally.
**K.-C. Yang and A.M. Glazer contributed equally.
For Sources of Funding and Disclosures, see page 131.
© 2026 American Heart Association, Inc.
Circulation: Genomic and Precision Medicine is available at www.ahajournals.org/journal/circgen

Nonstandard Abbreviations and Acronyms

AFM	atomic force microscopy
CRISPR-Cas9	clustered regularly interspaced short palindromic repeats/clustered regularly interspaced short palindromic repeat–associated 9
CVD	cardiovascular disease
EHT	engineered heart tissue
GFP	green fluorescent protein
gRNA	guide RNA
HCM	hypertrophic cardiomyopathy
hiPSC-CM	human induced pluripotent stem cell–derived cardiomyocyte
MAVE	multiplexed assay of variant effect
nCas9	Cas9 nickase
TUNEL	terminal deoxynucleotidyl transferase dUTP nick end labeling
VUS	variant of uncertain significance
WT	wild type
β-MHC	β-myosin heavy chain

associations and reassess genes previously implicated in Mendelian cardiac disorders.[4,5] While functional data are not used in isolation, they can serve as critical evidence in supporting these relationships.

We divide high-throughput variant studies into 2 main types: arrayed assays and pooled assays (Figure 1; Table 1). Arrayed assays represent a more traditional approach, where a series of variants are tested sequentially or in multiwell plates, where each well contains cells expressing a single variant.[37] Platforms such as automated patch-clamp devices, modified traction force microscopy instruments, and plate readers can accommodate experiments using 96- or 384-well plates; certain plate readers support up to 1536 wells. Although the throughput of these assays can be high, it is typically limited by the number of generated variants, the number of available wells, and the need for replicates. Quantitative data can be collected for various cellular properties, including ion influx, contractile force, or patch-clamp measurements, enabling nuanced mechanistic studies with direct functional comparisons between variants.[14,21]

Pooled assays, often referred to as multiplexed assays of variant effect (MAVEs), enable the simultaneous study of variant effects on molecular phenotypes such as protein function, gene expression, or mRNA splicing.[38] In an MAVE experiment, comprehensive libraries of variants are expressed in cells, subjected to multiplexed assays of properties such as fluorescence intensity or cell survival, and then deep sequenced to quantify variant-specific functional effects. Multiple MAVE assays for a single gene can provide key information about variant properties or identify mechanisms of variant pathogenicity. Deep mutational scanning is a subtype of MAVE focused on protein-coding variants. Standard methods include survival assays or fluorescence-based assays to separate cells based on their variant effects.[7] Pooled assays measure variant effects in identical conditions, reducing environmental variation. While pooled assays provide high-throughput data sets, array-based assays currently offer a wider range of functional assay types. Notably, certain assays are not yet compatible with pooled assays with current technologies. The choice between the 2 approaches depends on specific experimental goals.

Regardless of the assay type, a central challenge is efficiently generating cells that express genetic variants. In the first section, we review current strategies for introducing single or pools of genetic variants into cells. We then discuss a variety of arrayed assays used to study cardiac phenotypes. Last, we discuss pooled MAVE assays and their emerging role in characterizing hundreds to thousands of variants in cardiac disease genes.

GENERATION OF VARIANT-EXPRESSING CELLS

A key aspect of high-throughput variant assays is the efficient generation of cells expressing genetic variants. While transient transfections are efficient and fast, this approach is limited due to the inability to rigorously standardize expression levels.[39] Instead, creating cell lines that stably and consistently express different variants at the endogenous genomic locus or a landing pad site can be a more reliable approach. Figure 2 depicts commonly used methods to generate single-variant–expressing cells or pools of variant-expressing cells.

Classic CRISPR-Cas9 (clustered regularly interspaced short palindromic repeats/clustered regularly interspaced short palindromic repeat–associated 9) gene editing involves introducing a guide RNA (gRNA) with Cas9 that causes a double-strand break at a specific locus. If a repair template is provided, the homology-directed repair pathway repairs the DNA lesion and introduces a variant of interest.[40] Saturation genome editing uses CRISPR-Cas9 to generate a double-strand break in a gene of interest, followed by homology-directed repair from a pool of single-stranded oligodeoxynucleotide sequences containing a library of variants. This approach has been used in haploid cells, typically the Human Aneuploidy Project 1 (HAP1) cell line, to generate pools of cells expressing all possible variants at a locus.[41] Prime editing and base editing are 2 variants of CRISPR-Cas9 that can lead to enhanced genome editing efficiency and versatility.[42] Prime editing uses an nCas9 (Cas9 nickase) enzyme to make a single-stranded nick at a site directed by the prime editing gRNA. A reverse transcriptase enzyme fused to

Downloaded from http://ahajournals.org by on February 27, 2026



Figure 1. Overview of pooled and arrayed cardiac functional assays.
A, Pooled assays. A library of variants is transfected into pooled cells and subjected to multiplexed assays that determine function scores for each variant, which are displayed on a heatmap. These assays include cell survival assays, fluorescence assays, and splicing assays. **B**, Arrayed assays. In arrayed assays, for example, automated patch clamping and engineered heart tissues, variants are serially measured in separate wells.

nCas9 then uses the prime editing gRNA as a template for variant synthesis.[43] This approach can also be used to introduce many genetic variants simultaneously.[44,45] In recent work, a library of prime editing gRNAs was delivered to cells, and prime editing was used to generate thousands of variants in the epithelial growth factor receptor gene, which is implicated in cardiac remodeling and atrial fibrillation.[44,45] Base editing uses a catalytically dead Cas9 fused to a deaminase to chemically alter bases. Although no single base editing system can generate every possible modification, available tools can generate all 4 purine-to-purine and pyrimidine-to-pyrimidine transition modifications (C→T, A→G, G→A, and T→C).[46,47] Prime and base editing methods can reduce off- and on-target genomic changes by eliminating the need for double-strand breaks. An important limitation for all families of CRISPR gene-editing methods is the generation of undesired off-target edits that can confound studies of individual variants. However, in addition to off-target editing, base editing approaches can also introduce bystander changes by modifying unintended bases in an editing window.[48] While base editing systems are limited by the variants that they can generate, hundreds of variants can be made in the native genomic context. For example, recent studies have used base editing to screen over 400 variants in *MYBPC3* and 300 variants in *PARP1*.[49,50] A limitation of base and prime editing is a dependence on unique gRNA binding sites proximal to the desired site of mutation, whereas traditional CRISPR-Cas9 paired with homology-directed repair or serine integrase-based approaches (see below) can modify DNA more distantly.

An alternative to CRISPR-based editing is the use of serine integrases for genomic insertion of transgenes (Figure 2). Serine integrases are site-specific recombinases that mediate the recombination of short (≈50 bp) attachment site B and P (attB and attP) DNA sequences.[51] Plasmids containing an attB sequence can be integrated next to engineered attP sites in the genome, allowing for the insertion of large DNA sequences into a targeted location.[52,53] The attP site can be placed either at the native locus of a gene of interest or in a safe

Downloaded from http://ahajournals.org by on February 27, 2026

Table 1. Pooled and Arrayed Functional Assays to Interrogate Cardiac Variants

Assay method	Gene type	Genes	Functional alteration	Throughput
MAVE/pooled assays				
VAMP-seq	Any protein-coding gene	*MYH7*[6]	Protein abundance	High
Surface trafficking	Transmembrane proteins	*KCNE1*[7] and *KCNH2*[8]	Protein surface abundance	High
Survival	Protein with an activity that impacts cell viability	*MYH7*,[6] *LMNA*,[9] *SCN5A*,[10] and *KCNE1*[7]	Protein function	High
Splicing	Genes with a loss-of-function mechanism	*SCN5A*,[11] *LMNA*,[12] *MYBPC3*,[12] and *TTN*[13]	Splicing patterns	High
Arrayed assays				
Automated patch clamping	Voltage-gated ion channels	*SCN5A*,[14–17] *KCNQ1*,[16,18,19] *KCNH2*,[20] and *KCNE1*[18]	Ion current amplitude and activation/inactivation properties	Medium-high
Fluorescent plate reader	Ion channels	*KCNH2*,[21] *KCNQ2*,[22] and *TRPM4*[23]	Ion flux changes (eg, Ca^{2+} K^+, and Na^+) via fluorescence indicators	Medium-high
Fractional shortening	Sarcomeric and cytoskeletal genes	*MYH7*,[24] *TNNT2*,[25] *TTN*,[26] and *MYBPC3*[25]	Contraction/relaxation dynamics	Low-medium
Atomic force microscopy	Sarcomeric, cytoskeletal, and nuclear envelope	*LMNA*,[27,28] *DSP*,[29] and *DMPK*[30]	Cell stiffness, contractile force, and viscoelastic properties	Low-medium
Traction force microscopy	Sarcomeric genes	*MYH7*[31]	Force generation against the substrate	Low-medium
Engineered heart tissues	Sarcomeric and cytoskeletal genes	*MYH7*[32–34] and *TNNT2*[34]	Force generation based on post-deflection	Low-medium
Electrical impedance (MEA)	Sarcomeric and signaling genes	*MYH7*,[35] *MYBPC3*,[35] and *TTN*[36]	Contractile kinetics via impedance changes	Low-medium

Low: 1–10; medium: 10–100; and high: >100 variants. MAVE indicates multiplexed assay of variant effect; MEA, microelectrode array; and VAMP-seq, variant abundance by massively parallel sequencing.

harbor location engineered with flanking elements that facilitate expression and genetic selection.[54] One commonly used system is a genomic landing pad containing a tetracycline-inducible promoter and an attP site, as well as additional features such as a fluorescent marker or a caspase for genetic selection.[53] AttB-containing plasmids can be integrated into the landing pad, replacing expression of the landing pad genes with a gene of interest and a fluorescent marker from the plasmid. This system enables the identification and selection of on-target integrated versus nonintegrated cells. This system has been used to insert and express thousands of genetic variants in multiplexed analyses.[7,10,53]

Each of the methods described above can generate individual variants for single-variant or arrayed assays or a pool of variants for MAVE experiments. In their multiplexed forms, these methods involve the generation of pools of gRNAs (prime editing and base editing), pools of variant-containing DNA repair templates (saturation genome editing), or pools of variant cDNAs for genomic integration (CRISPR-Cas9 and serine integrases). Pools of bespoke DNAs can be readily synthesized or obtained commercially. Libraries of larger DNAs can also be obtained commercially or can be constructed through polymerase chain reaction using modified primers or by synthesizing short oligonucleotide pools that are subsequently subcloned into the full-length gene.[55,56] Recently developed methods, such as Scalable and Uniform Nicking Mutagenesis and Deep Indel Missense Programmable Library Engineering, promise to increase the efficiency and cost-effectiveness of variant library generation.[57,58]

ARRAYED ASSAYS

Automated Patch Clamping

Genes associated with inherited cardiac arrhythmias primarily encode voltage-gated ion channels.[59] These channels regulate the flux of sodium, potassium, and calcium ions across the cell membrane, generating action potentials that lead to coordinated cardiac contractions. Genetic variants that alter ion flow can disrupt this process, leading to arrhythmias such as long-QT syndrome or Brugada syndrome.[60] By understanding how a variant affects a channel's ability to regulate ion currents, researchers can better predict the pathogenicity of that variant.[61]

Patch clamping is the gold-standard method for characterizing ion channel function and can measure detailed channel properties. For voltage-gated channels, these properties can include peak current and the voltage dependence of channel activation and inactivation.[62] Patch-clamping data of variants with altered functional properties have been widely used to support the pathogenic classification of variants in arrhythmia-related genes.[63] Manual patch clamping is a technique that requires a highly skilled practitioner and is time-consuming.[64] Despite the valuable data provided by

Downloaded from http://ahajournals.org by on February 27, 2026



Figure 2. Commonly used techniques to introduce variants into cells.
A, CRISPR-Cas9 (clustered regularly interspaced short palindromic repeats/clustered regularly interspaced short palindromic repeat–associated 9). Traditional CRISPR systems can use a library of guide RNAs (gRNAs) to target different locations in the genome and a library of templates to generate various variants at each location. **B**, Prime editing systems have a single prime editing gRNA (pegRNA) that determines the location and alteration of the variant. Therefore, a library of pegRNAs can generate a diverse library of variants. **C**, Base editing systems will generate variants based on the modifying enzyme attached. A library of gRNAs can generate a variety of variants in different locations. **D**, Serine integrase recombination. A serine integration system inserts cDNA into a specific part of the genome via recombination. A library of plasmids can, therefore, generate a library of inserted sequences. RT indicates reverse transciptase.

Downloaded from http://ahajournals.org by on February 27, 2026

manual patch clamping, the time-consuming nature of the procedure has limited its use in high-throughput studies.[64,65]

Automated patch-clamping devices enable the simultaneous testing of many cells.[66] In this planar patch-clamp method, a single cell is suctioned into a small cavity in each well of a 96- or 384-well plate. The voltage across the cell membrane is manipulated to activate voltage-gated channels, and the resulting currents are measured.[66] This technique can efficiently characterize dozens or hundreds of variants while collecting data of similar quality to that obtained through manual patch clamping.[64] Automated patch clamping has been deployed to characterize over 1300 variants in cardiac ion channel genes, including *KCNQ1*, *KCNH2*, *KCNE1*, and *SCN5A* (Table 2). The high-throughput nature of automated patch clamping is exemplified by 2 recent studies, which tested 533 and 488 *KCNH2* variants.[69,71]

Fluorescent Plate Reader Assays of Ion Channel Variants

High-throughput screening using fluorescent plate readers offers a means of rapidly characterizing the function of ion channel variants.[73] This method can simultaneously record ion channel function across wells in a 96- or

Table 2. Automated Patch-Clamping Studies of Cardiac Ion Channel Variants

Study	Gene	# Variants	Summary
O'Neill et al (2025)[67]	*SCN5A*	252	Studied all variants found in a large Brugada Syndrome cohort
Vanoye et al (2025)[68]	*KCNE1*	95	Determined population carrier frequency of Jervell and Lange-Nielsen syndrome type 2
O'Neill et al (2024)[69]	*KCNH2*	533	Identified 331 variants with loss-of-function
Ma et al (2024)[15]	*SCN5A*	53	Validated assay with 25 benign and 24 pathogenic controls
Thomson et al (2024)[70]	*KCNH2*	78	Validated assay with 60 controls and evaluated 18 missense variants
Glazer et al (2022)[16]	*SCN5A*, *KCNQ1*, and *KCNH2*	50	Studied 50 variants found from population screening
Ng et al (2022)[71]	*KCNH2*	458	Studied every single nucleotide polymorphism in a hotspot exon
Jiang et al (2022)[72]	*KCNH2*	75	Calibrated assay with 31 controls before testing 44 undefined variants
O'Neill et al (2022)[17]	*SCN5A*	50	Tested dominant negative effects of 35 total and 15 partial loss-of-function variants
Ng et al (2020)[20]	*KCNH2*	30	Evaluated dominant negative effects of 13 pore-domain variants
Kuenze et al (2020)[18]	*KCNQ1* and *KCNE1*	16	Utilized modified cell lines to test dominant negative effects
Glazer et al (2020)[14]	*SCN5A*	83	Validated assay with 30 control variants and then tested 48 VUSs
Vanoye et al (2018)[19]	*KCNQ1*	78	Reclassified 23 of 35 VUSs

Studies with at least 10 variants are included. VUS indicates variant of uncertain significance.

384-well plate. The use of plate readers to measure ion channel activity relies on linking the channel's function to fluorescence changes within each well.[62] This is achieved using ion-sensitive fluorescent indicators.[74] The general method involves (1) distributing cells expressing ion channel variants or drug conditions into different wells of a plate, (2) adding ion-sensitive indicators, (3) inducing ion channel activation, and (4) observing fluorescence changes as a proxy for ion concentration.[75] Variants that alter activity compared with the wild type (WT) will cause altered dynamics of cytosolic ion concentrations and altered fluorescence.[75] These methods depend on the use of suitable fluorescent dyes, which are commercially available for all ions relevant to cardiac ion channels. This approach has been deployed for studies of all major ion types (eg, potassium,[22] calcium,[75] and sodium[23] channels), and extensive reviews comparing the different ionic indicators have been written for each ion.[76–78] In certain cases, ions other than the native ion can permeate a channel, such as thallium or rubidium ions passing through open potassium channels.[22] This property is utilized in thallium flux assays, which measure potassium channel activity by detecting thallium ion accumulation with a fluorescent dye.[62,79] A similar technique is applied in rubidium assays, where the passage of rubidium ions through potassium channels is quantified using spectrophotometry.[22]

RYR2 encodes a sarcoplasmic reticulum–localized calcium channel that plays a critical role in excitation-contraction coupling. Gain-of-function variants in *RYR2* are associated with catecholaminergic polymorphic ventricular tachycardia, whereas loss-of-function variants can cause the recently described calcium release deficiency syndrome.[80] *RYR2* function has been studied using fluorescence resonance energy transfer assays to analyze its interactions with calmodulin and FK506-binding protein 12.6 kDa (FKBP12.6).[81] In addition, *RYR2* activity has been investigated using R-CEPIA1er, a calcium sensor localized to the endoplasmic reticulum.[82] To date, these approaches have been used mainly for drug screening in plate reader formats. However, because differentiating between loss-of-function and gain-of-function *RYR2* variants is critical for understanding *RYR2*-related arrhythmias, we anticipate the growing use of high-throughput approaches to study *RYR2* variants.

Fluorescent plate reader measurements of ion channel function are more widely available than automated patch-clamping machines. However, a limitation of this method is that it typically does not determine precise ion channel properties such as activation or inactivation voltages.[62] While most high-throughput studies of this type are utilized for drug screening, these methods can be easily adapted for testing the effects of different variants; instead of each well containing a different channel modulator, each well can include a different variant. In summary, fluorescent plate reader assays offer a versatile and accessible method for assessing ion channel function, with applications ranging from drug screening to variant effect testing.

CARDIOMYOCYTE MECHANICS

Cardiomyocyte contractile dysfunction is closely associated with cardiac disease. Mechanical properties of cardiomyocytes altered in cardiomyopathy include active and passive force, contraction and relaxation time, and cell adhesion.[83] In this section, we describe a series of methods to assess contractile function, which can aid in characterizing variant pathogenicity and phenotype.[84] These methods can be adapted for single cells, cardiomyocyte monolayers, cardiac embryoid bodies, and engineered heart tissues (EHTs), with varying throughputs

Downloaded from http://ahajournals.org by on February 27, 2026

Table 3. Functional Mechanical Readouts

Technique	Throughput	Single cells	Monolayers	Tissues	# Variants reported	References
AFM	High	Yes	Yes	No	3	28–30
TFM	High	Yes	Yes	No	26	31,85,86
Sarcomere fractional shortening	High	Yes	Yes	No	3	87
Microposts	Medium	Yes	No	No	2	88,89
Electrical impedance	Medium	No	Yes	No	6	35
Isolated myofibrils	Low	Yes	No	No	2	90
EHTs	Low	No	No	Yes	5	91

AFM indicates atomic force microscopy; EHT, engineered heart tissue; and TFM, traction force microscopy.

and applications (Table 3; Figure 3). Typically, single-cell and monolayer methods offer higher throughput compared with tissue-based methods due to the larger cell numbers and hands-on time required to generate EHTs. However, recent advances in tissue miniaturization have increased the throughput of tissue mechanics testing, enabling their use for variant effect assessments.[92]

Fractional Shortening

As a cardiomyocyte contracts, both cell length and sarcomere length cyclically decrease. Fractional shortening measures the change in cell or sarcomere length normalized to the resting length. Both cell and sarcomere length can be detected with brightfield microscopy, enabling simple measurement across hundreds of cells.[93] Sarcomeres can also be tracked by incorporating fluorescent proteins using gene editing, small molecules, or viral transduction.[60,94] Cell membranes and sarcomeres can be tracked across video frames using tools such as SotaTool,[95] SarcTrack,[96] and SarcGraph,[97] or in proprietary software from IonOptix.[98] Notably, fractional shortening is an indirect measure of contractile force and can be a poor predictor of force in certain disease contexts.[87]

Atomic Force Microscopy

Atomic force microscopy (AFM) is a microscopy technique that uses a nanometer-scale metal tip attached to a flexible cantilever. When the tip makes contact with a cardiomyocyte, the cantilever is deflected by a certain distance. This deflection corresponds to piconewton-scale force measurements, determined by the cantilever's spring constant. AFM can, thus, capture the mechanical



Figure 3. Methods for mechanical assessment of cardiomyocytes.
A, Fractional shortening measures the relative change in cell length or sarcomere length in contracting cardiomyocytes. **B**, Atomic force microscopy uses a flexible cantilever in contact with a contracting cardiomyocyte to estimate local contractile force based on the deflection of the cantilever and the cantilever's stiffness. **C**, Traction force microscopy uses a hydrogel with embedded fluorescent beads. As cardiomyocytes contract, force can be estimated based on the displacement of beads beneath the cardiomyocyte as a function of hydrogel stiffness. **D**, Cardiomyocytes can be seeded on flexible microposts of defined stiffness, enabling the measurement of force based on post-deflection. **E**, Engineered heart tissues use a 3-dimensional tissue seeded across one flexible and one rigid post. Tissue contraction deflects the flexible post, permitting force measurement. **F**, In impedance assays, cardiomyocytes are seeded on top of microelectrode arrays. As cardiomyocytes contract, the electrical resistance decreases, a proxy for contractile function.

Downloaded from http://ahajournals.org by on February 27, 2026

properties of cardiomyocytes, such as cell stiffness and stress relaxation. Beat rate, contractile force, and contractile kinetics can be measured by locking the tip at the cell surface and measuring its deflection as the cell contracts. AFM was first applied to rat atrial myocytes in 1995[99] and to human induced pluripotent stem cell–derived cardiomyocytes (hiPSC-CMs) in 2012.[100] AFM is well-suited for studying contractile dynamics of single cells with high resolution.[90,101,102] The technique has been used to investigate the effects of variants in cardiomyopathy-related genes such as *LMNA*, *DSP*, and *DMPK* on cardiomyocyte mechanics.[28–30] Approximately 100 force tip indentations can be measured per hour, enabling rapid assessment of single variants in an arrayed format.[103]

Traction Force Microscopy

Traction force microscopy is a fluorescence microscopy technique used to measure the contractile mechanics of cardiomyocytes. In traction force microscopy, fluorescent spheres ranging from 100 to 1000 nm in diameter are embedded in a polymer hydrogel with adjustable stiffness. An attachment protein matrix is then patterned on the hydrogel using polydimethylsiloxane stamps or photolithography. Depending on the pattern, cells can attach to the coated surface as single cells, a monolayer, or a tissue.[104,105] When cardiomyocytes contract, they exert force on the underlying hydrogel, causing displacement of the embedded fluorescent beads. By capturing videos of these displacements, researchers can quantify traction force and contractile kinetics as a function of the hydrogel's Young modulus. This approach has been used to interrogate the effect that cardiomyopathy-associated *MYH7* variants have on hiPSC-CMs mechanics.[31,85,86]

Microposts

In the micropost technique, cardiomyocytes are cultured on arrays of elastomeric posts coated with extracellular matrix.[88,89,106–108] These posts deflect under cell tension, which can be converted to force measurement based on the physics of the bending posts. This principle has been adapted to measure the mechanical properties of cardiomyocytes.[88,89] However, a significant limitation is the inability to electrically stimulate the cardiomyocytes when they are cultured on the microposts.[88]

Engineered Heart Tissues

In vitro tissue approaches attempt to mimic the native cardiac architecture to capture macroscopic properties that are absent in single-cell or monolayer contexts. The most established method in this category is the EHT.[91] The basic principle involves coculturing cardiomyocytes with stromal cells in a slurry, which is then seeded onto hydrogels or scaffolds, such as flexible posts, to form tissue constructs. As the tissues contract, the attached posts deform, enabling measurement of contractile force. Contractile function of large EHTs can also be assessed by directly connecting the tissues to a force transducer. Early strategies using EHTs relied on tissues comprised of 10^5 to 10^6 cardiomyocytes per tissue, significantly limiting the scale of throughput across variants.[109,110] Recently, miniaturized or micro-EHTs have been developed, which mimic the contractile properties of larger EHTs while requiring as few as 16 000 cardiomyocytes per tissue.[92,111] These micro-EHTs offer scalability across variants, as they can be cast into individual wells of a 96-well plate, enabling parallel assessment of many tissues at once. The Curi Bio Mantarray system uses EHTs attached to posts with embedded magnets for label-free, electromagnetic measurement of contractile kinetics in response to electrical stimulation, enabling simultaneous measurement of up to 24 EHTs.[112]

Electrical Impedance With Microelectrode Array

Label-free electrical impedance is a well-established method for sensing cell behavior. Microelectrode arrays embedded in tissue culture plates can detect changes in impedance and correlate these changes to cardiomyocyte mechanical output, as described by a power law relationship.[113] Impedance measurements have been used with hiPSC-CMs to assess cardiotoxicity in response to putative therapeutics.[114] Furthermore, impedance measurements have recently been used to assess the efficacy of myosin ATPase inhibitors in restoring normal contractile function in hiPSC-CMs with HCM-causing *MYH7*, *MYBPC3*, and *TNNI3* variants.[35] This approach shows promise for high-throughput mechanical assessment of variant-expressing cells in a 96-well plate format.

POOLED ASSAYS

In contrast to arrayed assays where each variant is studied individually, MAVEs involve the pooled creation and joint characterization of a library of variants. The general approach involves (1) saturation mutagenesis of a target gene or region, (2) expression of pools of mutants in an appropriate cell model, (3) en masse multiplexed selection for cellular phenotypes that reflect variant effects on relevant protein function(s), (4) next-generation sequencing to compare variant abundance in preselection and postselection pools, and (5) data analysis and display. The most critical step of an MAVE is to develop an assay that faithfully models the disease process of interest. Several assay types are described below.

Flow Cytometry–Based Assays

Flow cytometry combined with cell sorting enables the rapid and high-throughput separation of cells into bins spanning

Downloaded from http://ahajournals.org by on February 27, 2026

a broad range of phenotypes. Each bin, characterized by a distinct phenotypic range, undergoes deep sequencing to quantify the frequency of each variant within that bin. By analyzing variant frequencies across all bins, researchers can reconstruct an estimated phenotypic profile for each variant, allowing for the detection of variant profiles that significantly diverge from controls. Flow cytometry–based assays for multiplexed functional assays rely on measuring single-cell phenotypes. These include cell size, expression of a fluorescent reporter, or an epitope prelabeled with fluorophore-conjugated antibodies. Figure 4 illustrates a general workflow for 3 different fluorescence-based flow cytometry MAVEs. Examples include (1) variant abundance by massively parallel sequencing, where the intensity of GFP (green fluorescent protein) fused to a protein of interest serves as a readout of protein expression,[115] (2) trafficking assays using fluorescent antibody staining to assess surface protein abundance,[116] and (3) assays measuring protein function that use fluorescent markers, where fluorescence changes occur as a downstream response to activity of the protein of interest.[117]

Arguably, the most fundamental way to assess the effect of a genetic variant is to determine whether it alters the abundance of its associated protein product. Pathogenic variants in many genes can reduce protein abundance by destabilizing the mRNA or protein produced, triggering nonsense-mediated decay of the mRNA, or through other mechanisms.[118] Matreyek et al[115] utilized this principle to develop a flow cytometry–based multiplexed assay technique called variant abundance by massively parallel sequencing. This method can detect deleterious variants that alter protein abundance.

Variant abundance by massively parallel sequencing involves tagging the protein of interest with a fluorescent marker, allowing a cell's fluorescence intensity, measured via flow cytometry, to serve as an indicator of the protein's abundance. A pooled library of tagged protein variants can be binned by fluorescence intensity using flow sorting, and then, each bin can be sequenced to quantify the variant's impact on protein abundance. This method enables high-throughput assessment of protein levels using flow cytometry for fluorescence-based cell sorting, followed by next-generation sequencing and abundance calculations. Friedman et al[6] applied variant abundance by massively parallel sequencing to evaluate a library of 113 heterozygous *MYH7* variants



Figure 4. Fluorescence-based multiplexed assays of variant effect.
A, Generate a pool of cells expressing different genetic variants. **B**, An assay is performed where cellular fluorescence is linked to protein function. **C**, Flow sorting is used to separate cells based on their fluorescence intensity. **D**, Each bin is deep sequenced to generate functional scores for each variant, which are presented as a heatmap. GFP indicates green fluorescent protein; POI, protein of interest; and VAMP-seq, variant abundance by massively parallel sequencing.

Downloaded from http://ahajournals.org by on February 27, 2026

in hiPSC-CMs, by measuring the abundance of β-MHC (β-myosin heavy chain), the protein encoded by *MYH7*. This approach facilitated the identification of pathogenic variants associated with low β-MHC abundance. Similarly, Yamamoto et al[49] used direct antibody-labeling of a library of *MYBPC3* variants in hiPSC-CMs, followed by flow sorting and sequencing, to identify *MYBPC3* variants disrupting protein abundance.

Additional CVD phenotypes, such as enlarged cardiomyocyte size or increased expression of the heart failure marker B-type natriuretic peptide (BNP), can be detected using flow cytometry–based assays. For instance, Pettinato et al[119] used CRISPR/Cas9 to knock in a tandem dimer Tomato (tdTomato) fluorescent reporter into the endogenous *NPPB* locus of a *TNNT2* knockout hiPSC line. They then transduced hiPSC-CMs derived from this line with 51 distinct *TNNT2* variant lentiviruses and used flow cytometry to measure tdTomato as an indicator of *NPPB* expression. *TNNT2* variants known to cause HCM exhibited significantly increased *NPPB* expression, whereas variants causing dilated cardiomyopathy showed decreased expression. In the recent *MYBPC3 MAVE* study, the authors also used a BNP reporter to assess cardiomyocyte hypertrophic signaling across the pool of variants.[49]

To enhance flow cytometry–based functional assessments of CVD-associated variants, additional pathogenic markers that can be fluorescently labeled, either through immunostaining or genetically encoded reporters, should be identified. These markers would enable evaluations in an arrayed format or, if the marker's dynamic range is sufficiently broad, in a pooled format. Alternatively, simple morphological parameters such as cell size could be assessed by flow cytometry, eliminating the need for fluorescent labeling. Indeed, several groups have used flow cytometric measurement of forward scatter as a proxy for cell area to evaluate variants causing HCM and dilated cardiomyopathy.[33,119] However, forward scatter assessments have only revealed minor (<10%) differences in relative cell size between pathogenic and WT controls. This observation is corroborated by absolute, flow cytometry–based volumetric assessments of hiPSC-CMs, which show a modest increase in cell volume (12%–51%).[110] Collectively, these findings suggest that enhancing hypertrophy phenotypes through various growth condition modifications would be necessary to expand the dynamic range for pooled forward scatter assays.

Trafficking Assays

Of the cardiac ion channels, *KCNH2* variants have been the most extensively studied using solubility and glycosylation assays, which serve as proxies for evaluating protein abundance and surface expression.[120,121] These studies established that most loss-of-function *KCNH2* variants impair protein stability and localization. They further led to a general appreciation that variants in transmembrane proteins that disrupt overall protein folding, stability, and trafficking to the membrane are usually the major mechanism of loss-of-function.[122] Building on this work, MAVE-compatible fluorescent assays of surface abundance were developed (Figure 4).[8] This approach uses fluorophore-conjugated antibodies targeting extracellular epitopes (native or epitope tagged) to label only proteins at the cell surface, leaving intracellular pools unlabeled. Each variant's surface abundance is quantified by fluorescence-activated cell sorter-sorting cells into fluorescence bins and sequencing the barcodes or variants from each bin.[123] Surface abundance scores are assigned to each variant based on its frequency in low-, WT-like, or high-fluorescence bins. This assay has been deployed to generate surface abundance scores for thousands of variants across cardiac ion channel genes, *KCNE1*,[7] *KCNJ2*,[124] and *KCNH2*.[69] Notably, O'Neill et al[69] performed an MAVE analysis of surface expression of over 17 000 *KCNH2* variants; this data set was also correlated with a large automated patch-clamp data set and clinical arrhythmia outcomes. As an alternative method for measuring cell surface expression, researchers assessed ion channel trafficking using a pH-sensitive fluorescent tag, pHluorin, which distinguishes between internal and plasma membrane–localized channels based on fluorescence changes at different pH levels. Lowering the extracellular pH from 7.4 to 6.0 caused a significant reduction in fluorescence for variant proteins located at the cell surface, while internal variant proteins remained unaffected.[125] We note that it is possible for a variant to be correctly trafficked to the surface yet still hinder function. Although a surface abundance assay alone may not identify this class of variants, pairing this assay with a functional assay can allow researchers to gain insights into the mechanisms underlying a variant's pathogenicity.[7]

Key considerations for this assay are that each cell should only express a single variant and that the overall expression of the ion channel must be relatively consistent. This can be achieved using specialized cell lines that allow for controlled cDNA integration or appropriate CRISPR systems.[43,53] Alternatively, coexpressing a fluorescent protein with the ion channel gene can provide a means to normalize trafficking scores to transgene expression.

Survival Assays

Survival assays that compare variant frequencies within a pooled variant library before and after a selection pressure are perhaps the most accessible high-throughput phenotyping strategy. Many cardiac diseases, such as dilated cardiomyopathy, inherited arrhythmia syndromes, and HCM, feature reduced cardiomyocyte survival relative to WT cardiomyocytes.[126] Pathogenic variants can

Downloaded from http://ahajournals.org by on February 27, 2026

exert damaging effects through mechanisms beyond cell death; however, cells that cannot survive cannot function.

Pathogenic variants in numerous CVD-related genes have been shown to cause reduced cardiomyocyte survival, priming these genes for survival-based MAVEs. For example, heterozygous premature truncating variants in the *DSP*, which encodes an intercalated disk protein, are known to cause a subtype of arrhythmogenic cardiomyopathy referred to as desmoplakin cardiomyopathy, often characterized by cell death.[127] Indeed, cardiomyocyte-specific deletion of *Dsp* in mice causes a robust PANoptosis phenotype (apoptosis, necroptosis, and pyroptosis).[127,128] Interestingly, the location of premature truncating variants in the *DSP* gene plays a role in the development of ventricular arrhythmias,[129] suggesting that other *DSP* phenotypes, such as reduced cell survival, could also be domain-dependent and benefit from high-throughput survival-based assessments in the future. In addition, heterozygous missense variants in the nuclear inner membrane gene *LMNA* cause a highly penetrant dilated cardiomyopathy subtype, with associated arrhythmias. This syndrome is often referred to as laminopathy and also features cell death.[130] For example, cardiomyocyte-specific expression of the p.Asp300Asn pathogenic variant of *LMNA* in mice causes a robust increase in TUNEL (terminal deoxynucleotidyl transferase dUTP nick end labeling)-positive cells and upregulation of *Tp53* signaling cascades,[130] suggesting that a high-throughput survival-based assay could help to map variant effect across *LMNA*. A comprehensive study of 178 *LMNA* missense variants utilized aggregation and solubility assays, with further validation in iPSC-CMs. Notably, the study revealed that pathogenic variants often lead to lamin A aggregation in the nucleus, indicating that these diseases are predominantly driven by protein misfolding.[9]

Pathogenic missense variants in the sarcomeric thick filament gene, *MYH7*, are a leading cause of HCM. Previous reports have shown that single pathogenic *MYH7* variants (p.Arg723Cys and p.Glu848Gly) can cause reduced survival in hiPSC-CMs.[33,131] Friedman et al[6] leveraged this observation to establish a multiplexed cardiomyocyte survival assay to functionally assess heterozygous *MYH7* missense variants in hiPSC-CMs. This assay involved performing next-generation sequencing of the *MYH7* variant library in hiPSCs and hiPSC-CMs. Frequencies of each variant were then calculated and compared between hiPSC-CMs and hiPSCs to discover which variants became prevalent after 47 days of culture in vitro. This analysis revealed that all 3 pathogenic variants, p.Glu848Gly, p.Met852Thr, and p.Met852Lys, had lower survival than 3 synonymous variants, p.Ser866=, p.Ala850=, and p.Glu848=(assumed to be functionally normal); a similar analysis of day 25 hiPSC-CMs revealed a similar outcome.[6] In addition, 2 pathogenic variants were lost from the day 47 library hiPSC-CMs (p.Lys865Arg and p.Met852Arg), indicating more severe cases of this survival phenotype.[6] This study serves as proof of concept that multiplexed survival assays can be leveraged for accurate functional annotation of additional *MYH7* (and other genes) variants in hiPSC-CMs in the future.

Survival assays have been deployed in ion channel MAVEs to obtain functional scores in Human Embryonic Kidney (HEK293T) cells for a 36-nucleotide region in *SCN5A* and nearly every variant in *KCNE1*.[7,10] Glazer et al[10] modified a previously developed assay that exposes cells to the sodium channel agonists, veratridine and brevetoxin, and the Na^+/K^+ exchanger inhibitor ouabain, to assess the function of a library of *SCN5A* variants; loss-of-function *SCN5A* variants resisted sodium influx and sensitivity to sodium ion overload, leading to increased (or unchanged) survival, whereas WT-like *SCN5A* variants showed reduced viability relative to pretreatment. A similar loss-of-function assay developed by Muhammad et al[7] tested the hypothesis that long-term (20-day) expression of WT *KCNE1* (in the context of a *KCNQ1* gain-of-function variant p.Ser140Gly) would decrease HEK293T viability relative to preexposure due to increased potassium ion efflux. Application of this assay to a library of *KCNE1* variants revealed 401 partial- and 173 loss-of-function variants with unchanged survival.[7] Other ion channel genes could likely be studied with similar strategies.

Splicing Assays

Up to 10% of pathogenic variants in Mendelian diseases disrupt RNA splicing, highlighting the importance of studying this type of variant.[132] However, assays that measure variant effects often use plasmids or DNA sequences containing only the coding sequences of a gene, which do not undergo RNA splicing, thereby preventing the identification of variants that alter RNA splicing. One approach to studying splice-disrupting variants is the use of minigene assays, in which DNA containing an exon or exons of interest, along with surrounding intronic sequences, is transfected into cells.[133] Normal or abnormal splicing can then be measured by RNA sequencing or reverse transcription polymerase chain reaction and gel electrophoresis.[12,134,135]

Recent efforts have scaled up this type of minigene assay to allow for multiplexed studies of pooled variants. Variant exon sequencing is a method in which a barcoded minigene system containing a library of variants is transfected into cells and given time for RNA processing to occur.[136] RNA is then extracted from the cells, reverse-transcribed into cDNA, and subjected to deep sequencing to determine the impact of each variant on splicing. In the original study, this system measured 2059 variants spanning 110 exons.[136] ParSE-seq is a similar process that utilizes a minigene system with barcoded RNA to measure variant effects via RNA sequencing

Downloaded from http://ahajournals.org by on February 27, 2026

after transfection.[134] This method was used to study 224 *SCN5A* variants in hiPSC-CMs. A similar pooled minigene method was used to measure 196 candidate splice-altering variants in *LMNA* and *MYBPC3*, which are associated with dilated cardiomyopathy and HCM, as well as *TTN* truncating variants that may contribute to dilated cardiomyopathy.[12,13] The scalability of these approaches has enabled the comprehensive testing of all possible splicing variants in targeted exons of cardiac genes, such as *POU1F1*, *MYBPC3*, and *FAS*.[49,137,138] An alternate minigene system relies on the introns and exon of interest being flanked by incomplete sequences of GFP. If the exon is skipped during splicing, cells will transcribe and translate GFP, allowing them to fluoresce. Fluorescent cells can then be isolated via flow sorting and subjected to deep sequencing. This method was used to measure splicing in more than 25 000 pooled variants.[139]

Evidence Strength Calculations

If functional assays are to be used for clinical variant classification, current ACMG guidelines recommend that assays be calibrated using a large set of known benign and pathogenic variants.[140] The assay's evidence strength is determined by the number of control variants and the assay's ability to distinguish benign from pathogenic variants. A Bayesian calculation of the odds of pathogenicity (termed OddsPath) can then be converted into ACMG evidence strengths (eg, supporting, moderate, or strong) for variant classification.[140,141] To improve OddsPath estimates for functional assays, we recommend incorporating as many benign and pathogenic controls as is feasible. A key advantage of MAVEs is that they generate functional data for large numbers of control variants, enabling the possibility of stronger OddsPath evidence codes.

Assays of the Future

This review has highlighted current methods for assessing variant effects in large classes of cardiac disorders, including inherited arrhythmia syndromes and HCM. We now outline future directions for high-throughput assays to enhance our understanding of genetic variation in CVDs.

Broader Contexts for Arrhythmia Assays

As outlined above, arrhythmia-causing variants in voltage-gated ion channels have been effectively studied using automated patch-clamping and MAVE experiments. However, much of this work has relied on heterologous expression systems, such as HEK293 cells with exogenous expression of a single cDNA isoform of each gene. While these approaches have been highly successful in identifying channel-altering variants, an important question remains: to what extent do accurate assessments of variant effects require a disease-relevant cellular context and endogenous gene expression? Future studies focusing on arrhythmia-related genes in more native contexts, such as gene-edited iPSC-CMs, and follow-up investigations of variants with discordant results between functional data and patient phenotypes, could help address this gap. Multiple gene-editing approaches in iPSC-CMs have already been applied to study dozens to hundreds of variants in genes such as *MYH7*,[6,86] *MYBPC3*,[49] and even *KCNH2*,[54] demonstrating the potential to expand functional models for arrhythmia-associated genes. In addition, studies in heterologous systems could be extended to a broader set of variant contexts, incorporating features such as coexpression of additional binding partners, heterozygous expression, common polymorphisms, splice isoforms, or response to drug exposures.

Scaling Up Mechanical Assays

Conventional wisdom holds that cardiomyopathy genes, which intricately alter contractile force dynamics, are best studied in cardiomyocytes with functional sarcomeres. As we outlined above, a broad set of mechanical assays can be deployed to assess variant effects and elucidate disease mechanisms. However, most studies utilizing these assays have been conducted in arrayed formats, which are often limited in throughput. Therefore, a critical future direction is to increase the throughput of arrayed assays and to develop innovative, MAVE-compatible assays for cardiomyopathy genes that capture key disease processes. In addition, efforts are underway to enhance the efficiency and scalability of gene editing in iPSC-derived cardiomyocytes. Finally, it may be possible to investigate some functions of cardiomyopathy-related genes in non-cardiomyocyte contexts, enabling the use of simpler heterologous cellular models.

PERSPECTIVES AND CONCLUSIONS

High-throughput methods for functional assessment have transformed the study of genetic variants implicated in cardiac diseases, enabling unprecedented insights into variant-disease relationships and the molecular mechanisms of conditions such as inherited arrhythmias and cardiomyopathies. While scalable techniques such as MAVEs and automated patch clamping have addressed critical challenges such as the classification of VUS, the field remains constrained by the need for disease-relevant models such as hiPSC-derived cardiomyocytes and pooled assays for physiologically relevant phenotypes. Expanding the range of phenotypes studied and linking these functional insights to clinically actionable outcomes will be essential for improving diagnostic accuracy and therapeutic decision-making. Despite these challenges, continued innovation promises to further advance variant classification, refine precision medicine approaches, and ultimately improve patient care in cardiovascular genetics.

Downloaded from http://ahajournals.org by on February 27, 2026

ARTICLE INFORMATION

Affiliations

Division of Genetic Medicine and Clinical Pharmacology, Department of Medicine, Vanderbilt Center for Arrhythmia Research and Therapeutics, Vanderbilt University Medical Center, Nashville, TN (R.E.D., A.M.G.). Department of Medicine/Cardiology, Center for Cardiovascular Biology, Institute for Stem Cell and Regenerative Medicine, University of Washington School of Medicine, University of Washington, Seattle (C.E.F., A.M.L., K.-C.Y.). Cardiology Section/Hospital Specialty Medicine Service Line, VA Puget Sound Health Care System, Seattle, WA (K.-C.Y.).

Acknowledgments

The content is solely the responsibility of the authors and does not necessarily represent the official views of the Department of Veterans Affairs or the US Government.

Sources of Funding

The authors acknowledge funding from the National Institutes of Health grants R01HL171174 (to Dr Yang), F32HL164108 (to Dr Loiben), K99HL177347 (to Dr Friedman), R00HG010904 (to Dr Glazer), R35GM150465 (to Dr Glazer), and R01HL164675 (to Dr Glazer); the American Heart Association grant 25PRE1360565 (to R.E. Dolder III); and the Department of Veterans Affairs grant 1I01 BX006428 (to Dr Yang).

Disclosures

None.

REFERENCES

1. Lindstrom M, DeCleene N, Dorsey H, Fuster V, Johnson CO, LeGrand KE, Mensah GA, Razo C, Stark B, Varieur Turco J, et al. Global burden of cardiovascular diseases and risks collaboration, 1990-2021. *J Am Coll Cardiol*. 2022;80:2372–2425. doi: 10.1016/j.jacc.2022.11.001
2. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M, Lyon E, Spector E, et al; ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17:405–424. doi: 10.1038/gim.2015.30
3. Miller DT, Lee K, Abul-Husn NS, Amendola LM, Brothers K, Chung WK, Gollob MH, Gordon AS, Harrison SM, Hershberger RE, et al; ACMG Secondary Findings Working Group. Electronic address: documents@acmg.net. ACMG SF v3.2 list for reporting of secondary findings in clinical exome and genome sequencing: a policy statement of the American College of Medical Genetics and Genomics (ACMG). *Genet Med*. 2023;25:100866. doi: 10.1016/j.gim.2023.100866
4. Hosseini SM, Kim R, Udupa S, Costain G, Jobling R, Liston E, Jamal SM, Szybowska M, Morel CF, Bowdin S, et al; National Institutes of Health Clinical Genome Resource Consortium. Reappraisal of reported genes for sudden arrhythmic death: Evidence-based evaluation of gene validity for Brugada syndrome. *Circulation*. 2018;138:1195–1205. doi: 10.1161/CIRCULATIONAHA.118.035070
5. Hespe S, Waddell A, Asatryan B, Owens E, Thaxton C, Adduru M-L, Anderson K, Brown EE, Hoffman-Andrews L, Jordan E, et al. Genes associated with hypertrophic cardiomyopathy: a reappraisal by the ClinGen hereditary cardiovascular disease gene curation expert panel. *J Am Coll Cardiol*. 2025;85:727–740. doi: 10.1016/j.jacc.2024.12.010
6. Friedman CE, Fayer S, Pendyala S, Chien W-M, Loiben A, Tran L, Chao LS, McKinstry A, Ahmed D, Farris SD, et al. Multiplexed functional assessments of MYH7 variants in human cardiomyocytes. *Circ Genom Precis Med*. 2024;17:e004377. doi: 10.1161/CIRCGEN.123.004377
7. Muhammad A, Calandranis ME, Li B, Yang T, Blackwell DJ, Harvey ML, Smith JE, Daniel ZA, Chew AE, Capra JA, et al. High-throughput functional mapping of variants in an arrhythmia gene, KCNE1, reveals novel biology. *Genome Med*. 2024;16:73. doi: 10.1186/s13073-024-01340-5
8. Kozek KA, Glazer AM, Ng C-A, Blackwell D, Egly CL, Vanags LR, Blair M, Mitchell D, Matreyek KA, Fowler DM, et al. High-throughput discovery of trafficking-deficient variants in the cardiac potassium channel KV11.1. *Heart Rhythm*. 2020;17:2180–2189. doi: 10.1016/j.hrthm.2020.05.041
9. Anderson CL, Langer ER, Routes TC, McWilliams SF, Bereslavskyy I, Kamp TJ, Eckhardt LL. Most myopathic lamin variants aggregate: a functional genomics approach for assessing variants of uncertain significance. *NPJ Genom Med*. 2021;6:103. doi: 10.1038/s41525-021-00265-x
10. Glazer AM, Kroncke BM, Matreyek KA, Yang T, Wada Y, Shields T, Salem J-E, Fowler DM, Roden DM. Deep mutational scan of an SCN5A voltage sensor. *Circ Genom Precis Med*. 2020;13:e002786. doi: 10.1161/CIRCGEN.119.002786
11. O'Neill MJ, Yang T, Laudeman J, Calandranis M, Solus J, Roden DM, Glazer AM. ParSE-seq: a calibrated multiplexed assay to facilitate the clinical classification of putative splice-altering variants. *Nat Commun*. 2023;15:8320. doi: 10.1101/2023.09.04.23295019
12. Ito K, Patel PN, Gorham JM, McDonough B, DePalma SR, Adler EE, Lam L, MacRae CA, Mohiuddin SM, Fatkin D, et al. Identification of pathogenic gene mutations in LMNA and MYBPC3 that alter RNA splicing. *Proc Natl Acad Sci USA*. 2017;114:7689–7694. doi: 10.1073/pnas.1707741114
13. Patel PN, Ito K, Willcox JAL, Haghighi A, Jang MY, Gorham JM, DePalma SR, Lam L, McDonough B, Johnson R, et al. Contribution of noncanonical splice variants to TTN truncating variant cardiomyopathy. *Circ Genom Precis Med*. 2021;14:e003389. doi: 10.1161/CIRCGEN.121.003389
14. Glazer AM, Wada Y, Li B, Muhammad A, Kalash OR, O'Neill MJ, Shields T, Hall L, Short L, Blair MA, et al. High-throughput reclassification of SCN5A variants. *Am J Hum Genet*. 2020;107:111–123. doi: 10.1016/j.ajhg.2020.05.015
15. Ma JG, O'Neill MJ, Richardson E, Thomson KL, Ingles J, Muhammad A, Solus JF, Davogustto G, Anderson KC, Shoemaker MB, et al. Multisite validation of a functional assay to adjudicate SCN5A Brugada syndrome-associated variants. *Circ Genom Precis Med*. 2024;17:e004569. doi: 10.1161/CIRCGEN.124.004569
16. Glazer AM, Davogustto G, Shaffer CM, Vanoye CG, Desai RR, Farber-Eger EH, Dikilitas O, Shang N, Pacheco JA, Yang T, et al; eMERGE Network. Arrhythmia variant associations and reclassifications in the eMERGE-III sequencing study. *Circulation*. 2022;145:877–891. doi: 10.1161/CIRCULATIONAHA.121.055562
17. O'Neill MJ, Muhammad A, Li B, Wada Y, Hall L, Solus JF, Short L, Roden DM, Glazer AM. Dominant negative effects of SCN5A missense variants. *Genet Med*. 2022;24:1238–1248. doi: 10.1016/j.gim.2022.02.010
18. Kuenze G, Vanoye CG, Desai RR, Adusumilli S, Brewer KR, Woods H, McDonald EF, Sanders CR, George AL Jr, Meiler J. Allosteric mechanism for KCNE1 modulation of KCNQ1 potassium channel activation. *Elife*. 2020;9:e57680. doi: 10.7554/eLife.57680
19. Vanoye CG, Desai RR, Fabre KL, Gallagher SL, Potet F, DeKeyser J-M, Macaya D, Meiler J, Sanders CR, George AL Jr. High-throughput functional evaluation of KCNQ1 decrypts variants of unknown significance. *Circ Genom Precis Med*. 2018;11:e002345. doi: 10.1161/CIRCGEN.118.002345
20. Ng C-A, Perry MD, Liang W, Smith NJ, Foo B, Shrier A, Lukacs GL, Hill AP, Vandenberg JI. High-throughput phenotyping of heteromeric human ether-à-go-go-related gene potassium channel variants can discriminate pathogenic from rare benign variants. *Heart Rhythm*. 2020;17:492–500. doi: 10.1016/j.hrthm.2019.09.020
21. Beacham DW, Blackmer T, O' Grady M, Hanson GT. Cell-based potassium ion channel screening using the FluxOR assay. *J Biomol Screen*. 2010;15:441–446. doi: 10.1177/1087057109359807
22. Scott CW, Wilkins DE, Trivedi S, Crankshaw DJ. A medium-throughput functional assay of KCNQ2 potassium channels using rubidium efflux and atomic absorption spectrometry. *Anal Biochem*. 2003;319:251–257. doi: 10.1016/s0003-2697(03)00328-2
23. Ozhathil LC, Delalande C, Bianchi B, Nemeth G, Kappel S, Thomet U, Ross-Kaschitza D, Simonin C, Rubin M, Gertsch J, et al. Identification of potent and selective small molecule inhibitors of the cation channel TRPM4. *Br J Pharmacol*. 2018;175:2504–2519. doi: 10.1111/bph.14220
24. Yang K-C, Breitbart A, De Lange WJ, Hofsteen P, Futakuchi-Tsuchida A, Xu J, Schopf C, Razumova MV, Jiao A, Boucek R, et al. Novel adult-onset systolic cardiomyopathy due to MYH7 E848G mutation in patient-derived induced pluripotent stem cells. *JACC Basic Transl Sci*. 2018;3:728–740. doi: 10.1016/j.jacbts.2018.08.008
25. Meisner JK, Renberg A, Smith ED, Tsan Y-C, Elder B, Bullard A, Merritt OL, Zheng SL, Lakdawala NK, Owens AT, et al. Low penetrance sarcomere variants contribute to additive risk in hypertrophic cardiomyopathy. *Circulation*. 2025;151:783–798. doi: 10.1161/CIRCULATIONAHA.124.069398
26. Chopra A, Kutys ML, Zhang K, Polacheck WJ, Sheng CC, Luu RJ, Eyckmans J, Hinson JT, Seidman JG, Seidman CE, et al. Force generation via β-cardiac myosin, titin, and α-actinin drives cardiac sarcomere assembly from cell-matrix adhesions. *Dev Cell*. 2018;44:87–96.e5. doi: 10.1016/j.devcel.2017.12.012
27. Shah PP, Lv W, Rhoades JH, Poleshko A, Abbey D, Caporizzo MA, Linares-Saldana R, Heffler JG, Sayed N, Thomas D, et al. Pathogenic LMNA variants disrupt cardiac lamina-chromatin interactions

Downloaded from http://ahajournals.org by on February 27, 2026

and de-repress alternative fate genes. *Cell Stem Cell*. 2021;28:938–954.e9. doi: 10.1016/j.stem.2020.12.016
28. Laurini E, Martinelli V, Lanzicher T, Puzzi L, Borin D, Chen SN, Long CS, Lee P, Mestroni L, Taylor MRG, et al. Biomechanical defects and rescue of cardiomyocytes expressing pathologic nuclear lamins. *Cardiovasc Res*. 2018;114:846–857. doi: 10.1093/cvr/cvy040
29. Puzzi L, Borin D, Martinelli V, Mestroni L, Kelsell DP, Sbaizero O. Cellular biomechanics impairment in keratinocytes is associated with a C-terminal truncated desmoplakin: An atomic force microscopy investigation. *Micron*. 2018;106:27–33. doi: 10.1016/j.micron.2017.12.005
30. Dinarelli S, Girasole M, Spitalieri P, Talarico RV, Murdocca M, Botta A, Novelli G, Mango R, Sangiuolo F, Longo G. AFM nano-mechanical study of the beating profile of hiPSC-derived cardiomyocytes beating bodies WT and DM1. *J Mol Recognit*. 2018;31:e2725. doi: 10.1002/jmr.2725
31. Dainis A, Zaleta-Rivera K, Ribeiro A, Chang ACH, Shang C, Lan F, Burridge PW, Liu WR, Wu JC, Chang ACY, et al. Silencing of MYH7 ameliorates disease phenotypes in human iPSC-cardiomyocytes. *Physiol Genomics*. 2020;52:293–303. doi: 10.1152/physiolgenomics.00021.2020
32. Lee S, Vander Roest AS, Blair CA, Kao K, Bremner SB, Childers MC, Pathak D, Heinrich P, Lee D, Chirikian O, et al. Incomplete-penetrant hypertrophic cardiomyopathy MYH7 G256E mutation causes hypercontractility and elevated mitochondrial respiration. *Proc Natl Acad Sci USA*. 2024;121:e2318413121. doi: 10.1073/pnas.2318413121
33. Loiben AM, Chien W-M, Friedman CE, Chao LS-L, Weber G, Goldstein A, Sniadecki NJ, Murry CE, Yang K-C. Cardiomyocyte apoptosis is associated with contractile dysfunction in stem cell model of MYH7 E848G hypertrophic cardiomyopathy. *Int J Mol Sci*. 2023;24:4909. doi: 10.3390/ijms24054909
34. Wijnker PJM, Dinani R, van der Laan NC, Algül S, Knollmann BC, Verkerk AO, Remme CA, Zuurbier CJ, Kuster DWD, van der Velden J. Hypertrophic cardiomyopathy dysfunction mimicked in human engineered heart tissue and improved by sodium-glucose cotransporter 2 inhibitors. *Cardiovasc Res*. 2024;120:301–317. doi: 10.1093/cvr/cvae004
35. Kinnear C, Said A, Meng G, Zhao Y, Wang EY, Rafatian N, Parmar N, Wei W, Billia F, Simmons CA, et al. Myosin inhibitor reverses hypertrophic cardiomyopathy in genotypically diverse pediatric iPSC-cardiomyocytes to mirror variant correction. *Cell Rep Med*. 2024;5:101520. doi: 10.1016/j.xcrm.2024.101520
36. Huang G, Bisaria A, Wakefield DL, Yamawaki TM, Luo X, Zhang JA, Vigneault P, Wang J, Reagan JD, Oliverio O, et al. Titin-truncating variants in hiPSC cardiomyocytes induce pathogenic proteinopathy and sarcomere defects with preserved core contractile machinery. *Stem Cell Rep*. 2023;18:220–236. doi: 10.1016/j.stemcr.2022.11.008
37. Ryoo H, Kimmel H, Rondo E, Underhill GH. Advances in high throughput cell culture technologies for therapeutic screening and biological discovery applications. *Bioeng Transl Med*. 2024;9:e10627. doi: 10.1002/btm2.10627
38. Fowler DM, Adams DJ, Gloyn AL, Hahn WC, Marks DS, Muffley LA, Neal JT, Roth FP, Rubin AF, Starita LM, et al. An atlas of variant effects to understand the genome at nucleotide resolution. *Genome Biol*. 2023;24:147. doi: 10.1186/s13059-023-02986-x
39. Chong ZX, Yeap SK, Ho WY. Transfection types, methods and strategies: a technical review. *PeerJ*. 2021;9:e11165. doi: 10.7717/peerj.11165
40. Adli M. The CRISPR tool kit for genome editing and beyond. *Nat Commun*. 2018;9:1911. doi: 10.1038/s41467-018-04252-2
41. Findlay GM, Daza RM, Martin B, Zhang MD, Leith AP, Gasperini M, Janizek JD, Huang X, Starita LM, Shendure J. Accurate classification of BRCA1 variants with saturation genome editing. *Nature*. 2018;562:217–222. doi: 10.1038/s41586-018-0461-z
42. Kim HS, Kweon J, Kim Y. Recent advances in CRISPR-based functional genomics for the study of disease-associated genetic variants. *Exp Mol Med*. 2024;56:861–869. doi: 10.1038/s12276-024-01212-3
43. Anzalone AV, Randolph PB, Davis JR, Sousa AA, Koblan LW, Levy JM, Chen PJ, Wilson C, Newby GA, Raguram A, et al. Search-and-replace genome editing without double-strand breaks or donor DNA. *Nature*. 2019;576:149–157. doi: 10.1038/s41586-019-1711-4
44. Belli O, Karava K, Farouni R, Platt RJ. Multimodal scanning of genetic variants with base and prime editing. *Nat Biotechnol*. 2024;43:1458–1470. doi: 10.1038/s41587-024-02439-1
45. Feng M, Xiang J-Z, Ming Z-Y, Fu Q, Ma R, Zhang Q-F, Dun Y-Y, Yang L, Liu H. Activation of epidermal growth factor receptor mediates reperfusion arrhythmias in anaesthetized rats. *Cardiovasc Res*. 2012;93:60–68. doi: 10.1093/cvr/cvr281
46. Anzalone AV, Koblan LW, Liu DR. Genome editing with CRISPR-Cas nucleases, base editors, transposases and prime editors. *Nat Biotechnol*. 2020;38:824–844. doi: 10.1038/s41587-020-0561-9
47. Kantor A, McClements ME, MacLaren RE. CRISPR-Cas9 DNA base-editing and prime-editing. *Int J Mol Sci*. 2020;21:6240. doi: 10.3390/ijms21176240
48. Jeong YK, Song B, Bae S. Current status and challenges of DNA base editing tools. *Mol Ther*. 2020;28:1938–1952. doi: 10.1016/j.ymthe.2020.07.021
49. Yamamoto Y, Chua K, Ferrasse A, Kirilova A, De Jong HN, Floyd BJ, Cadisch C, Wiel L, Wang Q, O'Neill MJ, et al. Scaled multidimensional assays of variant effect identify sequence-function relationships in hypertrophic cardiomyopathy. *bioRxivorg*. 2025. doi: 10.1101/2025.05.23.655878
50. Hanna RE, Hegde M, Fagre CR, DeWeirdt PC, Sangree AK, Szegletes Z, Griffith A, Feeley MN, Sanson KR, Baidi Y, et al. Massively parallel assessment of human variants with base editor screens. *Cell*. 2021;184:1064–1080.e20. doi: 10.1016/j.cell.2021.01.012
51. Xu Z, Thomas L, Davies B, Chalmers R, Smith M, Brown W. Accuracy and efficiency define Bxb1 integrase as the best of fifteen candidate serine recombinases for the integration of DNA into the human genome. *BMC Biotechnol*. 2013;13:87. doi: 10.1186/1472-6750-13-87
52. Brown WRA, Lee NCO, Xu Z, Smith MCM. Serine recombinases as tools for genome engineering. *Methods*. 2011;53:372–379. doi: 10.1016/j.ymeth.2010.12.031
53. Matreyek KA, Stephany JJ, Fowler DM. A platform for functional assessment of large variant libraries in mammalian cells. *Nucleic Acids Res*. 2017;45:e102. doi: 10.1093/nar/gkx183
54. Blanch-Asensio A, Grandela C, Brandão KO, de Korte T, Mei H, Ariyurek Y, Yiangou L, Mol MPH, van Meer BJ, Kloet SL, et al. STRAIGHT-IN enables high-throughput targeting of large DNA payloads in human pluripotent stem cells. *Cell Rep Methods*. 2022;2:100300. doi: 10.1016/j.crmeth.2022.100300
55. Li A, Acevedo-Rocha CG, Reetz MT. Boosting the efficiency of site-saturation mutagenesis for a difficult-to-randomize gene by a two-step PCR strategy. *Appl Microbiol Biotechnol*. 2018;102:6095–6103. doi: 10.1007/s00253-018-9041-2
56. Melnikov A, Rogov P, Wang L, Gnirke A, Mikkelsen TS. Comprehensive mutational scanning of a kinase in vivo reveals substrate-dependent fitness landscapes. *Nucleic Acids Res*. 2014;42:e112. doi: 10.1093/nar/gku511
57. Mighell TL, Toledano I, Lehner B. SUNi mutagenesis: scalable and uniform nicking for efficient generation of variant libraries. *PLoS One*. 2023;18:e0288158. doi: 10.1371/journal.pone.0288158
58. Macdonald CB, Nedrud D, Grimes PR, Trinidad D, Fraser JS, Coyote-Maestas W. DIMPLE: deep insertion, deletion, and missense mutation libraries for exploring protein variation in evolution, disease, and biology. *Genome Biol*. 2023;24:36. doi: 10.1186/s13059-023-02880-6
59. Beckmann B-M, Pfeufer A, Kääb S. Inherited cardiac arrhythmias: diagnosis, treatment, and prevention. *Dtsch Arztebl Int*. 2011;108:623–33; quiz 634. doi: 10.3238/arztebl.2011.0623
60. Dehghani-Samani A, Madreseh-Ghahfarokhi S, Dehghani-Samani A. Mutations of voltage-gated ionic channels and risk of severe cardiac arrhythmias. *Acta Cardiol Sin*. 2019;35:99–110. doi: 10.6515/ACS.201903_35(2).20181028A
61. Giudicessi JR, Ackerman MJ. Potassium-channel mutations and cardiac arrhythmias--diagnosis and therapy. *Nat Rev Cardiol*. 2012;9:319–332. doi: 10.1038/nrcardio.2012.3
62. Yu H-B, Li M, Wang W-P, Wang X-L. High throughput screening technologies for ion channels. *Acta Pharmacol Sin*. 2016;37:34–43. doi: 10.1038/aps.2015.108
63. Sakla M, Breitinger U, Breitinger H-G, Mansour S, Nabil Tammam S. A review of current and prospective treatments for channelopathies, with a focus on gene and protein therapy. *Curr Pharm Des*. 2023;29:1341–1360. doi: 10.2174/1381612829666230601122846
64. Yajuan X, Xin L, Zhiyuan L. A comparison of the performance and application differences between manual and automated patch-clamp techniques. *Curr Chem Genomics*. 2012;6:87–92. doi: 10.2174/1875397301206010087
65. Ma JG, Vandenberg JI, Ng C-A. Development of automated patch clamp assays to overcome the burden of variants of uncertain significance in inheritable arrhythmia syndromes. *Front Physiol*. 2023;14:1294741. doi: 10.3389/fphys.2023.1294741
66. Annecchino LA, Schultz SR. Progress in automating patch clamp cellular physiology. *Brain Neurosci Adv*. 2018;2:2398212818776561. doi: 10.1177/2398212818776561
67. O'Neill MJ, Ma JG, Aldridge JL, Solus JF, Harvey GR, Roberson PH, Barc J, Bezzina CR, Roden DM, Walsh R, et al. Automated patch clamp data improve variant classification and penetrance stratification for SCN5A-Brugada syndrome. *Eur Heart J*. 2025. doi: 10.1093/eurheartj/ehaf874
68. Vanoye CG, Desai RR, John JD, Hoffman SC, Fink N, Zhang Y, Venkatesh OG, Roe J, Adusumilli S, Jairam NP, et al. Functional profiling of KCNE1 variants informs population carrier frequency of Jervell and Lange-Nielsen

Downloaded from http://ahajournals.org by on February 27, 2026

syndrome type 2. *Circ Genom Precis Med*. 2025;18:e005219. doi: 10.1161/CIRCGEN.125.005219
69. O'Neill MJ, Ng C-A, Aizawa T, Sala L, Bains S, Winbo A, Ullah R, Shen Q, Tan C-Y, Kozek K, et al. Multiplexed assays of variant effect and automated patch clamping improve KCNH2-LQTS variant classification and cardiac event risk stratification. *Circulation*. 2024;150:1869–1881. doi: 10.1161/CIRCULATIONAHA.124.069828
70. Thomson KL, Jiang C, Richardson E, Westphal DS, Burkard T, Wolf CM, Vatta M, Harrison SM, Ingles J, Bezzina CR, et al. Clinical interpretation of KCNH2 variants using a robust PS3/BS3 functional patch-clamp assay. *HGG Adv*. 2024;5:100270. doi: 10.1016/j.xhgg.2024.100270
71. Ng C-A, Ullah R, Farr J, Hill AP, Kozek KA, Vanags LR, Mitchell DW, Kroncke BM, Vandenberg JI. A massively parallel assay accurately discriminates between functionally normal and abnormal variants in a hotspot domain of KCNH2. *Am J Hum Genet*. 2022;109:1208–1216. doi: 10.1016/j.ajhg.2022.05.003
72. Jiang C, Richardson E, Farr J, Hill AP, Ullah R, Kroncke BM, Harrison SM, Thomson KL, Ingles J, Vandenberg JI, et al. A calibrated functional patch-clamp assay to enhance clinical variant interpretation in KCNH2-related long QT syndrome. *Am J Hum Genet*. 2022;109:1199–1207. doi: 10.1016/j.ajhg.2022.05.002
73. González JE, Oades K, Leychkis Y, Harootunian A, Negulescu PA. Cell-based assays and instrumentation for screening ion-channel targets. *Drug Discov Today*. 1999;4:431–439. doi: 10.1016/s1359-6446(99)01383-5
74. Vetter I, Carter D, Bassett J, Deuis JR, Tay B, Jami S, Robinson SD. High-throughput fluorescence assays for ion channels and GPCRs. *Adv Exp Med Biol*. 2020;1131:27–72. doi: 10.1007/978-3-030-12457-1_3
75. Zhang Y-L, Moran SP, Allen A, Baez-Nieto D, Xu Q, Wang LA, Martenis WE, Sacher JR, Gale JP, Weïwer M, et al. Novel fluorescence-based high-throughput FLIPR assay utilizing membrane-tethered genetic calcium sensors to identify T-type calcium channel modulators. *ACS Pharmacol Transl Sci*. 2022;5:156–168. doi: 10.1021/acsptsci.1c00233
76. Iamshanova O, Mariot P, Lehen'kyi V'yacheslav, Prevarskaya N. Comparison of fluorescence probes for intracellular sodium imaging in prostate cancer cell lines. *Eur Biophys J*. 2016;45:765–777. doi: 10.1007/s00249-016-1173-7
77. Paredes RM, Etzler JC, Watts LT, Zheng W, Lechleiter JD. Chemical calcium indicators. *Methods*. 2008;46:143–151. doi: 10.1016/j.ymeth.2008.09.025
78. Shen Y, Wu S-Y, Rancic V, Aggarwal A, Qian Y, Miyashita S-I, Ballanyi K, Campbell RE, Dong M. Genetically encoded fluorescent indicators for imaging intracellular potassium ion concentration. *Commun Biol*. 2019;2:18. doi: 10.1038/s42003-018-0269-2
79. Soga O, Kinoshita H, Ueda M, Tanaka A. Evaluation of peroxisomal heme in yeast. *J Biochem*. 1997;121:25–28. doi: 10.1093/oxfordjournals.jbchem.a021564
80. Roston TM, Wei J, Guo W, Li Y, Zhong X, Wang R, Estillore JP, Peltenburg PJ, Noguer FRI, Till J, et al. Clinical and functional characterization of ryanodine receptor 2 variants implicated in calcium-release deficiency syndrome. *JAMA Cardiol*. 2022;7:84–92. doi: 10.1001/jamacardio.2021.4458
81. Rebbeck RT, Essawy MM, Nitu FR, Grant BD, Gillispie GD, Thomas DD, Bers DM, Cornea RL. High-throughput screens to discover small-molecule modulators of ryanodine receptor calcium release channels. *SLAS Discov*. 2017;22:176–186. doi: 10.1177/1087057116674312
82. Takenaka M, Kodama M, Murayama T, Ishigami-Yuasa M, Mori S, Ishida R, Suzuki J, Kanemaru K, Sugihara M, Iino M, et al. Screening for novel type 2 ryanodine receptor inhibitors by endoplasmic reticulum Ca2+ monitoring. *Mol Pharmacol*. 2023;104:275–286. doi: 10.1124/molpharm.123.000720
83. Lyon RC, Zanella F, Omens JH, Sheikh F. Mechanotransduction in cardiac hypertrophy and failure. *Circ Res*. 2015;116:1462–1476. doi: 10.1161/CIRCRESAHA.116.304937
84. Swiatlowska P, Iskratsch T. Tools for studying and modulating (cardiac muscle) cell mechanics and mechanosensing across the scales. *Biophys Rev*. 2021;13:611–623. doi: 10.1007/s12551-021-00837-2
85. Loiben AM, Chien W-M, McKinstry A, Ahmed D, Childers MC, Regnier M, Murry CE, Yang K-C. Functional interrogation of contextually correct MYH7 variants using CRaTER-flox gene editing and contractility profiling. *J Clin Invest*. 2026;136. doi: 10.1172/JCI192057
86. Friedman CE, Fayer S, Pendyala S, Chien W-M, Loiben A, Tran L, Chao LS, Mckinstry A, Ahmed D, Karbassi E, et al. CRaTER enrichment for on-target gene editing enables generation of variant libraries in hiPSCs. *J Mol Cell Cardiol*. 2023;179:60–71. doi: 10.1016/j.yjmcc.2023.03.017
87. Lipshultz SE, Orav EJ, Sanders SP, McIntosh K, Colan SD. Limitations of fractional shortening as an index of contractility in pediatric patients infected with human immunodeficiency virus. *J Pediatr*. 1994;125:563–570. doi: 10.1016/s0022-3476(94)70008-7
88. Rodriguez ML, Graham BT, Pabon LM, Han SJ, Murry CE, Sniadecki NJ. Measuring the contractile forces of human induced pluripotent stem cell-derived cardiomyocytes with arrays of microposts. *J Biomech Eng*. 2014;136:051005. doi: 10.1115/1.4027145
89. Beussman KM, Rodriguez ML, Leonard A, Taparia N, Thompson CR, Sniadecki NJ. Micropost arrays for measuring stem cell-derived cardiomyocyte contractility. *Methods*. 2016;94:43–50. doi: 10.1016/j.ymeth.2015.09.005
90. Jannati S, Maaref Y, Tibbits GF, Chiao M. Investigating viscoelastic properties of myofibrils isolated from hiPSC-CMs using atomic force microscopy and quasi-linear viscoelastic model. *J Appl Mech*. 2024;91. doi: 10.1115/1.4063141
91. Bremner S, Goldstein AJ, Higashi T, Sniadecki NJ. Engineered heart tissues for contractile, structural, and transcriptional assessment of human pluripotent stem cell-derived cardiomyocytes in a three-dimensional, auxotonic environment. *Methods Mol Biol*. 2022;2485:87–97. doi: 10.1007/978-1-0716-2261-2_6
92. Windt LM, Wiendels M, Dostanić M, Bellin M, Sarro PM, Mastrangeli M, Mummery CL, van Meer BJ. Miniaturized engineered heart tissues from hiPSC-derived triple cell type co-cultures to study human cardiac function. *Biochem Biophys Res Commun*. 2023;681:200–211. doi: 10.1016/j.bbrc.2023.09.034
93. Kamgoué A, Ohayon J, Usson Y, Riou L, Tracqui P. Quantification of cardiomyocyte contraction based on image correlation analysis. *Cytometry A*. 2009;75:298–308. doi: 10.1002/cyto.a.20700
94. Sharma A, Toepfer CN, Ward T, Wasson L, Agarwal R, Conner DA, Hu JH, Seidman CE. CRISPR/Cas9-mediated fluorescent tagging of endogenous proteins in human pluripotent stem cells. *Curr Protoc Hum Genet*. 2018;96:21.11.1–21.11.20. doi: 10.1002/cphg.52
95. Fassina L, Assenza MR, Miragoli M, Isidori AM, Naro F, Barbagallo F. Cell shortening and calcium homeostasis analysis in adult cardiomyocytes via a new software tool. *Biomedicines*. 2022;10:640. doi: 10.3390/biomedicines10030640
96. Toepfer CN, Sharma A, Cicconet M, Garfinkel AC, Mücke M, Neyazi M, Willcox JAL, Agarwal R, Schmid M, Rao J, et al. SarcTrack. *Circ Res*. 2019;124:1172–1183. doi: 10.1161/CIRCRESAHA.118.314505
97. Zhao B, Zhang K, Chen CS, Lejeune E. Sarc-Graph: automated segmentation, tracking, and analysis of sarcomeres in hiPSC-derived cardiomyocytes. *PLoS Comput Biol*. 2021;17:e1009443. doi: 10.1371/journal.pcbi.1009443
98. Henton A. Calcium and contractility system. IonOptix. 2019. https://www.ionoptix.com/products/systems/calcium-and-contractility-system/
99. Shroff SG, Saner DR, Lal R. Dynamic micromechanical properties of cultured rat atrial myocytes measured by atomic force microscopy. *Am J Physiol*. 1995;269:C286–C292. doi: 10.1152/ajpcell.1995.269.1.C286
100. Liu J, Sun N, Bruce MA, Wu JC, Butte MJ. Atomic force mechanobiology of pluripotent stem cell-derived cardiomyocytes. *PLoS One*. 2012;7:e37559. doi: 10.1371/journal.pone.0037559
101. Kit-Anan W, Mazo MM, Wang BX, Leonardo V, Pence IJ, Gopal S, Gelmi A, Nagelkerke A, Becce M, Chiappini C, et al. Multiplexing physical stimulation on single human induced pluripotent stem cell-derived cardiomyocytes for phenotype modulation. *Biofabrication*. 2021;13:025004. doi: 10.1088/1758-5090/abce0a
102. Nagarajan N, Vyas V, Huey BD, Zorlutuna P. Modulation of the contractility of micropatterned myocardial cells with nanoscale forces using atomic force microscopy. *Nanobiomedicine (Rij)*. 2016;3:1849543516675348. doi: 10.1177/1849543516675348
103. Peña B, Adbel-Hafiz M, Cavasin M, Mestroni L, Sbaizero O. Atomic force microscopy (AFM) applications in arrhythmogenic cardiomyopathy. *Int J Mol Sci*. 2022;23:3700. doi: 10.3390/ijms23073700
104. Ribeiro AJS, Ang Y-S, Fu J-D, Rivas RN, Mohamed TMA, Higgs GC, Srivastava D, Pruitt BL. Contractility of single cardiomyocytes differentiated from pluripotent stem cells depends on physiological shape and substrate stiffness. *Proc Natl Acad Sci USA*. 2015;112:12705–12710. doi: 10.1073/pnas.1508073112
105. Pasqualini FS, Agarwal A, O'Connor BB, Liu Q, Sheehy SP, Parker KK. Traction force microscopy of engineered cardiac tissues. *PLoS One*. 2018;13:e0194706. doi: 10.1371/journal.pone.0194706
106. Tanaka Y, Morishima K, Shimizu T, Kikuchi A, Yamato M, Okano T, Kitamori T. Demonstration of a PDMS-based bio-microactuator using cultured cardiomyocytes to drive polymer micropillars. *Lab Chip*. 2006;6:230–235. doi: 10.1039/b512099c
107. Ribeiro AJS, Zaleta-Rivera K, Ashley EA, Pruitt BL. Stable, covalent attachment of laminin to microposts improves the contractility of mouse neonatal

Downloaded from http://ahajournals.org by on February 27, 2026

cardiomyocytes. *ACS Appl Mater Interfaces*. 2014;6:15516–15526. doi: 10.1021/am5042324
108. Kim K, Taylor R, Sim JY, Park S-J, Norman J, Fajardo G, Bernstein D, Pruitt BL. Calibrated micropost arrays for biomechanical characterisation of cardiomyocytes. *Micro Nano Lett*. 2011;6:317–322. doi: 10.1049/mnl.2011.0031
109. Mannhardt I, Breckwoldt K, Letuffe-Brenière D, Schaaf S, Schulz H, Neuber C, Benzin A, Werner T, Eder A, Schulze T, et al. Human engineered heart tissue: analysis of contractile force. *Stem Cell Rep*. 2016;7:29–42. doi: 10.1016/j.stemcr.2016.04.011
110. Mosqueira D, Mannhardt I, Bhagwan JR, Lis-Slimak K, Katili P, Scott E, Hassan M, Prondzynski M, Harmer SC, Tinker A, et al. CRISPR/Cas9 editing in human pluripotent stem cell-cardiomyocytes highlights arrhythmias, hypocontractility, and energy depletion as potential therapeutic targets for hypertrophic cardiomyopathy. *Eur Heart J*. 2018;39:3879–3892. doi: 10.1093/eurheartj/ehy249
111. Cofiño-Fabres C, Boonen T, Rivera-Arbeláez JM, Rijpkema M, Blauw L, Rensen PCN, Schwach V, Ribeiro MC, Passier R. Micro-engineered heart tissues on-chip with heterotypic cell composition display self-organization and improved cardiac function. *Adv Healthc Mater*. 2024;13:e2303664. doi: 10.1002/adhm.202303664
112. Smith AS, Luttrell SM, Dupont J-B, Gray K, Lih D, Fleming JW, Cunningham NJ, Jepson S, Hesson J, Mathieu J, et al. High-throughput, real-time monitoring of engineered skeletal muscle function using magnetic sensing. *J Tissue Eng*. 2022;13:20417314221122127. doi: 10.1177/20417314221122127
113. Wang X, Wang L, Dou W, Huang Z, Zhao Q, Malhi M, Maynes JT, Sun Y. Electrical impedance-based contractile stress measurement of human iPSC-Cardiomyocytes. *Biosens Bioelectron*. 2020;166:112399. doi: 10.1016/j.bios.2020.112399
114. Koci B, Luerman G, Duenbostell A, Kettenhofen R, Bohlen H, Coyle L, Knight B, Ku W, Volberg W, Woska JR Jr, et al. An impedance-based approach using human iPSC-derived cardiomyocytes significantly improves in vitro prediction of in vivo cardiotox liabilities. *Toxicol Appl Pharmacol*. 2017;329:121–127. doi: 10.1016/j.taap.2017.05.023
115. Matreyek KA, Starita LM, Stephany JJ, Martin B, Chiasson MA, Gray VE, Kircher M, Khechaduri A, Dines JN, Hause RJ, et al. Multiplex assessment of protein variant abundance by massively parallel sequencing. *Nat Genet*. 2018;50:874–882. doi: 10.1038/s41588-018-0122-z
116. Kanner SA, Jain A, Colecraft HM. Development of a high-throughput flow cytometry assay to monitor defective trafficking and rescue of long QT2 mutant hERG channels. *Front Physiol*. 2018;9:397. doi: 10.3389/fphys.2018.00397
117. Schmidt R, Ward CC, Dajani R, Armour-Garb Z, Ota M, Allain V, Hernandez R, Layeghi M, Xing G, Goudy L, et al. Base-editing mutagenesis maps alleles to tune human T cell functions. *Nature*. 2024;625:805–812. doi: 10.1038/s41586-023-06835-6
118. Medina-Carmona E, Betancor-Fernández I, Santos J, Mesa-Torres N, Grottelli S, Batlle C, Naganathan AN, Oppici E, Cellini B, Ventura S, et al. Insight into the specificity and severity of pathogenic mechanisms associated with missense mutations through experimental and structural perturbation analyses. *Hum Mol Genet*. 2019;28:1–15. doi: 10.1093/hmg/ddy323
119. Pettinato AM, Ladha FA, Mellert DJ, Legere N, Cohn R, Romano R, Thakar K, Chen Y-S, Hinson JT. Development of a cardiac sarcomere functional genomics platform to enable scalable interrogation of human TNNT2 variants. *Circulation*. 2020;142:2262–2275. doi: 10.1161/CIRCULATIONAHA.120.047999
120. Anderson CL, Routes TC, Eckhardt LL, Delisle BP, January CT, Kamp TJ. A rapid solubility assay of protein domain misfolding for pathogenicity assessment of rare DNA sequence variants. *Genet Med*. 2020;22:1642–1652. doi: 10.1038/s41436-020-0842-1
121. Anderson CL, Kuzmicki CE, Childs RR, Hintz CJ, Delisle BP, January CT. Large-scale mutational analysis of Kv11.1 reveals molecular insights into type 2 long QT syndrome. *Nat Commun*. 2014;5:5535. doi: 10.1038/ncomms6535
122. Blandin CE, Gravez BJ, Hatem SN, Balse E. Remodeling of ion channel trafficking and cardiac arrhythmias. *Cells*. 2021;10:2417. doi: 10.3390/cells10092417
123. Telford WG. Flow cytometry and cell sorting. *Front Med (Lausanne)*. 2023;10:1287884. doi: 10.3389/fmed.2023.1287884
124. Coyote-Maestas W, Nedrud D, He Y, Schmidt D. Determinants of trafficking, conduction, and disease within a K+ channel revealed through multiparametric deep mutational scanning. *Elife*. 2022;11:e76903. doi: 10.7554/eLife.76903
125. Oliveira-Mendes B, Feliciangeli S, Ménard M, Chatelain F, Alameh M, Montnach J, Nicolas S, Ollivier B, Barc J, Baró I, et al. A standardised hERG phenotyping pipeline to evaluate KCNH2 genetic variant pathogenicity. *Clin Transl Med*. 2021;11:e609. doi: 10.1002/ctm2.609
126. Sheng S-Y, Li J-M, Hu X-Y, Wang Y. Regulated cell death pathways in cardiomyopathy. *Acta Pharmacol Sin*. 2023;44:1521–1535. doi: 10.1038/s41401-023-01068-9
127. Olcum M, Rouhi L, Fan S, Gonzales MM, Jeong H-H, Zhao Z, Gurha P, Marian AJ. PANoptosis is a prominent feature of desmoplakin cardiomyopathy. *J Cardiovasc Aging*. 2023;3:3. doi: 10.20517/jca.2022.34
128. Olcum M, Fan S, Rouhi L, Cheedipudi S, Cathcart B, Jeong H-H, Zhao Z, Gurha P, Marian AJ. Genetic inactivation of β-catenin is salubrious, whereas its activation is deleterious in desmoplakin cardiomyopathy. *Cardiovasc Res*. 2023;119:2712–2728. doi: 10.1093/cvr/cvad137
129. Hoorntje ET, Burns C, Marsili L, Corden B, Parikh VN, Te Meerman GJ, Gray B, Adiyaman A, Bagnall RD, Barge-Schaapveld DQCM, et al. Variant location is a novel risk factor for individuals with arrhythmogenic cardiomyopathy due to a desmoplakin (DSP) truncating variant. *Circ Genom Precis Med*. 2023;16:e003672. doi: 10.1161/CIRCGEN.121.003672
130. Chen SN, Lombardi R, Karmouch J, Tsai J-Y, Czernuszewicz G, Taylor MRG, Mestroni L, Coarfa C, Gurha P, Marian AJ. DNA damage response/TP53 pathway is activated and contributes to the pathogenesis of dilated cardiomyopathy associated with LMNA (lamin A/C) mutations. *Circ Res*. 2019;124:856–873. doi: 10.1161/CIRCRESAHA.118.314238
131. Hsieh J, Becklin KL, Givens S, Komosa ER, Lloréns JEA, Kamdar F, Moriarity BS, Webber BR, Singh BN, Ogle BM. Myosin heavy chain converter domain mutations drive early-stage changes in extracellular matrix dynamics in hypertrophic cardiomyopathy. *Front Cell Dev Biol*. 2022;10:894635. doi: 10.3389/fcell.2022.894635
132. Jaganathan K, Kyriazopoulou Panagiotopoulou S, McRae JF, Darbandi SF, Knowles D, Li YI, Kosmicki JA, Arbelaez J, Cui W, Schwartz GB, et al. Predicting splicing from primary sequence with deep learning. *Cell*. 2019;176:535–548.e24. doi: 10.1016/j.cell.2018.12.015
133. Zhang X, Cai M, Ma Y, Chen J, Huang S, Cai M, Ding Y, Ma D, Gao Q, Hu X, et al. Minigene assay as an effective molecular diagnostic strategy in determining the pathogenicity of noncanonical splice-site variants in FLCN. *J Mol Diagn*. 2023;25:110–120. doi: 10.1016/j.jmoldx.2022.10.005
134. O'Neill MJ, Yang T, Laudeman J, Calandranis ME, Harvey ML, Solus JF, Roden DM, Glazer AM. ParSE-seq: a calibrated multiplexed assay to facilitate the clinical classification of putative splice-altering variants. *Nat Commun*. 2024;15:8320. doi: 10.1038/s41467-024-52474-4
135. O'Neill MJ, Wada Y, Hall LD, Mitchell DW, Glazer AM, Roden DM. Functional assays reclassify suspected splice-altering variants of uncertain significance in Mendelian channelopathies. *Circ Genom Precis Med*. 2022;15:e003782. doi: 10.1161/CIRCGEN.122.003782
136. Adamson SI, Zhan L, Graveley BR. Vex-seq: high-throughput identification of the impact of genetic variation on pre-mRNA splicing efficiency. *Genome Biol*. 2018;19:71. doi: 10.1186/s13059-018-1437-x
137. Gergics P, Smith C, Bando H, Jorge AAL, Rockstroh-Lippold D, Vishnopolska SA, Castinetti F, Maksutova M, Carvalho LRS, Hoppmann J, et al. High-throughput splicing assays identify missense and silent splice-disruptive POU1F1 variants underlying pituitary hormone deficiency. *Am J Hum Genet*. 2021;108:1526–1539. doi: 10.1016/j.ajhg.2021.06.013
138. Julien P, Miñana B, Baeza-Centurion P, Valcárcel J, Lehner B. The complete local genotype-phenotype landscape for the alternative splicing of a human exon. *Nat Commun*. 2016;7:11558. doi: 10.1038/ncomms11558
139. Chong R, Insigne KD, Yao D, Burghard CP, Wang J, Hsiao Y-HE, Jones EM, Goodman DB, Xiao X, Kosuri S. A multiplexed assay for Exon recognition reveals that an unappreciated fraction of rare genetic variants cause large-effect splicing disruptions. *Mol Cell*. 2019;73:183–194.e8. doi: 10.1016/j.molcel.2018.10.037
140. Brnich SE, Abou Tayoun AN, Couch FJ, Cutting GR, Greenblatt MS, Heinen CD, Kanavy DM, Luo X, McNulty SM, Starita LM, et al; Clinical Genome Resource Sequence Variant Interpretation Working Group. Recommendations for application of the functional evidence PS3/BS3 criterion using the ACMG/AMP sequence variant interpretation framework. *Genome Med*. 2019;12:3. doi: 10.1186/s13073-019-0690-2
141. Fayer S, Horton C, Dines JN, Rubin AF, Richardson ME, McGoldrick K, Hernandez F, Pesaran T, Karam R, Shirts BH, et al. Closing the gap: Systematic integration of multiplexed functional data resolves variants of uncertain significance in BRCA1, TP53, and PTEN. *Am J Hum Genet*. 2021;108:2248–2258. doi: 10.1016/j.ajhg.2021.11.001

Downloaded from http://ahajournals.org by on February 27, 2026