

# Experimental Data Analysis

*in ©MATLAB*

## **Lecture 7:** Factorial ANOVA, ANCOVA, MANOVA Example of research project solution

Jan Rusz  
Czech Technical University in Prague



## Example

The measure of maximum phonation time through sustained vowel phonation well reflects subjects' aerodynamic efficiency of the vocal tract. Normal speaker should be able to perform sustained vowel phonation for more than 15 second. The aim of the research was to verify if there is some impairment of respiratory control across patients with Huntington's disease and Parkinson's disease as compared to healthy control subjects.

## Maximum phonation time

Groups

Single factor  
„group“


Random sample  
within each group.

Huntington's disease	Parkinson's disease	Healthy controls
5	8	29
7	17	22
8	11	18
5	10	18
5	15	13
4	10	11

## Why Factorial ANOVA?

- In a One-way ANOVA, we selected a random sample for each column/treatment group
- A factorial ANOVA compare means across two or more independent variables
- A two-way ANOVA allows us to “account for variation” at the ROW level due to some other factor or grouping
- This allows greater focus on COLUMN or GROUP differences making it easier to detect group differences
- We are attempting to minimize the ERROR variance due to the additional variance in the ROWS
- We will now have 5 types of Sum of Squares
  - Total Sum of Squares (SST)
  - Column (Between) Sum of Squares (SSC)
  - Error (Within) Sum of Squares (SSE)
  - Rows (Blocks) Sum of Squares (SSB)
  - Column  $\times$  Blocks Sum of Squares (SSCB)

## Maximum phonation time



Sex	Group		
	Huntington's disease	Parkinson's disease	Healthy controls
F	5	8	29
F	7	17	22
F	8	11	18
M	5	10	18
M	5	15	13
M	4	10	11

ANOVA  
One-way vs. Two-way  
Group

Huntington's disease
5
7
8
5
5
4
$\bar{x}_1 = 5.7$

Parkinson's disease
8
17
11
10
15
10
$\bar{x}_2 = 11.8$

Healthy controls
29
22
18
18
13
11
$\bar{x}_3 = 18.5$

Total average  $\bar{\bar{x}} = 12$


# ANOVA

## One-way vs. **Two-way**

### **Group & Sex**


	Huntington's disease	Parkinson's disease	Healthy controls	
Female	5	8	29	$\bar{x}_{R1} = 13.9$
	7	17	22	
	8	11	18	
	6.7	8.7	23	
Male	5	10	18	$\bar{x}_{R2} = 10.1$
	5	15	13	
	4	10	11	
	4.7	11.7	14	
	$\bar{x}_{C1} = 5.7$	$\bar{x}_{C2} = 11.8$	$\bar{x}_{C3} = 18.5$	$\bar{\bar{x}} = 12$

## Partitioning Sum of Squares: One-way ANOVA





## Partitioning Sum of Squares: Two-way ANOVA



## Established hypothesis?

- 1)  $H_0$ : Type of disease will have no significant effect on maximum phonation time;
- 2)  $H_0$ : Gender will have no significant effect on maximum phonation time;
- 3)  $H_0$ : Interaction of type of disease and gender will have no significant effect on maximum phonation time; in other words, maximum phonation time is not gender-dependent across different types of neurological diseases.

	Sum of Squares	d.f.	Mean Square	F score
1 <sup>st</sup> Factor (Group) Column	$SSC$	$C-1$	$\frac{SSC}{df_{columns}}$	$\frac{MSC}{MSE}$
2 <sup>nd</sup> Factor (Sex) Row	$SSB$	$B-1$	$\frac{SSB}{df_{blocks}}$	$\frac{MSB}{MSE}$
Both Factors (Group*Sex)	$SSCB$	$(C-1)(B-1)$	$\frac{SSCB}{df_{interaction}}$	$\frac{MSI}{MSE}$
Within (Error)	$SSE$	$N-C*B$	$\frac{SSE}{df_{error}}$	
Total	$SST$	$N-1$		

$N$  = total observations;  $C$  = number of columns;  $B$  = number of rows

# Sum of Squares 1st Factor (Group)

Sex	Huntington's disease	Parkinson's disease	Healthy controls	Average
Female	6.7	12.0	23.0	13.9
Male	4.7	11.7	14.0	10.1
Average	5.7	11.8	18.5	12.0

$$SSC = \sum r(\bar{x} - \bar{\bar{x}})^2 + \sum r(\bar{x} - \bar{\bar{x}})^2 + \sum r(\bar{x} - \bar{\bar{x}})^2$$

$r$  = number of data in each column

Squared		
Female	Group mean - Grand mean	
5	$(5.7 - 12)^2$	40.1
7	$(5.7 - 12)^2$	40.1
8	$(5.7 - 12)^2$	40.1
8	$(11.8 - 12)^2$	0.03
17	$(11.8 - 12)^2$	0.03
11	$(11.8 - 12)^2$	0.03
29	$(18.5 - 12)^2$	42.3
22	$(18.5 - 12)^2$	42.3
18	$(18.5 - 12)^2$	42.3
	<b>Average</b>	<b>247.2</b>

Squared		
Male	Group mean - Grand mean	
5	$(5.7 - 12)^2$	40.1
5	$(5.7 - 12)^2$	40.1
4	$(5.7 - 12)^2$	40.1
10	$(11.8 - 12)^2$	0.03
15	$(11.8 - 12)^2$	0.03
10	$(11.8 - 12)^2$	0.03
18	$(18.5 - 12)^2$	42.3
13	$(18.5 - 12)^2$	42.3
11	$(18.5 - 12)^2$	42.3
	<b>Average</b>	<b>247.2</b>

$$SSC = 247.2 + 247.2 = 494.4$$

	Sum of Squares	d.f.	Mean Square	F score
1 <sup>st</sup> Factor (Group) Column	494.4	C-1	$\frac{SSC}{df_{columns}}$	$\frac{MSC}{MSE}$
2 <sup>nd</sup> Factor (Sex) Row	$SSB$	$B-1$	$\frac{SSB}{df_{blocks}}$	$\frac{MSB}{MSE}$
Both Factors (Group*Sex)	$SSCB$	$(C-1)(B-1)$	$\frac{SSCB}{df_{interaction}}$	$\frac{SSI}{MSE}$
Within (Error)	$SSE$	$N-C*B$	$\frac{SSE}{df_{error}}$	
Total	$SST$	$N-1$		

$N$  = total observations;  $C$  = number of columns;  $B$  = number of rows

Sum of Squares 2nd Factor (Sex)

Sex	Huntington's disease	Parkinson's disease	Healthy controls	Average
Female	6.7	12.0	23.0	13.9
Male	4.7	11.7	14.0	10.1
Average	5.7	11.8	18.5	12.0

Female

Male

$$SSB = \sum r(\bar{x} - \bar{\bar{x}})^2 + \sum r(\bar{x} - \bar{\bar{x}})^2$$

$r$  = number of data in each column

Squared		
Female	Female mean - Grand mean	
5	$(13.9 - 12)^2$	3.6
7	$(13.9 - 12)^2$	3.6
8	$(13.9 - 12)^2$	3.6
8	$(13.9 - 12)^2$	3.6
17	$(13.9 - 12)^2$	3.6
11	$(13.9 - 12)^2$	3.6
29	$(13.9 - 12)^2$	3.6
22	$(13.9 - 12)^2$	3.6
18	$(13.9 - 12)^2$	3.6
	Average	32.1

Squared		
Male	Male mean - Grand mean	
5	$(10.1 - 12)^2$	3.6
5	$(10.1 - 12)^2$	3.6
4	$(10.1 - 12)^2$	3.6
10	$(10.1 - 12)^2$	3.6
15	$(10.1 - 12)^2$	3.6
10	$(10.1 - 12)^2$	3.6
18	$(10.1 - 12)^2$	3.6
13	$(10.1 - 12)^2$	3.6
11	$(10.1 - 12)^2$	3.6
	Average	32.1

**$SSB = 32.1 + 32.1 = 64.2$**

	Sum of Squares	d.f.	Mean Square	F score
1 <sup>st</sup> Factor (Group) Column	494.4	C-1	$\frac{SSC}{df_{columns}}$	$\frac{MSC}{MSE}$
2 <sup>nd</sup> Factor (Sex) Row	64.2	B-1	$\frac{SSB}{df_{blocks}}$	$\frac{MSB}{MSE}$
Both Factors (Group*Sex)	SSCB	(C-1)(B-1)	$\frac{SSCB}{df_{interaction}}$	$\frac{MSI}{MSE}$
Within (Error)	SSE	N-C*B	$\frac{SSE}{df_{error}}$	
Total	SST	N-1		

N = total observations; C = number of columns; B = number of rows

Sum of Squares Within (Error)

Sex	Huntington's disease	Parkinson's disease	Healthy controls	Average
Female	6.7	12.0	23.0	13.9
Male	4.7	11.7	14.0	10.1
Average	5.7	11.8	18.5	12.0

$SSE = \sum (x - \bar{x})^2$

Squared		
Female	Female score - Female mean	
5	$(5 - 6.7)^2$	2.8
7	$(7 - 6.7)^2$	0.1
8	$(8 - 6.7)^2$	1.8
8	$(8 - 12)^2$	16.0
17	$(17 - 12)^2$	25.0
11	$(11 - 12)^2$	1.0
29	$(29 - 23)^2$	36.0
22	$(22 - 23)^2$	1.0
18	$(18 - 23)^2$	25.0
	Average	108.7

Squared		
Male	Male score - Male mean	
5	$(5 - 4.7)^2$	0.1
5	$(5 - 4.7)^2$	0.1
4	$(4 - 4.7)^2$	0.4
10	$(10 - 11.7)^2$	2.8
15	$(15 - 11.7)^2$	11.1
10	$(10 - 11.7)^2$	2.8
18	$(18 - 14)^2$	16.0
13	$(13 - 14)^2$	1.0
11	$(11 - 14)^2$	9.0
	Average	43.3

**$SSE = 108.7 + 43.3 = 152.0$**



	Sum of Squares	d.f.	Mean Square	F score
1 <sup>st</sup> Factor (Group) Column	494.4	C-1	$\frac{SSC}{df_{columns}}$	$\frac{MSC}{MSE}$
2 <sup>nd</sup> Factor (Sex) Row	64.2	B-1	$\frac{SSB}{df_{blocks}}$	$\frac{MSB}{MSE}$
Both Factors (Group*Sex)	SSCB	(C-1)(B-1)	$\frac{SSCB}{df_{interaction}}$	$\frac{MSI}{MSE}$
Within (Error)	152.0	N-C*B	$\frac{SSE}{df_{error}}$	
Total	SST	N-1		

*N* = total observations; *C* = number of columns; *B* = number of rows

Sum of Squares Total


$SST = \sum (x - \bar{x})^2$

***SST* = 774**

Squared Each score - Grand mean		
5	$(5 - 12)^2$	49.0
7	$(7 - 12)^2$	25.0
8	$(8 - 12)^2$	16.0
5	$(5 - 12)^2$	49.0
5	$(5 - 12)^2$	49.0
4	$(4 - 12)^2$	64.0
8	$(8 - 12)^2$	16.0
17	$(17 - 12)^2$	25.0
11	$(11 - 12)^2$	1.0
10	$(10 - 12)^2$	4.0
15	$(15 - 12)^2$	9.0
10	$(10 - 12)^2$	4.0
29	$(29 - 12)^2$	289.0
22	$(22 - 12)^2$	100.0
18	$(18 - 12)^2$	36.0
18	$(18 - 12)^2$	36.0
13	$(13 - 12)^2$	1.0
11	$(11 - 12)^2$	1.0
Average		774.0

	Sum of Squares	d.f.	Mean Square	F score
1 <sup>st</sup> Factor (Group) Column	494.4	$C-1$	$\frac{SSC}{df_{columns}}$	$\frac{MSC}{MSE}$
2 <sup>nd</sup> Factor (Sex) Row	64.2	$B-1$	$\frac{SSB}{df_{blocks}}$	$\frac{MSB}{MSE}$
Both Factors (Group*Sex)	$SSCB$	$(C-1)(B-1)$	$\frac{SSCB}{df_{interaction}}$	$\frac{MSI}{MSE}$
Within (Error)	152.0	$N-C*B$	$\frac{SSE}{df_{error}}$	
Total	774.0	$N-1$		

$N$  = total observations;  $C$  = number of columns;  $B$  = number of rows



$$\text{SSCB} = 774.0 - 494.4 - 64.2 - 152.0 = \mathbf{63.4}$$

	Sum of Squares	d.f.	Mean Square	F score
1 <sup>st</sup> Factor (Group) Column	494.4	2	247.2	19.5
2 <sup>nd</sup> Factor (Sex) Row	64.2	1	64.2	5.1
Both Factors (Group*Sex)	63.4	2	31.7	2.5
Within (Error)	152.0	12	12.7	
Total	774.0	17		

$N$  = total observations;  $C$  = number of columns;  $B$  = number of rows

## Matlab output to replicate

### Matlab example 1

$p = 1 - \text{fcddc}(F, \text{df\_factor}, \text{df\_error});$

#### Analysis of Variance

Source	Sum Sq.	d.f.	Mean Sq.	F	Prob>F
group	494.333	2	247.167	19.51	0.0002
sex	64.222	1	64.222	5.07	0.0439
group*sex	63.444	2	31.722	2.5	0.1233
Error	152	12	12.667		
Total	774	17			

Group	Sex	Sample
1	1	5
1	1	7
1	1	8
1	2	5
1	2	5
1	2	4
2	1	8
2	1	17
2	1	11
2	2	10
2	2	15
2	2	10
3	1	29
3	1	22
3	1	18
3	2	18
3	2	13
3	2	11

## Answers to established hypotheses

- 1)  $H_0$ : Type of disease will have no significant effect on maximum phonation time;

$$F(2,12) = 19.5, p < 0.001$$

**REJECT**

- 2)  $H_0$ : Gender will have no significant effect on maximum phonation time;

$$F(1,12) = 5.1, p = 0.04$$

**REJECT**

- 3)  $H_0$ : Interaction of type of disease and gender will have no significant effect on maximum phonation time; in other words, maximum phonation time is not gender-dependent across different types of neurological diseases.

$$F(2,12) = 2.5, p = 0.12$$

**FAIL TO REJECT**

## Analysis of covariance (ANCOVA)

- Extension of ANOVA, in which main effects and interactions are assessed on scores of dependent variables (DV) **after DV has been adjusted for the relationship of DV with one or more covariates (CV)**
- CV is a variable that is related to the DV, which you can't manipulate, but you want to account for its relationship with DV
- CV should be collected before analysis is started



## Applications of ANCOVA


- Increase test sensitivity (main effects and interactions) by using the CV to account for more of the error variance and therefore making the error term smaller
- In other words, using **CV** we **can show a larger effect** or **CV can eliminate the effect**

## Assumptions using ANCOVA


- Normality of sampling distributions of the DV and each CV
- Absence of outliers
- Homogeneity of variance
- Linearity – we have to consider a linear relationship between each CV and DV as well as each pair of CV

## Matlab output to replicate

`[p,a,s] = anova1(Sample,Group)`



`[p,a,s] = aocool(Age,Sample,Group)`



Covariate

Group	Age	Sample
1	80	5
1	75	7
1	75	8
1	72	5
1	68	5
1	71	4
2	69	8
2	55	17
2	60	11
2	61	10
2	58	15
2	62	10
3	41	29
3	50	22
3	48	18
3	52	18
3	60	13
3	61	11

## Multivariate analysis of variance (MANOVA)


- Extension of ANOVA with the key difference that we are dealing with **many dependent variables** (not a single DV as in the case of ANOVA)
- Combination of DV is called “joint distribution”
- MANOVA gives us answer to question if joint distribution of 2 or more DV is significantly related to one or more factors
- The results of MANOVA simply tell us that a difference exists (or not) across groups

## Assumptions using MANOVA


- At least 2 DV
- Multivariate normality of sampling distributions of the DV
- Multivariate homogeneity of variance
- May be correlated, **DV should be related**

## Matlab output to replicate

`[p,a,s] = anova1(Sample1,Group)`



`[p,a,s] = anova1(Sample2,Group)`



`[d,p] = manova1([Sample1 Sample2],Group)`

$d = 1$  ... estimate of dimension of space containing group means

$p = [0.0051 \ 0.8946]$  ...  $p(d)$

$(r = 0.89, p < 0.001)$

Group	Sample1	Sample2
1	5	3
1	7	7
1	8	8
1	5	3
1	5	5
1	4	4
2	8	8
2	17	14
2	11	10
2	10	10
2	15	10
2	10	8
3	29	29
3	22	10
3	18	12
3	18	17
3	13	12
3	11	10

# Research project

ACOUSTIC ANALYSIS OF SPEECH  
PATTERNS IN PARKINSON'S DISEASE

## Why to conduct such research?

- Patients with PD have limited ability to express contrastive stress
- Impaired stress production is one the most common affected dimensions of speech in PD
- Very limited research regarding stress production in PD





## General motivation

- Early and accurate diagnosis
- Monitoring progression of disease
- Monitoring treatment efficacy
- Feedback for speech therapy



## Principle of stress expression

- Word prolongation (Time)
- Increase of pitch (Fo)
- Increase of loudness (Intensity)



Hypothesis: what happen during stress expression?

Parameter	Helthy controls	Parkinson's disease
Word length	↑↑↑	↑
Pitch	↑↑↑	↑
Loudness	↑↑↑	↑

## Methods: speech data

- The aim was to unnaturally emphasize certain keyword marked in the text


*Dnes jsme to již nestihli,*  
možná ZÍTRA navštívíme všechny své známé.

*Příbuzné jsme již navštívili,*  
možná zítra navštívíme všechny své ZNÁMÉ.

## Methods: speech data

5 words & 2 versions of reading

stressed




non-stressed



## Methods: variables







- Mean value of Fo
- Range of Fo (min-max)
- Mean value of intensity of loudness
- Range of intensity of loudness
- Length of word
- Stress pattern index (combining effects of all three fundamental aspects including pitch, loudness, and length)

# Results



PD = Parkinson's disease  
 HC = Healthy controls

# Results



—●— HC - stressed production  
 -○- HC - non-stressed production

—▲— PD - stressed production  
 -△- PD - non-stressed production

- 2 groups (PD, HC)
  - 6 variables
  - 5 words
  - 2 conditions (stressed, non-stressed)
- +1 condition (difference between stressed and non-stressed)

For each PD vs. HC comparison:

$$6 \times 5 \times 3 = 90 \text{ tests}$$

# Results

MEAN F0		STRESSED	NORMAL	ABS. STRESS
ZNAME	PN x HC	NS	NS	NS
NAVSTIVIME	PN x HC	NS	NS	NS
SVE	PN x HC	NS	NS	NS
VSECHNY	PN x HC	NS	NS	NS
ZITRA	PN x HC	NS	NS	NS

MEAN INTENSITY		STRESSED	NORMAL	ABS. STRESS
ZNAME	PN x HC	NS	NS	NS
NAVSTIVIME	PN x HC	NS	NS	NS
SVE	PN x HC	NS	NS	NS
VSECHNY	PN x HC	NS	NS	NS
ZITRA	PN x HC	NS	NS	NS

DURATION		STRESSED	NORMAL	ABS. STRESS
ZNAME	PN x HC	NS	p < 0.01	NS
NAVSTIVIME	PN x HC	p < 0.05	p < 0.01	NS
SVE	PN x HC	p < 0.05	p < 0.05	p < 0.05
VSECHNY	PN x HC	NS	p < 0.05	NS
ZITRA	PN x HC	p < 0.01	p < 0.01	NS

RANGE INTENSITY		STRESSED	NORMAL	ABS. STRESS
ZNAME	PN x HC	p < 0.05	p < 0.001	NS
NAVSTIVIME	PN x HC	p < 0.001	p < 0.001	NS
SVE	PN x HC	p < 0.001	p < 0.01	NS
VSECHNY	PN x HC	p < 0.05	p < 0.01	NS
ZITRA	PN x HC	p < 0.001	p < 0.001	NS

RANGE F0		STRESSED	NORMAL	ABS. STRESS
ZNAME	PN x HC	p < 0.001	p < 0.01	p < 0.01
NAVSTIVIME	PN x HC	p < 0.05	NS	NS
SVE	PN x HC	NS	NS	NS
VSECHNY	PN x HC	NS	NS	NS
ZITRA	PN x HC	p < 0.01	p < 0.001	NS

STRESS PATTERN INDEX		STRESSED	NORMAL	ABS. STRESS
ZNAME	PN x HC	p < 0.05	p < 0.001	NS
NAVSTIVIME	PN x HC	p < 0.001	p < 0.001	p < 0.05
SVE	PN x HC	p < 0.001	p < 0.05	p < 0.01
VSECHNY	PN x HC	NS	p < 0.01	NS
ZITRA	PN x HC	p < 0.001	p < 0.001	p < 0.01

- Very hard to **interpret** the results
- **Type I Error**
- Corrected:  $p = 0.05/90 \Rightarrow p < 0.00056$
- $\Rightarrow$  **no result is significant**

NS = not significant




## Solution: Three-way ANOVA

- $2 \times 2 \times 5$  ANOVA
- 3 factors
  - **SPEAKERS GROUP** (PD, HC)
  - **STRESS CONDITION** (stressed, non-stressed)
  - **KEYWORD ORDER** (first, second, third, fourth, fifth)

# Three-way ANOVA: Interpretation

## SPEAKERS GROUP

PD patients will manifest affected speech




## Three-way ANOVA: Results

	mean F0	mean Intensity	duration	range F0	range Intensity	SPI
<b>Speakers group</b>	<b>n.s.</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Stress condition	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Keyword order	<0.001	n.s.	<0.001	<0.001	<0.001	<0.001
Speakers group x Stress condition	n.s.	n.s.	n.s.	0.0014	n.s.	<0.001
Keyword order x Speakers group	n.s.	n.s.	n.s.	n.s.	<0.001	<0.001
Keyword order x Stress condition	n.s.	n.s.	<0.001	n.s.	n.s.	<0.001

# Three-way ANOVA: Interpretation

## STRESS CONDITION

Variable reflects effect of the stress condition




## Three-way ANOVA: Results

	mean F0	mean Intensity	duration	range F0	range Intensity	SPI
Speakers group	n.s.	<0.001	<0.001	<0.001	<0.001	<0.001
<b>Stress condition</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Keyword order	<0.001	n.s.	<0.001	<0.001	<0.001	<0.001
Speakers group x Stress condition	n.s.	n.s.	n.s.	0.0014	n.s.	<0.001
Keyword order x Speakers group	n.s.	n.s.	n.s.	n.s.	<0.001	<0.001
Keyword order x Stress condition	n.s.	n.s.	<0.001	n.s.	n.s.	<0.001

# Three-way ANOVA: Interpretation

## KEYWORD ORDER

Variable reflects the keyword order




## Three-way ANOVA: Results

	mean F0	mean Intensity	duration	range F0	range Intensity	SPI
Speakers group	n.s.	<0.001	<0.001	<0.001	<0.001	<0.001
Stress condition	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
<b>Keyword order</b>	<b>&lt;0.001</b>	<b>n.s.</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Speakers group x Stress condition	n.s.	n.s.	n.s.	0.0014	n.s.	<0.001
Keyword order x Speakers group	n.s.	n.s.	n.s.	n.s.	<0.001	<0.001
Keyword order x Stress condition	n.s.	n.s.	<0.001	n.s.	n.s.	<0.001

# Three-way ANOVA: Interpretation

**SPEAKERS GROUP ×  
STRESS CONDITION**

PD patients are not able to  
express stress as  
effectively as HC






## Three-way ANOVA: Results

	mean F0	mean Intensity	duration	range F0	range Intensity	SPI
Speakers group	n.s.	<0.001	<0.001	<0.001	<0.001	<0.001
Stress condition	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Keyword order	<0.001	n.s.	<0.001	<0.001	<0.001	<0.001
<b>Speakers group x Stress condition</b>	<b>n.s.</b>	<b>n.s.</b>	<b>n.s.</b>	<b>0.0014</b>	<b>n.s.</b>	<b>&lt;0.001</b>
Keyword order x Speakers group	n.s.	n.s.	n.s.	n.s.	<0.001	<0.001
Keyword order x Stress condition	n.s.	n.s.	<0.001	n.s.	n.s.	<0.001

# Three-way ANOVA: Interpretation

KEYWORD ORDER ×  
SPEAKERS GROUP

Certain word is more  
effective in the separation  
of groups




## Three-way ANOVA: Results

	mean F0	mean Intensity	duration	range F0	range Intensity	SPI
Speakers group	n.s.	<0.001	<0.001	<0.001	<0.001	<0.001
Stress condition	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Keyword order	<0.001	n.s.	<0.001	<0.001	<0.001	<0.001
Speakers group x Stress condition	n.s.	n.s.	n.s.	0.0014	n.s.	<0.001
<b>Keyword order x Speakers group</b>	<b>n.s.</b>	<b>n.s.</b>	<b>n.s.</b>	<b>n.s.</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Keyword order x Stress condition	n.s.	n.s.	<0.001	n.s.	n.s.	<0.001

# Three-way ANOVA: Interpretation

KEYWORD ORDER ×  
STRESS CONDITION

Certain word is more  
suitable for expression of  
stress condition



## Three-way ANOVA: Results

	mean F0	mean Intensity	duration	range F0	range Intensity	SPI
Speakers group	n.s.	<0.001	<0.001	<0.001	<0.001	<0.001
Stress condition	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Keyword order	<0.001	n.s.	<0.001	<0.001	<0.001	<0.001
Speakers group x Stress condition	n.s.	n.s.	n.s.	0.0014	n.s.	<0.001
Keyword order x Speakers group	n.s.	n.s.	n.s.	n.s.	<0.001	<0.001
<b>Keyword order x Stress condition</b>	<b>n.s.</b>	<b>n.s.</b>	<b>&lt;0.001</b>	<b>n.s.</b>	<b>n.s.</b>	<b>&lt;0.001</b>

## Three-way ANOVA: Interpretation

**SPEAKERS GROUP** – PD patients will manifest affected speech

**STRESS CONDITION** – variable reflects effect of the stress condition

**KEYWORD ORDER** – variable reflects the keyword order

**SPEAKERS GROUP × STRESS CONDITION** – PD patients are not able to express stress as effectively as HC

**KEYWORD ORDER × SPEAKERS GROUP** – certain word is more effective in the separation of groups

**KEYWORD ORDER × STRESS CONDITION** – certain word is more suitable for expression of stress condition

# Matlab output to replicate

Analysis of Variance					
Source	Sum Sq.	d.f.	Mean Sq.	F	Prob>F
group	66.628	1	66.6284	106.67	0
stress	95.182	1	95.182	152.38	0
keyword	165.214	4	41.3036	66.13	0
group*stress	8.844	1	8.8439	14.16	0.0002
group*keyword	26.217	4	6.5542	10.49	0
stress*keyword	38.71	4	9.6774	15.49	0
Error	211.123	338	0.6246		
Total	595.155	353			

## Main conclusions regarding our findings

- PD patients have decreased ability to express stress (**SPEAKERS GROUP × STRESS CONDITION**)
- PD patients are able to consciously improve their speech performance (**MAIN EFFECTS**)



## Some notes regarding interactions

- Always look for significant interaction first
- If it is significant, you do not analyze main effects
- Variables are interconnected to allows look at them individually