

Contents

| | |
|--|-------------|
| Glossary | xiii |
| Acronyms | xiv |
| 1 Introduction | 1 |
| 1.1 Cancer Research in the Post-Genomic Era | 1 |
| 1.1.1 Cancer as a Global Health Concern | 2 |
| 1.1.1.1 The Genetics and Molecular Biology of Cancers | 3 |
| 1.1.2 The Human Genome Revolution | 6 |
| 1.1.2.1 The First Human Genome Sequence | 6 |
| 1.1.2.2 Impact of Genomics | 7 |
| 1.1.3 Technologies to Enable Genetics Research | 7 |
| 1.1.3.1 DNA Sequencing and Genotyping Technologies | 7 |
| 1.1.3.2 Microarrays and Quantitative Technologies | 8 |
| 1.1.3.3 Massively Parallel “Next Generation” Sequencing | 9 |
| 1.1.3.3.1 Molecular Profiling with Genomics Technology | 11 |
| 1.1.3.3.2 Sequencing Technologies | 11 |
| 1.1.3.4 Bioinformatics as Interdisciplinary Genomic Analysis | 12 |
| 1.1.4 Follow-up Large-Scale Genomics Projects | 13 |
| 1.1.5 Cancer Genomes | 14 |
| 1.1.5.1 The Cancer Genome Atlas Project | 15 |
| 1.1.5.1.1 Findings from Cancer Genomes | 15 |
| 1.1.5.1.2 Genomic Comparisons Across Cancer Tissues | 17 |
| 1.1.5.1.3 Cancer Genomic Data Resources | 18 |
| 1.1.6 Genomic Cancer Medicine | 18 |
| 1.1.6.1 Cancer Genes and Driver Mutations | 18 |
| 1.1.6.2 Personalised or Precision Cancer Medicine | 19 |
| 1.1.6.2.1 Molecular Diagnostics and Pan-Cancer Medicine | 20 |
| 1.1.6.3 Targeted Therapeutics and Pharmacogenomics | 21 |
| 1.1.6.3.1 Targeting Oncogenic Driver Mutations | 21 |
| 1.1.6.4 Systems and Network Biology | 22 |
| 1.1.6.4.1 Network Medicine, and Polypharmacology | 24 |
| 1.2 A Synthetic Lethal Approach to Cancer Medicine | 25 |
| 1.2.1 Synthetic Lethal Genetic Interactions | 26 |
| 1.2.2 Synthetic Lethal Concepts in Genetics | 26 |
| 1.2.3 Studies of Synthetic Lethality | 27 |

| | | |
|-----------|---|-----------|
| 1.2.3.1 | Synthetic Lethal Pathways and Networks | 28 |
| 1.2.3.1.1 | Evolution of Synthetic Lethality | 29 |
| 1.2.4 | Synthetic Lethal Concepts in Cancer | 29 |
| 1.2.5 | Clinical Impact of Synthetic Lethality in Cancer | 31 |
| 1.2.6 | High-throughput Screening for Synthetic Lethality | 33 |
| 1.2.6.1 | Synthetic Lethal Screens | 34 |
| 1.2.7 | Computational Prediction of Synthetic Lethality | 37 |
| 1.2.7.1 | Bioinformatics Approaches to Genetic Interactions . . | 37 |
| 1.2.7.2 | Comparative Genomics | 38 |
| 1.2.7.3 | Analysis and Modelling of Protein Data | 41 |
| 1.2.7.4 | Differential Gene Expression | 43 |
| 1.2.7.5 | Data Mining and Machine Learning | 44 |
| 1.2.7.6 | Bimodality | 47 |
| 1.2.7.7 | Rationale for Further Development | 48 |
| 1.3 | E-cadherin as a Synthetic Lethal Target | 48 |
| 1.3.1 | The <i>CDH1</i> gene and it's Biological Functions | 48 |
| 1.3.1.1 | Cytoskeleton | 49 |
| 1.3.1.2 | Extracellular and Tumour Micro-Environment | 49 |
| 1.3.1.3 | Cell-Cell Adhesion and Signalling | 49 |
| 1.3.2 | <i>CDH1</i> as a Tumour (and Invasion) Suppressor | 50 |
| 1.3.2.1 | Breast Cancers and Invasion | 50 |
| 1.3.3 | Hereditary Diffuse Gastric Cancer and Lobular Breast Cancer . | 50 |
| 1.3.4 | Somatic Mutations | 52 |
| 1.3.4.1 | Mutation Rate | 52 |
| 1.3.4.2 | Co-occurring Mutations | 52 |
| 1.3.5 | Models of <i>CDH1</i> loss in cell lines | 53 |
| 1.4 | Summary and Research Direction of Thesis | 54 |
| 2 | Methods and Resources | 58 |
| 2.1 | Bioinformatics Resources for Genomics Research | 58 |
| 2.1.1 | Public Data and Software Packages | 58 |
| 2.1.1.1 | Cancer Genome Atlas Data | 59 |
| 2.1.1.2 | Reactome and Annotation Data | 60 |
| 2.2 | Data Handling | 60 |
| 2.2.1 | Normalisation | 60 |
| 2.2.2 | Sample Triage | 61 |
| 2.2.3 | Metagenes and the Singular Value Decomposition | 63 |
| 2.2.3.1 | Candidate Triage and Integration with Screen Data . . | 63 |
| 2.3 | Techniques | 64 |
| 2.3.1 | Statistical Procedures and Tests | 64 |
| 2.3.2 | Gene Set Over-representation Analysis | 65 |
| 2.3.3 | Clustering | 65 |
| 2.3.4 | Heatmap | 66 |
| 2.3.5 | Modeling and Simulations | 66 |
| 2.3.5.1 | Receiver Operating Characteristic (Performance) . . . | 67 |
| 2.3.6 | Resampling Analysis | 67 |

| | | |
|----------|---|------------|
| 2.4 | Pathway Structure Methods | 68 |
| 2.4.1 | Network and Graph Analysis | 68 |
| 2.4.2 | Sourcing Graph Structure Data | 69 |
| 2.4.3 | Constructing Pathway Subgraphs | 70 |
| 2.4.4 | Network Analysis Metrics | 70 |
| 2.5 | Implementation | 71 |
| 2.5.1 | Computational Resources and Linux Utilities | 71 |
| 2.5.2 | R Language and Packages | 72 |
| 2.5.3 | High Performance and Parallel Computing | 75 |
| 3 | Methods Developed During Thesis | 77 |
| 3.1 | A Synthetic Lethal Detection Methodology | 77 |
| 3.2 | Synthetic Lethal Simulation and Modelling | 80 |
| 3.2.1 | A Model of Synthetic Lethality in Expression Data | 80 |
| 3.2.2 | Simulation Procedure | 84 |
| 3.3 | Detecting Simulated Synthetic Lethal Partners | 87 |
| 3.3.1 | Binomial Simulation of Synthetic lethality | 87 |
| 3.3.2 | Multivariate Normal Simulation of Synthetic lethality | 89 |
| 3.3.2.1 | Multivariate Normal Simulation with Correlated Genes | 92 |
| 3.3.2.2 | Specificity with Query-Correlated Pathways | 99 |
| 3.3.2.3 | Importance of Directional Testing | 99 |
| 3.4 | Graph Structure Methods | 101 |
| 3.4.1 | Upstream and Downstream Gene Detection | 101 |
| 3.4.1.1 | Permutation Analysis for Statistical Significance | 102 |
| 3.4.1.2 | Hierarchy Based on Biological Context | 103 |
| 3.4.2 | Simulating Gene Expression from Graph Structures | 104 |
| 3.5 | Customised Functions and Packages Developed | 108 |
| 3.5.1 | Synthetic Lethal Interaction Prediction Tool | 108 |
| 3.5.2 | Data Visualisation | 109 |
| 3.5.3 | Extensions to the iGraph Package | 112 |
| 3.5.3.1 | Sampling Simulated Data from Graph Structures | 112 |
| 3.5.3.2 | Plotting Directed Graph Structures | 112 |
| 3.5.3.3 | Computing Information Centrality | 113 |
| 3.5.3.4 | Testing Pathway Structure with Permutation Testing | 113 |
| 3.5.3.5 | Metapackage to Install iGraph Functions | 114 |
| 4 | Synthetic Lethal Analysis of Gene Expression Data | 115 |
| 4.1 | Synthetic Lethal Genes in Breast Cancer | 116 |
| 4.1.1 | Synthetic Lethal Pathways in Breast Cancer | 118 |
| 4.1.2 | Expression Profiles of Synthetic Lethal Partners | 119 |
| 4.1.2.1 | Subgroup Pathway Analysis | 122 |
| 4.2 | Comparing Synthetic Lethal Gene Candidates | 125 |
| 4.2.1 | Primary siRNA Screen Candidates | 125 |
| 4.2.2 | Comparison with Correlation | 126 |
| 4.2.3 | Comparison with Primary Screen Viability | 128 |
| 4.2.4 | Comparison with Secondary siRNA Screen Validation | 129 |

| | | |
|----------|--|------------|
| 4.2.5 | Comparison to Primary Screen at Pathway Level | 131 |
| 4.2.5.1 | Resampling Genes for Pathway Enrichment | 133 |
| 4.2.6 | Integrating Synthetic Lethal Pathways and Screens | 136 |
| 4.3 | Metagene Analysis | 138 |
| 4.3.1 | Pathway Expression | 139 |
| 4.3.2 | Somatic Mutation | 141 |
| 4.3.3 | Synthetic Lethal Pathway Metagenes | 145 |
| 4.3.4 | Synthetic Lethality in Breast Cancer | 146 |
| 4.4 | Replication in Stomach Cancer | 147 |
| 4.5 | Discussion | 147 |
| 4.5.1 | Strengths of the SLIPT Methodology | 147 |
| 4.5.2 | Synthetic Lethal Pathways for E-cadherin | 148 |
| 4.5.3 | Replication and Validation | 150 |
| 4.5.3.1 | Integration with siRNA Screening | 150 |
| 4.5.3.2 | Replication across Tissues and Cell lines | 151 |
| 4.6 | Summary | 151 |
| 5 | Synthetic Lethal Pathway Structure | 155 |
| 5.1 | Synthetic Lethal Genes in Reactome Pathways | 155 |
| 5.1.1 | The PI3K/AKT Pathway | 156 |
| 5.1.2 | The Extracellular Matrix | 158 |
| 5.1.3 | G Protein Coupled Receptors | 161 |
| 5.1.4 | Gene Regulation and Translation | 161 |
| 5.2 | Network Analysis of Synthetic Lethal Genes | 162 |
| 5.2.1 | Gene Connectivity and Vertex Degree | 163 |
| 5.2.2 | Gene Importance and Centrality | 164 |
| 5.2.2.1 | Information Centrality | 164 |
| 5.2.2.2 | PageRank Centrality | 166 |
| 5.3 | Relationships between Synthetic Lethal Genes | 168 |
| 5.3.1 | Hierarchical Pathway Structure | 168 |
| 5.3.1.1 | Contextual Hierarchy of PI3K | 168 |
| 5.3.1.2 | Testing Contextual Hierarchy of Synthetic Lethal Genes | 168 |
| 5.3.2 | Upstream or Downstream Synthetic Lethality | 172 |
| 5.3.2.1 | Measuring Structure of Candidates within PI3K | 172 |
| 5.3.2.2 | Resampling for Synthetic Lethal Pathway Structure | 174 |
| 5.4 | Discussion | 176 |
| 5.5 | Summary | 178 |
| 6 | Simulation and Modeling of Synthetic Lethal Pathways | 181 |
| 6.1 | Comparing methods | 182 |
| 6.1.1 | Performance of SLIPT and χ^2 across Quantiles | 183 |
| 6.1.1.1 | Correlated Query Genes affects Specificity | 186 |
| 6.1.2 | Alternative Synthetic Lethal Detection Strategies | 188 |
| 6.1.2.1 | Correlation for Synthetic Lethal Detection | 188 |
| 6.1.2.2 | Testing for Bimodality with BiSEp | 190 |
| 6.2 | Simulations with Graph Structures | 192 |

| | | |
|----------|--|------------|
| 6.2.1 | Performance over a Graph Structure | 193 |
| 6.2.1.1 | Simple Graph Structures | 193 |
| 6.2.1.2 | Constructed Graph Structures | 195 |
| 6.2.2 | Performance with Inhibitions | 199 |
| 6.2.3 | Synthetic Lethality across Graph Structures | 205 |
| 6.2.4 | Performance within a Simulated Human Genome | 209 |
| 6.3 | Simulations over pathway-based graphs | 215 |
| 6.3.1 | Pathway Structures in a Simulated Human Genome | 217 |
| 6.4 | Discussion | 220 |
| 6.4.1 | Simulation Procedure | 220 |
| 6.4.2 | Design and Performance of SLIPT | 221 |
| 6.4.3 | Simulations from Graph Structures | 223 |
| 6.5 | Summary | 224 |
| 7 | Discussion | 227 |
| 7.1 | Synthetic Lethality and <i>CDH1</i> Biology | 227 |
| 7.1.1 | Established Functions of <i>CDH1</i> | 228 |
| 7.1.2 | The Molecular Role of <i>CDH1</i> in Cancer | 228 |
| 7.2 | Significance | 229 |
| 7.2.1 | Synthetic Lethality in the Genomic Era | 229 |
| 7.2.2 | Clinical Interventions based on Synthetic Lethality | 231 |
| 7.3 | Evaluating the Synthetic Lethality Prediction Tool | 232 |
| 7.3.1 | Strength of the Synthetic Lethality Prediction Tool | 232 |
| 7.3.2 | Limitations of the Synthetic Lethality Prediction Tool | 232 |
| 7.3.3 | Comparisons to Alternative Methods | 232 |
| 7.3.3.1 | Combined with Experimental Screening | 232 |
| 7.3.3.2 | Differences to Computational Methods | 232 |
| 7.4 | Future Directions | 232 |
| 7.4.1 | Refinements Synthetic Lethality Prediction Methods | 234 |
| 7.4.1.1 | Wider Use of Synthetic Lethality Prediction | 234 |
| 7.4.2 | Validation of Synthetic Lethal Genes and Pathways | 234 |
| 7.4.2.1 | Pre-clinical and Clinical Testing | 234 |
| 7.4.3 | Application to Further Genes and Pathways | 234 |
| 8 | Conclusion | 235 |
| | References | 239 |
| A | Sample Quality | 265 |
| A.1 | Sample Correlation | 265 |
| A.2 | Replicate Samples in TCGA Breast | 268 |
| B | Software Used for Thesis | 272 |

| | | |
|----------|---|------------|
| C | Mutation Analysis in Breast Cancer | 281 |
| C.1 | Synthetic Lethal Genes and Pathways | 281 |
| C.2 | Synthetic Lethal Expression Profiles | 284 |
| C.3 | Comparison to Primary Screen | 287 |
| C.3.1 | Resampling Analysis | 289 |
| C.4 | Compare SLIPT genes | 291 |
| C.5 | Metagene Analysis | 293 |
| C.6 | Expression of Somatic Mutations | 294 |
| C.7 | Metagene Expression Profiles | 297 |
| D | Intrinsic Subtyping | 300 |
| E | Stomach Expression Analysis | 302 |
| E.1 | Synthetic Lethal Genes and Pathways | 302 |
| E.2 | Synthetic Lethal Expression Profiles | 305 |
| E.3 | Comparison to Primary Screen | 308 |
| E.3.1 | Resampling Analysis | 310 |
| E.4 | Metagene Analysis | 312 |
| F | Synthetic Lethal Genes in Pathways | 313 |
| G | Pathway Connectivity for Mutation SLIPT | 321 |
| H | Information Centrality for Gene Essentiality | 325 |
| I | Pathway Structure for Mutation SLIPT | 328 |
| J | Performance of SLIPT and χ^2 | 331 |
| J.0.1 | Correlated Query Genes affects Specificity | 337 |
| K | Graph Structures | 343 |
| K.1 | Simulations from Graph Structures | 349 |
| K.2 | Simulations from Inhibiting Graph Structures | 354 |
| K.3 | Simulation across Graph Structures | 364 |
| K.4 | Graph Structure Simulations with 20K genes | 368 |
| K.4.1 | Inhibiting Graph Structure Simulations with 20K genes | 375 |
| K.5 | Simations from Pathway Graph Structures | 387 |

List of Figures

| | | |
|------|--|-----|
| 1.1 | Synthetic genetic interactions | 27 |
| 1.2 | Synthetic lethality in cancer | 30 |
| 2.1 | Read count density | 62 |
| 2.2 | Read count sample mean | 62 |
| 3.1 | Framework for synthetic lethal prediction | 78 |
| 3.2 | Synthetic lethal prediction adapted for mutation | 79 |
| 3.3 | A model of synthetic lethal gene expression | 81 |
| 3.4 | Modeling synthetic lethal gene expression | 82 |
| 3.5 | Synthetic lethality with multiple genes | 83 |
| 3.6 | Simulating gene function | 85 |
| 3.7 | Simulating synthetic lethal gene function | 85 |
| 3.8 | Simulating synthetic lethal gene expression | 86 |
| 3.9 | Performance of binomial simulations | 88 |
| 3.10 | Comparison of statistical performance | 88 |
| 3.11 | Performance of multivariate normal simulations | 90 |
| 3.12 | Simulating expression with correlated gene blocks | 93 |
| 3.13 | Simulating expression with correlated gene blocks | 94 |
| 3.14 | Synthetic lethal prediction across simulations | 95 |
| 3.15 | Performance with correlations | 96 |
| 3.16 | Comparison of statistical performance with correlation structure | 97 |
| 3.17 | Performance with query correlations | 98 |
| 3.18 | Statistical evaluation of directional criteria | 99 |
| 3.19 | Performance of directional criteria | 100 |
| 3.20 | Simulated graph structures | 104 |
| 3.21 | Simulating expression from a graph structure | 106 |
| 3.22 | Simulating expression from graph structure with inhibitions | 107 |
| 3.23 | Demonstration of violin plots with custom features | 110 |
| 3.24 | Demonstration of annotated heatmap | 110 |
| 3.25 | Simulating graph structures | 113 |
| 4.1 | Synthetic lethal expression profiles of analysed samples | 121 |
| 4.2 | Comparison of SLIPT to siRNA | 125 |
| 4.3 | Compare SLIPT and siRNA genes with correlation | 126 |
| 4.4 | Compare SLIPT and siRNA genes with correlation | 127 |
| 4.5 | Compare SLIPT and siRNA genes with viability | 128 |

| | | |
|------|---|-----|
| 4.6 | Compare SLIPT genes with siRNA viability | 129 |
| 4.7 | Resampled intersection of SLIPT and siRNA candidates | 133 |
| 4.8 | Pathway metagene expression profiles | 140 |
| 4.9 | Expression profiles for constituent genes of PI3K | 142 |
| 4.10 | Expression profiles for estrogen receptor related genes | 143 |
| 4.11 | Somatic mutation against the PI3K metagene | 144 |
| 5.1 | Synthetic Lethality in the PI3K Cascade | 157 |
| 5.2 | Synthetic Lethality in the Elastic Fibre Formation Pathway | 159 |
| 5.3 | Synthetic Lethality in the Fibrin Clot Formation | 160 |
| 5.4 | Synthetic Lethality and Vertex Degree | 163 |
| 5.5 | Synthetic Lethality and Centrality | 166 |
| 5.6 | Synthetic Lethality and PageRank | 167 |
| 5.7 | Hierarchical Structure of PI3K | 169 |
| 5.8 | Hierarchy Score in PI3K against Synthetic Lethality in PI3K | 170 |
| 5.9 | Structure of Synthetic Lethality in PI3K | 172 |
| 5.10 | Structure of Synthetic Lethality Resampling in PI3K | 173 |
| 6.1 | Performance of χ^2 and SLIPT across quantiles | 184 |
| 6.2 | Performance of χ^2 and SLIPT across quantiles with more genes | 185 |
| 6.3 | Performance of χ^2 and SLIPT across quantiles with query correlation | 186 |
| 6.4 | Performance of χ^2 and SLIPT across quantiles with query correlation and more genes | 187 |
| 6.5 | Performance of negative correlation and SLIPT | 189 |
| 6.6 | Performance of simulations on a simple graph | 194 |
| 6.7 | Performance of simulations is similar in simple graphs | 195 |
| 6.8 | Performance of simulations on a constructed graph | 196 |
| 6.9 | Performance of simulations on a large graph | 198 |
| 6.10 | Performance of simulations on a simple graph with inhibition | 200 |
| 6.11 | Performance is higher on a simple inhibiting graph | 201 |
| 6.12 | Performance of simulations on a constructed graph with inhibition | 203 |
| 6.13 | Performance is affected by inhibition in graphs | 204 |
| 6.14 | Detection of Synthetic Lethality within a Graph Structure | 206 |
| 6.15 | Detection of Synthetic Lethality within a Graph Structure with Inhibitions | 208 |
| 6.16 | Performance of simulations including a simple graph | 210 |
| 6.17 | Performance on a simple graph improves with more genes | 211 |
| 6.18 | Performance on an inhibiting graph with more genes | 212 |
| 6.19 | Performance on an inhibiting graph improves with more genes | 214 |
| 6.20 | Performance of simulations on the PI3K cascade | 216 |
| 6.21 | Performance of simulations including the PI3K cascade | 218 |
| 6.22 | Performance on pathways improves with more genes | 219 |
| A.1 | Correlation profiles of removed samples | 266 |
| A.2 | Correlation analysis and sample removal | 267 |
| A.3 | Replicate excluded samples | 268 |
| A.4 | Replicate samples with all remaining | 269 |

| | | |
|------|---|-----|
| A.5 | Replicate samples with some excluded | 270 |
| C.1 | Synthetic lethal expression profiles of analysed samples | 285 |
| C.2 | Comparison of mtSLIPT to siRNA | 287 |
| C.3 | Compare mtSLIPT and siRNA genes with correlation | 291 |
| C.4 | Compare mtSLIPT and siRNA genes with correlation | 291 |
| C.5 | Compare mtSLIPT and siRNA genes with siRNA viability | 292 |
| C.6 | Somatic mutation against PIK3CA metagene | 294 |
| C.7 | Somatic mutation against PI3K protein | 295 |
| C.8 | Somatic mutation against AKT protein | 296 |
| C.9 | Pathway metagene expression profiles | 297 |
| C.10 | Expression profiles for p53 related genes | 298 |
| C.11 | Expression profiles for BRCA related genes | 299 |
| E.1 | Synthetic lethal expression profiles of stomach samples | 307 |
| E.2 | Comparison of SLIPT in stomach to siRNA | 308 |
| F.1 | Synthetic Lethality in the PI3K/AKT Pathway | 313 |
| F.2 | Synthetic Lethality in the PI3K/AKT Pathway in Cancer | 314 |
| F.3 | Synthetic Lethality in the Extracellular Matrix | 315 |
| F.4 | Synthetic Lethality in the GPCRs | 316 |
| F.5 | Synthetic Lethality in the GPCR Downstream | 317 |
| F.6 | Synthetic Lethality in the Translation Elongation | 318 |
| F.7 | Synthetic Lethality in the Nonsense-mediated Decay | 319 |
| F.8 | Synthetic Lethality in the 3' UTR | 320 |
| G.1 | Synthetic Lethality and Vertex Degree | 321 |
| G.2 | Synthetic Lethality and Centrality | 322 |
| G.3 | Synthetic Lethality and PageRank | 323 |
| H.1 | Information centrality distribution | 327 |
| I.1 | Synthetic Lethality and Heirarchy Score in PI3K | 328 |
| I.2 | Heirarchy Score in PI3K against Synthetic Lethality in PI3K | 329 |
| I.3 | Structure of Synthetic Lethality in PI3K | 329 |
| I.4 | Structure of Synthetic Lethality Resampling | 330 |
| J.1 | Performance of χ^2 and SLIPT across quantiles | 331 |
| J.2 | Performance of χ^2 and SLIPT across quantiles | 333 |
| J.3 | Performance of χ^2 and SLIPT across quantiles with more genes | 335 |
| J.4 | Performance of χ^2 and SLIPT across quantiles with query correlation | 337 |
| J.5 | Performance of χ^2 and SLIPT across quantiles with query correlation | 339 |
| J.6 | Performance of χ^2 and SLIPT across quantiles with query correlation and more genes | 341 |
| K.1 | Simple graph structures | 343 |
| K.2 | Simple graph structure | 344 |
| K.3 | Constructed graph structure | 344 |

| | | |
|------|--|-----|
| K.4 | Large constructed graph structure. | 345 |
| K.5 | Branching constructed graph structure | 345 |
| K.6 | Complex constructed graph structure | 347 |
| K.7 | Performance of simulations on a simple graph | 350 |
| K.8 | Performance of simulations on a constructed graph | 351 |
| K.9 | Performance of simulations on a branching graph | 352 |
| K.10 | Performance of simulations on a complex graph | 353 |
| K.11 | Performance of simulations on a simple graph with inhibition | 355 |
| K.12 | Performance of simulations on a simple graph with inhibition | 356 |
| K.13 | Performance of simulations on a constructed graph with inhibition | 357 |
| K.14 | Performance of simulations on a large constructed graph with inhibition | 358 |
| K.15 | Performance of simulations on a large constructed graph with inhibition | 359 |
| K.16 | Performance of simulations on a branching graph with inhibition | 360 |
| K.17 | Performance of simulations on a branching graph with inhibition | 361 |
| K.18 | Performance of simulations on a complex graph with inhibition | 362 |
| K.19 | Performance of simulations on a complex graph with inhibition | 363 |
| K.20 | Detection of Synthetic Lethality within a Graph Structure | 364 |
| K.21 | Detection of Synthetic Lethality within an Inhibiting Graph Structure | 366 |
| K.22 | Detection of Synthetic Lethality within an Inhibiting Graph Structure | 367 |
| K.23 | Performance of simulations on a simple graph with more genes | 369 |
| K.24 | Performance of simulations including a simple graph | 370 |
| K.25 | Performance of simulations including a constructed graph | 371 |
| K.26 | Performance of simulations including a large graph | 372 |
| K.27 | Performance of simulations including a branching graph | 373 |
| K.28 | Performance of simulations including a complex graph | 374 |
| K.29 | Performance of simulations including a simple graph with inhibition | 376 |
| K.30 | Performance of simulations including a simple graph with inhibition | 377 |
| K.31 | Performance of simulations including a simple graph with inhibition | 378 |
| K.32 | Performance of simulations including a constructed graph with inhibition | 379 |
| K.33 | Performance of simulations including a constructed graph with inhibition | 380 |
| K.34 | Performance of simulations including a large graph with inhibition | 381 |
| K.35 | Performance of simulations including a large graph with inhibition | 382 |
| K.36 | Performance of simulations including a branching graph with inhibition | 383 |
| K.37 | Performance of simulations including a branching graph with inhibition | 384 |
| K.38 | Performance of simulations including a complex graph with inhibition | 385 |
| K.39 | Performance of simulations including a complex graph with inhibition | 386 |
| K.40 | Performance of simulations on the $G_{\alpha i}$ signalling pathway | 387 |
| K.41 | Performance of simulations including the $G_{\alpha i}$ signalling pathway | 388 |

List of Tables

| | | |
|-----|--|-----|
| 1.1 | Methods for Predicting Genetic Interactions | 38 |
| 1.2 | Methods for Predicting Synthetic Lethality in Cancer | 39 |
| 1.3 | Methods used by Wu <i>et al.</i> (2014) | 40 |
| 2.1 | Excluded Samples by Batch and Clinical Characteristics. | 61 |
| 2.2 | Computers used during Thesis | 72 |
| 2.3 | Linux Utilities and Applications used during Thesis | 72 |
| 2.4 | R Installations used during Thesis | 73 |
| 2.5 | R Packages used during Thesis | 73 |
| 2.6 | R Packages Developed during Thesis | 75 |
| 4.1 | Candidate synthetic lethal gene partners of <i>CDH1</i> from SLIPT | 117 |
| 4.2 | Pathways for <i>CDH1</i> partners from SLIPT | 119 |
| 4.3 | Pathway composition for clusters of <i>CDH1</i> partners from SLIPT | 123 |
| 4.4 | Analysis of variance (ANOVA) for Synthetic Lethality and Correlation with <i>CDH1</i> | 127 |
| 4.5 | Comparing SLIPT genes against secondary siRNA screen in breast cancer | 130 |
| 4.6 | Pathway composition for <i>CDH1</i> partners from SLIPT and siRNA screen- ing | 132 |
| 4.7 | Pathways for <i>CDH1</i> partners from SLIPT | 135 |
| 4.8 | Pathways for <i>CDH1</i> partners from SLIPT and siRNA primary screen . | 137 |
| 4.9 | Candidate synthetic lethal metagenes against <i>CDH1</i> from SLIPT | 146 |
| 5.1 | ANOVA for Synthetic Lethality and Vertex Degree | 164 |
| 5.2 | ANOVA for Synthetic Lethality and Information Centrality | 166 |
| 5.3 | ANOVA for Synthetic Lethality and PageRank Centrality | 168 |
| 5.4 | ANOVA for Synthetic Lethality and PI3K Hierarchy | 171 |
| 5.5 | Resampling for pathway structure of synthetic lethal detection methods | 175 |
| B.1 | R Packages used during Thesis | 272 |
| C.1 | Candidate synthetic lethal gene partners of <i>CDH1</i> from mtSLIPT | 282 |
| C.2 | Pathways for <i>CDH1</i> partners from mtSLIPT | 283 |
| C.3 | Pathway composition for clusters of <i>CDH1</i> partners from mtSLIPT . . . | 286 |
| C.4 | Pathway composition for <i>CDH1</i> partners from mtSLIPT and siRNA . . . | 288 |
| C.5 | Pathways for <i>CDH1</i> partners from mtSLIPT | 289 |
| C.6 | Pathways for <i>CDH1</i> partners from mtSLIPT and siRNA primary screen | 290 |
| C.7 | Candidate synthetic lethal metagenes against <i>CDH1</i> from mtSLIPT . . . | 293 |

| | | |
|-----|--|-----|
| D.1 | Comparison of Intrinsic Subtypes | 300 |
| E.1 | Synthetic lethal gene partners of <i>CDH1</i> from SLIPT in stomach cancer | 303 |
| E.2 | Pathways for <i>CDH1</i> partners from SLIPT in stomach cancer | 304 |
| E.3 | Pathway composition for clusters of <i>CDH1</i> partners in stomach SLIPT | 306 |
| E.4 | Pathway composition for <i>CDH1</i> partners from SLIPT and siRNA screen- ing | 309 |
| E.5 | Pathways for <i>CDH1</i> partners from SLIPT in stomach cancer | 310 |
| E.6 | Pathways for <i>CDH1</i> partners from SLIPT in stomach and siRNA screen | 311 |
| E.7 | Candidate synthetic lethal metagenes against <i>CDH1</i> from SLIPT in stomach cancer | 312 |
| G.1 | ANOVA for Synthetic Lethality and Vertex Degree | 324 |
| G.2 | ANOVA for Synthetic Lethality and Information Centrality | 324 |
| G.3 | ANOVA for Synthetic Lethality and PageRank Centrality | 324 |
| H.1 | Information centrality for genes and molecules in the Reactome network | 326 |
| I.1 | ANOVA for Synthetic Lethality and PI3K Hierarchy | 328 |
| I.2 | Resampling for pathway structure of synthetic lethal detection methods | 330 |

Glossary

| | |
|------------------|--|
| RNA-Seq | Transcriptome data from sequencing RNA. |
| synthetic lethal | Genetic interactions where inactivation of multiple genes is inviable (or deleterious) when they are viable if inactivated separately. |

Acronyms

ANOVA Analysis of Variance.

PAM50 Prediction Analysis of Microarray 50.

SLIPT Synthetic lethal interaction prediction tool.

TCGA The Cancer Genome Atlas (genomics project).

UCSC University of California, Santa Cruz.

References

- Aarts, M., Bajrami, I., Herrera-Abreu, M.T., Elliott, R., Brough, R., Ashworth, A., Lord, C.J., and Turner, N.C. (2015) Functional genetic screen identifies increased sensitivity to weel inhibition in cells with defects in fanconi anemia and hr pathways. *Mol Cancer Ther*, **14**(4): 865–76.
- Abeshouse, A., Ahn, J., Akbani, R., Ally, A., Amin, S., Andry, C.D., Annala, M., Aprikian, A., Armenia, J., Arora, A., *et al.* (2015) The Molecular Taxonomy of Primary Prostate Cancer. *Cell*, **163**(4): 1011–1025.
- Adamski, M.G., Gumann, P., and Baird, A.E. (2014) A method for quantitative analysis of standard and high-throughput qPCR expression data based on input sample quantity. *PLoS ONE*, **9**(8): e103917.
- Adler, D. (2005) *vioplot: Violin plot*. R package version 0.2.
- Agarwal, S., Deane, C.M., Porter, M.A., and Jones, N.S. (2010) Revisiting date and party hubs: Novel approaches to role assignment in protein interaction networks. *PLoS Comput Biol*, **6**(6): e1000817.
- Agrawal, N., Akbani, R., Aksoy, B.A., Ally, A., Arachchi, H., Asa, S.L., Auman, J.T., Balasundaram, M., Balu, S., Baylin, S.B., *et al.* (2014) Integrated genomic characterization of papillary thyroid carcinoma. *Cell*, **159**(3): 676–690.
- Akbani, R., Akdemir, K.C., Aksoy, B.A., Albert, M., Ally, A., Amin, S.B., Arachchi, H., Arora, A., Auman, J.T., Ayala, B., *et al.* (2015) Genomic Classification of Cutaneous Melanoma. *Cell*, **161**(7): 1681–1696.
- Akobeng, A.K. (2007) Understanding diagnostic tests 3: receiver operating characteristic curves. *Acta Pdiatrica*, **96**(5): 644–647.
- American Cancer Society (2017) Genetics and cancer. <https://www.cancer.org/cancer/cancer-causes/genetics.html>. Accessed: 22/03/2017.

- American Society for Clinical Oncology (ASCO) (2017) The genetics of cancer. <http://www.cancer.net/navigating-cancer-care/cancer-basics/genetics/genetics-cancer>. Accessed: 22/03/2017.
- Anjomshoaa, A., Lin, Y.H., Black, M.A., McCall, J.L., Humar, B., Song, S., Fukuzawa, R., Yoon, H.S., Holzmann, B., Friederichs, J., *et al.* (2008) Reduced expression of a gene proliferation signature is associated with enhanced malignancy in colon cancer. *Br J Cancer*, **99**(6): 966–973.
- Araki, H., Knapp, C., Tsai, P., and Print, C. (2012) GeneSetDB: A comprehensive meta-database, statistical and visualisation framework for gene set analysis. *FEBS Open Bio*, **2**: 76–82.
- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., *et al.* (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet*, **25**(1): 25–29.
- Ashworth, A. (2008) A synthetic lethal therapeutic approach: poly(adp) ribose polymerase inhibitors for the treatment of cancers deficient in dna double-strand break repair. *J Clin Oncol*, **26**(22): 3785–90.
- Audeh, M.W., Carmichael, J., Penson, R.T., Friedlander, M., Powell, B., Bell-McGuinn, K.M., Scott, C., Weitzel, J.N., Oaknin, A., Loman, N., *et al.* (2010) Oral poly(adp-ribose) polymerase inhibitor olaparib in patients with *BRCA1* or *BRCA2* mutations and recurrent ovarian cancer: a proof-of-concept trial. *Lancet*, **376**(9737): 245–51.
- Babyak, M.A. (2004) What you see may not be what you get: a brief, nontechnical introduction to overfitting in regression-type models. *Psychosom Med*, **66**(3): 411–21.
- Bamford, S., Dawson, E., Forbes, S., Clements, J., Pettett, R., Dogan, A., Flanagan, A., Teague, J., Futreal, P.A., Stratton, M.R., *et al.* (2004) The COSMIC (Catalogue of Somatic Mutations in Cancer) database and website. *Br J Cancer*, **91**(2): 355–358.
- Barabási, A.L. and Albert, R. (1999) Emergence of scaling in random networks. *Science*, **286**(5439): 509–12.

- Barabási, A.L., Gulbahce, N., and Loscalzo, J. (2011) Network medicine: a network-based approach to human disease. *Nat Rev Genet*, **12**(1): 56–68.
- Barabási, A.L. and Oltvai, Z.N. (2004) Network biology: understanding the cell’s functional organization. *Nat Rev Genet*, **5**(2): 101–13.
- Barrat, A. and Weigt, M. (2000) On the properties of small-world network models. *The European Physical Journal B - Condensed Matter and Complex Systems*, **13**(3): 547–560.
- Barretina, J., Caponigro, G., Stransky, N., Venkatesan, K., Margolin, A.A., Kim, S., Wilson, C.J., Lehar, J., Kryukov, G.V., Sonkin, D., *et al.* (2012) The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. *Nature*, **483**(7391): 603–607.
- Barry, W.T. (2016) *safe: Significance Analysis of Function and Expression*. R package version 3.14.0.
- Baryshnikova, A., Costanzo, M., Dixon, S., Vizeacoumar, F.J., Myers, C.L., Andrews, B., and Boone, C. (2010a) Synthetic genetic array (sga) analysis in *saccharomyces cerevisiae* and *schizosaccharomyces pombe*. *Methods Enzymol*, **470**: 145–79.
- Baryshnikova, A., Costanzo, M., Kim, Y., Ding, H., Koh, J., Toufighi, K., Youn, J.Y., Ou, J., San Luis, B.J., Bandyopadhyay, S., *et al.* (2010b) Quantitative analysis of fitness and genetic interactions in yeast on a genome scale. *Nat Meth*, **7**(12): 1017–1024.
- Bass, A.J., Thorsson, V., Shmulevich, I., Reynolds, S.M., Miller, M., Bernard, B., Hinoue, T., Laird, P.W., Curtis, C., Shen, H., *et al.* (2014) Comprehensive molecular characterization of gastric adenocarcinoma. *Nature*, **513**(7517): 202–209.
- Bates, D. and Maechler, M. (2016) *Matrix: Sparse and Dense Matrix Classes and Methods*. R package version 1.2-7.1.
- Bateson, W. and Mendel, G. (1909) *Mendel’s principles of heredity*, by W. Bateson. University Press, Cambridge [Eng.].
- Beck, T.F., Mullikin, J.C., and Biesecker, L.G. (2016) Systematic Evaluation of Sanger Validation of Next-Generation Sequencing Variants. *Clin Chem*, **62**(4): 647–654.

- Becker, K.F., Atkinson, M.J., Reich, U., Becker, I., Nekarda, H., Siewert, J.R., and Hfler, H. (1994) E-cadherin gene mutations provide clues to diffuse type gastric carcinomas. *Cancer Research*, **54**(14): 3845–3852.
- Bell, D., Berchuck, A., Birrer, M., Chien, J., Cramer, D., Dao, F., Dhir, R., DiSaia, P., Gabra, H., Glenn, P., *et al.* (2011) Integrated genomic analyses of ovarian carcinoma. *Nature*, **474**(7353): 609–615.
- Benjamini, Y. and Hochberg, Y. (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B (Methodological)*, **57**(1): 289–300.
- Berx, G., Cleton-Jansen, A.M., Nollet, F., de Leeuw, W.J., van de Vijver, M., Cornelisse, C., and van Roy, F. (1995) E-cadherin is a tumour/invasion suppressor gene mutated in human lobular breast cancers. *EMBO J*, **14**(24): 6107–15.
- Berx, G., Cleton-Jansen, A.M., Strumane, K., de Leeuw, W.J., Nollet, F., van Roy, F., and Cornelisse, C. (1996) E-cadherin is inactivated in a majority of invasive human lobular breast cancers by truncation mutations throughout its extracellular domain. *Oncogene*, **13**(9): 1919–25.
- Berx, G. and van Roy, F. (2009) Involvement of members of the cadherin superfamily in cancer. *Cold Spring Harb Perspect Biol*, **1**: a003129.
- Bitler, B.G., Aird, K.M., Garipov, A., Li, H., Amatangelo, M., Kossenkov, A.V., Schultz, D.C., Liu, Q., Shih Ie, M., Conejo-Garcia, J.R., *et al.* (2015) Synthetic lethality by targeting ezh2 methyltransferase activity in arid1a-mutated cancers. *Nat Med*, **21**(3): 231–8.
- Blake, J.A., Christie, K.R., Dolan, M.E., Drabkin, H.J., Hill, D.P., Ni, L., Sitnikov, D., Burgess, S., Buza, T., Gresham, C., *et al.* (2015) Gene Ontology Consortium: going forward. *Nucleic Acids Res*, **43**(Database issue): D1049–1056.
- Boettcher, M., Lawson, A., Ladenburger, V., Fredebohm, J., Wolf, J., Hoheisel, J.D., Frezza, C., and Shlomi, T. (2014) High throughput synthetic lethality screen reveals a tumorigenic role of adenylate cyclase in fumarate hydratase-deficient cancer cells. *BMC Genomics*, **15**: 158.
- Boone, C., Bussey, H., and Andrews, B.J. (2007) Exploring genetic interactions and networks with yeast. *Nat Rev Genet*, **8**(6): 437–49.

- Borgatti, S.P. (2005) Centrality and network flow. *Social Networks*, **27**(1): 55 – 71.
- Boucher, B. and Jenna, S. (2013) Genetic interaction networks: better understand to better predict. *Front Genet*, **4**: 290.
- Breiman, L. (2001) Random forests. *Machine Learning*, **45**(1): 5–32.
- Brin, S. and Page, L. (1998) The anatomy of a large-scale hypertextual web search engine. *Computer Networks and ISDN Systems*, **30**(1): 107 – 117.
- Bryant, H.E., Schultz, N., Thomas, H.D., Parker, K.M., Flower, D., Lopez, E., Kyle, S., Meuth, M., Curtin, N.J., and Helleday, T. (2005) Specific killing of *BRCA2*-deficient tumours with inhibitors of polyadprribose polymerase. *Nature*, **434**(7035): 913–7.
- Burk, R.D., Chen, Z., Saller, C., Tarvin, K., Carvalho, A.L., Scapulatempo-Neto, C., Silveira, H.C., Fregnani, J.H., Creighton, C.J., Anderson, M.L., *et al.* (2017) Integrated genomic and molecular characterization of cervical cancer. *Nature*, **543**(7645): 378–384.
- Bussey, H., Andrews, B., and Boone, C. (2006) From worm genetic networks to complex human diseases. *Nat Genet*, **38**(8): 862–3.
- Butland, G., Babu, M., Diaz-Mejia, J.J., Bohdana, F., Phanse, S., Gold, B., Yang, W., Li, J., Gagarinova, A.G., Pogoutse, O., *et al.* (2008) esga: E. coli synthetic genetic array analysis. *Nat Methods*, **5**(9): 789–95.
- Cancer Research UK (2017) Family history and cancer genes. <http://www.cancerresearchuk.org/about-cancer/causes-of-cancer/inherited-cancer-genes-and-increased-cancer-risk/family-history-and-inherited-cancer-genes>. Accessed: 22/03/2017.
- cBioPortal for Cancer Genomics (cBioPortal) (2017) cBioPortal for Cancer Genomics. <http://www.cbioportal.org/>. Accessed: 26/03/2017.
- Cerami, E.G., Gross, B.E., Demir, E., Rodchenkov, I., Babur, O., Anwar, N., Schultz, N., Bader, G.D., and Sander, C. (2011) Pathway Commons, a web resource for biological pathway data. *Nucleic Acids Res*, **39**(Database issue): D685–690.
- Chen, A., Beetham, H., Black, M.A., Priya, R., Telford, B.J., Guest, J., Wiggins, G.A.R., Godwin, T.D., Yap, A.S., and Guilford, P.J. (2014) E-cadherin loss alters

- cytoskeletal organization and adhesion in non-malignant breast cells but is insufficient to induce an epithelial-mesenchymal transition. *BMC Cancer*, **14**(1): 552.
- Chen, K., Yang, D., Li, X., Sun, B., Song, F., Cao, W., Brat, D.J., Gao, Z., Li, H., Liang, H., *et al.* (2015) Mutational landscape of gastric adenocarcinoma in Chinese: implications for prognosis and therapy. *Proc Natl Acad Sci USA*, **112**(4): 1107–1112.
- Chen, S. and Parmigiani, G. (2007) Meta-analysis of BRCA1 and BRCA2 penetrance. *J Clin Oncol*, **25**(11): 1329–1333.
- Chen, X. and Tompa, M. (2010) Comparative assessment of methods for aligning multiple genome sequences. *Nat Biotechnol*, **28**(6): 567–572.
- Cherniack, A.D., Shen, H., Walter, V., Stewart, C., Murray, B.A., Bowlby, R., Hu, X., Ling, S., Soslow, R.A., Broaddus, R.R., *et al.* (2017) Integrated Molecular Characterization of Uterine Carcinosarcoma. *Cancer Cell*, **31**(3): 411–423.
- Chipman, K. and Singh, A. (2009) Predicting genetic interactions with random walks on biological networks. *BMC Bioinformatics*, **10**(1): 17.
- Christofori, G. and Semb, H. (1999) The role of the cell-adhesion molecule E-cadherin as a tumour-suppressor gene. *Trends in Biochemical Sciences*, **24**(2): 73 – 76.
- Ciriello, G., Gatza, M.L., Beck, A.H., Wilkerson, M.D., Rhie, S.K., Pastore, A., Zhang, H., McLellan, M., Yau, C., Kandoth, C., *et al.* (2015) Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. *Cell*, **163**(2): 506–519.
- Clark, M.J. (2004) Endogenous Regulator of G Protein Signaling Proteins Suppress G o-Dependent μ -Opioid Agonist-Mediated Adenylyl Cyclase Supersensitization. *Journal of Pharmacology and Experimental Therapeutics*, **310**(1): 215–222.
- Clough, E. and Barrett, T. (2016) The Gene Expression Omnibus Database. *Methods Mol Biol*, **1418**: 93–110.
- Collingridge, D.S. (2013) A primer on quantitized data analysis and permutation testing. *Journal of Mixed Methods Research*, **7**(1): 81–97.
- Collins, F.S. and Barker, A.D. (2007) Mapping the cancer genome. Pinpointing the genes involved in cancer will help chart a new course across the complex landscape of human malignancies. *Sci Am*, **296**(3): 50–57.

- Collins, F.S., Morgan, M., and Patrinos, A. (2003) The Human Genome Project: lessons from large-scale biology. *Science*, **300**(5617): 286–290.
- Collisson, E., Campbell, J., Brooks, A., Berger, A., Lee, W., Chmielecki, J., Beer, D., Cope, L., Creighton, C., Danilova, L., *et al.* (2014) Comprehensive molecular profiling of lung adenocarcinoma. *Nature*, **511**(7511): 543–550.
- Corcoran, R.B., Ebi, H., Turke, A.B., Coffee, E.M., Nishino, M., Cogdill, A.P., Brown, R.D., Della Pelle, P., Dias-Santagata, D., Hung, K.E., *et al.* (2012) Egfr-mediated re-activation of mapk signaling contributes to insensitivity of *BRAF*-mutant colorectal cancers to raf inhibition with vemurafenib. *Cancer Discovery*, **2**(3): 227–235.
- Costanzo, M., Baryshnikova, A., Bellay, J., Kim, Y., Spear, E.D., Sevier, C.S., Ding, H., Koh, J.L., Toufighi, K., Mostafavi, S., *et al.* (2010) The genetic landscape of a cell. *Science*, **327**(5964): 425–31.
- Costanzo, M., Baryshnikova, A., Myers, C.L., Andrews, B., and Boone, C. (2011) Charting the genetic interaction map of a cell. *Curr Opin Biotechnol*, **22**(1): 66–74.
- Courtney, K.D., Corcoran, R.B., and Engelman, J.A. (2010) The PI3K pathway as drug target in human cancer. *J Clin Oncol*, **28**(6): 1075–1083.
- Creighton, C.J., Morgan, M., Gunaratne, P.H., Wheeler, D.A., Gibbs, R.A., Robertson, A., Chu, A., Beroukhim, R., Cibulskis, K., Signoretti, S., *et al.* (2013) Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature*, **499**(7456): 43–49.
- Croft, D., Mundo, A.F., Haw, R., Milacic, M., Weiser, J., Wu, G., Caudy, M., Garapati, P., Gillespie, M., Kamdar, M.R., *et al.* (2014) The Reactome pathway knowledge-base. *Nucleic Acids Res*, **42**(database issue): D472D477.
- Crunkhorn, S. (2014) Cancer: Predicting synthetic lethal interactions. *Nat Rev Drug Discov*, **13**(11): 812.
- Csardi, G. and Nepusz, T. (2006) The igraph software package for complex network research. *InterJournal*, **Complex Systems**: 1695.
- Curtis, C., Shah, S.P., Chin, S.F., Turashvili, G., Rueda, O.M., Dunning, M.J., Speed, D., Lynch, A.G., Samarajiwa, S., Yuan, Y., *et al.* (2012) The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature*, **486**(7403): 346–352.

- Dai, X., Li, T., Bai, Z., Yang, Y., Liu, X., Zhan, J., and Shi, B. (2015) Breast cancer intrinsic subtype classification, clinical use and future trends. *Am J Cancer Res*, **5**(10): 2929–2943.
- Davierwala, A.P., Haynes, J., Li, Z., Brost, R.L., Robinson, M.D., Yu, L., Mnaimneh, S., Ding, H., Zhu, H., Chen, Y., *et al.* (2005) The synthetic genetic interaction spectrum of essential genes. *Nat Genet*, **37**(10): 1147–1152.
- De Leeuw, W.J., Berx, G., Vos, C.B., Peterse, J.L., Van de Vijver, M.J., Litvinov, S., Van Roy, F., Cornelisse, C.J., and Cleton-Jansen, A.M. (1997) Simultaneous loss of E-cadherin and catenins in invasive lobular breast cancer and lobular carcinoma in situ. *J Pathol*, **183**(4): 404–11.
- Demir, E., Babur, O., Rodchenkov, I., Aksoy, B.A., Fukuda, K.I., Gross, B., Sumer, O.S., Bader, G.D., and Sander, C. (2013) Using biological pathway data with Paxtools. *PLoS Comput Biol*, **9**(9): e1003194.
- Deshpande, R., Asiedu, M.K., Klebig, M., Sutor, S., Kuzmin, E., Nelson, J., Piotrowski, J., Shin, S.H., Yoshida, M., Costanzo, M., *et al.* (2013) A comparative genomic approach for identifying synthetic lethal interactions in human cancer. *Cancer Res*, **73**(20): 6128–36.
- Dickson, D. (1999) Wellcome funds cancer database. *Nature*, **401**(6755): 729.
- Dienstmann, R. and Tabernero, J. (2011) *BRAF* as a target for cancer therapy. *Anti-cancer Agents Med Chem*, **11**(3): 285–95.
- Dijkstra, E.W. (1959) A note on two problems in connexion with graphs. *Numerische Mathematik*, **1**(1): 269–271.
- Dixon, S.J., Andrews, B.J., and Boone, C. (2009) Exploring the conservation of synthetic lethal genetic interaction networks. *Commun Integr Biol*, **2**(2): 78–81.
- Dixon, S.J., Fedyszyn, Y., Koh, J.L., Prasad, T.S., Chahwan, C., Chua, G., Toufighi, K., Baryshnikova, A., Hayles, J., Hoe, K.L., *et al.* (2008) Significant conservation of synthetic lethal genetic interaction networks between distantly related eukaryotes. *Proc Natl Acad Sci U S A*, **105**(43): 16653–8.
- Dorogovtsev, S.N. and Mendes, J.F. (2003) *Evolution of networks: From biological nets to the Internet and WWW*. Oxford University Press, USA.

- Dorsam, R.T. and Gutkind, J.S. (2007) G-protein-coupled receptors and cancer. *Nat Rev Cancer*, **7**(2): 79–94.
- Erdős, P. and Rényi, A. (1959) On random graphs I. *Publ Math Debrecen*, **6**: 290–297.
- Erdős, P. and Rényi, A. (1960) On the evolution of random graphs. In *Publ. Math. Inst. Hung. Acad. Sci*, volume 5, 17–61.
- Eroles, P., Bosch, A., Perez-Fidalgo, J.A., and Lluch, A. (2012) Molecular biology in breast cancer: intrinsic subtypes and signaling pathways. *Cancer Treat Rev*, **38**(6): 698–707.
- Ezkurdia, I., Juan, D., Rodriguez, J.M., Frankish, A., Diekhans, M., Harrow, J., Vazquez, J., Valencia, A., and Tress, M.L. (2014) Multiple evidence strands suggest that there may be as few as 19 000 human protein-coding genes. *Human Molecular Genetics*, **23**(22): 5866.
- Farmer, H., McCabe, N., Lord, C.J., Tutt, A.N., Johnson, D.A., Richardson, T.B., Santarosa, M., Dillon, K.J., Hickson, I., Knights, C., *et al.* (2005) Targeting the dna repair defect in BRCA mutant cells as a therapeutic strategy. *Nature*, **434**(7035): 917–21.
- Fawcett, T. (2006) An introduction to ROC analysis. *Pattern Recognition Letters*, **27**(8): 861 – 874. {ROC} Analysis in Pattern Recognition.
- Fece de la Cruz, F., Gapp, B.V., and Nijman, S.M. (2015) Synthetic lethal vulnerabilities of cancer. *Annu Rev Pharmacol Toxicol*, **55**: 513–531.
- Ferlay, J., Soerjomataram, I., Dikshit, R., Eser, S., Mathers, C., Rebelo, M., Parkin, D.M., Forman, D., and Bray, F. (2015) Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*, **136**(5): E359–386.
- Fisher, R.A. (1919) Xv.the correlation between relatives on the supposition of mendelian inheritance. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, **52**(02): 399–433.
- Fong, P.C., Boss, D.S., Yap, T.A., Tutt, A., Wu, P., Mergui-Roelvink, M., Mortimer, P., Swaisland, H., Lau, A., O’Connor, M.J., *et al.* (2009) Inhibition of poly(adp-ribose) polymerase in tumors from BRCA mutation carriers. *N Engl J Med*, **361**(2): 123–34.

- Fong, P.C., Yap, T.A., Boss, D.S., Carden, C.P., Mergui-Roelvink, M., Gourley, C., De Greve, J., Lubinski, J., Shanley, S., Messiou, C., *et al.* (2010) Poly(adp)-ribose polymerase inhibition: frequent durable responses in BRCA carrier ovarian cancer correlating with platinum-free interval. *J Clin Oncol*, **28**(15): 2512–9.
- Forbes, S.A., Beare, D., Gunasekaran, P., Leung, K., Bindal, N., Boutselakis, H., Ding, M., Bamford, S., Cole, C., Ward, S., *et al.* (2015) COSMIC: exploring the world’s knowledge of somatic mutations in human cancer. *Nucleic Acids Res*, **43**(Database issue): D805–811.
- Fraser, A. (2004) Towards full employment: using RNAi to find roles for the redundant. *Oncogene*, **23**(51): 8346–52.
- Futreal, P.A., Coin, L., Marshall, M., Down, T., Hubbard, T., Wooster, R., Rahman, N., and Stratton, M.R. (2004) A census of human cancer genes. *Nat Rev Cancer*, **4**(3): 177–183.
- Futreal, P.A., Kasprzyk, A., Birney, E., Mullikin, J.C., Wooster, R., and Stratton, M.R. (2001) Cancer and genomics. *Nature*, **409**(6822): 850–852.
- Gao, B. and Roux, P.P. (2015) Translational control by oncogenic signaling pathways. *Biochimica et Biophysica Acta*, **1849**(7): 753–65.
- Gatza, M.L., Kung, H.N., Blackwell, K.L., Dewhirst, M.W., Marks, J.R., and Chi, J.T. (2011) Analysis of tumor environmental response and oncogenic pathway activation identifies distinct basal and luminal features in HER2-related breast tumor subtypes. *Breast Cancer Res*, **13**(3): R62.
- Gatza, M.L., Lucas, J.E., Barry, W.T., Kim, J.W., Wang, Q., Crawford, M.D., Datto, M.B., Kelley, M., Mathey-Prevot, B., Potti, A., *et al.* (2010) A pathway-based classification of human breast cancer. *Proc Natl Acad Sci USA*, **107**(15): 6994–6999.
- Gatza, M.L., Silva, G.O., Parker, J.S., Fan, C., and Perou, C.M. (2014) An integrated genomics approach identifies drivers of proliferation in luminal-subtype human breast cancer. *Nat Genet*, **46**(10): 1051–1059.
- Gentleman, R.C., Carey, V.J., Bates, D.M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B., Gautier, L., Ge, Y., Gentry, J., *et al.* (2004) Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol*, **5**(10): R80.

- Genz, A. and Bretz, F. (2009) Computation of multivariate normal and t probabilities. In *Lecture Notes in Statistics*, volume 195. Springer-Verlag, Heidelberg.
- Genz, A., Bretz, F., Miwa, T., Mi, X., Leisch, F., Scheipl, F., and Hothorn, T. (2016) *mvtnorm: Multivariate Normal and t Distributions*. R package version 1.0-5. URL.
- Gilbert, W. and Maxam, A. (1973) The nucleotide sequence of the lac operator. *Proceedings of the National Academy of Sciences*, **70**(12): 3581–3584.
- Git, A., Dvinge, H., Salmon-Divon, M., Osborne, M., Kutter, C., Hadfield, J., Bertone, P., and Caldas, C. (2010) Systematic comparison of microarray profiling, real-time PCR, and next-generation sequencing technologies for measuring differential microRNA expression. *RNA*, **16**(5): 991–1006.
- Globus (Globus) (2017) Research data management simplified. <https://www.globus.org/>. Accessed: 25/03/2017.
- Graziano, F., Humar, B., and Guilford, P. (2003) The role of the E-cadherin gene (*CDH1*) in diffuse gastric cancer susceptibility: from the laboratory to clinical practice. *Annals of Oncology*, **14**(12): 1705–1713.
- Güell, O., Sagus, F., and Serrano, M. (2014) Essential plasticity and redundancy of metabolism unveiled by synthetic lethality analysis. *PLoS Comput Biol*, **10**(5): e1003637.
- Guilford, P. (1999) E-cadherin downregulation in cancer: fuel on the fire? *Molecular Medicine Today*, **5**(4): 172 – 177.
- Guilford, P., Hopkins, J., Harraway, J., McLeod, M., McLeod, N., Harawira, P., Taite, H., Scoular, R., Miller, A., and Reeve, A.E. (1998) E-cadherin germline mutations in familial gastric cancer. *Nature*, **392**(6674): 402–5.
- Guilford, P., Humar, B., and Blair, V. (2010) Hereditary diffuse gastric cancer: translation of *CDH1* germline mutations into clinical practice. *Gastric Cancer*, **13**(1): 1–10.
- Guilford, P.J., Hopkins, J.B., Grady, W.M., Markowitz, S.D., Willis, J., Lynch, H., Rajput, A., Wiesner, G.L., Lindor, N.M., Burgart, L.J., *et al.* (1999) E-cadherin germline mutations define an inherited cancer syndrome dominated by diffuse gastric cancer. *Hum Mutat*, **14**(3): 249–55.

- Guo, J., Liu, H., and Zheng, J. (2016) SynLethDB: synthetic lethality database toward discovery of selective and sensitive anticancer drug targets. *Nucleic Acids Res*, **44**(D1): D1011–1017.
- Hajian-Tilaki, K. (2013) Receiver Operating Characteristic (ROC) Curve Analysis for Medical Diagnostic Test Evaluation. *Caspian J Intern Med*, **4**(2): 627–635.
- Hall, M., Frank, E., Holmes, G., Pfahringer, B., Reutemann, P., and Witten, I.H. (2009) The weka data mining software: an update. *SIGKDD Explor Newsl*, **11**(1): 10–18.
- Hammerman, P.S., Lawrence, M.S., Voet, D., Jing, R., Cibulskis, K., Sivachenko, A., Stojanov, P., McKenna, A., Lander, E.S., Gabriel, S., *et al.* (2012) Comprehensive genomic characterization of squamous cell lung cancers. *Nature*, **489**(7417): 519–525.
- Han, J.D.J., Bertin, N., Hao, T., Goldberg, D.S., Berriz, G.F., Zhang, L.V., Dupuy, D., Walhout, A.J.M., Cusick, M.E., Roth, F.P., *et al.* (2004) Evidence for dynamically organized modularity in the yeast protein-protein interaction network. *Nature*, **430**(6995): 88–93.
- Hanahan, D. and Weinberg, R.A. (2000) The hallmarks of cancer. *Cell*, **100**(1): 57–70.
- Hanahan, D. and Weinberg, R.A. (2011) Hallmarks of cancer: the next generation. *Cell*, **144**(5): 646–674.
- Hanna, S. (2003) Cancer incidence in new zealand (2003-2007). In D. Forman, D. Bray F Brewster, C. Gombe Mbalawa, B. Kohler, M. Piñeros, E. Steliarova-Foucher, R. Swaminathan, and J. Ferlay (editors), *Cancer Incidence in Five Continents*, volume X, 902–907. International Agency for Research on Cancer, Lyon, France. Electronic version <http://ci5.iarc.fr> Accessed 22/03/2017.
- Heiskanen, M., Bian, X., Swan, D., and Basu, A. (2014) caArray microarray database in the cancer biomedical informatics gridTM (caBIGTM). *Cancer Research*, **67**(9 Supplement): 3712–3712.
- Heiskanen, M.A. and Aittokallio, T. (2012) Mining high-throughput screens for cancer drug targets-lessons from yeast chemical-genomic profiling and synthetic lethality. *Wiley Interdisciplinary Reviews: Data Mining and Knowledge Discovery*, **2**(3): 263–272.

- Hell, P. (1976) Graphs with given neighbourhoods i. problèmes combinatorics at theorie des graphes. *Proc Coil Int CNRS, Orsay*, **260**: 219–223.
- Herschkowitz, J.I., Simin, K., Weigman, V.J., Mikaelian, I., Usary, J., Hu, Z., Rasmussen, K.E., Jones, L.P., Assefnia, S., Chandrasekharan, S., *et al.* (2007) Identification of conserved gene expression features between murine mammary carcinoma models and human breast tumors. *Genome Biol*, **8**(5): R76.
- Hillenmeyer, M.E. (2008) The chemical genomic portrait of yeast: uncovering a phenotype for all genes. *Science*, **320**: 362–365.
- Hoadley, K.A., Yau, C., Wolf, D.M., Cherniack, A.D., Tamborero, D., Ng, S., Leiserson, M.D., Niu, B., McLellan, M.D., Uzunangelov, V., *et al.* (2014) Multiplatform analysis of 12 cancer types reveals molecular classification within and across tissues of origin. *Cell*, **158**(4): 929–944.
- Hoehndorf, R., Hardy, N.W., Osumi-Sutherland, D., Tweedie, S., Schofield, P.N., and Gkoutos, G.V. (2013) Systematic analysis of experimental phenotype data reveals gene functions. *PLoS ONE*, **8**(4): e60847.
- Holm, S. (1979) A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, **6**(2): 65–70.
- Holme, P. and Kim, B.J. (2002) Growing scale-free networks with tunable clustering. *Physical Review E*, **65**(2): 026107.
- Hopkins, A.L. (2008) Network pharmacology: the next paradigm in drug discovery. *Nat Chem Biol*, **4**(11): 682–690.
- Hu, Z., Fan, C., Oh, D.S., Marron, J.S., He, X., Qaqish, B.F., Livasy, C., Carey, L.A., Reynolds, E., Dressler, L., *et al.* (2006) The molecular portraits of breast tumors are conserved across microarray platforms. *BMC Genomics*, **7**: 96.
- Huang, E., Cheng, S., Dressman, H., Pittman, J., Tsou, M., Horng, C., Bild, A., Iversen, E., Liao, M., Chen, C., *et al.* (2003) Gene expression predictors of breast cancer outcomes. *Lancet*, **361**: 1590–1596.
- Illumina, Inc (Illumina) (2017) Sequencing and array-based solutions for genetic research. <https://www.illumina.com/>. Accessed: 26/03/2017.

- International HapMap 3 Consortium (HapMap) (2003) The International HapMap Project. *Nature*, **426**(6968): 789–796.
- International Human Genome Sequencing Consortium (IHGSC) (2004) Finishing the euchromatic sequence of the human genome. *Nature*, **431**(7011): 931–945.
- Jerby-Arnon, L., Pfetzer, N., Waldman, Y., McGarry, L., James, D., Shanks, E., Seashore-Ludlow, B., Weinstock, A., Geiger, T., Clemons, P., *et al.* (2014) Predicting cancer-specific vulnerability via data-driven detection of synthetic lethality. *Cell*, **158**(5): 1199–1209.
- Joachims, T. (1999) Making large-scale support vector machine learning practical. In S. Bernhard, I. Kopr, J.C.B. Christopher, and J.S. Alexander (editors), *Advances in kernel methods*, 169–184. MIT Press.
- Ju, Z., Liu, W., Roebuck, P.L., Siwak, D.R., Zhang, N., Lu, Y., Davies, M.A., Akbani, R., Weinstein, J.N., Mills, G.B., *et al.* (2015) Development of a robust classifier for quality control of reverse-phase protein arrays. *Bioinformatics*, **31**(6): 912.
- Kaelin, Jr, W. (2005) The concept of synthetic lethality in the context of anticancer therapy. *Nat Rev Cancer*, **5**(9): 689–98.
- Kaelin, Jr, W. (2009) Synthetic lethality: a framework for the development of wiser cancer therapeutics. *Genome Med*, **1**: 99.
- Kakiuchi, M., Nishizawa, T., Ueda, H., Gotoh, K., Tanaka, A., Hayashi, A., Yamamoto, S., Tatsuno, K., Katoh, H., Watanabe, Y., *et al.* (2014) Recurrent gain-of-function mutations of RHOA in diffuse-type gastric carcinoma. *Nat Genet*, **46**(6): 583–587.
- Kamada, T. and Kawai, S. (1989) An algorithm for drawing general undirected graphs. *Information Processing Letters*, **31**(1): 7–15.
- Kandoth, C., Schultz, N., Cherniack, A.D., Akbani, R., Liu, Y., Shen, H., Robertson, A.G., Pashtan, I., Shen, R., Benz, C.C., *et al.* (2013) Integrated genomic characterization of endometrial carcinoma. *Nature*, **497**(7447): 67–73.
- Kawai, J., Shinagawa, A., Shibata, K., Yoshino, M., Itoh, M., Ishii, Y., Arakawa, T., Hara, A., Fukunishi, Y., Konno, H., *et al.* (2001) Functional annotation of a full-length mouse cDNA collection. *Nature*, **409**(6821): 685–690.

- Kelley, R. and Ideker, T. (2005) Systematic interpretation of genetic interactions using protein networks. *Nat Biotech*, **23**(5): 561–566.
- Kelly, S., Chen, A., Guilford, P., and Black, M. (2017a) Synthetic lethal interaction prediction of target pathways in E-cadherin deficient breast cancers. Submitted to *BMC Genomics*.
- Kelly, S.T. (2013) *Statistical Predictions of Synthetic Lethal Interactions in Cancer*. Dissertation, University of Otago.
- Kelly, S.T., Single, A.B., Telford, B.J., Beetham, H.G., Godwin, T.D., Chen, A., Black, M.A., and Guilford, P.J. (2017b) Towards HDGC chemoprevention: vulnerabilities in E-cadherin-negative cells identified by genome-wide interrogation of isogenic cell lines and whole tumors. Submitted to *Cancer Prev Res*.
- Kozlov, K.N., Gursky, V.V., Kulakovskiy, I.V., and Samsonova, M.G. (2015) Sequence-based model of gap gene regulation network. *BMC Genomics*, **15**(Suppl 12): S6.
- Kranthi, S., Rao, S., and Manimaran, P. (2013) Identification of synthetic lethal pairs in biological systems through network information centrality. *Mol BioSyst*, **9**(8): 2163–2167.
- Lander, E.S. (2011) Initial impact of the sequencing of the human genome. *Nature*, **470**(7333): 187–197.
- Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., *et al.* (2001) Initial sequencing and analysis of the human genome. *Nature*, **409**(6822): 860–921.
- Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009) Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol*, **10**(3): R25.
- Latora, V. and Marchiori, M. (2001) Efficient behavior of small-world networks. *Phys Rev Lett*, **87**: 198701.
- Laufer, C., Fischer, B., Billmann, M., Huber, W., and Boutros, M. (2013) Mapping genetic interactions in human cancer cells with RNAi and multiparametric phenotyping. *Nat Methods*, **10**(5): 427–31.

- Law, C.W., Chen, Y., Shi, W., and Smyth, G.K. (2014) voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol*, **15**(2): R29.
- Lawrence, M.S., Sougnez, C., Lichtenstein, L., Cibulskis, K., Lander, E., Gabriel, S.B., Getz, G., Ally, A., Balasundaram, M., Birol, I., *et al.* (2015) Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*, **517**(7536): 576–582.
- Le Meur, N. and Gentleman, R. (2008) Modeling synthetic lethality. *Genome Biol*, **9**(9): R135.
- Le Meur, N., Jiang, Z., Liu, T., Mar, J., and Gentleman, R.C. (2014) Slgi: Synthetic lethal genetic interaction. r package version 1.26.0.
- Lee, A.Y., Perreault, R., Harel, S., Boulier, E.L., Suderman, M., Hallett, M., and Jenna, S. (2010a) Searching for signaling balance through the identification of genetic interactors of the rab guanine-nucleotide dissociation inhibitor gdi-1. *PLoS ONE*, **5**(5): e10624.
- Lee, I., Lehner, B., Vavouri, T., Shin, J., Fraser, A.G., and Marcotte, E.M. (2010b) Predicting genetic modifier loci using functional gene networks. *Genome Research*, **20**(8): 1143–1153.
- Lee, I. and Marcotte, E.M. (2009) Effects of functional bias on supervised learning of a gene network model. *Methods Mol Biol*, **541**: 463–75.
- Lee, M.J., Ye, A.S., Gardino, A.K., Heijink, A.M., Sorger, P.K., MacBeath, G., and Yaffe, M.B. (2012) Sequential application of anticancer drugs enhances cell death by rewiring apoptotic signaling networks. *Cell*, **149**(4): 780–94.
- Lehner, B., Crombie, C., Tischler, J., Fortunato, A., and Fraser, A.G. (2006) Systematic mapping of genetic interactions in *caenorhabditis elegans* identifies common modifiers of diverse signaling pathways. *Nat Genet*, **38**(8): 896–903.
- Li, X.J., Mishra, S.K., Wu, M., Zhang, F., and Zheng, J. (2014) Syn-lethality: An integrative knowledge base of synthetic lethality towards discovery of selective anticancer therapies. *Biomed Res Int*, **2014**: 196034.
- Linehan, W.M., Spellman, P.T., Ricketts, C.J., Creighton, C.J., Fei, S.S., Davis, C., Wheeler, D.A., Murray, B.A., Schmidt, L., Vocke, C.D., *et al.* (2016) Comprehen-

- sive Molecular Characterization of Papillary Renal-Cell Carcinoma. *N Engl J Med*, **374**(2): 135–145.
- Lokody, I. (2014) Computational modelling: A computational crystal ball. *Nature Reviews Cancer*, **14**(10): 649–649.
- Lord, C.J., Tutt, A.N., and Ashworth, A. (2015) Synthetic lethality and cancer therapy: lessons learned from the development of PARP inhibitors. *Annu Rev Med*, **66**: 455–470.
- Lu, X., Kensche, P.R., Huynen, M.A., and Notebaart, R.A. (2013) Genome evolution predicts genetic interactions in protein complexes and reveals cancer drug targets. *Nat Commun*, **4**: 2124.
- Lu, X., Megchelenbrink, W., Notebaart, R.A., and Huynen, M.A. (2015) Predicting human genetic interactions from cancer genome evolution. *PLoS One*, **10**(5): e0125795.
- Lum, P.Y., Armour, C.D., Stepaniants, S.B., Cavet, G., Wolf, M.K., Butler, J.S., Hinshaw, J.C., Garnier, P., Prestwich, G.D., Leonardson, A., *et al.* (2004) Discovering modes of action for therapeutic compounds using a genome-wide screen of yeast heterozygotes. *Cell*, **116**(1): 121–137.
- Luo, J., Solimini, N.L., and Elledge, S.J. (2009) Principles of Cancer Therapy: Oncogene and Non-oncogene Addiction. *Cell*, **136**(5): 823–837.
- Machado, J., Olivera, C., Carvalh, R., Soares, P., Berx, G., Caldas, C., Sercuca, R., Carneiro, F., and Sorbrinho-Simoes, M. (2001) E-cadherin gene (*CDH1*) promoter methylation as the second hit in sporadic diffuse gastric carcinoma. *Oncogene*, **20**: 1525–1528.
- Masciari, S., Larsson, N., Senz, J., Boyd, N., Kaurah, P., Kandel, M.J., Harris, L.N., Pinheiro, H.C., Troussard, A., Miron, P., *et al.* (2007) Germline E-cadherin mutations in familial lobular breast cancer. *J Med Genet*, **44**(11): 726–31.
- Mattison, J., van der Weyden, L., Hubbard, T., and Adams, D.J. (2009) Cancer gene discovery in mouse and man. *Biochim Biophys Acta*, **1796**(2): 140–161.
- Maxam, A.M. and Gilbert, W. (1977) A new method for sequencing DNA. *Proceedings of the National Academy of Science*, **74**(2): 560–564.

- McCourt, C.M., McArt, D.G., Mills, K., Catherwood, M.A., Maxwell, P., Waugh, D.J., Hamilton, P., O’Sullivan, J.M., and Salto-Tellez, M. (2013) Validation of next generation sequencing technologies in comparison to current diagnostic gold standards for BRAF, EGFR and KRAS mutational analysis. *PLoS ONE*, **8**(7): e69604.
- McLachlan, J., George, A., and Banerjee, S. (2016) The current status of parp inhibitors in ovarian cancer. *Tumori*, **102**(5): 433–440.
- McLendon, R., Friedman, A., Bigner, D., Van Meir, E.G., Brat, D.J., Mastrogiannis, G.M., Olson, J.J., Mikkelsen, T., Lehman, N., Aldape, K., *et al.* (2008) Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature*, **455**(7216): 1061–1068.
- Miles, D.W. (2001) Update on HER-2 as a target for cancer therapy: herceptin in the clinical setting. *Breast Cancer Res*, **3**(6): 380–384.
- Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L., and Wold, B. (2008) Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods*, **5**(7): 621–628.
- Muzny, D.M., Bainbridge, M.N., Chang, K., Dinh, H.H., Drummond, J.A., Fowler, G., Kovar, C.L., Lewis, L.R., Morgan, M.B., Newsham, I.F., *et al.* (2012) Comprehensive molecular characterization of human colon and rectal cancer. *Nature*, **487**(7407): 330–337.
- Nagalla, S., Chou, J.W., Willingham, M.C., Ruiz, J., Vaughn, J.P., Dubey, P., Lash, T.L., Hamilton-Dutoit, S.J., Bergh, J., Sotiriou, C., *et al.* (2013) Interactions between immunity, proliferation and molecular subtype in breast cancer prognosis. *Genome Biol*, **14**(4): R34.
- Neeley, E.S., Kornblau, S.M., Coombes, K.R., and Baggerly, K.A. (2009) Variable slope normalization of reverse phase protein arrays. *Bioinformatics*, **25**(11): 1384.
- Novomestky, F. (2012) *matrixcalc: Collection of functions for matrix calculations*. R package version 1.0-3.
- Oliveira, C., Senz, J., Kaurah, P., Pinheiro, H., Sanges, R., Haegert, A., Corso, G., Schouten, J., Fitzgerald, R., Vogelsang, H., *et al.* (2009) Germline *CDH1* deletions in hereditary diffuse gastric cancer families. *Human Molecular Genetics*, **18**(9): 1545–1555.

- Oliveira, C., Seruca, R., Hoogerbrugge, N., Ligtenberg, M., and Carneiro, F. (2013) Clinical utility gene card for: Hereditary diffuse gastric cancer (HDGC). *Eur J Hum Genet*, **21**(8).
- Pandey, G., Zhang, B., Chang, A.N., Myers, C.L., Zhu, J., Kumar, V., and Schadt, E.E. (2010) An integrative multi-network and multi-classifier approach to predict genetic interactions. *PLoS Comput Biol*, **6**(9).
- Parker, J., Mullins, M., Cheung, M., Leung, S., Voduc, D., Vickery, T., Davies, S., Fauron, C., He, X., Hu, Z., *et al.* (2009) Supervised risk predictor of breast cancer based on intrinsic subtypes. *Journal of Clinical Oncology*, **27**(8): 1160–1167.
- Peltonen, L. and McKusick, V.A. (2001) Genomics and medicine. Dissecting human disease in the postgenomic era. *Science*, **291**(5507): 1224–1229.
- Pereira, B., Chin, S.F., Rueda, O.M., Vollan, H.K., Provenzano, E., Bardwell, H.A., Pugh, M., Jones, L., Russell, R., Sammut, S.J., *et al.* (2016) Erratum: The somatic mutation profiles of 2,433 breast cancers refine their genomic and transcriptomic landscapes. *Nat Commun*, **7**: 11908.
- Perou, C.M., Sørlie, T., Eisen, M.B., van de Rijn, M., Jeffrey, S.S., Rees, C.A., Pollack, J.R., Ross, D.T., Johnsen, H., Akslen, L.A., *et al.* (2000) Molecular portraits of human breast tumours. *Nature*, **406**(6797): 747–752.
- Pleasance, E.D., Cheetham, R.K., Stephens, P.J., McBride, D.J., Humphray, S.J., Greenman, C.D., Varela, I., Lin, M.L., Ordonez, G.R., Bignell, G.R., *et al.* (2010) A comprehensive catalogue of somatic mutations from a human cancer genome. *Nature*, **463**(7278): 191–196.
- Polyak, K. and Weinberg, R.A. (2009) Transitions between epithelial and mesenchymal states: acquisition of malignant and stem cell traits. *Nat Rev Cancer*, **9**(4): 265–73.
- Prahallad, A., Sun, C., Huang, S., Di Nicolantonio, F., Salazar, R., Zecchin, D., Beijersbergen, R.L., Bardelli, A., and Bernards, R. (2012) Unresponsiveness of colon cancer to *BRAF*(v600e) inhibition through feedback activation of egfr. *Nature*, **483**(7387): 100–3.
- R Core Team (2016) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. R version 3.3.2.

- Ravnan, M.C. and Matalka, M.S. (2012) Vemurafenib in patients with *BRAF* v600e mutation-positive advanced melanoma. *Clin Ther*, **34**(7): 1474–86.
- Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research*, **43**(7): e47.
- Robin, J.D., Ludlow, A.T., LaRanger, R., Wright, W.E., and Shay, J.W. (2016) Comparison of DNA Quantification Methods for Next Generation Sequencing. *Sci Rep*, **6**: 24067.
- Robinson, M.D. and Oshlack, A. (2010) A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol*, **11**(3): R25.
- Roguev, A., Bandyopadhyay, S., Zofall, M., Zhang, K., Fischer, T., Collins, S.R., Qu, H., Shales, M., Park, H.O., Hayles, J., *et al.* (2008) Conservation and rewiring of functional modules revealed by an epistasis map in fission yeast. *Science*, **322**(5900): 405–10.
- Rung, J. and Brazma, A. (2013) Reuse of public genome-wide gene expression data. *Nat Rev Genet*, **14**(2): 89–99.
- Rustici, G., Kolesnikov, N., Brandizi, M., Burdett, T., Dylag, M., Emam, I., Farne, A., Hastings, E., Ison, J., Keays, M., *et al.* (2013) ArrayExpress update—trends in database growth and links to data analysis tools. *Nucleic Acids Res*, **41**(Database issue): D987–990.
- Ryan, C., Lord, C., and Ashworth, A. (2014) Daisy: Picking synthetic lethals from cancer genomes. *Cancer Cell*, **26**(3): 306–308.
- Sander, J.D. and Joung, J.K. (2014) Crispr-cas systems for editing, regulating and targeting genomes. *Nat Biotechnol*, **32**(4): 347–55.
- Sanger, F. and Coulson, A. (1975) A rapid method for determining sequences in dna by primed synthesis with dna polymerase. *Journal of Molecular Biology*, **94**(3): 441 – 448.
- Scheuer, L., Kauff, N., Robson, M., Kelly, B., Barakat, R., Satagopan, J., Ellis, N., Hensley, M., Boyd, J., Borgen, P., *et al.* (2002) Outcome of preventive surgery and screening for breast and ovarian cancer in BRCA mutation carriers. *J Clin Oncol*, **20**(5): 1260–1268.

- Semb, H. and Christofori, G. (1998) The tumor-suppressor function of E-cadherin. *Am J Hum Genet*, **63**(6): 1588–93.
- Sing, T., Sander, O., Beerenwinkel, N., and Lengauer, T. (2005) Rocr: visualizing classifier performance in r. *Bioinformatics*, **21**(20): 7881.
- Slurm development team (Slurm) (2017) Slurm workload manager. <https://slurm.schedmd.com/>. Accessed: 25/03/2017.
- Sørbye, T., Perou, C.M., Tibshirani, R., Aas, T., Geisler, S., Johnsen, H., Hastie, T., Eisen, M.B., van de Rijn, M., Jeffrey, S.S., *et al.* (2001) Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci USA*, **98**(19): 10869–10874.
- Stajich, J.E. and Lapp, H. (2006) Open source tools and toolkits for bioinformatics: significance, and where are we? *Brief Bioinformatics*, **7**(3): 287–296.
- Stratton, M.R., Campbell, P.J., and Futreal, P.A. (2009) The cancer genome. *Nature*, **458**(7239): 719–724.
- Ström, C. and Helleday, T. (2012) Strategies for the use of poly(adenosine diphosphate ribose) polymerase (parp) inhibitors in cancer therapy. *Biomolecules*, **2**(4): 635–649.
- Sun, C., Wang, L., Huang, S., Heynen, G.J.J.E., Prahallad, A., Robert, C., Haanen, J., Blank, C., Wesseling, J., Willems, S.M., *et al.* (2014) Reversible and adaptive resistance to *BRAF*(v600e) inhibition in melanoma. *Nature*, **508**(7494): 118–122.
- Taylor, I.W., Linding, R., Warde-Farley, D., Liu, Y., Pesquita, C., Faria, D., Bull, S., Pawson, T., Morris, Q., and Wrana, J.L. (2009) Dynamic modularity in protein interaction networks predicts breast cancer outcome. *Nat Biotechnol*, **27**(2): 199–204.
- Telford, B.J., Chen, A., Beetham, H., Frick, J., Brew, T.P., Gould, C.M., Single, A., Godwin, T., Simpson, K.J., and Guilford, P. (2015) Synthetic lethal screens identify vulnerabilities in gpcr signalling and cytoskeletal organization in E-cadherin-deficient cells. *Mol Cancer Ther*, **14**(5): 1213–1223.
- The 1000 Genomes Project Consortium (1000 Genomes) (2010) A map of human genome variation from population-scale sequencing. *Nature*, **467**(7319): 1061–1073.

- The Cancer Genome Atlas Research Network (TCGA) (2012) Comprehensive molecular portraits of human breast tumours. *Nature*, **490**(7418): 61–70.
- The Cancer Genome Atlas Research Network (TCGA) (2017a) The Cancer Genome Atlas Project. <https://cancergenome.nih.gov/>. Accessed: 26/03/2017.
- The Cancer Genome Atlas Research Network (TCGA) (2017b) The Cancer Genome Atlas Project Data Portal. <https://tcga-data.nci.nih.gov/>. Accessed: 06/02/2017 (via cBioPortal).
- The Cancer Society of New Zealand (Cancer Society of NZ) (2017) What is cancer? <https://otago-southland.cancernz.org.nz/en/cancer-information/other-links/what-is-cancer-3/>. Accessed: 22/03/2017.
- The Catalogue Of Somatic Mutations In Cancer (COSMIC) (2016) Cosmic: The catalogue of somatic mutations in cancer. <http://cancer.sanger.ac.uk/cosmic>. Release 79 (23/08/2016), Accessed: 05/02/2017.
- The Comprehensive R Archive Network (CRAN) (2017) Cran. <https://cran.r-project.org/>. Accessed: 24/03/2017.
- The ENCODE Project Consortium (ENCODE) (2004) The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science*, **306**(5696): 636–640.
- The International Cancer Genome Consortium (ICGC) (2017) ICGC Data Portal. <https://dcc.icgc.org/>. Accessed: 06/02/2017.
- The National Cancer Institute (NCI) (2015) The genetics of cancer. <https://www.cancer.gov/about-cancer/causes-prevention/genetics>. Published: 22/04/2015, Accessed: 22/03/2017.
- The New Zealand eScience Infrastructure (NeSI) (2017) NeSI. <https://www.nesi.org.nz/>. Accessed: 25/03/2017.
- The Pharmaceutical Management Agency (PHARMAC) (2016) Approval of multi-product funding proposal with roche.
- Tierney, L., Rossini, A.J., Li, N., and Sevcikova, H. (2015) *snow: Simple Network of Workstations*. R package version 0.4-2.

- Tiong, K.L., Chang, K.C., Yeh, K.T., Liu, T.Y., Wu, J.H., Hsieh, P.H., Lin, S.H., Lai, W.Y., Hsu, Y.C., Chen, J.Y., *et al.* (2014) Csnk1e/ctnnb1 are synthetic lethal to tp53 in colorectal cancer and are markers for prognosis. *Neoplasia*, **16**(5): 441–50.
- Tischler, J., Lehner, B., and Fraser, A.G. (2008) Evolutionary plasticity of genetic interaction networks. *Nat Genet*, **40**(4): 390–391.
- Tomasetti, C. and Vogelstein, B. (2015) Cancer etiology. Variation in cancer risk among tissues can be explained by the number of stem cell divisions. *Science*, **347**(6217): 78–81.
- Tong, A.H., Evangelista, M., Parsons, A.B., Xu, H., Bader, G.D., Page, N., Robinson, M., Raghibizadeh, S., Hogue, C.W., Bussey, H., *et al.* (2001) Systematic genetic analysis with ordered arrays of yeast deletion mutants. *Science*, **294**(5550): 2364–8.
- Tong, A.H., Lesage, G., Bader, G.D., Ding, H., Xu, H., Xin, X., Young, J., Berriz, G.F., Brost, R.L., Chang, M., *et al.* (2004) Global mapping of the yeast genetic interaction network. *Science*, **303**(5659): 808–13.
- Travers, J. and Milgram, S. (1969) An experimental study of the small world problem. *Sociometry*, **32**(4): 425–443.
- Tsai, H.C., Li, H., Van Neste, L., Cai, Y., Robert, C., Rassool, F.V., Shin, J.J., Harbom, K.M., Beaty, R., Pappou, E., *et al.* (2012) Transient low doses of dna-demethylating agents exert durable antitumor effects on hematological and epithelial tumor cells. *Cancer Cell*, **21**(3): 430–46.
- Tutt, A., Robson, M., Garber, J.E., Domchek, S.M., Audeh, M.W., Weitzel, J.N., Friedlander, M., Arun, B., Loman, N., Schmutzler, R.K., *et al.* (2010) Oral poly(adp-ribose) polymerase inhibitor olaparib in patients with *BRCA1* or *BRCA2* mutations and advanced breast cancer: a proof-of-concept trial. *Lancet*, **376**(9737): 235–44.
- van der Meer, R., Song, H.Y., Park, S.H., Abdulkadir, S.A., and Roh, M. (2014) RNAi screen identifies a synthetic lethal interaction between PIM1 overexpression and PLK1 inhibition. *Clinical Cancer Research*, **20**(12): 3211–3221.
- van Steen, K. (2012) Travelling the world of genegene interactions. *Briefings in Bioinformatics*, **13**(1): 1–19.
- van Steen, M. (2010) *Graph Theory and Complex Networks: An Introduction*. Maarten van Steen, VU Amsterdam.

- Vapnik, V.N. (1995) *The nature of statistical learning theory*. Springer-Verlag New York, Inc.
- Vargas, J.J., Gusella, G., Najfeld, V., Klotman, M., and Cara, A. (2004) Novel integrase-defective lentiviral episomal vectors for gene transfer. *Hum Gene Ther*, **15**: 361–372.
- Vizeacoumar, F.J., Arnold, R., Vizeacoumar, F.S., Chandrashekhar, M., Buzina, A., Young, J.T., Kwan, J.H., Sayad, A., Mero, P., Lawo, S., *et al.* (2013) A negative genetic interaction map in isogenic cancer cell lines reveals cancer cell vulnerabilities. *Mol Syst Biol*, **9**: 696.
- Vogelstein, B., Papadopoulos, N., Velculescu, V.E., Zhou, S., Diaz, L.A., and Kinzler, K.W. (2013) Cancer genome landscapes. *Science*, **339**(6127): 1546–1558.
- Vos, C.B., Cleton-Jansen, A.M., Berx, G., de Leeuw, W.J., ter Haar, N.T., van Roy, F., Cornelisse, C.J., Peterse, J.L., and van de Vijver, M.J. (1997) E-cadherin inactivation in lobular carcinoma in situ of the breast: an early event in tumorigenesis. *Br J Cancer*, **76**(9): 1131–3.
- Wang, K., Singh, D., Zeng, Z., Coleman, S.J., Huang, Y., Savich, G.L., He, X., Mieczkowski, P., Grimm, S.A., Perou, C.M., *et al.* (2010) MapSplice: accurate mapping of RNA-seq reads for splice junction discovery. *Nucleic Acids Res*, **38**(18): e178.
- Wang, K., Yuen, S.T., Xu, J., Lee, S.P., Yan, H.H., Shi, S.T., Siu, H.C., Deng, S., Chu, K.M., Law, S., *et al.* (2014) Whole-genome sequencing and comprehensive molecular profiling identify new driver mutations in gastric cancer. *Nat Genet*, **46**(6): 573–582.
- Wang, X. and Simon, R. (2013) Identification of potential synthetic lethal genes to p53 using a computational biology approach. *BMC Medical Genomics*, **6**(1): 30.
- Wappett, M. (2014) Bisep: Toolkit to identify candidate synthetic lethality. r package version 2.0.
- Wappett, M., Dulak, A., Yang, Z.R., Al-Watban, A., Bradford, J.R., and Dry, J.R. (2016) Multi-omic measurement of mutually exclusive loss-of-function enriches for candidate synthetic lethal gene pairs. *BMC Genomics*, **17**: 65.

- Warnes, G.R., Bolker, B., Bonebakker, L., Gentleman, R., Liaw, W.H.A., Lumley, T., Maechler, M., Magnusson, A., Moeller, S., Schwartz, M., *et al.* (2015) *gplots: Various R Programming Tools for Plotting Data*. R package version 2.17.0.
- Watts, D.J. and Strogatz, S.H. (1998) Collective dynamics of 'small-world' networks. *Nature*, **393**(6684): 440–2.
- Weinstein, I.B. (2000) Disorders in cell circuitry during multistage carcinogenesis: the role of homeostasis. *Carcinogenesis*, **21**(5): 857–864.
- Weinstein, J.N., Akbani, R., Broom, B.M., Wang, W., Verhaak, R.G., McConkey, D., Lerner, S., Morgan, M., Creighton, C.J., Smith, C., *et al.* (2014) Comprehensive molecular characterization of urothelial bladder carcinoma. *Nature*, **507**(7492): 315–322.
- Weinstein, J.N., Collisson, E.A., Mills, G.B., Shaw, K.R., Ozenberger, B.A., Ellrott, K., Shmulevich, I., Sander, C., Stuart, J.M., Chang, K., *et al.* (2013) The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet*, **45**(10): 1113–1120.
- Wickham, H. and Chang, W. (2016) *devtools: Tools to Make Developing R Packages Easier*. R package version 1.12.0.
- Wickham, H., Danenberg, P., and Eugster, M. (2017) *roxygen2: In-Line Documentation for R*. R package version 6.0.1.
- Wong, S.L., Zhang, L.V., Tong, A.H.Y., Li, Z., Goldberg, D.S., King, O.D., Lesage, G., Vidal, M., Andrews, B., Bussey, H., *et al.* (2004) Combining biological networks to predict genetic interactions. *Proceedings of the National Academy of Sciences of the United States of America*, **101**(44): 15682–15687.
- World Health Organization (WHO) (2017) Fact sheet: Cancer. <http://www.who.int/mediacentre/factsheets/fs297/en/>. Updated February 2017, Accessed: 22/03/2017.
- Wu, M., Li, X., Zhang, F., Li, X., Kwoh, C.K., and Zheng, J. (2014) In silico prediction of synthetic lethality by meta-analysis of genetic interactions, functions, and pathways in yeast and human cancer. *Cancer Inform*, **13**(Suppl 3): 71–80.
- Yu, H. (2002) Rmpi: Parallel statistical computing in r. *R News*, **2**(2): 10–14.

- Zhang, F., Wu, M., Li, X.J., Li, X.L., Kwoh, C.K., and Zheng, J. (2015) Predicting essential genes and synthetic lethality via influence propagation in signaling pathways of cancer cell fates. *J Bioinform Comput Biol*, **13**(3): 1541002.
- Zhang, J., Baran, J., Cros, A., Guberman, J.M., Haider, S., Hsu, J., Liang, Y., Rivkin, E., Wang, J., Whitty, B., *et al.* (2011) International cancer genome consortium data portala one-stop shop for cancer genomics data. *Database: The Journal of Biological Databases and Curation*, **2011**: bar026.
- Zhong, W. and Sternberg, P.W. (2006) Genome-wide prediction of c. elegans genetic interactions. *Science*, **311**(5766): 1481–1484.
- Zweig, M.H. and Campbell, G. (1993) Receiver-operating characteristic (roc) plots: a fundamental evaluation tool in clinical medicine. *Clinical Chemistry*, **39**(4): 561–577.

Appendix C

Mutation Analysis in Breast Cancer

C.1 Synthetic Lethal Genes and Pathways

SLIPT expression analysis (described in Section 3.1) on TCGA breast cancer data ($n = 969$) found the following genes and pathways, described in sections 4.1 and 4.1.1.

Table C.1: Candidate synthetic lethal gene partners of *CDH1* from mtSLIPT

| Gene | Observed | Expected | χ^2 value | p-value | p-value (FDR) |
|-----------------|----------|----------|----------------|------------------------|------------------------|
| <i>TFAP2B</i> | 8 | 36.7 | 89.5 | 3.60×10^{-20} | 8.37×10^{-17} |
| <i>ZNF423</i> | 15 | 36.7 | 78.8 | 7.89×10^{-18} | 1.22×10^{-14} |
| <i>CALCOCO1</i> | 11 | 36.7 | 76.8 | 2.09×10^{-17} | 2.59×10^{-14} |
| <i>RBM5</i> | 13 | 36.7 | 75.7 | 3.65×10^{-17} | 4.00×10^{-14} |
| <i>BTG2</i> | 7 | 36.7 | 71.7 | 2.72×10^{-16} | 1.81×10^{-13} |
| <i>RXRA</i> | 6 | 36.7 | 70.5 | 5.00×10^{-16} | 2.97×10^{-13} |
| <i>SLC27A1</i> | 11 | 36.7 | 70.3 | 5.42×10^{-16} | 2.97×10^{-13} |
| <i>MEF2D</i> | 12 | 36.7 | 69.6 | 7.86×10^{-16} | 3.95×10^{-13} |
| <i>NISCH</i> | 12 | 36.7 | 69.6 | 7.86×10^{-16} | 3.95×10^{-13} |
| <i>AVPR2</i> | 9 | 36.7 | 69.2 | 9.36×10^{-16} | 4.58×10^{-13} |
| <i>CRY2</i> | 13 | 36.7 | 68.9 | 1.07×10^{-15} | 4.98×10^{-13} |
| <i>RAPGEF3</i> | 13 | 36.7 | 68.9 | 1.07×10^{-15} | 4.98×10^{-13} |
| <i>NRIP2</i> | 10 | 36.7 | 68.2 | 1.58×10^{-15} | 7.18×10^{-13} |
| <i>DARC</i> | 12 | 36.7 | 66.4 | 3.76×10^{-15} | 1.54×10^{-12} |
| <i>SFRS5</i> | 12 | 36.7 | 66.4 | 3.76×10^{-15} | 1.54×10^{-12} |
| <i>NOSTRIN</i> | 5 | 36.7 | 65.1 | 7.40×10^{-15} | 2.70×10^{-12} |
| <i>KIF13B</i> | 12 | 36.7 | 63.4 | 1.69×10^{-14} | 5.16×10^{-12} |
| <i>TENC1</i> | 10 | 36.7 | 62.5 | 2.67×10^{-14} | 7.40×10^{-12} |
| <i>MFAP4</i> | 12 | 36.7 | 60.5 | 7.17×10^{-14} | 1.67×10^{-11} |
| <i>ELN</i> | 13 | 36.7 | 59.7 | 1.07×10^{-13} | 2.32×10^{-11} |
| <i>SGK223</i> | 14 | 36.7 | 59 | 1.51×10^{-13} | 3.05×10^{-11} |
| <i>KIF12</i> | 11 | 36.7 | 58.8 | 1.74×10^{-13} | 3.34×10^{-11} |
| <i>SELP</i> | 11 | 36.7 | 58.8 | 1.74×10^{-13} | 3.34×10^{-11} |
| <i>CIRBP</i> | 9 | 36.7 | 58.7 | 1.83×10^{-13} | 3.41×10^{-11} |
| <i>CTDSP1</i> | 9 | 36.7 | 58.7 | 1.83×10^{-13} | 3.41×10^{-11} |

Strongest candidate SL partners for *CDH1* by mtSLIPT with observed and expected numbers of *CDH1* mutant The Cancer Genome Atlas (TCGA) breast tumours with low expression of partner genes.

Table C.2: Pathways for *CDH1* partners from mtSLIPT

| Pathways Over-represented | Pathway Size | SL Genes | p-value (FDR) |
|---|--------------|----------|------------------------|
| Eukaryotic Translation Elongation | 86 | 60 | 2.0×10^{-128} |
| Peptide chain elongation | 83 | 59 | 2.0×10^{-128} |
| Eukaryotic Translation Termination | 83 | 58 | 2.3×10^{-125} |
| Viral mRNA Translation | 81 | 57 | 2.5×10^{-124} |
| Nonsense Mediated Decay independent of the Exon Junction Complex | 88 | 59 | 8.6×10^{-124} |
| Nonsense-Mediated Decay | 103 | 61 | 5.2×10^{-117} |
| Nonsense Mediated Decay enhanced by the Exon Junction Complex | 103 | 61 | 5.2×10^{-117} |
| Formation of a pool of free 40S subunits | 93 | 58 | 1.6×10^{-116} |
| L13a-mediated translational silencing of Ceruloplasmin expression | 103 | 59 | 1.3×10^{-111} |
| 3' -UTR-mediated translational regulation | 103 | 59 | 1.3×10^{-111} |
| GTP hydrolysis and joining of the 60S ribosomal subunit | 104 | 59 | 6.2×10^{-111} |
| SRP-dependent cotranslational protein targeting to membrane | 104 | 58 | 2.9×10^{-108} |
| Eukaryotic Translation Initiation | 111 | 59 | 3.0×10^{-106} |
| Cap-dependent Translation Initiation | 111 | 59 | 3.0×10^{-106} |
| Influenza Viral RNA Transcription and Replication | 108 | 57 | 5.1×10^{-103} |
| Influenza Infection | 117 | 59 | 1.5×10^{-102} |
| Translation | 141 | 64 | 3.7×10^{-101} |
| Influenza Life Cycle | 112 | 57 | 1.4×10^{-100} |
| GPCR downstream signalling | 472 | 116 | 1.0×10^{-80} |
| Hemostasis | 422 | 105 | 1.4×10^{-78} |

Gene set over-representation analysis (hypergeometric test) for Reactome pathways in mtSLIPT partners for *CDH1*.

The genes and pathways identified in Tables C.1 and C.2 were derived from comparing the expression profiles of potential partners to the mutation status of *CDH1* (as shown in Figure 3.2). Thus the following analysis is only limited the samples for which TCGA provides both expression and somatic mutation data.

C.2 Synthetic Lethal Expression Profiles

Similar to the analysis of synthetic lethal partners against low *CDH1* expression in 4.1.2, the partners detected from *CDH1* mutation were also examined for their expression profiles and the pathway composition of gene clusters. Hierarchical clustering was performed on mtSLIPT partners for *CDH1* as showing in Figure C.1. Overrepresentation for Reactome pathways for each of the gene clusters identified is given in Table C.3.




Figure C.1: Synthetic lethal expression profiles of analysed samples. Gene expression profile heatmap (correlation distance) of all samples (separated by *CDH1* somatic mutation status) analysed in TCGA breast cancer dataset for gene expression of 3,743 candidate partners of E-cadherin (*CDH1*) from mtSLIPT prediction (with significant FDR adjusted $p < 0.05$). Deeply clustered, inter-correlated genes form several main groups, each containing genes that were SL candidates or toxic in an siRNA screen Telford *et al.* (2015). Clusters had different sample groups highly expressing the synthetic lethal candidates in *CDH1* mutant samples and often lowly expressing *CDH1* wildtype samples (which were not tested for), although many of the *CDH1* mutant samples had among the lowest *CDH1* expression. In contrast to the expression analysis the (predominantly *CDH1* wildtype) basal subtype and estrogen receptor negative samples have depleted expression among most candidate synthetic lethal partners.

Table C.3: Pathway composition for clusters of *CDH1* partners from mtSLIPT

| Pathways Over-represented in Cluster 1 | Pathway Size | Cluster Genes | p-value (FDR) |
|---|--------------|---------------|------------------------|
| Olfactory Signalling Pathway | 57 | 8 | 7.1×10^{-9} |
| Assembly of the primary cilium | 149 | 14 | 8.0×10^{-9} |
| Sphingolipid metabolism | 62 | 8 | 9.6×10^{-9} |
| Signalling by ERBB4 | 133 | 12 | 5.1×10^{-8} |
| PI3K Cascade | 65 | 7 | 4.9×10^{-7} |
| Circadian Clock | 33 | 5 | 4.9×10^{-7} |
| Nuclear signalling by ERBB4 | 34 | 5 | 4.9×10^{-7} |
| Intraflagellar transport | 35 | 5 | 4.9×10^{-7} |
| PI3K events in ERBB4 signalling | 87 | 8 | 4.9×10^{-7} |
| PIP3 activates AKT signalling | 87 | 8 | 4.9×10^{-7} |
| PI3K events in ERBB2 signalling | 87 | 8 | 4.9×10^{-7} |
| PI-3K cascade:FGFR1 | 87 | 8 | 4.9×10^{-7} |
| PI-3K cascade:FGFR2 | 87 | 8 | 4.9×10^{-7} |
| PI-3K cascade:FGFR3 | 87 | 8 | 4.9×10^{-7} |
| PI-3K cascade:FGFR4 | 87 | 8 | 4.9×10^{-7} |
| Deadenylation of mRNA | 22 | 4 | 5.6×10^{-7} |
| PI3K/AKT activation | 90 | 8 | 5.6×10^{-7} |
| Cargo trafficking to the periciliary membrane | 38 | 5 | 5.6×10^{-7} |
| Pathways Over-represented in Cluster 2 | Pathway Size | Cluster Genes | p-value (FDR) |
| G _{αs} signalling events | 83 | 19 | 5.1×10^{-25} |
| Extracellular matrix organization | 238 | 30 | 1.4×10^{-18} |
| Hemostasis | 422 | 46 | 2.7×10^{-16} |
| Aquaporin-mediated transport | 32 | 9 | 2.7×10^{-16} |
| Transcriptional regulation of white adipocyte differentiation | 56 | 11 | 1.7×10^{-15} |
| Degradation of the extracellular matrix | 102 | 15 | 1.7×10^{-15} |
| Integration of energy metabolism | 84 | 13 | 8.8×10^{-15} |
| GPCR downstream signalling | 472 | 48 | 2.8×10^{-14} |
| G _{αz} signalling events | 15 | 6 | 5.0×10^{-14} |
| Molecules associated with elastic fibres | 33 | 8 | 5.4×10^{-14} |
| Phase 1 - Functionalization of compounds | 67 | 11 | 5.6×10^{-14} |
| Platelet activation, signalling and aggregation | 179 | 20 | 5.6×10^{-14} |
| Vasopressin regulates renal water homeostasis via Aquaporins | 24 | 7 | 6.1×10^{-14} |
| Elastic fibre formation | 37 | 8 | $.03 \times 10^{-13}$ |
| Calmodulin induced events | 27 | 7 | 3.3×10^{-13} |
| CaM pathway | 27 | 7 | 3.3×10^{-13} |
| cGMP effects | 18 | 6 | 3.6×10^{-13} |
| G _{αi} signalling events | 167 | 18 | 6.3×10^{-13} |
| Pathways Over-represented in Cluster 3 | Pathway Size | Cluster Genes | p-value (FDR) |
| Eukaryotic Translation Elongation | 86 | 55 | 1.1×10^{-112} |
| Peptide chain elongation | 83 | 54 | 1.3×10^{-112} |
| Viral mRNA Translation | 81 | 53 | 1.6×10^{-111} |
| Eukaryotic Translation Termination | 83 | 53 | 7.1×10^{-110} |
| Nonsense Mediated Decay independent of the Exon Junction Complex | 88 | 54 | 1.0×10^{-108} |
| Formation of a pool of free 40S subunits | 93 | 53 | 4.1×10^{-102} |
| Nonsense-Mediated Decay | 103 | 54 | 3.9×10^{-98} |
| Nonsense Mediated Decay enhanced by the Exon Junction Complex | 103 | 54 | 3.9×10^{-98} |
| LI3a-mediated translational silencing of Ceruloplasmin expression | 103 | 53 | 1.2×10^{-95} |
| 3' -UTR-mediated translational regulation | 103 | 53 | 1.2×10^{-95} |
| SRP-dependent cotranslational protein targeting to membrane | 104 | 53 | 4.3×10^{-95} |
| GTP hydrolysis and joining of the 60S ribosomal subunit | 104 | 53 | 4.3×10^{-95} |
| Influenza Viral RNA Transcription and Replication | 108 | 53 | 9.6×10^{-93} |
| Eukaryotic Translation Initiation | 111 | 53 | 4.2×10^{-91} |
| Cap-dependent Translation Initiation | 111 | 53 | 4.2×10^{-91} |
| Influenza Life Cycle | 112 | 53 | 1.4×10^{-90} |
| Influenza Infection | 117 | 53 | 6.2×10^{-88} |
| Translation | 141 | 55 | 3×10^{-81} |
| Pathways Over-represented in Cluster 4 | Pathway Size | Cluster Genes | p-value (FDR) |
| ECM proteoglycans | 66 | 10 | 2.9×10^{-11} |
| deactivation of the beta-catenin transactivating complex | 38 | 7 | 5.1×10^{-10} |
| Arachidonic acid metabolism | 41 | 7 | 1.1×10^{-9} |
| G _{αq} signalling events | 149 | 14 | 4.0×10^{-9} |
| HS-GAG degradation | 21 | 5 | 4.5×10^{-9} |
| Uptake and actions of bacterial toxins | 22 | 5 | 6.1×10^{-9} |
| Gastrin-CREB signalling pathway via PKC and MAPK | 170 | 15 | 6.1×10^{-9} |
| RNA Polymerase I, RNA Polymerase III, and Mitochondrial Transcription | 64 | 8 | 6.1×10^{-9} |
| Non-integrin membrane-ECM interactions | 53 | 7 | 1.5×10^{-8} |
| Syndecan interactions | 25 | 5 | 1.5×10^{-8} |
| NOTCH1 Intracellular Domain Regulates Transcription | 40 | 6 | 2.3×10^{-8} |
| Synthesis of Leukotrienes and Eoxins | 15 | 4 | 3.2×10^{-8} |
| Signalling by NOTCH1 | 59 | 7 | 5.3×10^{-8} |
| Regulation of insulin secretion | 44 | 6 | 6.0×10^{-8} |
| Metabolism of lipids and lipoproteins | 471 | 37 | 8.2×10^{-8} |
| Signalling by NOTCH | 80 | 8 | 1.2×10^{-7} |
| Platelet activation, signalling and aggregation | 179 | 14 | 1.2×10^{-7} |
| Recruitment of mitotic centrosome proteins and complexes | 64 | 7 | 1.2×10^{-7} |

Pathway over-representation analysis for Reactome pathways with the number of genes in each pathway (Pathway Size), number of genes within the pathway identified (Cluster Genes), and the pathway over-representation p-value (adjusted by FDR) from the hypergeometric test.

C.3 Comparison to Primary Screen

The mutation synthetic lethal partners with *CDH1* were also compared to siRNA primary screen data (Telford *et al.*, 2015), as performed in Section 4.2.1. These are expected to be more concordant with the experimental results performed on a null mutant, however this not the case at the gene level: less genes overlapped with experimental candidates in Figure C.2. This may be affected by lower sample size for mutations in TCGA data or lower frequency (expected value) of *CDH1* mutations compared to low expression.




Figure C.2: **Comparison of mtSLIPT to siRNA.** Testing the overlap of gene candidates for E-cadherin synthetic lethal partners between computational (SLIPT) and experimental screening (siRNA) approaches. The χ^2 test suggests that the overlap is no more than would be expected by chance ($p = 0.281$).

Despite a lower sample size (and low number of a predicted partners) for mutation analysis, the pathway composition (Tables C.2 and C.4) is similar to expression analysis, as described in Section 4.2.5. In particular, the resampling analysis (Section C.3.1) supported many of the results of expression analysis (Section 4.2.5.1) with Tables C.5 and C.6 detecting many of the same or functionally-related pathways.

Table C.4: Pathway composition for *CDH1* partners from mtSLIPT and siRNA

| Predicted only by SLIPT (2901 genes) | Pathway Size | Genes Identified | p-value (FDR) |
|---|--------------|------------------|------------------------|
| Eukaryotic Translation Elongation | 87 | 57 | 2.8×10^{-120} |
| Peptide chain elongation | 84 | 56 | 3.1×10^{-120} |
| Eukaryotic Translation Termination | 84 | 55 | 2.8×10^{-117} |
| Viral mRNA Translation | 82 | 54 | 4.1×10^{-116} |
| Nonsense Mediated Decay independent of the Exon Junction Complex | 89 | 55 | 3.7×10^{-113} |
| Formation of a pool of free 40S subunits | 94 | 55 | 2.8×10^{-109} |
| Nonsense-Mediated Decay | 104 | 57 | 8.4×10^{-108} |
| Nonsense Mediated Decay enhanced by the Exon Junction Complex | 104 | 57 | 8.4×10^{-108} |
| L13a-mediated translational silencing of Ceruloplasmin expression | 104 | 56 | 3.4×10^{-105} |
| 3' -UTR-mediated translational regulation | 104 | 56 | 3.4×10^{-105} |
| GTP hydrolysis and joining of the 60S ribosomal subunit | 105 | 56 | 1.4×10^{-104} |
| Eukaryotic Translation Initiation | 112 | 56 | 2.8×10^{-100} |
| Cap-dependent Translation Initiation | 112 | 56 | 2.8×10^{-100} |
| SRP-dependent cotranslational protein targeting to membrane | 105 | 54 | 2.2×10^{-99} |
| Influenza Viral RNA Transcription and Replication | 109 | 54 | 5.3×10^{-97} |
| Influenza Life Cycle | 113 | 54 | 9.6×10^{-95} |
| Influenza Infection | 118 | 55 | 1.7×10^{-94} |
| Translation | 142 | 60 | 3.5×10^{-94} |
| Infectious disease | 349 | 77 | 5.9×10^{-62} |
| Extracellular matrix organization | 241 | 54 | 3.0×10^{-52} |

| Detected only by siRNA screen (1752 genes) | Pathway Size | Genes Identified | p-value (FDR) |
|--|--------------|------------------|-----------------------|
| Class A/1 (Rhodopsin-like receptors) | 282 | 69 | 1.9×10^{-59} |
| GPCR ligand binding | 363 | 78 | 2.7×10^{-54} |
| Peptide ligand-binding receptors | 175 | 41 | 1.5×10^{-42} |
| $G_{\alpha i}$ signalling events | 184 | 41 | 1.1×10^{-40} |
| Gastrin-CREB signalling pathway via PKC and MAPK | 180 | 37 | 1.5×10^{-35} |
| $G_{\alpha q}$ signalling events | 159 | 34 | 3.7×10^{-35} |
| DAP12 interactions | 159 | 27 | 1.1×10^{-24} |
| VEGFA-VEGFR2 Pathway | 91 | 19 | 1.0×10^{-23} |
| Downstream signal transduction | 146 | 24 | 1.9×10^{-22} |
| Signalling by VEGF | 99 | 19 | 2.6×10^{-22} |
| DAP12 signalling | 149 | 24 | 4.2×10^{-22} |
| Organelle biogenesis and maintenance | 264 | 34 | 4.3×10^{-20} |
| Downstream signalling of activated FGFR1 | 134 | 21 | 4.3×10^{-20} |
| Downstream signalling of activated FGFR2 | 134 | 21 | 4.3×10^{-20} |
| Downstream signalling of activated FGFR3 | 134 | 21 | 4.3×10^{-20} |
| Downstream signalling of activated FGFR4 | 134 | 21 | 4.3×10^{-20} |
| Signalling by ERBB2 | 146 | 22 | 5.3×10^{-20} |
| Signalling by FGFR | 146 | 22 | 5.3×10^{-20} |
| Signalling by FGFR1 | 146 | 22 | 5.3×10^{-20} |
| Signalling by FGFR2 | 146 | 22 | 5.3×10^{-20} |

| Intersection of SLIPT and siRNA screen (450 genes) | Pathway Size | Genes Identified | p-value (FDR) |
|--|--------------|------------------|----------------------|
| HS-GAG degradation | 21 | 4 | 4.9×10^{-6} |
| Retinoid metabolism and transport | 39 | 5 | 4.9×10^{-6} |
| Platelet activation, signalling and aggregation | 186 | 13 | 4.9×10^{-6} |
| Signalling by NOTCH4 | 11 | 3 | 4.9×10^{-6} |
| $G_{\alpha s}$ signalling events | 100 | 8 | 5.0×10^{-6} |
| Defective EXT2 causes exostoses 2 | 12 | 3 | 5.0×10^{-6} |
| Defective EXT1 causes exostoses 1, TRPS2 and CHDS | 12 | 3 | 5.0×10^{-6} |
| Class A/1 (Rhodopsin-like receptors) | 289 | 18 | 2.2×10^{-5} |
| Signalling by PDGF | 173 | 11 | 2.9×10^{-5} |
| Circadian Clock | 34 | 4 | 2.9×10^{-5} |
| Signalling by ERBB4 | 139 | 9 | 4.3×10^{-5} |
| Role of LAT2/NTAL/LAB on calcium mobilization | 99 | 7 | 4.4×10^{-5} |
| Peptide ligand-binding receptors | 181 | 11 | 4.5×10^{-5} |
| Defective B4GALT7 causes EDS, progeroid type | 19 | 3 | 4.5×10^{-5} |
| Defective B3GAT3 causes JDSSDHD | 19 | 3 | 4.5×10^{-5} |
| Signalling by NOTCH | 80 | 6 | 4.5×10^{-5} |
| $G_{\alpha q}$ signalling events | 164 | 10 | 5.1×10^{-5} |
| Response to elevated platelet cytosolic Ca^{2+} | 84 | 6 | 7.1×10^{-5} |
| Signalling by ERBB2 | 148 | 9 | 7.1×10^{-5} |
| Signalling by SCF-KIT | 129 | 8 | 8.3×10^{-5} |

C.3.1 Resampling Analysis

Table C.5: Pathways for *CDH1* partners from mtSLIPT

| Reactome Pathway | Over-representation | Permutation |
|---|------------------------|--------------------------|
| Eukaryotic Translation Elongation | 3.2×10^{-128} | $< 7.035 \times 10^{-4}$ |
| Peptide chain elongation | 3.2×10^{-128} | $< 7.035 \times 10^{-4}$ |
| Eukaryotic Translation Termination | 3.7×10^{-125} | $< 7.035 \times 10^{-4}$ |
| Viral mRNA Translation | 4.1×10^{-124} | $< 7.035 \times 10^{-4}$ |
| Nonsense Mediated Decay independent of the Exon Junction Complex | 1.4×10^{-123} | $< 7.035 \times 10^{-4}$ |
| Nonsense-Mediated Decay | 8.4×10^{-117} | $< 7.035 \times 10^{-4}$ |
| Nonsense Mediated Decay enhanced by the Exon Junction Complex | 8.4×10^{-117} | $< 7.035 \times 10^{-4}$ |
| Formation of a pool of free 40S subunits | 2.6×10^{-116} | $< 7.035 \times 10^{-4}$ |
| L13a-mediated translational silencing of Ceruloplasmin expression | 2.0×10^{-111} | $< 7.035 \times 10^{-4}$ |
| 3' -UTR-mediated translational regulation | 2.0×10^{-111} | $< 7.035 \times 10^{-4}$ |
| GTP hydrolysis and joining of the 60S ribosomal subunit | 9.9×10^{-111} | $< 7.035 \times 10^{-4}$ |
| SRP-dependent cotranslational protein targeting to membrane | 4.7×10^{-108} | $< 7.035 \times 10^{-4}$ |
| Eukaryotic Translation Initiation | 4.8×10^{-106} | $< 7.035 \times 10^{-4}$ |
| Cap-dependent Translation Initiation | 4.8×10^{-106} | $< 7.035 \times 10^{-4}$ |
| Influenza Viral RNA Transcription and Replication | 8.1×10^{-103} | $< 7.035 \times 10^{-4}$ |
| Influenza Infection | 2.4×10^{-102} | $< 7.035 \times 10^{-4}$ |
| Translation | 6.0×10^{-101} | $< 7.035 \times 10^{-4}$ |
| Influenza Life Cycle | 2.2×10^{-100} | $< 7.035 \times 10^{-4}$ |
| Disease | 2.1×10^{-90} | 0.013347 |
| GPCR downstream signalling | 1.6×10^{-80} | 0.095478 |
| Hemostasis | 2.1×10^{-78} | 0.2671 |
| Signalling by GPCR | 1.2×10^{-73} | 0.44939 |
| <i>Extracellular matrix organization</i> | 2.2×10^{-67} | 0.054008 |
| Metabolism of proteins | 1.4×10^{-66} | 0.9607 |
| Signal Transduction | 2.1×10^{-66} | 0.48184 |
| Developmental Biology | 2.5×10^{-66} | 0.54075 |
| Innate Immune System | 5.3×10^{-66} | 0.9589 |
| Infectious disease | 9.6×10^{-66} | 0.21075 |
| Signalling by NGF | 1.1×10^{-62} | 0.43356 |
| Immune System | 2.8×10^{-62} | 0.23052 |

Over-representation (hypergeometric test) and Permutation p-values adjusted for multiple tests across pathways (FDR). Significant pathways are marked in bold (FDR < 0.05) and italics (FDR < 0.1).

Table C.6: Pathways for *CDH1* partners from mtSLIPT and siRNA primary screen

| Reactome Pathway | Over-representation | Permutation |
|---|----------------------|-------------|
| Visual phototransduction | 1.2×10^{-9} | 0.86279 |
| G_{as} signalling events | 2.9×10^{-7} | 0.023066 |
| Retinoid metabolism and transport | 2.9×10^{-7} | 0.299 |
| Acyl chain remodelling of PS | 1.1×10^{-5} | 0.42584 |
| Transcriptional regulation of white adipocyte differentiation | 1.1×10^{-5} | 0.53928 |
| Chemokine receptors bind chemokines | 1.1×10^{-5} | 0.95259 |
| <i>Signalling by NOTCH4</i> | 1.2×10^{-5} | 0.079229 |
| Defective EXT2 causes exostoses 2 | 1.2×10^{-5} | 0.22292 |
| Defective EXT1 causes exostoses 1, TRPS2 and CHDS | 1.2×10^{-5} | 0.22292 |
| Platelet activation, signalling and aggregation | 1.2×10^{-5} | 0.48853 |
| Serotonin receptors | 1.4×10^{-5} | 0.34596 |
| Nicotinamide salvaging | 1.4×10^{-5} | 0.70881 |
| Phase 1 - Functionalization of compounds | 2×10^{-5} | 0.31142 |
| Amine ligand-binding receptors | 2.5×10^{-5} | 0.34934 |
| Acyl chain remodelling of PE | 3.8×10^{-5} | 0.42615 |
| Signalling by GPCR | 3.8×10^{-5} | 0.93888 |
| Molecules associated with elastic fibres | 3.9×10^{-5} | 0.017982 |
| DAP12 interactions | 3.9×10^{-5} | 0.71983 |
| Beta defensins | 3.9×10^{-5} | 0.91458 |
| Cytochrome P ₄₅₀ - arranged by substrate type | 4.7×10^{-5} | 0.83493 |
| GPCR ligand binding | 5.7×10^{-5} | 0.95258 |
| Acyl chain remodelling of PC | 6.1×10^{-5} | 0.42584 |
| Response to elevated platelet cytosolic Ca ²⁺ | 6.4×10^{-5} | 0.54046 |
| Arachidonic acid metabolism | 6.7×10^{-5} | 0.026696 |
| Defective B4GALT7 causes EDS, progeroid type | 7.3×10^{-5} | 0.24921 |
| Defective B3GAT3 causes JDSSDHD | 7.3×10^{-5} | 0.24921 |
| Hydrolysis of LPC | 7.3×10^{-5} | 0.80663 |
| Elastic fibre formation | 7.4×10^{-5} | 0.0058768 |
| HS-GAG degradation | 9.4×10^{-5} | 0.0083179 |
| <i>Bile acid and bile salt metabolism</i> | 9.4×10^{-5} | 0.079905 |
| Netrin-1 signalling | 0.00011 | 0.92216 |
| Integration of energy metabolism | 0.00011 | 0.011152 |
| Dectin-2 family | 0.00012 | 0.10385 |
| Platelet sensitization by LDL | 0.00012 | 0.34596 |
| DAP12 signalling | 0.00012 | 0.62787 |
| Defensins | 0.00012 | 0.77542 |
| GPCR downstream signalling | 0.00012 | 0.79454 |
| <i>Diseases associated with glycosaminoglycan metabolism</i> | 0.00013 | 0.065927 |
| <i>Diseases of glycosylation</i> | 0.00013 | 0.065927 |
| Signalling by Retinoic Acid | 0.00013 | 0.22292 |
| Signalling by Leptin | 0.00013 | 0.34596 |
| Signalling by SCF-KIT | 0.00013 | 0.70881 |
| Opioid Signalling | 0.00013 | 0.96053 |
| Signalling by NOTCH | 0.00015 | 0.26884 |
| Platelet homeostasis | 0.00015 | 0.4878 |
| Signalling by NOTCH1 | 0.00016 | 0.13043 |
| Class B/2 (Secretin family receptors) | 0.00016 | 0.13994 |
| <i>Diseases of Immune System</i> | 0.0002 | 0.0795 |
| <i>Diseases associated with the TLR signalling cascade</i> | 0.0002 | 0.0795 |
| A tetrasaccharide linker sequence is required for GAG synthesis | 0.0002 | 0.42615 |

Over-representation (hypergeometric test) and Permutation p-values adjusted for multiple tests across pathways (FDR). Significant pathways are marked in bold (FDR < 0.05) and italics (FDR < 0.1).

C.4 Compare SLIPT genes

The mutation synthetic lethal partners with *CDH1* were also compared to siRNA primary screen data (Telford *et al.*, 2015), by correlation and siRNA viability as described in sections 4.2.2 and 4.2.3.




Figure C.3: **Compare mtSLIPT and siRNA genes with correlation.** The mtSLIPT p-values were compared against Pearson's correlation of expression with *CDH1*. Genes detected by SLIPT or siRNA are coloured according to the legend.




Figure C.4: **Compare mtSLIPT and siRNA genes with correlation.** Genes detected by mtSLIPT against *CDH1* mutation and siRNA screening were compared against Pearson's correlation of expression with *CDH1*. There were no differences in correlation between the gene groups.




Figure C.5: **Compare mtSLIPT and siRNA genes with siRNA viability.** Genes detected as candidate synthetic lethal partners by mtSLIPT (in TCGA breast cancer) expression analysis against *CDH1* mutation and experimental screening (with siRNA) were compared against the viability ratio of *CDH1* mutant and wildtype cells in the primary siRNA screen. There were clear no differences in viability between genes detected by mtSLIPT and those not with the differences being primarily due to viability thresholds being used to detect synthetic lethality by Telford *et al.* (2015).

C.5 Metagene Analysis

Metagene analysis was also performed for synthetic lethal candidates for *CDH1* mutation. These are described and compared to expression analysis in Section 4.3.3.

Table C.7: Candidate synthetic lethal metagenes against *CDH1* from mtSLIPT

| Pathway | ID | Observed | Expected | χ^2 value | p-value | p-value (FDR) |
|--|---------|----------|----------|----------------|------------------------|------------------------|
| Neurotoxicity of clostridium toxins | 168799 | 8 | 36.7 | 79.4 | 5.71×10^{-18} | 3.14×10^{-15} |
| Aquaporin-mediated transport | 445717 | 8 | 36.7 | 76.3 | 2.73×10^{-17} | 9.01×10^{-15} |
| Toxicity of botulinum toxin type G (BoNT/G) | 5250989 | 8 | 36.7 | 76.3 | 2.73×10^{-17} | 9.01×10^{-15} |
| ABC-family proteins mediated transport | 382556 | 10 | 36.7 | 68.2 | 1.58×10^{-15} | 1.86×10^{-13} |
| G α_s signalling events | 418597 | 10 | 36.7 | 59.9 | 9.97×10^{-14} | 5.48×10^{-12} |
| Regulation of IGF transport and uptake by IGFBPs | 381426 | 9 | 36.7 | 56.3 | 5.88×10^{-13} | 2.11×10^{-11} |
| GP1b-IX-V activation signalling | 430116 | 8 | 36.7 | 55.7 | 8.20×10^{-13} | 2.76×10^{-11} |
| GABA receptor activation | 977443 | 12 | 36.7 | 55.1 | 1.07×10^{-12} | 3.26×10^{-11} |
| Vasopressin regulates renal water homeostasis via Aquaporins | 432040 | 9 | 36.7 | 54.1 | 1.77×10^{-12} | 4.88×10^{-11} |
| Toxicity of botulinum toxin type D (BoNT/D) | 5250955 | 14 | 36.7 | 53.4 | 2.54×10^{-12} | 6.64×10^{-11} |
| Toxicity of botulinum toxin type F (BoNT/F) | 5250981 | 14 | 36.7 | 53.4 | 2.54×10^{-12} | 6.64×10^{-11} |
| STAT6-mediated induction of chemokines | 3249367 | 16 | 36.7 | 52.2 | 4.72×10^{-12} | 1.13×10^{-10} |
| Toxicity of botulinum toxin type B (BoNT/B) | 5250958 | 14 | 36.7 | 50.8 | 9.5×10^{-12} | 1.98×10^{-10} |
| S6K1 signalling | 165720 | 12 | 36.7 | 50.2 | 1.24×10^{-11} | 2.5×10^{-10} |
| G α_s signalling events | 418555 | 11 | 36.7 | 49.2 | 2.08×10^{-11} | 3.85×10^{-10} |
| RHO GTPases activate CIT | 5625900 | 14 | 36.7 | 48.2 | 3.34×10^{-11} | 5.9×10^{-10} |
| NADE modulates death signalling | 205025 | 15 | 36.7 | 47.4 | 5.00×10^{-11} | 8.32×10^{-10} |
| Keratan sulfate degradation | 2022857 | 10 | 36.7 | 46.6 | 7.5×10^{-11} | 1.15×10^{-9} |
| Signalling by Retinoic Acid | 5362517 | 10 | 36.7 | 46.6 | 7.5×10^{-11} | 1.15×10^{-9} |
| Adenylate cyclase inhibitory pathway | 170670 | 14 | 36.7 | 45.9 | 1.11×10^{-10} | 1.59×10^{-9} |
| Inhibition of adenylate cyclase pathway | 997269 | 14 | 36.7 | 45.9 | 1.11×10^{-10} | 1.59×10^{-9} |
| Fatty acids | 211935 | 6 | 36.7 | 45.7 | 1.21×10^{-10} | 1.72×10^{-9} |
| Ionotropic activity of Kainate Receptors | 451306 | 13 | 36.7 | 44.6 | 2.03×10^{-10} | 2.58×10^{-9} |
| Activation of Ca-permeable Kainate Receptor | 451308 | 13 | 36.7 | 44.6 | 2.03×10^{-10} | 2.58×10^{-9} |
| RA biosynthesis pathway | 5365859 | 13 | 36.7 | 44.6 | 2.03×10^{-10} | 2.58×10^{-9} |

Strongest candidate SL partners for *CDH1* by mtSLIPT with observed and expected numbers of mutant *CDH1* TCGA breast cancer tumours with low expression of partner metagenes.

C.6 Expression of Somatic Mutations







Figure C.6: **Somatic mutation against PIK3CA metagene.** Mutations in *PIK3CA*, *PIK3R1*, *CDH1*, and *TP53* were examined in TCGA breast cancer for their effect on the PIK3CA (Gatza *et al.*, 2014) pathway metagene. The tumour suppressors *CDH1* and *TP53* showed an increase and decrease in the metagene respectively, whereas *PIK3CA* and *PIK3R1* mutations weaker evidence of decrease in metagene levels.




(a) *PIK3CA* mutation



(b) *PIK3CA* or *PIK3R1* mutation




(c) *CDH1* mutation




(d) *TP53* mutation


Figure C.7: Somatic mutation against PI3K protein. Mutations in *PIK3CA*, *PIK3R1*, *CDH1*, and *TP53* were examined in TCGA breast cancer for their effect on the expression of the p110 α protein (encoded by *PIK3CA*). Protein levels were significantly elevated in samples with *PIK3CA* or *PIK3R1* mutations and lower in samples with *TP53* mutations.




(a) *PIK3CA* mutation



(b) *PIK3CA* or *PIK3R1* mutation



(c) *CDH1* mutation



(d) *TP53* mutation

Figure C.8: **Somatic mutation against AKT protein.** Mutations in *PIK3CA*, *PIK3R1*, *CDH1*, and *TP53* were examined in TCGA breast cancer for their effect on the expression of the AKT protein (a downstream target of *PIK3CA*). Protein levels were not significantly different in samples mutations in any of these cancer genes.

C.7 Metagene Expression Profiles




Figure C.9: **Pathway metagene expression profiles.** Expression profiles for metagene signatures from Gatzia *et al.* (2014) in TCGA breast data, annotated for clinical factors and cancer gene mutations.




Figure C.10: **Expression profiles for p53 related genes.** Expression profiles the genes contained in the *TP53* gene signature from Gatza *et al.* (2011) in TCGA breast data, annotated for clinical factors and cancer gene mutations. Samples are separated by *CDH1* expression status and sorted by the metagene. In both cases, the majority of genes were consistent with the direction of the metagene, with few very exceptions. *TP53* mutant samples had low metagene expression, consistent with loss of tumour suppressor functions, and were less likely to have *CDH1* or *PIK3CA* mutations.




Figure C.11: **Expression profiles for BRCA related genes.** Expression profiles the genes contained in the gene signature related to *BRCA1* and *BRCA2* functions from Gatza *et al.* (2014) in TCGA breast data, annotated for clinical factors and cancer gene mutations. Samples are separated by *CDH1* expression status and sorted by the metagene. In both cases, the majority of genes were consistent with the direction of the metagene, with few very exceptions. *BRCA1* and *BRCA2* mutant samples had higher metagene expression than most samples for the ductal subtype, although this was not the case (for the lobular samples for which the metagene was lower). However, the metagene was higher for basal subtype and estrogen receptor negative samples.

Appendix D

Intrinsic Subtyping

The intrinsic subtypes for TCGA breast cancer samples provided by University of California, Santa Cruz (UCSC) (TCGA, 2012) that were derived from microarray analysis have been compared to the Prediction Analysis of Microarray 50 (PAM50) results for performing subtyping from RNA-Seq data (Parker *et al.*, 2009). As shown in Table D.1, these subtypes were highly concordant for samples which had both procedures performed upon them ($\chi^2 = 1305.9$, $p = 2.73 \times 10^{-268}$). The main exception were the luminal A samples some of which were reclassified as luminal B or “normal-like”.

Table D.1: Comparison of Intrinsic Subtypes

| UCSC Subtype | | | | | |
|--------------|------------|---------------|-----------|-----------|-------------|
| | Basal-like | HER2-enriched | Luminal A | Luminal B | Normal-like |
| | 100 | 58 | 232 | 128 | 30 |

| PAM50 Subtype | | | | | |
|---------------|------------|---------------|-----------|-----------|-------------|
| | Basal-like | HER2-enriched | Luminal A | Luminal B | Normal-like |
| | 208 | 94 | 314 | 334 | 227 |

| UCSC Subtype | | | | | |
|---------------|------------|---------------|-----------|-----------|-------------|
| PAM50 Subtype | Basal-like | HER2-enriched | Luminal A | Luminal B | Normal-like |
| Basal-like | 96 | 4 | 2 | 2 | 1 |
| HER2-enriched | 0 | 47 | 5 | 3 | 0 |
| Luminal A | 1 | 0 | 141 | 1 | 0 |
| Luminal B | 2 | 7 | 49 | 121 | 0 |
| Normal-like | 1 | 0 | 35 | 1 | 29 |

The intrinsic subtypes of TCGA breast samples were compared between those provided by UCSC (TCGA, 2012) from microarray expression to those derived from RNA-Seq data (Parker *et al.*, 2009). Comparisons between these were limited to samples for which both data types were available.

The PAM50 subtypes are potentially more accurate given similarity of these subtypes and that the remainder of the subtypes were accurately recapitulated with RNA-Seq data. Furthermore, UCSC subtypes correctly identified $^{22}/_{22}$ normal samples as “normal-like” and PAM50 subtyping in RNA-Seq data had a success rate of $^{112}/_{113}$ (including all of those identified from microarrays). Therefore the PAM50 subtypes (performed on a larger cohort of samples) are appropriate to use for further interpretation, superceding the UCSC subtypes available for a limited set of samples.

Appendix E

Stomach Expression Analysis

The following results are a replication of the TCGA results (in Chapter 4) with stomach cancer data, using synthetic lethality (SLIPT) against *CDH1* mutation.

E.1 Synthetic Lethal Genes and Pathways

Table E.1: Synthetic lethal gene partners of *CDH1* from SLIPT in stomach cancer

| Gene | Observed | Expected | χ^2 value | p-value | p-value (FDR) |
|------------------|----------|----------|----------------|------------------------|------------------------|
| <i>PRAF2</i> | 17 | 50.4 | 121 | 3.54×10^{-25} | 1.45×10^{-21} |
| <i>EMP3</i> | 17 | 50.4 | 115 | 5.06×10^{-24} | 1.48×10^{-20} |
| <i>PLEKHO1</i> | 22 | 50.4 | 112 | 2.14×10^{-23} | 4.75×10^{-20} |
| <i>SELM</i> | 20 | 50.4 | 111 | 5.13×10^{-23} | 8.09×10^{-20} |
| <i>GYPC</i> | 20 | 50.4 | 110 | 5.77×10^{-23} | 8.45×10^{-20} |
| <i>COX7A1</i> | 18 | 50.4 | 109 | 1.15×10^{-22} | 1.39×10^{-19} |
| <i>TNFSF12</i> | 20 | 50.4 | 106 | 4.06×10^{-22} | 4.38×10^{-19} |
| <i>SEPT4</i> | 17 | 50.4 | 106 | 6.58×10^{-22} | 5.91×10^{-19} |
| <i>LGALS1</i> | 19 | 50.4 | 105 | 6.64×10^{-22} | 5.91×10^{-19} |
| <i>RARRES2</i> | 27 | 50.4 | 105 | 8.02×10^{-22} | 6.85×10^{-19} |
| <i>VEGFB</i> | 16 | 50.4 | 104 | 1.19×10^{-21} | 9.74×10^{-19} |
| <i>PRR24</i> | 22 | 50.4 | 102 | 2.96×10^{-21} | 2.02×10^{-18} |
| <i>SYNC</i> | 19 | 50.4 | 102 | 3.73×10^{-21} | 2.39×10^{-18} |
| <i>MAGEH1</i> | 17 | 50.4 | 100 | 9.52×10^{-21} | 5.01×10^{-18} |
| <i>HSPB2</i> | 23 | 50.4 | 99.6 | 1.19×10^{-20} | 5.82×10^{-18} |
| <i>SMARCD3</i> | 19 | 50.4 | 99 | 1.59×10^{-20} | 7.57×10^{-18} |
| <i>CREM</i> | 13 | 50.4 | 98.1 | 2.48×10^{-20} | 1.13×10^{-17} |
| <i>GNG11</i> | 20 | 50.4 | 97.3 | 3.68×10^{-20} | 1.59×10^{-17} |
| <i>GNAI2</i> | 17 | 50.4 | 96.4 | 5.75×10^{-20} | 2.36×10^{-17} |
| <i>FUNDC2</i> | 22 | 50.4 | 95.9 | 7.39×10^{-20} | 2.91×10^{-17} |
| <i>CNRIP1</i> | 21 | 50.4 | 95.3 | 1.0×10^{-19} | 3.66×10^{-17} |
| <i>CALHM2</i> | 22 | 50.4 | 93.1 | 2.94×10^{-19} | 1.06×10^{-16} |
| <i>ARID5A</i> | 18 | 50.4 | 92.7 | 3.47×10^{-19} | 1.22×10^{-16} |
| <i>ST3GAL3</i> | 27 | 50.4 | 92.2 | 4.49×10^{-19} | 1.56×10^{-16} |
| <i>LOC339524</i> | 21 | 50.4 | 92.1 | 4.8×10^{-19} | 1.59×10^{-16} |

SLIPT partners of *CDH1* with observed and expected numbers of TCGA stomach cancer samples with low expression of both genes.

Table E.2: Pathways for *CDH1* partners from SLIPT in stomach cancer

| Pathways Over-represented | Pathway Size | SL Genes | p-value (FDR) |
|---|--------------|----------|------------------------|
| Extracellular matrix organization | 241 | 104 | 7.5×10^{-140} |
| Hemostasis | 445 | 138 | 1.8×10^{-121} |
| Developmental Biology | 432 | 125 | 9.2×10^{-107} |
| Axon guidance | 289 | 94 | 1.5×10^{-102} |
| Eukaryotic Translation Termination | 84 | 49 | 1.9×10^{-99} |
| GPCR ligand binding | 373 | 108 | 3.8×10^{-99} |
| Viral mRNA Translation | 82 | 48 | 3.3×10^{-98} |
| Formation of a pool of free 40S subunits | 94 | 51 | 3.3×10^{-98} |
| Eukaryotic Translation Elongation | 87 | 49 | 1.6×10^{-97} |
| Peptide chain elongation | 84 | 48 | 7.2×10^{-97} |
| Class A/1 (Rhodopsin-like receptors) | 289 | 90 | 2.7×10^{-96} |
| Nonsense Mediated Decay independent of the Exon Junction Complex | 89 | 49 | 3.0×10^{-96} |
| Infectious disease | 349 | 100 | 2.6×10^{-94} |
| GTP hydrolysis and joining of the 60S ribosomal subunit | 105 | 52 | 3.4×10^{-94} |
| L13a-mediated translational silencing of Ceruloplasmin expression | 104 | 51 | 2.8×10^{-92} |
| 3' -UTR-mediated translational regulation | 104 | 51 | 2.8×10^{-92} |
| Neuronal System | 272 | 84 | 8.4×10^{-92} |
| SRP-dependent cotranslational protein targeting to membrane | 105 | 51 | 9.5×10^{-92} |
| Eukaryotic Translation Initiation | 112 | 52 | 2.0×10^{-90} |
| Cap-dependent Translation Initiation | 112 | 52 | 2.0×10^{-90} |

Gene set over-representation analysis (hypergeometric test) for Reactome pathways in Synthetic Lethal Interaction Prediction Tool (SLIPT) partners for *CDH1*.

E.2 Synthetic Lethal Expression Profiles

Table E.3: Pathway composition for clusters of *CDH1* partners in stomach SLIPT

| Pathways Over-represented in Cluster 1 | Pathway Size | Cluster Genes | p-value (FDR) |
|--|--------------|---------------|------------------------|
| Viral mRNA Translation | 82 | 48 | 1.3×10^{-97} |
| Formation of a pool of free 40S subunits | 94 | 51 | 1.3×10^{-97} |
| Eukaryotic Translation Elongation | 87 | 49 | 4.8×10^{-97} |
| Peptide chain elongation | 84 | 48 | 1.4×10^{-96} |
| Eukaryotic Translation Termination | 84 | 48 | 1.4×10^{-96} |
| GTP hydrolysis and joining of the 60S ribosomal subunit | 105 | 52 | 7.9×10^{-94} |
| Nonsense Mediated Decay independent of the Exon Junction Complex | 89 | 48 | 3.1×10^{-93} |
| L13a-mediated translational silencing of Ceruloplasmin expression | 104 | 51 | 5.1×10^{-92} |
| 3' -UTR-mediated translational regulation | 104 | 51 | 5.1×10^{-92} |
| SRP-dependent cotranslational protein targeting to membrane | 105 | 51 | 1.7×10^{-91} |
| Eukaryotic Translation Initiation | 112 | 52 | 3.3×10^{-90} |
| Cap-dependent Translation Initiation | 112 | 52 | 3.3×10^{-90} |
| Translation | 142 | 56 | 3.6×10^{-85} |
| Nonsense-Mediated Decay | 104 | 48 | 1.2×10^{-84} |
| Nonsense Mediated Decay enhanced by the Exon Junction Complex | 104 | 48 | 1.2×10^{-84} |
| Influenza Viral RNA Transcription and Replication | 109 | 48 | 4.1×10^{-82} |
| Influenza Life Cycle | 113 | 48 | 3.4×10^{-80} |
| Influenza Infection | 118 | 48 | 6.4×10^{-78} |
| Pathways Over-represented in Cluster 2 | Pathway Size | Cluster Genes | p-value (FDR) |
| Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell | 65 | 12 | 1.3×10^{-15} |
| Phosphorylation of CD3 and TCR zeta chains | 18 | 6 | 1.7×10^{-12} |
| Generation of second messenger molecules | 29 | 7 | 2.7×10^{-12} |
| PD-1 signalling | 21 | 6 | 7.4×10^{-12} |
| TCR signalling | 62 | 9 | 4.3×10^{-11} |
| Translocation of ZAP-70 to Immunological synapse | 16 | 5 | 1.1×10^{-10} |
| Interferon alpha/beta signalling | 68 | 9 | 1.6×10^{-10} |
| Initial triggering of complement | 17 | 5 | 1.6×10^{-10} |
| IKK complex recruitment mediated by RIP1 | 19 | 5 | 5.1×10^{-10} |
| TRIF-mediated programmed cell death | 10 | 4 | 6.2×10^{-10} |
| Creation of C4 and C2 activators | 11 | 4 | 1.3×10^{-9} |
| RHO GTPases Activate NADPH Oxidases | 11 | 4 | 1.3×10^{-9} |
| Interferon Signalling | 175 | 15 | 2.3×10^{-9} |
| Chemokine receptors bind chemokines | 52 | 7 | 4.0×10^{-9} |
| Interferon gamma signalling | 74 | 8 | 1.6×10^{-8} |
| TRAF6 mediated induction of TAK1 complex | 15 | 4 | 1.6×10^{-8} |
| Activation of IRF3/IRF7 mediated by TBK1/IKK epsilon | 16 | 4 | 2.7×10^{-8} |
| Downstream TCR signalling | 45 | 6 | 3.5×10^{-8} |
| Pathways Over-represented in Cluster 3 | Pathway Size | Cluster Genes | p-value (FDR) |
| Uptake and actions of bacterial toxins | 22 | 4 | 3.5×10^{-6} |
| Neurotoxicity of clostridium toxins | 10 | 3 | 3.5×10^{-6} |
| Activation of PPARGC1A (PGC-1alpha) by phosphorylation | 10 | 3 | 3.5×10^{-6} |
| SMAD2/SMAD3:SMAD4 heterotrimer regulates transcription | 28 | 4 | 1.4×10^{-5} |
| Assembly of the primary cilium | 149 | 10 | 2.5×10^{-5} |
| Serotonin Neurotransmitter Release Cycle | 15 | 3 | 2.5×10^{-5} |
| Glycosaminoglycan metabolism | 114 | 8 | 3.3×10^{-5} |
| Platelet homeostasis | 54 | 5 | 3.3×10^{-5} |
| Norepinephrine Neurotransmitter Release Cycle | 17 | 3 | 3.3×10^{-5} |
| Acetylcholine Neurotransmitter Release Cycle | 17 | 3 | 3.3×10^{-5} |
| G _{12s} signalling events | 100 | 7 | 5.5×10^{-5} |
| GABA synthesis, release, reuptake and degradation | 19 | 3 | 5.6×10^{-5} |
| deactivation of the beta-catenin transactivating complex | 39 | 4 | 6.7×10^{-5} |
| Dopamine Neurotransmitter Release Cycle | 20 | 3 | 6.7×10^{-5} |
| IRS-related events triggered by IGF1R | 83 | 6 | 7.1×10^{-5} |
| Generic Transcription Pathway | 186 | 11 | 7.1×10^{-5} |
| Termination of O-glycan biosynthesis | 21 | 3 | 7.4×10^{-5} |
| Kinesins | 22 | 3 | 8.5×10^{-5} |
| Pathways Over-represented in Cluster 4 | Pathway Size | Cluster Genes | p-value (FDR) |
| Extracellular matrix organization | 241 | 97 | 8.8×10^{-126} |
| Axon guidance | 289 | 75 | 8.3×10^{-72} |
| Hemostasis | 445 | 101 | 8.3×10^{-72} |
| Developmental Biology | 432 | 95 | 3.0×10^{-67} |
| Response to elevated platelet cytosolic Ca ²⁺ | 84 | 37 | 5.8×10^{-67} |
| Platelet degranulation | 79 | 36 | 5.8×10^{-67} |
| Degradation of the extracellular matrix | 104 | 39 | 6.7×10^{-63} |
| Platelet activation, signalling and aggregation | 186 | 52 | 6.6×10^{-62} |
| ECM proteoglycans | 66 | 31 | 8.1×10^{-61} |
| Neuronal System | 272 | 64 | 5.1×10^{-60} |
| Signalling by PDGF | 173 | 47 | 9.7×10^{-57} |
| Integrin cell surface interactions | 82 | 31 | 1.9×10^{-53} |
| Collagen biosynthesis and modifying enzymes | 56 | 26 | 1.1×10^{-52} |
| Collagen formation | 67 | 28 | 1.4×10^{-52} |
| Class A/1 (Rhodopsin-like receptors) | 289 | 61 | 2.3×10^{-52} |
| GPCR ligand binding | 373 | 73 | 2.8×10^{-52} |
| Elastic fibre formation | 38 | 22 | 4.7×10^{-52} |
| Non-integrin membrane-ECM interactions | 53 | 24 | 7.0×10^{-49} |

Pathway over-representation analysis for Reactome pathways with the number of genes in each pathway (Pathway Size), number of genes within the pathway identified (Cluster Genes), and the pathway over-representation p-value (adjusted by FDR) from the hypergeometric test.




Figure E.1: **Synthetic lethal expression profiles of analysed samples.** Gene expression profile heatmap (correlation distance) of all samples (separated by the $1/3$ quantile of *CDH1* expression) analysed in TCGA stomach cancer dataset for gene expression of 4,365 candidate partners of E-cadherin (*CDH1*) from SLIPT prediction (with significant FDR adjusted $p < 0.05$). Deeply clustered, inter-correlated genes form several main groups, each containing genes that were SL candidates or toxic in an siRNA screen Telford *et al.* (2015). Clusters had different sample groups highly expressing the synthetic lethal candidates in *CDH1* low samples, notably diffuse and *CDH1* mutant samples have elevated expression in one or more distinct clusters, although there was less complexity and variation among candidate synthetic lethal partners than in breast data. *CDH1* low samples also contained most of samples with *CDH1* mutations.

E.3 Comparison to Primary Screen

The synthetic lethal partners with *CDH1* expression in stomach cancers were also compared to siRNA primary screen data (Telford *et al.*, 2015), as performed in Section 4.2.1. These are expected to be more concordant with the experimental results performed on a null mutant, however this not the case at the gene level: less genes overlapped with experimental candidates in Figure E.2. This may be affected by lower sample size for mutations in TCGA data or lower frequency (expected value) of *CDH1* mutations compared to low expression.




Figure E.2: **Comparison of SLIPT in stomach to siRNA.** Testing the overlap of gene candidates for E-cadherin synthetic lethal partners between computational (SLIPT) and experimental screening (siRNA) approaches. The χ^2 test suggests that the overlap is no more than would be expected by chance ($p = 0.281$).

Table E.4: Pathway composition for *CDH1* partners from SLIPT and siRNA screening

| Predicted only by SLIPT (3392 genes) | Pathway Size | Genes Identified | p-value (FDR) |
|---|--------------|------------------|------------------------|
| Extracellular matrix organization | 238 | 90 | 3.4×10^{-107} |
| Eukaryotic Translation Termination | 79 | 46 | 7.6×10^{-91} |
| Viral mRNA Translation | 77 | 45 | 1.2×10^{-89} |
| Eukaryotic Translation Elongation | 82 | 46 | 5.8×10^{-89} |
| Peptide chain elongation | 79 | 45 | 2.1×10^{-88} |
| Nonsense Mediated Decay independent of the Exon Junction Complex | 84 | 46 | 9.4×10^{-88} |
| Formation of a pool of free 40S subunits | 89 | 47 | 3.3×10^{-87} |
| GTP hydrolysis and joining of the 60S ribosomal subunit | 100 | 48 | 3.2×10^{-83} |
| Axon guidance | 284 | 84 | 3.9×10^{-82} |
| Developmental Biology | 426 | 111 | 4.2×10^{-82} |
| L13a-mediated translational silencing of Ceruloplasmin expression | 99 | 47 | 1.4×10^{-81} |
| 3' -UTR-mediated translational regulation | 99 | 47 | 1.4×10^{-81} |
| SRP-dependent cotranslational protein targeting to membrane | 99 | 47 | 1.4×10^{-81} |
| Nonsense-Mediated Decay | 99 | 47 | 1.4×10^{-81} |
| Nonsense Mediated Decay enhanced by the Exon Junction Complex | 99 | 47 | 1.4×10^{-81} |
| Hemostasis | 438 | 112 | 1.2×10^{-80} |
| Eukaryotic Translation Initiation | 107 | 48 | 8.0×10^{-80} |
| Cap-dependent Translation Initiation | 107 | 48 | 8.0×10^{-80} |
| Infectious disease | 338 | 90 | 1.6×10^{-76} |
| Neuronal System | 267 | 77 | 1.6×10^{-76} |

| Detected only by siRNA screen (1803 genes) | Pathway Size | Genes Identified | p-value (FDR) |
|--|--------------|------------------|-----------------------|
| Class A/1 (Rhodopsin-like receptors) | 282 | 62 | 8.1×10^{-50} |
| GPCR ligand binding | 363 | 71 | 4.9×10^{-46} |
| Peptide ligand-binding receptors | 175 | 38 | 7.9×10^{-38} |
| G _{αi} signalling events | 184 | 37 | 1.1×10^{-34} |
| Gastrin-CREB signalling pathway via PKC and MAPK | 180 | 35 | 1.4×10^{-32} |
| G _{αq} signalling events | 159 | 32 | 4.8×10^{-32} |
| DAP12 interactions | 159 | 29 | 1.4×10^{-27} |
| Downstream signal transduction | 146 | 26 | 2.4×10^{-25} |
| DAP12 signalling | 149 | 26 | 6.4×10^{-25} |
| VEGFA-VEGFR2 Pathway | 91 | 19 | 8.1×10^{-24} |
| Signalling by PDGF | 172 | 27 | 5.7×10^{-23} |
| Signalling by ERBB2 | 146 | 24 | 1.4×10^{-22} |
| Signalling by VEGF | 99 | 19 | 2.0×10^{-22} |
| Visual phototransduction | 85 | 17 | 1.3×10^{-21} |
| Downstream signalling of activated FGFR1 | 134 | 22 | 1.3×10^{-21} |
| Downstream signalling of activated FGFR2 | 134 | 22 | 1.3×10^{-21} |
| Downstream signalling of activated FGFR3 | 134 | 22 | 1.3×10^{-21} |
| Downstream signalling of activated FGFR4 | 134 | 22 | 1.3×10^{-21} |
| Signalling by FGFR | 146 | 23 | 2.0×10^{-21} |
| Signalling by FGFR1 | 146 | 23 | 2.0×10^{-21} |

| Intersection of SLIPT and siRNA screen (547 genes) | Pathway Size | Genes Identified | p-value (FDR) |
|--|--------------|------------------|----------------------|
| Class A/1 (Rhodopsin-like receptors) | 282 | 25 | 3.9×10^{-9} |
| Platelet activation, signalling and aggregation | 182 | 17 | 3.9×10^{-9} |
| Response to elevated platelet cytosolic Ca ²⁺ | 82 | 9 | 5.5×10^{-8} |
| Platelet homeostasis | 53 | 7 | 5.7×10^{-8} |
| Nucleotide-like (purinergic) receptors | 16 | 4 | 1.8×10^{-7} |
| Platelet degranulation | 77 | 8 | 2.8×10^{-7} |
| Peptide ligand-binding receptors | 175 | 14 | 3.8×10^{-7} |
| Molecules associated with elastic fibres | 34 | 5 | 7.1×10^{-7} |
| Amine ligand-binding receptors | 35 | 5 | 8.6×10^{-7} |
| G _{αi} signalling events | 184 | 14 | 9.8×10^{-7} |
| GPCR ligand binding | 363 | 27 | 1.1×10^{-6} |
| Elastic fibre formation | 38 | 5 | 1.5×10^{-6} |
| G _{αq} signalling events | 159 | 12 | 1.9×10^{-6} |
| Serotonin receptors | 12 | 3 | 3.8×10^{-6} |
| P2Y receptors | 12 | 3 | 3.8×10^{-6} |
| Signal amplification | 16 | 3 | 2.3×10^{-5} |
| Gastrin-CREB signalling pathway via PKC and MAPK | 180 | 12 | 2.3×10^{-5} |
| Complement cascade | 33 | 4 | 2.4×10^{-5} |
| Glycosaminoglycan metabolism | 110 | 8 | 2.5×10^{-5} |
| Glycogen breakdown (glycogenolysis) | 17 | 3 | 2.7×10^{-5} |

E.3.1 Resampling Analysis

Table E.5: Pathways for *CDH1* partners from SLIPT in stomach cancer

| Reactome Pathway | Over-representation | Permutation |
|--|------------------------|--------------------------|
| <i>Extracellular matrix organization</i> | 7.5×10^{-140} | 0.070215 |
| Hemostasis | 1.8×10^{-121} | 0.25804 |
| Developmental Biology | 9.2×10^{-107} | 0.53032 |
| Axon guidance | 1.5×10^{-102} | 0.6704 |
| Eukaryotic Translation Termination | 1.9×10^{-99} | $> 1.031 \times 10^{-5}$ |
| GPCR ligand binding | 3.8×10^{-99} | 0.54914 |
| Viral mRNA Translation | 3.3×10^{-98} | $> 1.031 \times 10^{-5}$ |
| Formation of a pool of free 40S subunits | 3.3×10^{-98} | $> 1.031 \times 10^{-5}$ |
| Eukaryotic Translation Elongation | 1.6×10^{-97} | $> 1.031 \times 10^{-5}$ |
| Peptide chain elongation | 7.2×10^{-97} | $> 1.031 \times 10^{-5}$ |
| Class A/1 (Rhodopsin-like receptors) | 2.7×10^{-96} | 0.58174 |
| Nonsense Mediated Decay independent of the Exon Junction Complex | 3×10^{-96} | $> 1.031 \times 10^{-5}$ |
| Infectious disease | 2.6×10^{-94} | 0.25484 |
| GTP hydrolysis and joining of the 60S ribosomal subunit | 3.4×10^{-94} | $> 1.031 \times 10^{-5}$ |
| L13a-mediated translational silencing of Ceruloplasmin expression | 2.8×10^{-92} | $> 1.031 \times 10^{-5}$ |
| 3' -UTR-mediated translational regulation | 2.8×10^{-92} | $> 1.031 \times 10^{-5}$ |
| Neuronal System | 8.4×10^{-92} | 0.53433 |
| SRP-dependent cotranslational protein targeting to membrane | 9.5×10^{-92} | $> 1.031 \times 10^{-5}$ |
| Eukaryotic Translation Initiation | 2.0×10^{-90} | $> 1.031 \times 10^{-5}$ |
| Cap-dependent Translation Initiation | 2.0×10^{-90} | $> 1.031 \times 10^{-5}$ |
| Nonsense-Mediated Decay | 7.4×10^{-90} | $> 1.031 \times 10^{-5}$ |
| Nonsense Mediated Decay enhanced by the Exon Junction Complex | 7.4×10^{-90} | $> 1.031 \times 10^{-5}$ |
| Adaptive Immune System | 8.1×10^{-88} | 0.14116 |
| Translation | 1.3×10^{-87} | $> 1.031 \times 10^{-5}$ |
| Platelet activation, signalling and aggregation | 1.3×10^{-86} | 0.28959 |
| Influenza Infection | 1×10^{-82} | $> 1.031 \times 10^{-5}$ |
| Influenza Viral RNA Transcription and Replication | 2.4×10^{-82} | $> 1.031 \times 10^{-5}$ |
| Influenza Life Cycle | 2×10^{-80} | $> 1.031 \times 10^{-5}$ |
| Response to elevated platelet cytosolic Ca^{2+} | 4.9×10^{-78} | 0.50817 |
| Signalling by NGF | 1.6×10^{-75} | 0.38518 |
| Rho GTPase cycle | 5.1×10^{-75} | 0.14864 |
| Signalling by PDGF | 7.4×10^{-74} | 0.40493 |
| <i>Signalling by Rho GTPases</i> | 5.1×10^{-73} | 0.077217 |
| Glycosaminoglycan metabolism | 1.4×10^{-68} | 0.52984 |
| $\text{G}_{\alpha i}$ signalling events | 1.8×10^{-66} | 0.9254 |
| Metabolism of carbohydrates | 1.1×10^{-65} | 0.39501 |
| $\text{G}_{\alpha s}$ signalling events | 2.7×10^{-65} | 0.0050293 |
| Potassium Channels | 2.7×10^{-65} | 0.53359 |
| Transmission across Chemical Synapses | 1.8×10^{-64} | 0.81833 |
| ECM proteoglycans | 3.4×10^{-64} | 0.083482 |
| Peptide ligand-binding receptors | 4.8×10^{-64} | 0.62817 |
| Degradation of the extracellular matrix | 1.1×10^{-63} | 0.80879 |
| Platelet homeostasis | 5.3×10^{-63} | 0.53134 |
| NGF signalling via TRKA from the plasma membrane | 6.1×10^{-63} | 0.5717 |
| Integration of energy metabolism | 4.5×10^{-61} | 0.10889 |
| Collagen formation | 5.4×10^{-61} | 0.29896 |
| Integrin cell surface interactions | 7×10^{-59} | 0.18167 |
| Collagen biosynthesis and modifying enzymes | 7×10^{-59} | 0.30208 |
| Neurotransmitter Receptor Binding And Downstream Transmission | 8.7×10^{-57} | 0.82522 |
| In The Postsynaptic Cell | 8.7×10^{-57} | 0.82522 |
| Signalling by Wnt | 8.7×10^{-57} | 0.25468 |

Over-representation (hypergeometric test) and Permutation p-values adjusted for multiple tests across pathways (FDR). Significant pathways are marked in bold (FDR < 0.05) and italics (FDR < 0.1).

Table E.6: Pathways for *CDH1* partners from SLIPT in stomach and siRNA screen

| Reactome Pathway | Over-representation | Permutation |
|---|----------------------|---------------------------|
| Platelet activation, signalling and aggregation | 3.9×10^{-9} | 0.49557 |
| Class A/1 (Rhodopsin-like receptors) | 3.9×10^{-9} | 0.98432 |
| Response to elevated platelet cytosolic Ca^{2+} | 5.5×10^{-8} | 0.54349 |
| Platelet homeostasis | 5.7×10^{-8} | 0.45017 |
| Nucleotide-like (purinergic) receptors | 1.8×10^{-7} | 0.36966 |
| Peptide ligand-binding receptors | 3.8×10^{-7} | 0.91294 |
| Molecules associated with elastic fibres | 7.1×10^{-7} | 0.0025868 |
| Amine ligand-binding receptors | 8.6×10^{-7} | 0.43303 |
| $G_{\alpha i}$ signalling events | 9.8×10^{-7} | 0.99626 |
| GPCR ligand binding | 1.1×10^{-6} | 0.97733 |
| Elastic fibre formation | 1.5×10^{-6} | 0.0025868 |
| $G_{\alpha q}$ signalling events | 1.9×10^{-6} | 0.86089 |
| P2Y receptors | 3.8×10^{-6} | 0.18795 |
| Serotonin receptors | 3.8×10^{-6} | 0.37853 |
| Signal amplification | 2.3×10^{-5} | 0.47856 |
| Gastrin-CREB signalling pathway via PKC and MAPK | 2.3×10^{-5} | 0.98567 |
| Complement cascade | 2.4×10^{-5} | $> 3.4628 \times 10^{-6}$ |
| Glycosaminoglycan metabolism | 2.5×10^{-5} | 0.38953 |
| Glycogen breakdown (glycogenolysis) | 2.7×10^{-5} | 0.83772 |
| Defective B4GALT7 causes EDS, progeroid type | 4.9×10^{-5} | 0.10792 |
| Defective B3GAT3 causes JDSSDHD | 4.9×10^{-5} | 0.10792 |
| Role of LAT2/NTAL/LAB on calcium mobilization | 5.6×10^{-5} | 0.35373 |
| Cell surface interactions at the vascular wall | 5.6×10^{-5} | 0.47642 |
| $G_{\alpha s}$ signalling events | 6×10^{-5} | 0.019858 |
| Signalling by NOTCH | 6×10^{-5} | 0.19008 |
| A tetrasaccharide linker sequence is required for GAG synthesis | 0.00017 | 0.47642 |
| Extracellular matrix organization | 0.00018 | 0.0047308 |
| Collagen formation | 0.00018 | 0.19245 |
| Effects of PIP2 hydrolysis | 0.0002 | 0.37779 |
| Syndecan interactions | 0.0002 | 0.37779 |
| Diseases associated with glycosaminoglycan metabolism | 0.00023 | 0.01028 |
| Diseases of glycosylation | 0.00023 | 0.01028 |
| <i>Chondroitin sulfate/dermatan sulfate metabolism</i> | 0.00023 | 0.085541 |
| Integrin alphaIIb beta3 signalling | 0.00028 | 0.76936 |
| Keratan sulfate biosynthesis | 0.00034 | 0.68744 |
| Rho GTPase cycle | 0.00034 | 0.15675 |
| Creation of C4 and C2 activators | 0.00035 | 0.12275 |
| Abacavir transport and metabolism | 0.00035 | 0.12443 |
| Amine compound SLC transporters | 0.00037 | 0.69773 |
| FCERI mediated NF-kB activation | 0.00037 | 0.69846 |
| Fc epsilon receptor (FCERI) signalling | 0.00056 | 0.43303 |
| Defective EXT2 causes exostoses 2 | 0.00067 | 0.16053 |
| Defective EXT1 causes exostoses 1, TRPS2 and CHDS | 0.00067 | 0.16053 |
| <i>Collagen biosynthesis and modifying enzymes</i> | 0.00071 | 0.052911 |
| Keratan sulfate/keratin metabolism | 0.00073 | 0.46533 |
| G alpha (12/13) signalling events | 0.00078 | 0.59164 |
| SEMA3A-Plexin repulsion signalling by inhibiting Integrin adhesion | 0.00084 | 0.038504 |
| Signal attenuation | 0.00084 | 0.37779 |
| Eicosanoid ligand-binding receptors | 0.0011 | 0.11117 |
| SOS-mediated signalling | 0.0011 | 0.25387 |

Over-representation (hypergeometric test) and Permutation p-values adjusted for multiple tests across pathways (FDR). Significant pathways are marked in bold (FDR < 0.05) and italics (FDR < 0.1).

E.4 Metagene Analysis

Metagene analysis was also performed for synthetic lethal candidates for *CDH1* expression in stomach cancer. These are described and compared to mutation analysis in Section ??.

Table E.7: Candidate synthetic lethal metagenes against *CDH1* from SLIPT in stomach cancer

| Pathway | ID | Observed | Expected | χ^2 value | p-value | p-value (FDR) |
|---|---------|----------|----------|----------------|------------------------|------------------------|
| Cell-Cell communication | 1500931 | 18 | 50.4 | 110 | 7.43×10^{-23} | 1.53×10^{-20} |
| VEGFR2 mediated vascular permeability | 5218920 | 19 | 50.4 | 109 | 1.36×10^{-22} | 2.49×10^{-20} |
| Sema4D in semaphorin signalling | 400685 | 20 | 50.4 | 104 | 1.62×10^{-21} | 2.12×10^{-19} |
| Ion transport by P-type ATPases | 936837 | 17 | 50.4 | 100 | 8.29×10^{-21} | 8.06×10^{-19} |
| Sialic acid metabolism | 4085001 | 19 | 50.4 | 95.3 | 9.95×10^{-20} | 7.82×10^{-18} |
| Synthesis of pyrophosphates in the cytosol | 1855167 | 26 | 50.4 | 94 | 1.86×10^{-19} | 1.23×10^{-17} |
| Keratan sulfate/keratin metabolism | 1638074 | 25 | 50.4 | 93.5 | 2.36×10^{-19} | 1.44×10^{-17} |
| Ion channel transport | 983712 | 19 | 50.4 | 92.8 | 3.37×10^{-19} | 1.99×10^{-17} |
| Keratan sulfate biosynthesis | 2022854 | 26 | 50.4 | 91.4 | 6.79×10^{-19} | 3.62×10^{-17} |
| Arachidonic acid metabolism | 2142753 | 22 | 50.4 | 90.6 | 9.81×10^{-19} | 5.07×10^{-17} |
| RHO GTPases activate CIT | 5625900 | 22 | 50.4 | 87 | 5.80×10^{-18} | 2.66×10^{-16} |
| Stimuli-sensing channels | 2672351 | 25 | 50.4 | 85.8 | 1.03×10^{-17} | 4.58×10^{-16} |
| Synthesis of PI | 1483226 | 19 | 50.4 | 85.6 | 1.15×10^{-17} | 4.89×10^{-16} |
| G-protein activation | 202040 | 19 | 50.4 | 85.3 | 1.34×10^{-17} | 5.53×10^{-16} |
| NrCAM interactions | 447038 | 22 | 50.4 | 84.3 | 2.1×10^{-17} | 8.27×10^{-16} |
| Inwardly rectifying K^+ channels | 1296065 | 24 | 50.4 | 83.5 | 3.19×10^{-17} | 1.22×10^{-15} |
| Calcitonin-like ligand receptors | 419812 | 20 | 50.4 | 82.2 | 6.07×10^{-17} | 2.13×10^{-15} |
| Prostacyclin signalling through prostacyclin receptor | 392851 | 24 | 50.4 | 81.8 | 7.27×10^{-17} | 2.5×10^{-15} |
| Presynaptic function of Kainate receptors | 500657 | 26 | 50.4 | 79.7 | 2.00×10^{-16} | 6.34×10^{-15} |
| ADP signalling through P2Y purinoceptor 12 | 392170 | 23 | 50.4 | 79.2 | 2.57×10^{-16} | 7.71×10^{-15} |
| regulation of FZD by ubiquitination | 4641263 | 22 | 50.4 | 78.8 | 3.15×10^{-16} | 9.3×10^{-15} |
| Toxicity of tetanus toxin (TeNT) | 5250982 | 27 | 50.4 | 78.7 | 3.36×10^{-16} | 9.75×10^{-15} |
| Gap junction degradation | 190873 | 21 | 50.4 | 78.5 | 3.66×10^{-16} | 1.04×10^{-14} |
| Nephrin interactions | 373753 | 25 | 50.4 | 78.2 | 4.21×10^{-16} | 1.14×10^{-14} |
| GABA synthesis, release, reuptake and degradation | 888590 | 26 | 50.4 | 77 | 7.69×10^{-16} | 1.95×10^{-14} |

Strongest candidate SL partners for *CDH1* by SLIPT with observed and expected numbers of TCGA stomach cancer samples with low expression of both genes.