

Contents

Glossary	xiii
Acronyms	xvi
1 Introduction and Literature Review	1
1.1 Cancer Research in the Post-Genomic Era	1
1.1.1 Cancer is a Global Health Issue	2
1.1.1.1 The Genetics and Molecular Biology of Cancers	3
1.1.2 The Genomics Revolution in Cancer Research	3
1.1.2.1 High-Throughput Technologies	4
1.1.2.2 Bioinformatics and Genomic Data	5
1.1.3 Genomics Projects	5
1.1.3.1 The Cancer Genome Project	6
1.1.3.2 The Cancer Genome Atlas Project	6
1.1.4 Genomic Cancer Medicine	8
1.1.4.1 Cancer Genes and Driver Mutations	8
1.1.4.2 Precision Cancer Medicine	9
1.1.4.3 Molecular Diagnostics and Pan-Cancer Medicine	9
1.1.4.4 Targeted Therapeutics and Pharmacogenomics	10
1.1.5 Systems and Network Biology	11
1.2 Synthetic Lethal Cancer Medicine	12
1.2.1 Synthetic Lethal Genetic Interactions	13
1.2.2 Synthetic Lethal Concepts in Genetics	13
1.2.3 Synthetic Lethality in Model Systems	15
1.2.3.1 Synthetic Lethal Pathways and Networks	15
1.2.3.2 Evolution of Synthetic Lethality	16
1.2.4 Synthetic Lethality in Cancer	17
1.2.5 Clinical Impact of Synthetic Lethality in Cancer	18
1.2.6 High-throughput Screening for Synthetic Lethality	20
1.2.6.1 Synthetic Lethal Screens	21
1.2.7 Computational Prediction of Synthetic Lethality	22
1.2.7.1 Bioinformatics Approaches to Genetic Interactions	22
1.2.7.2 Comparative Genomics	24
1.2.7.3 Analysis and Modelling of Protein Data	26
1.2.7.4 Differential Gene Expression	28
1.2.7.5 Data Mining and Machine Learning	29

1.2.7.6	Mutual Exclusivity and Bimodality	31
1.2.7.7	Rationale for Further Development	33
1.3	E-cadherin as a Synthetic Lethal Target	33
1.3.1	The <i>CDH1</i> gene and its Biological Functions	33
1.3.1.1	Cytoskeleton	34
1.3.1.2	Extracellular and Tumour Micro-environment	34
1.3.1.3	Cell-Cell Adhesion and Signalling	34
1.3.2	<i>CDH1</i> as a Tumour (and Invasion) Suppressor	35
1.3.2.1	Breast Cancers and Invasion	35
1.3.3	Hereditary Diffuse Gastric (and Lobular Breast) Cancer	35
1.3.4	Cell Line Models of <i>CDH1</i> Null Mutations	37
1.4	Summary and Research Direction of Thesis	37
1.4.1	Thesis Aims	39
2	Methods and Resources	40
2.1	Bioinformatics Resources for Genomics Research	40
2.1.1	Public Data and Software Packages	40
2.1.1.1	Cancer Genome Atlas Data	41
2.1.1.2	Reactome and Annotation Data	42
2.2	Data Handling	42
2.2.1	Normalisation	42
2.2.2	Sample Triage	43
2.2.3	Metagenes and the Singular Value Decomposition	43
2.2.4	Candidate Triage and Integration with Screen Data	45
2.3	Techniques	46
2.3.1	Statistical Procedures and Tests	46
2.3.2	Gene Set Over-representation Analysis	47
2.3.3	Clustering	47
2.3.4	Heatmap	47
2.3.5	Modelling and Simulations	48
2.3.5.1	Receiver Operating Characteristic Curves	49
2.3.6	Resampling Analysis	49
2.4	Pathway Structure Methods	50
2.4.1	Network and Graph Analysis	50
2.4.2	Sourcing Graph Structure Data	51
2.4.3	Constructing Pathway Subgraphs	51
2.4.4	Network Analysis Metrics	52
2.5	Implementation	53
2.5.1	Computational Resources and Linux Utilities	53
2.5.2	R Language and Packages	54
2.5.3	High Performance and Parallel Computing	57
3	Methods Developed During Thesis	59
3.1	A Synthetic Lethal Detection Methodology	59
3.2	Synthetic Lethal Simulation and Modelling	61
3.2.1	A Model of Synthetic Lethality in Expression Data	62

3.2.2	Simulation Procedure	66
3.3	Detecting Simulated Synthetic Lethal Partners	69
3.3.1	Binomial Simulation of Synthetic Lethality	69
3.3.2	Multivariate Normal Simulation of Synthetic Lethality	71
3.3.2.1	Multivariate Normal Simulation with Correlated Genes	73
3.3.2.2	Specificity with Query-Correlated Pathways	81
3.4	Graph Structure Methods	83
3.4.1	Upstream and Downstream Gene Detection	83
3.4.1.1	Permutation Analysis for Statistical Significance	84
3.4.1.2	Hierarchy Based on Biological Context	84
3.4.2	Simulating Gene Expression from Graph Structures	85
3.5	Customised Functions and Packages Developed	90
3.5.1	Synthetic Lethal Interaction Prediction Tool	90
3.5.2	Data Visualisation	91
3.5.3	Extensions to the iGraph Package	93
3.5.3.1	Sampling Simulated Data from Graph Structures	93
3.5.3.2	Plotting Directed Graph Structures	93
3.5.3.3	Computing Information Centrality	94
3.5.3.4	Testing Pathway Structure with Permutation Testing .	94
3.5.3.5	Metapackage to Install iGraph Functions	95
4	Synthetic Lethal Analysis of Gene Expression Data	96
4.1	Synthetic Lethal Genes in Breast Cancer	97
4.1.1	Synthetic Lethal Pathways in Breast Cancer	98
4.1.2	Expression Profiles of Synthetic Lethal Partners	100
4.1.2.1	Subgroup Pathway Analysis	103
4.2	Comparing Synthetic Lethal Gene Candidates	105
4.2.1	Primary siRNA Screen Candidates	105
4.2.2	Comparison with Correlation	105
4.2.3	Comparison with Primary Screen Viability	108
4.2.4	Comparison with Secondary siRNA Screen Validation	110
4.2.5	Comparison to Primary Screen at Pathway Level	111
4.2.5.1	Resampling Genes for Pathway Enrichment	113
4.2.6	Integrating Synthetic Lethal Pathways and Screens	118
4.3	Synthetic Lethal Pathway Metagenes	119
4.4	Replication in Stomach Cancer	121
4.5	Discussion	122
4.5.1	Strengths of the SLIPT Methodology	122
4.5.2	Synthetic Lethal Pathways for E-cadherin	123
4.5.3	Replication and Validation	125
4.5.3.1	Integration with siRNA Screening	125
4.5.3.2	Replication across Tissues	126
4.6	Summary	126

5 Synthetic Lethal Pathway Structure	128
5.1 Synthetic Lethal Genes in Reactome Pathways	128
5.1.1 The PI3K/AKT Pathway	129
5.1.2 The Extracellular Matrix	131
5.1.3 G Protein Coupled Receptors	134
5.1.4 Gene Regulation and Translation	134
5.2 Network Analysis of Synthetic Lethal Genes	135
5.2.1 Gene Connectivity and Vertex Degree	136
5.2.2 Gene Importance and Centrality	137
5.2.2.1 Information Centrality	137
5.2.2.2 PageRank Centrality	139
5.3 Relationships between Synthetic Lethal Genes	141
5.3.1 Hierarchical Pathway Structure	141
5.3.1.1 Contextual Hierarchy of PI3K	141
5.3.1.2 Testing Contextual Hierarchy of Synthetic Lethal Genes	141
5.3.2 Upstream or Downstream Synthetic Lethality	145
5.3.2.1 Measuring Structure of Candidates within PI3K . . .	145
5.3.2.2 Resampling for Synthetic Lethal Pathway Structure .	147
5.4 Discussion	149
5.5 Summary	151
6 Simulation and Modelling of Synthetic Lethal Pathways	152
6.1 Synthetic Lethal Detection Methods	153
6.1.1 Performance of SLIPT and χ^2 across Quantiles	154
6.1.1.1 Correlated Query Genes affects Specificity	157
6.1.2 Alternative Synthetic Lethal Detection Strategies	159
6.1.2.1 Correlation for Synthetic Lethal Detection	160
6.1.2.2 Testing for Bimodality with BiSEp	161
6.2 Simulations with Graph Structures	162
6.2.1 Performance over Graph Structures	163
6.2.1.1 Simple Graph Structures	163
6.2.1.2 Constructed Graph Structures	166
6.2.2 Performance with Inhibitions	168
6.2.3 Synthetic Lethality across Graph Structures	174
6.2.4 Performance within a Simulated Human Genome	177
6.3 Simulations in More Complex Graph Structures	182
6.3.1 Simulations over Pathway-based Graphs	183
6.3.2 Pathway Structures in a Simulated Human Genome	185
6.4 Discussion	188
6.4.1 Simulation Procedure	188
6.4.2 Comparing Methods with Simulated Data	189
6.4.3 Design and Performance of SLIPT	190
6.4.4 Simulations from Graph Structures	192
6.5 Summary	193

7 Discussion	195
7.1 Synthetic Lethality and <i>CDH1</i> Biology	195
7.1.1 Established Functions of <i>CDH1</i>	196
7.1.2 The Molecular Role of <i>CDH1</i> in Cancer	196
7.2 Significance	197
7.2.1 Synthetic Lethality in the Genomic Era	197
7.2.2 Clinical Interventions based on Synthetic Lethality	199
7.3 Future Directions	200
7.4 Conclusions	202
References	204
A Sample Quality	228
A.1 Sample Correlation	228
A.2 Replicate Samples in TCGA Breast Cancer Data	231
B Software Used for Thesis	235
C Mutation Analysis in Breast Cancer	244
C.1 Synthetic Lethal Genes and Pathways	244
C.2 Synthetic Lethal Expression Profiles	245
C.3 Comparison to Primary Screen	248
C.3.1 Resampling Analysis	250
C.4 Compare SLIPT genes	252
D Metagene Analysis	254
D.1 Pathway Signature Expression	254
D.2 Somatic Mutation	263
D.3 Synthetic Lethal Reactome Metagenes	264
D.4 Expression of Somatic Mutations	266
E Intrinsic Subtyping	269
F Stomach Expression Analysis	271
F.1 Synthetic Lethal Genes and Pathways	271
F.2 Comparison to Primary Screen	275
F.2.1 Resampling Analysis	277
F.3 Metagene Analysis	279
G Synthetic Lethal Genes in Pathways	279
H Pathway Connectivity for Mutation SLIPT	287
I Information Centrality for Gene Essentiality	291
J Pathway Structure for Mutation SLIPT	294

K Performance of SLIPT and χ^2	297
K.1 Correlated Query Genes affects Specificity	303
L Simulations on Graph Structures	309
L.0.1 Simulations from Inhibiting Graph Structures	310
L.1 Simulation across Graph Structures	313
L.2 Simulations from Complex Graph Structures	317
L.2.1 Simulations from Complex Inhibiting Graphs	320
L.3 Simulations from Pathway Graph Structures	326

List of Figures

1.1	Synthetic genetic interactions	14
1.2	Synthetic lethality in cancer	17
2.1	Read count density	44
2.2	Read count sample mean	44
3.1	Framework for synthetic lethal prediction	60
3.2	Synthetic lethal prediction adapted for mutation	61
3.3	A model of synthetic lethal gene expression	63
3.4	Modelling synthetic lethal gene expression	64
3.5	Synthetic lethality with multiple genes	65
3.6	Simulating gene function	67
3.7	Simulating synthetic lethal gene function	67
3.8	Simulating synthetic lethal gene expression	68
3.9	Performance of binomial simulations	70
3.10	Comparison of statistical performance	70
3.11	Performance of multivariate normal simulations	72
3.12	Simulating expression with correlated gene blocks	74
3.13	Simulating expression with correlated gene blocks	75
3.14	Synthetic lethal prediction across simulations	77
3.15	Performance with correlations	78
3.16	Comparison of statistical performance with correlation structure	79
3.17	Performance with query correlations	80
3.18	Statistical evaluation of directional criteria	81
3.19	Performance of directional criteria	82
3.20	Simulated graph structures	86
3.21	Simulating expression from a graph structure	87
3.22	Simulating expression from graph structure with inhibitions	88
3.23	Demonstration of violin plots with custom features	92
3.24	Demonstration of annotated heatmap	92
3.25	Simulating graph structures	94
4.1	Synthetic lethal expression profiles of analysed samples	101
4.2	Comparison of SLIPT with siRNA	106
4.3	Comparison of SLIPT and siRNA genes with correlation	106
4.4	Comparison of SLIPT and siRNA genes with correlation	108
4.5	Comparison of SLIPT and siRNA genes with screen viability	109

4.6	Comparison of SLIPT genes with siRNA screen viability	109
4.7	Resampled intersection of SLIPT and siRNA candidate genes	114
5.1	synthetic lethality in the PI3K cascade	130
5.2	synthetic lethality in Elastic Fibre Formation	132
5.3	Synthetic lethality in Fibrin Clot Formation	133
5.4	Synthetic lethality and vertex degree	136
5.5	Synthetic lethality and centrality	139
5.6	Synthetic lethality and PageRank	140
5.7	Hierarchical structure of PI3K	142
5.8	Hierarchy score in PI3K against synthetic lethality in PI3K	143
5.9	Structure of synthetic lethality in PI3K	145
5.10	Structure of synthetic lethality resampling in PI3K	146
6.1	Performance of χ^2 and SLIPT across quantiles	155
6.2	Performance of χ^2 and SLIPT across quantiles with more genes	156
6.3	Performance of χ^2 and SLIPT across quantiles with query correlation .	157
6.4	Performance of χ^2 and SLIPT across quantiles with query correlation and more genes	158
6.5	Performance of negative correlation and SLIPT	161
6.6	Simple graph structures	164
6.7	Performance of simulations on a simple graph	165
6.8	Performance of simulations is similar in simple graphs	166
6.9	Performance of simulations on a pathway	167
6.10	Performance of simulations on a simple graph with inhibition	169
6.11	Performance is higher on a simple inhibiting graph	171
6.12	Performance of simulations on a constructed graph with inhibition . .	172
6.13	Performance is affected by inhibition in graphs	173
6.14	Detection of synthetic lethality within a graph structure	175
6.15	Performance of simulations including a simple graph	179
6.16	Performance on a simple graph improves with more genes	180
6.17	Performance on an inhibiting graph improves with more genes	181
6.18	Performance of simulations on the PI3K cascade	184
6.19	Performance of simulations including the PI3K cascade	186
6.20	Performance on pathways improves with more genes	187
A.1	Correlation profiles of removed samples	229
A.2	Correlation analysis and sample removal	230
A.3	Replicate excluded samples	231
A.4	Replicate samples with all remaining	232
A.5	Replicate samples with some excluded	233
C.1	Synthetic lethal expression profiles of analysed samples	246
C.2	Comparison of mtSLIPT to short interfering RNA (siRNA)	248
C.3	Compare mtSLIPT and siRNA genes with correlation	252
C.4	Compare mtSLIPT and siRNA genes with correlation	252
C.5	Compare mtSLIPT and siRNA genes with siRNA viability	253

D.1	Pathway metagene expression profiles	256
D.2	Expression profiles for constituent genes of PI3K	258
D.3	Expression profiles for estrogen receptor related genes	259
D.4	Pathway metagene expression profiles	260
D.5	Expression profiles for p53 related genes	261
D.6	Expression profiles for BRCA related genes	262
D.7	Somatic mutation against the PI3K metagene	263
D.8	Somatic mutation against PIK3CA metagene	266
D.9	Somatic mutation against PI3K protein	267
D.10	Somatic mutation against AKT protein	268
F.1	Synthetic lethal expression profiles of stomach samples	273
F.2	Comparison of SLIPT in stomach to siRNA	275
G.1	Synthetic lethality in the PI3K/AKT pathway	279
G.2	Synthetic lethality in the PI3K/AKT pathway in cancer	280
G.3	Synthetic lethality in the Extracellular Matrix	281
G.4	Synthetic lethality in the GPCRs	282
G.5	Synthetic lethality in the GPCR Downstream	283
G.6	Synthetic lethality in the Translation Elongation	284
G.7	Synthetic lethality in the Nonsense-mediated Decay	285
G.8	Synthetic lethality in the 3' UTR	286
H.1	Synthetic lethality and vertex degree	287
H.2	Synthetic lethality and centrality	288
H.3	Synthetic lethality and PageRank	289
I.1	Information centrality distribution	293
J.1	Synthetic lethality and heirarchy score in PI3K	294
J.2	Heirarchy score in PI3K against synthetic lethality in PI3K	295
J.3	Structure of synthetic lethality in PI3K	295
J.4	Structure of synthetic lethality resampling	296
K.1	Performance of χ^2 and SLIPT across quantiles	297
K.2	Performance of χ^2 and SLIPT across quantiles	299
K.3	Performance of χ^2 and SLIPT across quantiles with more genes	301
K.4	Performance of χ^2 and SLIPT across quantiles with query correlation	303
K.5	Performance of χ^2 and SLIPT across quantiles with query correlation	305
K.6	Performance of χ^2 and SLIPT across quantiles with query correlation and more genes	307
L.1	Performance of simulations on a simple graph	309
L.2	Performance of simulations on an inhibiting graph	310
L.3	Performance of simulations on a constructed graph with inhibition	311
L.4	Performance of simulations on a constructed graph with inhibition	312
L.5	Detection of synthetic lethality within a graph structure	313
L.6	Detection of synthetic lethality within an inhibiting graph	315

L.7	Detection of synthetic lethality within an inhibiting graph	316
L.8	Performance of simulations on a branching graph	317
L.9	Performance of simulations on a complex graph	318
L.10	Performance of simulations on a large graph	319
L.11	Performance of simulations on a branching graph with inhibition	320
L.12	Performance of simulations on a branching graph with inhibition	321
L.13	Performance of simulations on a complex graph with inhibition	322
L.14	Performance of simulations on a complex graph with inhibition	323
L.15	Performance of simulations on a large constructed graph with inhibition	324
L.16	Performance of simulations on a large constructed graph with inhibition	325
L.17	Performance of simulations on the $G_{\alpha i}$ signalling pathway	326
L.18	Performance of simulations including the $G_{\alpha i}$ signalling pathway	327

List of Tables

1.1	Methods for predicting genetic interactions	23
1.2	Methods for predicting synthetic lethality in cancer	23
1.3	Methods used by Wu <i>et al.</i> (2014)	25
2.1	Excluded samples by batch and clinical characteristics.	43
2.2	Computers used during thesis	53
2.3	Linux utilities and applications used during thesis	54
2.4	R installations used during thesis	55
2.5	R Packages used during thesis	55
2.6	R packages developed during thesis	57
4.1	Candidate synthetic lethal gene partners of <i>CDH1</i> from SLIPT	98
4.2	Pathways for <i>CDH1</i> partners from SLIPT	99
4.3	Pathways for clusters of <i>CDH1</i> partners from SLIPT	104
4.4	ANOVA for synthetic lethality and correlation with <i>CDH1</i>	107
4.5	Comparison of Synthetic Lethal Interaction Prediction Tool (SLIPT) genes against secondary siRNA screen	111
4.6	Pathways for <i>CDH1</i> partners from SLIPT and siRNA	112
4.7	Pathways for <i>CDH1</i> partners from SLIPT	115
4.8	Pathways for <i>CDH1</i> partners from SLIPT and siRNA primary screen .	116
4.9	Examples of candidate metagenes synthetic lethal for <i>CDH1</i> from SLIPT	120
5.1	ANOVA for synthetic lethality and vertex degree	137
5.2	ANOVA for synthetic lethality and information centrality	139
5.3	ANOVA for synthetic lethality and PageRank centrality	141
5.4	ANOVA for synthetic lethality and PI3K hierarchy	144
5.5	Resampling for pathway structure of synthetic lethal detection methods	148
B.1	Complete list of R packages used during this thesis	235
C.1	Candidate synthetic lethal gene partners of <i>CDH1</i> from mtSLIPT	244
C.2	Pathways for <i>CDH1</i> partners from mtSLIPT	245
C.3	Pathways for clusters of <i>CDH1</i> partners from mtSLIPT	247
C.4	Pathways for <i>CDH1</i> partners from mtSLIPT and siRNA	249
C.5	Pathways for <i>CDH1</i> partners from mtSLIPT	250
C.6	Pathways for <i>CDH1</i> partners from mtSLIPT and siRNA primary screen	251
D.1	Candidate synthetic lethal metagenes against <i>CDH1</i> from mtSLIPT . .	265

E.1	Comparison of intrinsic subtypes	269
F.1	Synthetic lethal gene partners of <i>CDH1</i> from SLIPT in stomach cancer	271
F.2	Pathways for <i>CDH1</i> partners from SLIPT in stomach cancer	272
F.3	Pathways for clusters of <i>CDH1</i> partners in stomach SLIPT	274
F.4	Pathways for <i>CDH1</i> partners from SLIPT and siRNA	276
F.5	Pathways for <i>CDH1</i> partners from SLIPT in stomach cancer	277
F.6	Pathways for <i>CDH1</i> partners from SLIPT in stomach and siRNA . . .	278
F.7	Synthetic lethal metagenes against <i>CDH1</i> in stomach cancer	279
H.1	ANOVA for synthetic lethality and vertex degree	290
H.2	ANOVA for synthetic lethality and information centrality	290
H.3	ANOVA for synthetic lethality and PageRank centrality	290
I.1	Information centrality for genes and molecules in the Reactome network	292
J.1	ANOVA for synthetic lethality and PI3K hierarchy	294
J.2	Resampling for pathway structure of synthetic lethal detection methods	296

Glossary

allele	A gene variant with a specific sequence and phenotype.
centrality	A network metric which identifies important vertices.
driver mutation	A mutation which promotes cancer growth.
E-cadherin	Epithelial cadherin (calcium-dependent adhesion), a cell-adhesion protein encoded by <i>CDH1</i> .
essential	A gene which is required to be functional or expressed for a cell or organism to be viable, grow or develop.
gene expression	A measure of the relative expression of each gene from the mRNA extracted from (pooled) cells.
graph or network	A mathematical structure modelling or depicting the relationships between elements.
hereditary	A trait or disease which has a genetic cause and is inherited from family members.
information centrality	A network centrality metric which uses the impact of removing a vertex or node on connections in the network.
intrinsic subtype	Distinguishing cancer by molecular and genetic features.
metagene	A consistent signal of expression for a collection of genes such as a biological pathway, derived from singular value decomposition.

microarray	A high-throughput technique to measure presence or abundance of nucleic acid sequences from binding to probes.
mutant	A variant or dysfunctional phenotype arising from a mutation in a gene.
mutation	A change in DNA sequence that disrupts gene function.
oncogene	A gene that potentially causes cancer, typically by over-expression or mutant gene variants.
PageRank centrality	A network centrality metric which uses eigenvectors with a scaling factor (Brin and Page, 1998).
pathway	A series of biomolecules that produces a particular product or biological function.
recurrent mutation	The repeated occurrence of mutations in a particular gene across cancers.
RNA-Seq	The generation of transcriptome data from sequencing RNA.
somatic mutation	A mutation that occurs in somatic cells, during a patient's lifespan.
synthetic lethal	Genetic interactions where inactivation of multiple genes is inviable (or deleterious) which are viable if inactivated separately.
tumour suppressor	A gene potentially causes cancer, typically by disruption of functions which protect the cell from cancer.
vertex degree	A network metric of connectivity of vertices which uses the number of edges connected to each vertex or node.
vertex or node	An element of a graph structure or network.
wild-type	A natural phenotype of a trait or the normally functional allele which encodes it.

Acronyms

ANOVA	Analysis of Variance.
CRAN	comprehensive R archive network.
ER	Estrogen Receptor.
FDR	False Discovery Rate.
GPCR	G Crotein Coupled Receptor.
mRNA	Messenger RNA.
mtSLIPT	Synthetic Lethal Interaction Prediction Tool (against mutation).
NMD	Nonsense-Mediated Decay.
PAM50	Prediction Analysis of Microarray 50.
PI3K	Phosphoinositide 3-kinase.
PR	Progesterone Receptor.
RNA	Ribonucleic Acid.
ROC	Reciever Operating Characteristic (curve).
siRNA	Short Interfering RNA.
SLIPT	Synthetic Lethal Interaction Prediction Tool.
TCGA	The Cancer Genome Atlas (genomics project).
UCSC	University of California, Santa Cruz.
UTR	Untranslated Region (of mRNA).

References

- Abeshouse, A., Ahn, J., Akbani, R., Ally, A., Amin, S., Andry, C.D., Annala, M., Aprikian, A., Armenia, J., Arora, A., *et al.* (2015) The Molecular Taxonomy of Primary Prostate Cancer. *Cell*, **163**(4): 1011–1025.
- Adler, D. (2005) *vioplot: Violin plot*. R package version 0.2.
- Akbani, R., Akdemir, K.C., Aksoy, B.A., Albert, M., Ally, A., Amin, S.B., Arachchi, H., Arora, A., Auman, J.T., Ayala, B., *et al.* (2015) Genomic Classification of Cutaneous Melanoma. *Cell*, **161**(7): 1681–1696.
- Akobeng, A.K. (2007) Understanding diagnostic tests 3: receiver operating characteristic curves. *Acta Pdiatrica*, **96**(5): 644–647.
- American Cancer Society (2017) Genetics and cancer. <https://www.cancer.org/cancer/cancer-causes/genetics.html>. Accessed: 22/03/2017.
- Anjomshoaa, A., Lin, Y.H., Black, M.A., McCall, J.L., Humar, B., Song, S., Fukuzawa, R., Yoon, H.S., Holzmann, B., Friederichs, J., *et al.* (2008) Reduced expression of a gene proliferation signature is associated with enhanced malignancy in colon cancer. *Br J Cancer*, **99**(6): 966–973.
- Araki, H., Knapp, C., Tsai, P., and Print, C. (2012) GeneSetDB: A comprehensive meta-database, statistical and visualisation framework for gene set analysis. *FEBS Open Bio*, **2**: 76–82.
- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., *et al.* (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet*, **25**(1): 25–29.
- Ashworth, A. (2008) A synthetic lethal therapeutic approach: poly(adp) ribose polymerase inhibitors for the treatment of cancers deficient in dna double-strand break repair. *J Clin Oncol*, **26**(22): 3785–90.

- Ashworth, A., Lord, C.J., and Reis-Filho, J.S. (2011) Genetic interactions in cancer progression and treatment. *Cell*, **145**(1): 30–38.
- Audeh, M.W., Carmichael, J., Penson, R.T., Friedlander, M., Powell, B., Bell-McGuinn, K.M., Scott, C., Weitzel, J.N., Oaknin, A., Loman, N., *et al.* (2010) Oral poly(adp-ribose) polymerase inhibitor olaparib in patients with *BRCA1* or *BRCA2* mutations and recurrent ovarian cancer: a proof-of-concept trial. *Lancet*, **376**(9737): 245–51.
- Babyak, M.A. (2004) What you see may not be what you get: a brief, nontechnical introduction to overfitting in regression-type models. *Psychosom Med*, **66**(3): 411–21.
- Bamford, S., Dawson, E., Forbes, S., Clements, J., Pettett, R., Dogan, A., Flanagan, A., Teague, J., Futreal, P.A., Stratton, M.R., *et al.* (2004) The COSMIC (Catalogue of Somatic Mutations in Cancer) database and website. *Br J Cancer*, **91**(2): 355–358.
- Barabási, A.L. and Albert, R. (1999) Emergence of scaling in random networks. *Science*, **286**(5439): 509–12.
- Barabási, A.L., Gulbahce, N., and Loscalzo, J. (2011) Network medicine: a network-based approach to human disease. *Nat Rev Genet*, **12**(1): 56–68.
- Barabási, A.L. and Oltvai, Z.N. (2004) Network biology: understanding the cell's functional organization. *Nat Rev Genet*, **5**(2): 101–13.
- Barrat, A. and Weigt, M. (2000) On the properties of small-world network models. *The European Physical Journal B - Condensed Matter and Complex Systems*, **13**(3): 547–560.
- Barretina, J., Caponigro, G., Stransky, N., Venkatesan, K., Margolin, A.A., Kim, S., Wilson, C.J., Lehar, J., Kryukov, G.V., Sonkin, D., *et al.* (2012) The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. *Nature*, **483**(7391): 603–607.
- Barry, W.T. (2016) *safe: Significance Analysis of Function and Expression*. R package version 3.14.0.

- Baryshnikova, A., Costanzo, M., Dixon, S., Vizeacoumar, F.J., Myers, C.L., Andrews, B., and Boone, C. (2010a) Synthetic genetic array (sga) analysis in *saccharomyces cerevisiae* and *schizosaccharomyces pombe*. *Methods Enzymol*, **470**: 145–79.
- Baryshnikova, A., Costanzo, M., Kim, Y., Ding, H., Koh, J., Toufighi, K., Youn, J.Y., Ou, J., San Luis, B.J., Bandyopadhyay, S., *et al.* (2010b) Quantitative analysis of fitness and genetic interactions in yeast on a genome scale. *Nat Meth*, **7**(12): 1017–1024.
- Bass, A.J., Thorsson, V., Shmulevich, I., Reynolds, S.M., Miller, M., Bernard, B., Hinoue, T., Laird, P.W., Curtis, C., Shen, H., *et al.* (2014) Comprehensive molecular characterization of gastric adenocarcinoma. *Nature*, **513**(7517): 202–209.
- Bates, D. and Maechler, M. (2016) *Matrix: Sparse and Dense Matrix Classes and Methods*. R package version 1.2-7.1.
- Bateson, W. and Mendel, G. (1909) *Mendel's principles of heredity, by W. Bateson*. University Press, Cambridge [Eng.].
- Becker, K.F., Atkinson, M.J., Reich, U., Becker, I., Nekarda, H., Siewert, J.R., and Hfler, H. (1994) E-cadherin gene mutations provide clues to diffuse type gastric carcinomas. *Cancer Research*, **54**(14): 3845–3852.
- Bell, D., Berchuck, A., Birrer, M., Chien, J., Cramer, D., Dao, F., Dhir, R., DiSaia, P., Gabra, H., Glenn, P., *et al.* (2011) Integrated genomic analyses of ovarian carcinoma. *Nature*, **474**(7353): 609–615.
- Benjamini, Y. and Hochberg, Y. (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B (Methodological)*, **57**(1): 289–300.
- Berx, G., Cleton-Jansen, A.M., Nollet, F., de Leeuw, W.J., van de Vijver, M., Cornelisse, C., and van Roy, F. (1995) E-cadherin is a tumour/invasion suppressor gene mutated in human lobular breast cancers. *EMBO J*, **14**(24): 6107–15.
- Berx, G., Cleton-Jansen, A.M., Strumane, K., de Leeuw, W.J., Nollet, F., van Roy, F., and Cornelisse, C. (1996) E-cadherin is inactivated in a majority of invasive human lobular breast cancers by truncation mutations throughout its extracellular domain. *Oncogene*, **13**(9): 1919–25.

- Berx, G. and van Roy, F. (2009) Involvement of members of the cadherin superfamily in cancer. *Cold Spring Harb Perspect Biol*, **1**: a003129.
- Bitler, B.G., Aird, K.M., Garipov, A., Li, H., Amatangelo, M., Kossenkov, A.V., Schultz, D.C., Liu, Q., Shih Ie, M., Conejo-Garcia, J.R., *et al.* (2015) Synthetic lethality by targeting ezh2 methyltransferase activity in arid1a-mutated cancers. *Nat Med*, **21**(3): 231–8.
- Blake, J.A., Christie, K.R., Dolan, M.E., Drabkin, H.J., Hill, D.P., Ni, L., Sitnikov, D., Burgess, S., Buza, T., Gresham, C., *et al.* (2015) Gene Ontology Consortium: going forward. *Nucleic Acids Res*, **43**(Database issue): D1049–1056.
- Boone, C., Bussey, H., and Andrews, B.J. (2007) Exploring genetic interactions and networks with yeast. *Nat Rev Genet*, **8**(6): 437–49.
- Borgatti, S.P. (2005) Centrality and network flow. *Social Networks*, **27**(1): 55 – 71.
- Boucher, B. and Jenna, S. (2013) Genetic interaction networks: better understand to better predict. *Front Genet*, **4**: 290.
- Bozovic-Spasojevic, I., Azambuja, E., McCaskill-Stevens, W., Dinh, P., and Cardoso, F. (2012) Chemoprevention for breast cancer. *Cancer treatment reviews*, **38**(5): 329–339.
- Breiman, L. (2001) Random forests. *Machine Learning*, **45**(1): 5–32.
- Brin, S. and Page, L. (1998) The anatomy of a large-scale hypertextual web search engine. *Computer Networks and ISDN Systems*, **30**(1): 107 – 117.
- Brouxhon, S.M., Kyrkanides, S., Teng, X., Athar, M., Ghazizadeh, S., Simon, M., O'Banion, M.K., and Ma, L. (2014) Soluble E-cadherin: a critical oncogene modulating receptor tyrosine kinases, MAPK and PI3K/Akt/mTOR signaling. *Oncogene*, **33**(2): 225–235.
- Brückner, A., Polge, C., Lentze, N., Auerbach, D., and Schlattner, U. (2009) Yeast two-hybrid, a powerful tool for systems biology. *Int J Mol Sci*, **10**(6): 2763–2788.
- Bryant, H.E., Schultz, N., Thomas, H.D., Parker, K.M., Flower, D., Lopez, E., Kyle, S., Meuth, M., Curtin, N.J., and Helleday, T. (2005) Specific killing of *BRCA2*-deficient tumours with inhibitors of polyadprbose polymerase. *Nature*, **434**(7035): 913–7.

- Bussey, H., Andrews, B., and Boone, C. (2006) From worm genetic networks to complex human diseases. *Nat Genet*, **38**(8): 862–3.
- Butland, G., Babu, M., Diaz-Mejia, J.J., Bohdana, F., Phanse, S., Gold, B., Yang, W., Li, J., Gagarinova, A.G., Pogoutse, O., et al. (2008) esga: E. coli synthetic genetic array analysis. *Nat Methods*, **5**(9): 789–95.
- cBioPortal for Cancer Genomics (cBioPortal) (2017) cBioPortal for Cancer Genomics. <http://www.cbioportal.org/>. Accessed: 26/03/2017.
- Cerami, E.G., Gross, B.E., Demir, E., Rodchenkov, I., Babur, O., Anwar, N., Schultz, N., Bader, G.D., and Sander, C. (2011) Pathway Commons, a web resource for biological pathway data. *Nucleic Acids Res*, **39**(Database issue): D685–690.
- Chen, A., Beetham, H., Black, M.A., Priya, R., Telford, B.J., Guest, J., Wiggins, G.A.R., Godwin, T.D., Yap, A.S., and Guilford, P.J. (2014) E-cadherin loss alters cytoskeletal organization and adhesion in non-malignant breast cells but is insufficient to induce an epithelial-mesenchymal transition. *BMC Cancer*, **14**(1): 552.
- Chen, S. and Parmigiani, G. (2007) Meta-analysis of BRCA1 and BRCA2 penetrance. *J Clin Oncol*, **25**(11): 1329–1333.
- Chipman, K. and Singh, A. (2009) Predicting genetic interactions with random walks on biological networks. *BMC Bioinformatics*, **10**(1): 17.
- Christofori, G. and Semb, H. (1999) The role of the cell-adhesion molecule E-cadherin as a tumour-suppressor gene. *Trends in Biochemical Sciences*, **24**(2): 73 – 76.
- Ciriello, G., Gatza, M.L., Beck, A.H., Wilkerson, M.D., Rhie, S.K., Pastore, A., Zhang, H., McLellan, M., Yau, C., Kandoth, C., et al. (2015) Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. *Cell*, **163**(2): 506–519.
- Clark, M.J. (2004) Endogenous Regulator of G Protein Signaling Proteins Suppress G_o-Dependent μ -Opioid Agonist-Mediated Adenylyl Cyclase Supersensitization. *Journal of Pharmacology and Experimental Therapeutics*, **310**(1): 215–222.
- Collingridge, D.S. (2013) A primer on quantitized data analysis and permutation testing. *Journal of Mixed Methods Research*, **7**(1): 81–97.

- Collins, F.S. and Barker, A.D. (2007) Mapping the cancer genome. Pinpointing the genes involved in cancer will help chart a new course across the complex landscape of human malignancies. *Sci Am*, **296**(3): 50–57.
- Collisson, E., Campbell, J., Brooks, A., Berger, A., Lee, W., Chmielecki, J., Beer, D., Cope, L., Creighton, C., Danilova, L., *et al.* (2014) Comprehensive molecular profiling of lung adenocarcinoma. *Nature*, **511**(7511): 543–550.
- Costanzo, M., Baryshnikova, A., Bellay, J., Kim, Y., Spear, E.D., Sevier, C.S., Ding, H., Koh, J.L., Toufighi, K., Mostafavi, S., *et al.* (2010) The genetic landscape of a cell. *Science*, **327**(5964): 425–31.
- Costanzo, M., Baryshnikova, A., Myers, C.L., Andrews, B., and Boone, C. (2011) Charting the genetic interaction map of a cell. *Curr Opin Biotechnol*, **22**(1): 66–74.
- Courtney, K.D., Corcoran, R.B., and Engelman, J.A. (2010) The PI3K pathway as drug target in human cancer. *J Clin Oncol*, **28**(6): 1075–1083.
- Creighton, C.J., Morgan, M., Gunaratne, P.H., Wheeler, D.A., Gibbs, R.A., Robertson, A., Chu, A., Beroukhim, R., Cibulskis, K., Signoretti, S., *et al.* (2013) Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature*, **499**(7456): 43–49.
- Croft, D., Mundo, A.F., Haw, R., Milacic, M., Weiser, J., Wu, G., Caudy, M., Garapati, P., Gillespie, M., Kamdar, M.R., *et al.* (2014) The Reactome pathway knowledge-base. *Nucleic Acids Res*, **42**(database issue): D472D477.
- Crunkhorn, S. (2014) Cancer: Predicting synthetic lethal interactions. *Nat Rev Drug Discov*, **13**(11): 812.
- Csardi, G. and Nepusz, T. (2006) The igraph software package for complex network research. *InterJournal, Complex Systems*: 1695.
- Dai, X., Li, T., Bai, Z., Yang, Y., Liu, X., Zhan, J., and Shi, B. (2015) Breast cancer intrinsic subtype classification, clinical use and future trends. *Am J Cancer Res*, **5**(10): 2929–2943.
- Davierwala, A.P., Haynes, J., Li, Z., Brost, R.L., Robinson, M.D., Yu, L., Mnaimneh, S., Ding, H., Zhu, H., Chen, Y., *et al.* (2005) The synthetic genetic interaction spectrum of essential genes. *Nat Genet*, **37**(10): 1147–1152.

- De Leeuw, W.J., Berx, G., Vos, C.B., Peterse, J.L., Van de Vijver, M.J., Litvinov, S., Van Roy, F., Cornelisse, C.J., and Cleton-Jansen, A.M. (1997) Simultaneous loss of E-cadherin and catenins in invasive lobular breast cancer and lobular carcinoma in situ. *J Pathol*, **183**(4): 404–11.
- De Santis, G., Miotti, S., Mazzi, M., Canevari, S., and Tomassetti, A. (2009) E-cadherin directly contributes to PI3K/AKT activation by engaging the PI3K-p85 regulatory subunit to adherens junctions of ovarian carcinoma cells. *Oncogene*, **28**(9): 1206–1217.
- Demir, E., Babur, O., Rodchenkov, I., Aksoy, B.A., Fukuda, K.I., Gross, B., Sumer, O.S., Bader, G.D., and Sander, C. (2013) Using biological pathway data with Paxtools. *PLoS Comput Biol*, **9**(9): e1003194.
- Deshpande, R., Asiedu, M.K., Klebig, M., Sutor, S., Kuzmin, E., Nelson, J., Piotrowski, J., Shin, S.H., Yoshida, M., Costanzo, M., *et al.* (2013) A comparative genomic approach for identifying synthetic lethal interactions in human cancer. *Cancer Res*, **73**(20): 6128–36.
- Dickson, D. (1999) Wellcome funds cancer database. *Nature*, **401**(6755): 729.
- Dijkstra, E.W. (1959) A note on two problems in connexion with graphs. *Numerische Mathematik*, **1**(1): 269–271.
- Dixon, S.J., Andrews, B.J., and Boone, C. (2009) Exploring the conservation of synthetic lethal genetic interaction networks. *Commun Integr Biol*, **2**(2): 78–81.
- Dixon, S.J., Fedyshyn, Y., Koh, J.L., Prasad, T.S., Chahwan, C., Chua, G., Toufighi, K., Baryshnikova, A., Hayles, J., Hoe, K.L., *et al.* (2008) Significant conservation of synthetic lethal genetic interaction networks between distantly related eukaryotes. *Proc Natl Acad Sci U S A*, **105**(43): 16653–8.
- Dong, L.L., Liu, L., Ma, C.H., Li, J.S., Du, C., Xu, S., Han, L.H., Li, L., and Wang, X.W. (2012) E-cadherin promotes proliferation of human ovarian cancer cells in vitro via activating MEK/ERK pathway. *Acta Pharmacol Sin*, **33**(6): 817–822.
- Dorsam, R.T. and Gutkind, J.S. (2007) G-protein-coupled receptors and cancer. *Nat Rev Cancer*, **7**(2): 79–94.
- Erdős, P. and Rényi, A. (1959) On random graphs I. *Publ Math Debrecen*, **6**: 290–297.

- Erdős, P. and Rényi, A. (1960) On the evolution of random graphs. In *Publ. Math. Inst. Hung. Acad. Sci*, volume 5, 17–61.
- Eroles, P., Bosch, A., Perez-Fidalgo, J.A., and Lluch, A. (2012) Molecular biology in breast cancer: intrinsic subtypes and signaling pathways. *Cancer Treat Rev*, **38**(6): 698–707.
- Farmer, H., McCabe, N., Lord, C.J., Tutt, A.N., Johnson, D.A., Richardson, T.B., Santarosa, M., Dillon, K.J., Hickson, I., Knights, C., *et al.* (2005) Targeting the dna repair defect in BRCA mutant cells as a therapeutic strategy. *Nature*, **434**(7035): 917–21.
- Fawcett, T. (2006) An introduction to ROC analysis. *Pattern Recognition Letters*, **27**(8): 861 – 874. {ROC} Analysis in Pattern Recognition.
- Fece de la Cruz, F., Gapp, B.V., and Nijman, S.M. (2015) Synthetic lethal vulnerabilities of cancer. *Annu Rev Pharmacol Toxicol*, **55**: 513–531.
- Ferlay, J., Soerjomataram, I., Dikshit, R., Eser, S., Mathers, C., Rebelo, M., Parkin, D.M., Forman, D., and Bray, F. (2015) Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*, **136**(5): E359–386.
- Fisher, R.A. (1919) Xv.the correlation between relatives on the supposition of mendelian inheritance. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, **52**(02): 399–433.
- Fong, P.C., Boss, D.S., Yap, T.A., Tutt, A., Wu, P., Mergui-Roelvink, M., Mortimer, P., Swaisland, H., Lau, A., O'Connor, M.J., *et al.* (2009) Inhibition of poly(adp-ribose) polymerase in tumors from BRCA mutation carriers. *N Engl J Med*, **361**(2): 123–34.
- Fong, P.C., Yap, T.A., Boss, D.S., Carden, C.P., Mergui-Roelvink, M., Gourley, C., De Greve, J., Lubinski, J., Shanley, S., Messiou, C., *et al.* (2010) Poly(adp)-ribose polymerase inhibition: frequent durable responses in BRCA carrier ovarian cancer correlating with platinum-free interval. *J Clin Oncol*, **28**(15): 2512–9.
- Forbes, S.A., Beare, D., Gunasekaran, P., Leung, K., Bindal, N., Boutselakis, H., Ding, M., Bamford, S., Cole, C., Ward, S., *et al.* (2015) COSMIC: exploring the world's

knowledge of somatic mutations in human cancer. *Nucleic Acids Res*, **43**(Database issue): D805–811.

Fraser, A. (2004) Towards full employment: using RNAi to find roles for the redundant. *Oncogene*, **23**(51): 8346–52.

Fromental-Romain, C., Warot, X., Lakkaraju, S., Favier, B., Haack, H., Birling, C., Dierich, A., Doll e, P., and Chambon, P. (1996) Specific and redundant functions of the paralogous Hoxa-9 and Hoxd-9 genes in forelimb and axial skeleton patterning. *Development*, **122**(2): 461–472.

Futreal, P.A., Coin, L., Marshall, M., Down, T., Hubbard, T., Wooster, R., Rahman, N., and Stratton, M.R. (2004) A census of human cancer genes. *Nat Rev Cancer*, **4**(3): 177–183.

Futreal, P.A., Kasprzyk, A., Birney, E., Mullikin, J.C., Wooster, R., and Stratton, M.R. (2001) Cancer and genomics. *Nature*, **409**(6822): 850–852.

Gao, B. and Roux, P.P. (2015) Translational control by oncogenic signaling pathways. *Biochimica et Biophysica Acta*, **1849**(7): 753–65.

Gatza, M.L., Kung, H.N., Blackwell, K.L., Dewhirst, M.W., Marks, J.R., and Chi, J.T. (2011) Analysis of tumor environmental response and oncogenic pathway activation identifies distinct basal and luminal features in HER2-related breast tumor subtypes. *Breast Cancer Res*, **13**(3): R62.

Gatza, M.L., Lucas, J.E., Barry, W.T., Kim, J.W., Wang, Q., Crawford, M.D., Datto, M.B., Kelley, M., Mathey-Prevot, B., Potti, A., *et al.* (2010) A pathway-based classification of human breast cancer. *Proc Natl Acad Sci USA*, **107**(15): 6994–6999.

Gatza, M.L., Silva, G.O., Parker, J.S., Fan, C., and Perou, C.M. (2014) An integrated genomics approach identifies drivers of proliferation in luminal-subtype human breast cancer. *Nat Genet*, **46**(10): 1051–1059.

Gentleman, R.C., Carey, V.J., Bates, D.M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B., Gautier, L., Ge, Y., Gentry, J., *et al.* (2004) Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol*, **5**(10): R80.

Genz, A. and Bretz, F. (2009) Computation of multivariate normal and t probabilities. In *Lecture Notes in Statistics*, volume 195. Springer-Verlag, Heidelberg.

- Genz, A., Bretz, F., Miwa, T., Mi, X., Leisch, F., Scheipl, F., and Hothorn, T. (2016) *mvtnorm: Multivariate Normal and t Distributions*. R package version 1.0-5. URL.
- Glaire, M.A., Brown, M., Church, D.N., and Tomlinson, I. (2017) Cancer predisposition syndromes: lessons for truly precision medicine. *J Pathol*, **241**(2): 226–235.
- Globus (Globus) (2017) Research data management simplified. <https://www.globus.org/>. Accessed: 25/03/2017.
- Goodwin, S., McPherson, J.D., and McCombie, W.R. (2016) Coming of age: ten years of next-generation sequencing technologies. *Nat Rev Genet*, **17**(6): 333–351.
- Grady, W.M., Willis, J., Guilford, P.J., Dunbier, A.K., Toro, T.T., Lynch, H., Wiesner, G., Ferguson, K., Eng, C., Park, J.G., *et al.* (2000) Methylation of the CDH1 promoter as the second genetic hit in hereditary diffuse gastric cancer. *Nat Genet*, **26**(1): 16–17.
- Graziano, F., Humar, B., and Guilford, P. (2003) The role of the E-cadherin gene (*CDH1*) in diffuse gastric cancer susceptibility: from the laboratory to clinical practice. *Annals of Oncology*, **14**(12): 1705–1713.
- Guaragnella, N., Palermo, V., Galli, A., Moro, L., Mazzoni, C., and Giannattasio, S. (2014) The expanding role of yeast in cancer research and diagnosis: insights into the function of the oncosuppressors p53 and BRCA1/2. *FEMS Yeast Res*, **14**(1): 2–16.
- Güell, O., Sagus, F., and Serrano, M. (2014) Essential plasticity and redundancy of metabolism unveiled by synthetic lethality analysis. *PLoS Comput Biol*, **10**(5): e1003637.
- Guilford, P. (1999) E-cadherin downregulation in cancer: fuel on the fire? *Molecular Medicine Today*, **5**(4): 172 – 177.
- Guilford, P., Hopkins, J., Harraway, J., McLeod, M., McLeod, N., Harawira, P., Taite, H., Scouler, R., Miller, A., and Reeve, A.E. (1998) E-cadherin germline mutations in familial gastric cancer. *Nature*, **392**(6674): 402–5.
- Guilford, P., Humar, B., and Blair, V. (2010) Hereditary diffuse gastric cancer: translation of *CDH1* germline mutations into clinical practice. *Gastric Cancer*, **13**(1): 1–10.

- Guilford, P.J., Hopkins, J.B., Grady, W.M., Markowitz, S.D., Willis, J., Lynch, H., Rajput, A., Wiesner, G.L., Lindor, N.M., Burgart, L.J., *et al.* (1999) E-cadherin germline mutations define an inherited cancer syndrome dominated by diffuse gastric cancer. *Hum Mutat*, **14**(3): 249–55.
- Guo, J., Liu, H., and Zheng, J. (2016) SynLethDB: synthetic lethality database toward discovery of selective and sensitive anticancer drug targets. *Nucleic Acids Res*, **44**(D1): D1011–1017.
- Hajian-Tilaki, K. (2013) Receiver Operating Characteristic (ROC) Curve Analysis for Medical Diagnostic Test Evaluation. *Caspian J Intern Med*, **4**(2): 627–635.
- Hall, M., Frank, E., Holmes, G., Pfahringer, B., Reutemann, P., and Witten, I.H. (2009) The weka data mining software: an update. *SIGKDD Explor Newsl*, **11**(1): 10–18.
- Hamerman, P.S., Lawrence, M.S., Voet, D., Jing, R., Cibulskis, K., Sivachenko, A., Stojanov, P., McKenna, A., Lander, E.S., Gabriel, S., *et al.* (2012) Comprehensive genomic characterization of squamous cell lung cancers. *Nature*, **489**(7417): 519–525.
- Hanahan, D. and Weinberg, R.A. (2000) The hallmarks of cancer. *Cell*, **100**(1): 57–70.
- Hanahan, D. and Weinberg, R.A. (2011) Hallmarks of cancer: the next generation. *Cell*, **144**(5): 646–674.
- Hanna, S. (2003) Cancer incidence in new zealand (2003-2007). In D. Forman, D. Bray F Brewster, C. Gombe Mbalawa, B. Kohler, M. Piñeros, E. Steliarova-Foucher, R. Swaminathan, and J. Ferlay (editors), *Cancer Incidence in Five Continents*, volume X, 902–907. International Agency for Research on Cancer, Lyon, France. Electronic version <http://ci5.iarc.fr> Accessed 22/03/2017.
- Hansford, S., Kaurah, P., Li-Chang, H., Woo, M., Senz, J., Pinheiro, H., Schrader, K.A., Schaeffer, D.F., Shumansky, K., Zogopoulos, G., *et al.* (2015) Hereditary Diffuse Gastric Cancer Syndrome: CDH1 Mutations and Beyond. *JAMA Oncol*, **1**(1): 23–32.
- Heiskanen, M.A. and Aittokallio, T. (2012) Mining high-throughput screens for cancer drug targets-lessons from yeast chemical-genomic profiling and synthetic lethality.

Wiley Interdisciplinary Reviews: Data Mining and Knowledge Discovery, **2**(3): 263–272.

- Hell, P. (1976) Graphs with given neighbourhoods i. problèmes combinatoires at théorie des graphes. *Proc Coll Int CNRS, Orsay*, **260**: 219–223.
- Higgins, M.E., Claremont, M., Major, J.E., Sander, C., and Lash, A.E. (2007) CancerGenes: a gene selection resource for cancer genome projects. *Nucleic Acids Res*, **35**(Database issue): D721–726.
- Hillenmeyer, M.E. (2008) The chemical genomic portrait of yeast: uncovering a phenotype for all genes. *Science*, **320**: 362–365.
- Hoadley, K.A., Yau, C., Wolf, D.M., Cherniack, A.D., Tamborero, D., Ng, S., Leiserson, M.D., Niu, B., McLellan, M.D., Uzunangelov, V., et al. (2014) Multiplatform analysis of 12 cancer types reveals molecular classification within and across tissues of origin. *Cell*, **158**(4): 929–944.
- Hoehndorf, R., Hardy, N.W., Osumi-Sutherland, D., Tweedie, S., Schofield, P.N., and Gkoutos, G.V. (2013) Systematic analysis of experimental phenotype data reveals gene functions. *PLoS ONE*, **8**(4): e60847.
- Holm, S. (1979) A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, **6**(2): 65–70.
- Hopkins, A.L. (2008) Network pharmacology: the next paradigm in drug discovery. *Nat Chem Biol*, **4**(11): 682–690.
- Hu, Z., Fan, C., Oh, D.S., Marron, J.S., He, X., Qaqish, B.F., Livasy, C., Carey, L.A., Reynolds, E., Dressler, L., et al. (2006) The molecular portraits of breast tumors are conserved across microarray platforms. *BMC Genomics*, **7**: 96.
- Huang, E., Cheng, S., Dressman, H., Pittman, J., Tsou, M., Horng, C., Bild, A., Iversen, E., Liao, M., Chen, C., et al. (2003) Gene expression predictors of breast cancer outcomes. *Lancet*, **361**: 1590–1596.
- Hutchison, C.A., Chuang, R.Y., Noskov, V.N., Assad-Garcia, N., Deerinck, T.J., Elliston, M.H., Gill, J., Kannan, K., Karas, B.J., Ma, L., et al. (2016) Design and synthesis of a minimal bacterial genome. *Science*, **351**(6280): aad6253.

- International HapMap 3 Consortium (HapMap) (2003) The International HapMap Project. *Nature*, **426**(6968): 789–796.
- Jeanes, A., Gottardi, C.J., and Yap, A.S. (2008) Cadherins and cancer: how does cadherin dysfunction promote tumor progression? *Oncogene*, **27**(55): 6920–6929.
- Jerby-Arnon, L., Pfetzer, N., Waldman, Y., McGarry, L., James, D., Shanks, E., Seashore-Ludlow, B., Weinstock, A., Geiger, T., Clemons, P., *et al.* (2014) Predicting cancer-specific vulnerability via data-driven detection of synthetic lethality. *Cell*, **158**(5): 1199–1209.
- Joachims, T. (1999) Making large-scale support vector machine learning practical. In S. Bernhard, lkopf, J.C.B. Christopher, and J.S. Alexander (editors), *Advances in kernel methods*, 169–184. MIT Press.
- Ju, Z., Liu, W., Roebuck, P.L., Siwak, D.R., Zhang, N., Lu, Y., Davies, M.A., Akbani, R., Weinstein, J.N., Mills, G.B., *et al.* (2015) Development of a robust classifier for quality control of reverse-phase protein arrays. *Bioinformatics*, **31**(6): 912.
- Kaelin, Jr, W. (2005) The concept of synthetic lethality in the context of anticancer therapy. *Nat Rev Cancer*, **5**(9): 689–98.
- Kaelin, Jr, W. (2009) Synthetic lethality: a framework for the development of wiser cancer therapeutics. *Genome Med*, **1**: 99.
- Kamada, T. and Kawai, S. (1989) An algorithm for drawing general undirected graphs. *Information Processing Letters*, **31**(1): 7–15.
- Kawai, J., Shinagawa, A., Shibata, K., Yoshino, M., Itoh, M., Ishii, Y., Arakawa, T., Hara, A., Fukunishi, Y., Konno, H., *et al.* (2001) Functional annotation of a full-length mouse cDNA collection. *Nature*, **409**(6821): 685–690.
- Kelley, R. and Ideker, T. (2005) Systematic interpretation of genetic interactions using protein networks. *Nat Biotech*, **23**(5): 561–566.
- Kelly, S.T. (2013) *Statistical Predictions of Synthetic Lethal Interactions in Cancer*. Dissertation, University of Otago.
- Keshava Prasad, T.S., Goel, R., Kandasamy, K., Keerthikumar, S., Kumar, S., Mathivanan, S., Telikicherla, D., Raju, R., Shafreeen, B., Venugopal, A., *et al.* (2009)

Human Protein Reference Database—2009 update. *Nucleic Acids Res*, **37**(Database issue): D767–772.

Kim, N.G., Koh, E., Chen, X., and Gumbiner, B.M. (2011) E-cadherin mediates contact inhibition of proliferation through Hippo signaling-pathway components. *Proc Natl Acad Sci USA*, **108**(29): 11930–11935.

Koboldt, D.C., Fulton, R.S., McLellan, M.D., Schmidt, H., Kalicki-Veizer, J., McMichael, J.F., Fulton, L.L., Dooling, D.J., Ding, L., Mardis, E.R., *et al.* (2012) Comprehensive molecular portraits of human breast tumours. *Nature*, **490**(7418): 61–70.

Kockel, L., Zeitlinger, J., Staszewski, L.M., Mlodzik, M., and Bohmann, D. (1997) Jun in drosophila development: redundant and nonredundant functions and regulation by two mapk signal transduction pathways. *Genes & Development*, **11**(13): 1748–1758.

Kozlov, K.N., Gursky, V.V., Kulakovskiy, I.V., and Samsonova, M.G. (2015) Sequence-based model of gap gene regulation network. *BMC Genomics*, **15**(Suppl 12): S6.

Kranthi, S., Rao, S., and Manimaran, P. (2013) Identification of synthetic lethal pairs in biological systems through network information centrality. *Mol BioSyst*, **9**(8): 2163–2167.

Kroepil, F., Fluegen, G., Totikov, Z., Baldus, S.E., Vay, C., Schauer, M., Topp, S.A., Esch, J.S., Knoefel, W.T., and Stoecklein, N.H. (2012) Down-regulation of CDH1 is associated with expression of SNAI1 in colorectal adenomas. *PLoS ONE*, **7**(9): e46665.

Lander, E.S. (2011) Initial impact of the sequencing of the human genome. *Nature*, **470**(7333): 187–197.

Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., *et al.* (2001) Initial sequencing and analysis of the human genome. *Nature*, **409**(6822): 860–921.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009) Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol*, **10**(3): R25.

- Latora, V. and Marchiori, M. (2001) Efficient behavior of small-world networks. *Phys Rev Lett*, **87**: 198701.
- Laufer, C., Fischer, B., Billmann, M., Huber, W., and Boutros, M. (2013) Mapping genetic interactions in human cancer cells with RNAi and multiparametric phenotyping. *Nat Methods*, **10**(5): 427–31.
- Law, C.W., Chen, Y., Shi, W., and Smyth, G.K. (2014) voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol*, **15**(2): R29.
- Le Meur, N. and Gentleman, R. (2008) Modeling synthetic lethality. *Genome Biol*, **9**(9): R135.
- Le Meur, N., Jiang, Z., Liu, T., Mar, J., and Gentleman, R.C. (2014) Slgi: Synthetic lethal genetic interaction. r package version 1.26.0.
- Lee, A.Y., Perreault, R., Harel, S., Boulier, E.L., Suderman, M., Hallett, M., and Jenna, S. (2010a) Searching for signaling balance through the identification of genetic interactors of the rab guanine-nucleotide dissociation inhibitor gdi-1. *PLoS ONE*, **5**(5): e10624.
- Lee, I., Lehner, B., Vavouri, T., Shin, J., Fraser, A.G., and Marcotte, E.M. (2010b) Predicting genetic modifier loci using functional gene networks. *Genome Research*, **20**(8): 1143–1153.
- Lee, I. and Marcotte, E.M. (2009) Effects of functional bias on supervised learning of a gene network model. *Methods Mol Biol*, **541**: 463–75.
- Lee, M.J., Ye, A.S., Gardino, A.K., Heijink, A.M., Sorger, P.K., MacBeath, G., and Yaffe, M.B. (2012) Sequential application of anticancer drugs enhances cell death by rewiring apoptotic signaling networks. *Cell*, **149**(4): 780–94.
- Lehner, B., Crombie, C., Tischler, J., Fortunato, A., and Fraser, A.G. (2006) Systematic mapping of genetic interactions in *caenorhabditis elegans* identifies common modifiers of diverse signaling pathways. *Nat Genet*, **38**(8): 896–903.
- Li, B., Ruotti, V., Stewart, R.M., Thomson, J.A., and Dewey, C.N. (2010) RNA-Seq gene expression estimation with read mapping uncertainty. *Bioinformatics*, **26**(4): 493–500.

- Li, X.J., Mishra, S.K., Wu, M., Zhang, F., and Zheng, J. (2014) Syn-lethality: An integrative knowledge base of synthetic lethality towards discovery of selective anticancer therapies. *Biomed Res Int*, **2014**: 196034.
- Linehan, W.M., Spellman, P.T., Ricketts, C.J., Creighton, C.J., Fei, S.S., Davis, C., Wheeler, D.A., Murray, B.A., Schmidt, L., Vocke, C.D., *et al.* (2016) Comprehensive Molecular Characterization of Papillary Renal-Cell Carcinoma. *N Engl J Med*, **374**(2): 135–145.
- Lokody, I. (2014) Computational modelling: A computational crystal ball. *Nature Reviews Cancer*, **14**(10): 649–649.
- Lord, C.J., Tutt, A.N., and Ashworth, A. (2015) Synthetic lethality and cancer therapy: lessons learned from the development of PARP inhibitors. *Annu Rev Med*, **66**: 455–470.
- Lu, X., Kensche, P.R., Huynen, M.A., and Notebaart, R.A. (2013) Genome evolution predicts genetic interactions in protein complexes and reveals cancer drug targets. *Nat Commun*, **4**: 2124.
- Lu, X., Megchelenbrink, W., Notebaart, R.A., and Huynen, M.A. (2015) Predicting human genetic interactions from cancer genome evolution. *PLoS One*, **10**(5): e0125795.
- Lum, P.Y., Armour, C.D., Stepaniants, S.B., Cavet, G., Wolf, M.K., Butler, J.S., Hinchshaw, J.C., Garnier, P., Prestwich, G.D., Leonardson, A., *et al.* (2004) Discovering modes of action for therapeutic compounds using a genome-wide screen of yeast heterozygotes. *Cell*, **116**(1): 121–137.
- Luo, J., Solimini, N.L., and Elledge, S.J. (2009) Principles of Cancer Therapy: Oncogene and Non-oncogene Addiction. *Cell*, **136**(5): 823–837.
- Machado, J., Olivera, C., Carvalh, R., Soares, P., Berx, G., Caldas, C., Sercuca, R., Carneiro, F., and Sorbrinho-Simoes, M. (2001) E-cadherin gene (*CDH1*) promoter methylation as the second hit in sporadic diffuse gastric carcinoma. *Oncogene*, **20**: 1525–1528.
- Markowetz, F. (2017) All biology is computational biology. *PLoS Biol*, **15**(3): e2002050.

- Masciari, S., Larsson, N., Senz, J., Boyd, N., Kaurah, P., Kandel, M.J., Harris, L.N., Pinheiro, H.C., Troussard, A., Miron, P., *et al.* (2007) Germline E-cadherin mutations in familial lobular breast cancer. *J Med Genet*, **44**(11): 726–31.
- Mattison, J., van der Weyden, L., Hubbard, T., and Adams, D.J. (2009) Cancer gene discovery in mouse and man. *Biochim Biophys Acta*, **1796**(2): 140–161.
- McLachlan, J., George, A., and Banerjee, S. (2016) The current status of parp inhibitors in ovarian cancer. *Tumori*, **102**(5): 433–440.
- McLendon, R., Friedman, A., Bigner, D., Van Meir, E.G., Brat, D.J., Mastrogianakis, G.M., Olson, J.J., Mikkelsen, T., Lehman, N., Aldape, K., *et al.* (2008) Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature*, **455**(7216): 1061–1068.
- Miles, D.W. (2001) Update on HER-2 as a target for cancer therapy: herceptin in the clinical setting. *Breast Cancer Res*, **3**(6): 380–384.
- Muzny, D.M., Bainbridge, M.N., Chang, K., Dinh, H.H., Drummond, J.A., Fowler, G., Kovar, C.L., Lewis, L.R., Morgan, M.B., Newsham, I.F., *et al.* (2012) Comprehensive molecular characterization of human colon and rectal cancer. *Nature*, **487**(7407): 330–337.
- Nagalla, S., Chou, J.W., Willingham, M.C., Ruiz, J., Vaughn, J.P., Dubey, P., Lash, T.L., Hamilton-Dutoit, S.J., Bergh, J., Sotiriou, C., *et al.* (2013) Interactions between immunity, proliferation and molecular subtype in breast cancer prognosis. *Genome Biol*, **14**(4): R34.
- Neeley, E.S., Kornblau, S.M., Coombes, K.R., and Baggerly, K.A. (2009) Variable slope normalization of reverse phase protein arrays. *Bioinformatics*, **25**(11): 1384.
- Novomestky, F. (2012) *matrixcalc: Collection of functions for matrix calculations*. R package version 1.0-3.
- Nowak, M.A., Boerlijst, M.C., Cooke, J., and Smith, J.M. (1997) Evolution of genetic redundancy. *Nature*, **388**(6638): 167–171.
- Oliveira, C., Senz, J., Kaurah, P., Pinheiro, H., Sanges, R., Haegert, A., Corso, G., Schouten, J., Fitzgerald, R., Vogelsang, H., *et al.* (2009) Germline *CDH1* deletions in hereditary diffuse gastric cancer families. *Human Molecular Genetics*, **18**(9): 1545–1555.

- Oliveira, C., Seruca, R., Hoogerbrugge, N., Ligtenberg, M., and Carneiro, F. (2013) Clinical utility gene card for: Hereditary diffuse gastric cancer (HDGC). *Eur J Hum Genet*, **21**(8).
- Pandey, G., Zhang, B., Chang, A.N., Myers, C.L., Zhu, J., Kumar, V., and Schadt, E.E. (2010) An integrative multi-network and multi-classifier approach to predict genetic interactions. *PLoS Comput Biol*, **6**(9).
- Parker, J., Mullins, M., Cheung, M., Leung, S., Voduc, D., Vickery, T., Davies, S., Fauron, C., He, X., Hu, Z., et al. (2009) Supervised risk predictor of breast cancer based on intrinsic subtypes. *Journal of Clinical Oncology*, **27**(8): 1160–1167.
- Pereira, B., Chin, S.F., Rueda, O.M., Vollan, H.K., Provenzano, E., Bardwell, H.A., Pugh, M., Jones, L., Russell, R., Sammut, S.J., et al. (2016) Erratum: The somatic mutation profiles of 2,433 breast cancers refine their genomic and transcriptomic landscapes. *Nat Commun*, **7**: 11908.
- Perou, C.M., Sørlie, T., Eisen, M.B., van de Rijn, M., Jeffrey, S.S., Rees, C.A., Pollack, J.R., Ross, D.T., Johnsen, H., Akslen, L.A., et al. (2000) Molecular portraits of human breast tumours. *Nature*, **406**(6797): 747–752.
- Polyak, K. and Weinberg, R.A. (2009) Transitions between epithelial and mesenchymal states: acquisition of malignant and stem cell traits. *Nat Rev Cancer*, **9**(4): 265–73.
- R Core Team (2016) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. R version 3.3.2.
- Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research*, **43**(7): e47.
- Roguev, A., Bandyopadhyay, S., Zofall, M., Zhang, K., Fischer, T., Collins, S.R., Qu, H., Shales, M., Park, H.O., Hayles, J., et al. (2008) Conservation and rewiring of functional modules revealed by an epistasis map in fission yeast. *Science*, **322**(5900): 405–10.
- Roychowdhury, S. and Chinnaiyan, A.M. (2016) Translating cancer genomes and transcriptomes for precision oncology. *CA Cancer J Clin*, **66**(1): 75–88.
- Rung, J. and Brazma, A. (2013) Reuse of public genome-wide gene expression data. *Nat Rev Genet*, **14**(2): 89–99.

- Ryan, C., Lord, C., and Ashworth, A. (2014) Daisy: Picking synthetic lethals from cancer genomes. *Cancer Cell*, **26**(3): 306–308.
- Schena, M. (1996) Genome analysis with gene expression microarrays. *Bioessays*, **18**(5): 427–431.
- Scheuer, L., Kauff, N., Robson, M., Kelly, B., Barakat, R., Satagopan, J., Ellis, N., Hensley, M., Boyd, J., Borgen, P., *et al.* (2002) Outcome of preventive surgery and screening for breast and ovarian cancer in BRCA mutation carriers. *J Clin Oncol*, **20**(5): 1260–1268.
- Semb, H. and Christofori, G. (1998) The tumor-suppressor function of E-cadherin. *Am J Hum Genet*, **63**(6): 1588–93.
- Sing, T., Sander, O., Beerenwinkel, N., and Lengauer, T. (2005) RocR: visualizing classifier performance in r. *Bioinformatics*, **21**(20): 7881.
- Slurm development team (Slurm) (2017) Slurm workload manager. <https://slurm.schedmd.com/>. Accessed: 25/03/2017.
- Sørlie, T., Perou, C.M., Tibshirani, R., Aas, T., Geisler, S., Johnsen, H., Hastie, T., Eisen, M.B., van de Rijn, M., Jeffrey, S.S., *et al.* (2001) Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci USA*, **98**(19): 10869–10874.
- Srihari, S., Singla, J., Wong, L., and Ragan, M.A. (2015) Inferring synthetic lethal interactions from mutual exclusivity of genetic events in cancer. *Biology Direct*, **10**(1): 57.
- Stajich, J.E. and Lapp, H. (2006) Open source tools and toolkits for bioinformatics: significance, and where are we? *Brief Bioinformatics*, **7**(3): 287–296.
- Stratton, M.R., Campbell, P.J., and Futreal, P.A. (2009) The cancer genome. *Nature*, **458**(7239): 719–724.
- Ström, C. and Helleday, T. (2012) Strategies for the use of poly(adenosine diphosphate ribose) polymerase (parp) inhibitors in cancer therapy. *Biomolecules*, **2**(4): 635–649.
- Tarazona, S., Garcia-Alcalde, F., Dopazo, J., Ferrer, A., and Conesa, A. (2011) Differential expression in RNA-seq: a matter of depth. *Genome Res*, **21**(12): 2213–2223.

- Telford, B.J., Chen, A., Beetham, H., Frick, J., Brew, T.P., Gould, C.M., Single, A., Godwin, T., Simpson, K.J., and Guilford, P. (2015) Synthetic lethal screens identify vulnerabilities in gpcr signalling and cytoskeletal organization in E-cadherin-deficient cells. *Mol Cancer Ther*, **14**(5): 1213–1223.
- The 1000 Genomes Project Consortium (1000 Genomes) (2010) A map of human genome variation from population-scale sequencing. *Nature*, **467**(7319): 1061–1073.
- The Cancer Genome Atlas Research Network (TCGA) (2017) The Cancer Genome Atlas Project. <https://cancergenome.nih.gov/>. Accessed: 26/03/2017.
- The Catalogue Of Somatic Mutations In Cancer (COSMIC) (2016) Cosmic: The catalogue of somatic mutations in cancer. <http://cancer.sanger.ac.uk/cosmic>. Release 79 (23/08/2016), Accessed: 05/02/2017.
- The Comprehensive R Archive Network (CRAN) (2017) Cran. <https://cran.r-project.org/>. Accessed: 24/03/2017.
- The ENCODE Project Consortium (ENCODE) (2004) The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science*, **306**(5696): 636–640.
- The National Cancer Institute (NCI) (2015) The genetics of cancer. <https://www.cancer.gov/about-cancer/causes-prevention/genetics>. Published: 22/04/2015, Accessed: 22/03/2017.
- The New Zealand eScience Infrastructure (NeSI) (2017) NeSI. <https://www.nesi.org.nz/>. Accessed: 25/03/2017.
- Tierney, L., Rossini, A.J., Li, N., and Sevcikova, H. (2015) *snow: Simple Network of Workstations*. R package version 0.4-2.
- Tiong, K.L., Chang, K.C., Yeh, K.T., Liu, T.Y., Wu, J.H., Hsieh, P.H., Lin, S.H., Lai, W.Y., Hsu, Y.C., Chen, J.Y., et al. (2014) Csnk1e/ctnnb1 are synthetic lethal to tp53 in colorectal cancer and are markers for prognosis. *Neoplasia*, **16**(5): 441–50.
- Tischler, J., Lehner, B., and Fraser, A.G. (2008) Evolutionary plasticity of genetic interaction networks. *Nat Genet*, **40**(4): 390–391.
- Tomasetti, C. and Vogelstein, B. (2015) Cancer etiology. Variation in cancer risk among tissues can be explained by the number of stem cell divisions. *Science*, **347**(6217): 78–81.

- Tong, A.H., Evangelista, M., Parsons, A.B., Xu, H., Bader, G.D., Page, N., Robinson, M., Raghibizadeh, S., Hogue, C.W., Bussey, H., *et al.* (2001) Systematic genetic analysis with ordered arrays of yeast deletion mutants. *Science*, **294**(5550): 2364–8.
- Tong, A.H., Lesage, G., Bader, G.D., Ding, H., Xu, H., Xin, X., Young, J., Berriz, G.F., Brost, R.L., Chang, M., *et al.* (2004) Global mapping of the yeast genetic interaction network. *Science*, **303**(5659): 808–13.
- Tran, B., Dancey, J.E., Kamel-Reid, S., McPherson, J.D., Bedard, P.L., Brown, A.M., Zhang, T., Shaw, P., Onetto, N., Stein, L., *et al.* (2012) Cancer genomics: technology, discovery, and translation. *J Clin Oncol*, **30**(6): 647–660.
- Travers, J. and Milgram, S. (1969) An experimental study of the small world problem. *Sociometry*, **32**(4): 425–443.
- Tunggal, J.A., Helfrich, I., Schmitz, A., Schwarz, H., Gunzel, D., Fromm, M., Kemler, R., Krieg, T., and Niessen, C.M. (2005) E-cadherin is essential for in vivo epidermal barrier function by regulating tight junctions. *EMBO J*, **24**(6): 1146–1156.
- Tutt, A., Robson, M., Garber, J.E., Domchek, S.M., Audeh, M.W., Weitzel, J.N., Friedlander, M., Arun, B., Loman, N., Schmutzler, R.K., *et al.* (2010) Oral poly(adenylic ribose) polymerase inhibitor olaparib in patients with *BRCA1* or *BRCA2* mutations and advanced breast cancer: a proof-of-concept trial. *Lancet*, **376**(9737): 235–44.
- University of California, Santa Cruz (UCSC) (2012) Ucsc cancer browser. Accessed 29/03/2012.
- van der Post, R.S., Vogelaar, I.P., Carneiro, F., Guilford, P., Huntsman, D., Hoogerbrugge, N., Caldas, C., Schreiber, K.E., Hardwick, R.H., Ausems, M.G., *et al.* (2015) Hereditary diffuse gastric cancer: updated clinical guidelines with an emphasis on germline CDH1 mutation carriers. *J Med Genet*, **52**(6): 361–374.
- van Steen, K. (2012) Travelling the world of gene-gene interactions. *Briefings in Bioinformatics*, **13**(1): 1–19.
- van Steen, M. (2010) *Graph Theory and Complex Networks: An Introduction*. Maarten van Steen, VU Amsterdam.
- Vapnik, V.N. (1995) *The nature of statistical learning theory*. Springer-Verlag New York, Inc.

- Vizeacoumar, F.J., Arnold, R., Vizeacoumar, F.S., Chandrashekhar, M., Buzina, A., Young, J.T., Kwan, J.H., Sayad, A., Mero, P., Lawo, S., *et al.* (2013) A negative genetic interaction map in isogenic cancer cell lines reveals cancer cell vulnerabilities. *Mol Syst Biol*, **9**: 696.
- Vogelstein, B., Papadopoulos, N., Velculescu, V.E., Zhou, S., Diaz, L.A., and Kinzler, K.W. (2013) Cancer genome landscapes. *Science*, **339**(6127): 1546–1558.
- Vos, C.B., Cleton-Jansen, A.M., Berx, G., de Leeuw, W.J., ter Haar, N.T., van Roy, F., Cornelisse, C.J., Peterse, J.L., and van de Vijver, M.J. (1997) E-cadherin inactivation in lobular carcinoma in situ of the breast: an early event in tumorigenesis. *Br J Cancer*, **76**(9): 1131–3.
- Waldron, D. (2016) Cancer genomics: A multi-layer omics approach to cancer. *Nat Rev Genet*, **17**(8): 436–437.
- Wang, K., Singh, D., Zeng, Z., Coleman, S.J., Huang, Y., Savich, G.L., He, X., Mieczkowski, P., Grimm, S.A., Perou, C.M., *et al.* (2010) MapSplice: accurate mapping of RNA-seq reads for splice junction discovery. *Nucleic Acids Res*, **38**(18): e178.
- Wang, X. and Simon, R. (2013) Identification of potential synthetic lethal genes to p53 using a computational biology approach. *BMC Medical Genomics*, **6**(1): 30.
- Wappett, M. (2014) Bisep: Toolkit to identify candidate synthetic lethality. r package version 2.0.
- Wappett, M., Dulak, A., Yang, Z.R., Al-Watban, A., Bradford, J.R., and Dry, J.R. (2016) Multi-omic measurement of mutually exclusive loss-of-function enriches for candidate synthetic lethal gene pairs. *BMC Genomics*, **17**: 65.
- Warnes, G.R., Bolker, B., Bonebakker, L., Gentleman, R., Liaw, W.H.A., Lumley, T., Maechler, M., Magnusson, A., Moeller, S., Schwartz, M., *et al.* (2015) *gplots: Various R Programming Tools for Plotting Data*. R package version 2.17.0.
- Watts, D.J. and Strogatz, S.H. (1998) Collective dynamics of 'small-world' networks. *Nature*, **393**(6684): 440–2.
- Weinstein, I.B. (2000) Disorders in cell circuitry during multistage carcinogenesis: the role of homeostasis. *Carcinogenesis*, **21**(5): 857–864.

- Weinstein, J.N., Akbani, R., Broom, B.M., Wang, W., Verhaak, R.G., McConkey, D., Lerner, S., Morgan, M., Creighton, C.J., Smith, C., *et al.* (2014) Comprehensive molecular characterization of urothelial bladder carcinoma. *Nature*, **507**(7492): 315–322.
- Weinstein, J.N., Collisson, E.A., Mills, G.B., Shaw, K.R., Ozenberger, B.A., Ellrott, K., Shmulevich, I., Sander, C., Stuart, J.M., Chang, K., *et al.* (2013) The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet*, **45**(10): 1113–1120.
- Wickham, H. and Chang, W. (2016) *devtools: Tools to Make Developing R Packages Easier*. R package version 1.12.0.
- Wickham, H., Danenberg, P., and Eugster, M. (2017) *roxygen2: In-Line Documentation for R*. R package version 6.0.1.
- Wong, S.L., Zhang, L.V., Tong, A.H.Y., Li, Z., Goldberg, D.S., King, O.D., Lesage, G., Vidal, M., Andrews, B., Bussey, H., *et al.* (2004) Combining biological networks to predict genetic interactions. *Proceedings of the National Academy of Sciences of the United States of America*, **101**(44): 15682–15687.
- World Health Organization (WHO) (2017) Fact sheet: Cancer. <http://www.who.int/mediacentre/factsheets/fs297/en/>. Updated February 2017, Accessed: 22/03/2017.
- Wu, M., Li, X., Zhang, F., Li, X., Kwoh, C.K., and Zheng, J. (2014) In silico prediction of synthetic lethality by meta-analysis of genetic interactions, functions, and pathways in yeast and human cancer. *Cancer Inform*, **13**(Suppl 3): 71–80.
- Yu, H. (2002) Rmpi: Parallel statistical computing in r. *R News*, **2**(2): 10–14.
- Zhang, F., Wu, M., Li, X.J., Li, X.L., Kwoh, C.K., and Zheng, J. (2015) Predicting essential genes and synthetic lethality via influence propagation in signaling pathways of cancer cell fates. *J Bioinform Comput Biol*, **13**(3): 1541002.
- Zhang, J., Baran, J., Cros, A., Guberman, J.M., Haider, S., Hsu, J., Liang, Y., Rivkin, E., Wang, J., Whitty, B., *et al.* (2011) International cancer genome consortium data portal a one-stop shop for cancer genomics data. *Database: The Journal of Biological Databases and Curation*, **2011**: bar026.
- Zhong, W. and Sternberg, P.W. (2006) Genome-wide prediction of *c. elegans* genetic interactions. *Science*, **311**(5766): 1481–1484.

Zweig, M.H. and Campbell, G. (1993) Receiver-operating characteristic (roc) plots: a fundamental evaluation tool in clinical medicine. *Clinical Chemistry*, **39**(4): 561–577.

Appendix A

Sample Quality

A.1 Sample Correlation




Figure A.1: Correlation profiles of removed samples. Heatmap (Euclidean distance) of samples in The Cancer Genome Atlas (TCGA) breast cancer dataset (left) clustered for all samples against removed samples (top): tissue source site (TSS), sample type with reds for tumour and greens for normal, patient (A2QH in pink), with varied analyte and plate. Excluded samples clustered at the bottom and annotation (left) show shared properties between samples in the dataset.




Figure A.2: Correlation analysis and sample removal. Correlation matrix heatmap (Euclidean distance) of all samples in TCGA breast cancer dataset against each other annotated for sample clinical data: sample type, tissue type, tumour stage, Estrogen receptor (ER) and intrinsic subtype (from the PAM50 method). *CDH1* somatic mutation, gene expression, and status for SLIPT prediction were also annotated. Discrete variables were coloured as displayed in the legend and continuous variables on a blue–red scale as shown in the colour key. Trimmed samples cluster at the bottom of the heatmap and the colour bars of the left show which were removed for quality concerns.

A.2 Replicate Samples in TCGA Breast Cancer Data

Replicate samples were picked where possible from the TCGA breast cancer gene expression data to examine for sample quality. Independent samples of the same tumour were expected to have very high Pearson correlation between their expression profiles unless there were issues with sample collection or preparation and were thus an indicator of sample quality. The log-transformed raw read counts for replicate samples were examined in Figures A.3–A.5. These were examined before normalisation which would be expected to increase sample concordance.

Another consideration was the samples which were removed for quality concerns (in Section 2.2.2). While these were selected by unbiased hierarchical clustering (See Figure A.2), many of the excluded (tumour) samples were performed in replicate despite relatively few replicate samples in the overall dataset. These samples correlate poorly with the rest of the dataset, in addition to correlation with replicate samples.




Figure A.3: **Replicate excluded samples.** Both tumour samples of patient A2QH were excluded as they were poorly correlated with other samples, although they were highly similar to each other as shown by Pearson correlation of log-raw counts.




Figure A.4: **Replicate samples with all remaining.** Patient A26J was sampled 3 times and compared pairwise. Pairs of samples were also compared for other patients with replicate samples. In all cases, replicate samples remaining in the dataset were highly concordant, as shown by Pearson correlation of log-raw counts.




Figure A.5: **Replicate samples with some excluded.** (continued on next page)




Figure A.5: **Replicate samples with some excluded.** Patients A0DB, A13D, A13E, and A26E were each sampled 3 times and compared pairwise. Pairs of samples were also compared for other patients with replicate samples. In all cases, the replicate samples remaining in the dataset more were highly concordant than those excluded from the analysis, as shown by Pearson correlation of log-log counts.

Appendix B

Software Used for Thesis

Table B.1: Complete list of R packages used during this thesis

Package	Repository	Laptop	Lab	Server	NeSI
base	base	3.3.2	3.3.2	3.3.1	3.3.0
abind	CRAN		1.4-5		1.4-3
acepack	CRAN		1.4.1		1.3-3.3
ade4	CRAN		1.7-5		
annaffy	Bioconductor		1.46.0		
AnnotationDbi	Bioconductor		1.36.0	1.36.0	1.34.4
apComplex	CRAN		2.40.0		
ape	CRAN		4		3.4
arm	CRAN		1.9-3		
assertthat	CRAN	0.1	0.1	0.1	0.1
backports	CRAN	1.0.5	1.0.4	1.0.5	1.0.2
base64	CRAN			2	2
base64enc	CRAN		0.1-3		0.1-3
beanplot	CRAN		1.2	1.2	1.2
BH	CRAN	1.60.0-2	1.62.0-1	1.62.0-1	1.60.0-2
Biobase	Bioconductor		2.34.0	2.34.0	2.32.0
BiocGenerics	Bioconductor		0.20.0	0.20.0	0.18.0
BiocInstaller	Bioconductor		1.24.0	1.20.3	1.22.3
BiocParallel	Bioconductor		1.8.1	1.8.1	
Biostrings	Bioconductor		2.42.1	2.42.0	
BiSEp	Bioconductor		2.0.1	2.0.1	2.0.1

bitops	CRAN	1.0-6	1.0-6	1.0-6	1.0-6
boot	base	1.3-18	1.3-18	1.3-18	1.3-18
brew	CRAN	1.0-6	1.0-6	1.0-6	1.0-6
broom	CRAN	0.4.1			
caTools	CRAN	1.17.1	1.17.1	1.17.1	1.17.1
cgdssr	CRAN		1.2.5		
checkmate	CRAN		1.8.2		1.7.4
chron	CRAN	2.3-47	2.3-48	2.3-50	2.3-47
class	base	7.3-14	7.3-14	7.3-14	7.3-14
cluster	base	2.0.5	2.0.5	2.0.5	2.0.4
coda	CRAN		0.19-1		0.18-1
codetools	base	0.2-15	0.2-15	0.2-15	0.2-14
colorRamps	CRAN		2.3		
colorspace	CRAN	1.2-6	1.3-2	1.3-2	1.2-6
commonmark	CRAN	1.1		1.2	
compiler	base	3.3.2	3.3.2	3.3.1	3.3.0
corpcor	CRAN		1.6.8	1.6.8	1.6.8
Cprob	CRAN		1.2.4		
crayon	CRAN	1.3.2	1.3.2	1.3.2	1.3.2
crop	CRAN		0.0-2	0.0-2	
curl	CRAN	1.2	2.3	2.3	0.9.7
d3Network	CRAN		0.5.2.1		
data.table	CRAN	1.9.6	1.10.0	1.10.1	1.9.6
data.tree	CRAN		0.7.0	0.7.0	
datasets	base	3.3.2	3.3.2	3.3.1	3.3.0
DBI	CRAN	0.5-1	0.5-1	0.5-1	0.5-1
dendextend	CRAN	1.4.0	1.4.0	1.4.0	
DEoptimR	CRAN	1.0-8	1.0-8	1.0-8	1.0-4
desc	CRAN	1.1.0		1.1.0	
devtools	CRAN	1.12.0	1.12.0	1.12.0	1.12.0
DiagrammeR	CRAN		0.9.0	0.9.0	
dichromat	CRAN	2.0-0	2.0-0	2.0-0	2.0-0
digest	CRAN	0.6.10	0.6.11	0.6.12	0.6.9
diptest	CRAN	0.75-7	0.75-7	0.75-7	
doParallel	CRAN	1.0.10	1.0.10	1.0.10	1.0.10

dplyr	CRAN	0.5.0	0.5.0	0.5.0	0.5.0
ellipse	CRAN		0.3-8	0.3-8	0.3-8
evaluate	CRAN		0.1	0.1	0.9
fdrtool	CRAN		1.2.15		
fields	CRAN		8.1		
flexmix	CRAN	2.3-13	2.3-13	2.3-13	
forcats	CRAN	0.2.0			
foreach	CRAN	1.4.3	1.4.3	1.4.3	1.4.3
foreign	base	0.8-67	0.8-67	0.8-67	0.8-66
formatR	CRAN		1.4	1.4	1.4
Formula	CRAN		1.2-1		1.2-1
fpc	CRAN	2.1-10	2.1-10	2.1-10	
futile.logger	CRAN		1.4.3	1.4.3	1.4.1
futile.options	CRAN		1.0.0	1.0.0	1.0.0
gdata	CRAN	2.17.0	2.17.0	2.17.0	2.17.0
geepack	CRAN		1.2-1		
GenomeInfoDb	Bioconductor		1.10.2	1.10.1	
GenomicAlignments	Bioconductor		1.10.0	1.10.0	
GenomicRanges	Bioconductor		1.26.2	1.26.1	
ggm	CRAN		2.3		
ggplot2	CRAN	2.1.0	2.2.1	2.2.1	2.1.0
git2r	CRAN	0.15.0	0.18.0	0.16.0	0.15.0
glasso	CRAN		1.8		
GO.db	Bioconductor		3.4.0	3.2.2	3.3.0
GOSemSim	Bioconductor		2.0.3	1.28.2	1.30.3
gplots	CRAN	3.0.1	3.0.1	3.0.1	3.0.1
graph	Bioconductor		1.52.0		
graphics	base	3.3.2	3.3.2	3.3.1	3.3.0
graphsim	GitHub TomKellyGenetics	0.1.0	0.1.0	0.1.0	0.1.0
grDevices	base	3.3.2	3.3.2	3.3.1	3.3.0
grid	base	3.3.2	3.3.2	3.3.1	3.3.0
gridBase	CRAN	0.4-7	0.4-7	0.4-7	0.4-7
gridExtra	CRAN	2.2.1	2.2.1	2.2.1	2.2.1
gridGraphics	CRAN		0.1-5		

gtable	CRAN	0.2.0	0.2.0	0.2.0	0.2.0
gtools	CRAN	3.5.0	3.5.0	3.5.0	3.5.0
haven	CRAN	1.0.0			
heatmap.2x	GitHub		0.0.0.9000	0.0.0.9000	0.0.0.9000
	TomKellyGenetics				0.0.0.9000
hgu133plus2.db	Bioconductor		3.2.3		
highr	CRAN		0.6	0.6	0.6
Hmisc	CRAN		4.0-2	4.0-2	3.17-4
hms	CRAN	0.2	0.3		
htmlTable	CRAN		1.8	1.9	
htmltools	CRAN	0.3.5	0.3.5	0.3.5	0.3.5
htmlwidgets	CRAN		0.8	0.8	
httpuv	CRAN	1.3.3		1.3.3	
httr	CRAN	1.2.1	1.2.1	1.2.1	1.1.0
huge	CRAN		1.2.7		
hunspell	CRAN		2.3		2
hypergraph	CRAN		1.46.0		
igraph	CRAN	1.0.1	1.0.1	1.0.1	1.0.1
igraph.extensions	GitHub				
	TomKellyGenetics	0.1.0.9001	0.1.0.9001	0.1.0.9001	0.1.0.9001
influenceR	CRAN		0.1.0	0.1.0	
info.centrality	GitHub				
	TomKellyGenetics	0.1.0	0.1.0	0.1.0	0.1.0
IRanges	Bioconductor		2.8.1	2.8.1	2.6.1
irlba	CRAN	2.1.1	2.1.2	2.1.2	2.0.0
iterators	CRAN	1.0.8	1.0.8	1.0.8	1.0.8
jpeg	CRAN		0.1-8		
jsonlite	CRAN	1.1	1.2	1.3	0.9.20
KEGG.db	Bioconductor		3.2.3		
kernlab	CRAN	0.9-25	0.9-25	0.9-25	
KernSmooth	base	2.23-15	2.23-15	2.23-15	2.23-15
knitr	CRAN		1.15.1	1.15.1	1.14
labeling	CRAN	0.3	0.3	0.3	0.3
lambda.r	CRAN		1.1.9	1.1.9	1.1.7
lattice	base	0.20-34	0.20-34	0.20-34	0.20-33

latticeExtra	CRAN		0.6-28		0.6-28
lava	CRAN		1.4.6		
lavaan	CRAN		0.5-22		
lazyeval	CRAN	0.2.0	0.2.0	0.2.0	0.2.0
les	CRAN		1.24.0		
lgtdl	CRAN		1.1.3		
limma	Bioconductor		3.30.7	3.30.3	
lme4	CRAN		1.1-12		1.1-12
lubridate	CRAN	1.6.0			
magrittr	CRAN	1.5	1.5	1.5	1.5
maps	CRAN		3.1.1		
markdown	CRAN		0.7.7	0.7.7	0.7.7
MASS	base	7.3-45	7.3-45	7.3-45	7.3-45
Matrix	base	1.2-7.1	1.2-7.1	1.2-8	1.2-6
matrixcalc	CRAN	1.0-3	1.0-3	1.0-3	1.0-3
mclust	CRAN	5.2	5.2.1	5.2.2	5.2
memoise	CRAN	1.0.0	1.0.0	1.0.0	1.0.0
methods	base	3.3.2	3.3.2	3.3.1	3.3.0
mgcv	base	1.8-16	1.8-16	1.8-17	1.8-12
mi	CRAN		1		
mime	CRAN	0.5	0.5	0.5	0.4
minqa	CRAN		1.2.4		1.2.4
mnormt	CRAN	1.5-5	1.5-5		1.5-4
modelr	CRAN	0.1.0			
modeltools	CRAN	0.2-21	0.2-21	0.2-21	
multtest	Bioconductor		2.30.0	2.30.0	
munsell	CRAN	0.4.3	0.4.3	0.4.3	0.4.3
mvtnorm	CRAN	1.0-5	1.0-5	1.0-6	1.0-5
network	CRAN		1.13.0		
nlme	base	3.1-128	3.1-128	3.1-131	3.1-128
nloptr	CRAN		1.0.4		1.0.4
NMF	CRAN	0.20.6	0.20.6	0.20.6	0.20.6
nnet	base	7.3-12	7.3-12	7.3-12	7.3-12
numDeriv	CRAN		2016.8-1		2014.2-1
openssl	CRAN	0.9.4	0.9.6	0.9.6	0.9.4

org.Hs.eg.db	Bioconductor		3.1.2		3.3.0
org.Sc.sgd.db	Bioconductor		3.4.0		
parallel	base	3.3.2	3.3.2	3.3.1	3.3.0
pathway.structure	GitHub	0.1.0	0.1.0	0.1.0	0.1.0
.permutation	TomKellyGenetics				
pbivnorm	CRAN		0.6.0		
PGSEA	Bioconductor		1.48.0		
pkgmaker	CRAN	0.22	0.22	0.22	0.22
PKI	CRAN		0.1-3		
plogr	CRAN		0.1-1	0.1-1	
plot.igraph	GitHub	0.0.0.9001	0.0.0.9001	0.0.0.9001	0.0.0.9001
	TomKellyGenetics				
plotrix	CRAN		3.6-4		
plyr	CRAN	1.8.4	1.8.4	1.8.4	1.8.3
png	CRAN		0.1-7		0.1-7
prabclus	CRAN	2.2-6	2.2-6	2.2-6	
praise	CRAN	1.0.0	1.0.0		1.0.0
pROC	CRAN		1.8	1.9.1	
prodlim	CRAN		1.5.7		
prof.tree	CRAN		0.1.0		
protools	CRAN		0.99-2		
progress	CRAN			1.1.2	
psych	CRAN	1.6.12	1.6.12		
purrr	CRAN	0.2.2	0.2.2	0.2.2	0.2.2
qgraph	CRAN		1.4.1		
quadprog	CRAN		1.5-5	1.5-5	1.5-5
R.methodsS3	CRAN		1.7.1		1.7.1
R.oo	CRAN		1.21.0		1.20.0
R.utils	CRAN		2.5.0		
R6	CRAN	2.1.3	2.2.0	2.2.0	2.1.3
RBGL	CRAN		1.50.0		
RColorBrewer	CRAN	1.1-2	1.1-2	1.1-2	1.1-2
Rcpp	CRAN	0.12.7	0.12.9	0.12.9	0.12.7
RcppArmadillo	CRAN			0.7.700.0.0	0.6.700.6.0
RcppEigen	CRAN		0.3.2.9.0		0.3.2.8.1

RCurl	CRAN		1.95-4.8	1.95-4.8	1.95-4.8
reactome.db	Bioconductor		1.52.1	1.52.1	
reactometree	GitHub		0.1		
	TomKellyGenetics				
readr	CRAN	1.0.0	1.0.0		
readxl	CRAN	0.1.1			
registry	CRAN	0.3	0.3	0.3	0.3
reshape2	CRAN	1.4.1	1.4.2	1.4.2	1.4.1
rgeff	CRAN		0.15.3	0.15.3	
rgl	CRAN			0.97.0	0.95.1441
Rgraphviz	CRAN		2.18.0		
rjson	CRAN		0.2.15		
RJSONIO	CRAN		1.3-0		
rmarkdown	CRAN		1.3	1.3	1
Rmpi	CRAN		0.6-6		0.6-5
rngtools	CRAN	1.2.4	1.2.4	1.2.4	1.2.4
robustbase	CRAN	0.92-7	0.92-7	0.92-7	0.92-5
ROCR	CRAN	1.0-7	1.0-7	1.0-7	1.0-7
Rook	CRAN		1.1-1	1.1-1	
roxygen2	CRAN	6.0.1	5.0.1	6.0.1	5.0.1
rpart	base	4.1-10	4.1-10	4.1-10	4.1-10
rprojroot	CRAN	1.2	1.1	1.2	
Rsamtools	Bioconductor		1.26.1	1.26.1	
rsconnect	CRAN		0.7		
RSQLite	CRAN		1.1-2	1.1-2	1.0.0
rstudioapi	CRAN	0.6	0.6	0.6	0.6
rvest	CRAN	0.3.2			
S4Vectors	Bioconductor		0.12.1	0.12.0	0.10.3
safe	Bioconductor		3.14.0	3.10.0	
scales	CRAN	0.4.0	0.4.1	0.4.1	0.4.0
selectr	CRAN	0.3-1			
sem	CRAN		3.1-8		
shiny	CRAN	0.14		1.0.0	
slipt	GitHub	0.1.0	0.1.0	0.1.0	0.1.0
	TomKellyGenetics				

sm	CRAN	2.2-5.4	2.2-5.4		
sna	CRAN		2.4		
snow	CRAN	0.4-1	0.4-2	0.4-2	0.3-13
sourcetools	CRAN	0.1.5		0.1.5	
SparseM	CRAN		1.74		1.7
spatial	base	7.3-11	7.3-11	7.3-11	7.3-11
splines	base	3.3.2	3.3.2	3.3.1	3.3.0
statnet.common	CRAN		3.3.0		
stats	base	3.3.2	3.3.2	3.3.1	3.3.0
stats4	base	3.3.2	3.3.2	3.3.1	3.3.0
stringi	CRAN	1.1.1	1.1.2	1.1.2	1.0-1
stringr	CRAN	1.1.0	1.1.0	1.2.0	1.0.0
Summarized Experiment	Bioconductor		1.4.0	1.4.0	
survival	base	2.39-4	2.40-1	2.40-1	2.39-4
tcltk	base	3.3.2	3.3.2	3.3.1	3.3.0
testthat	CRAN	1.0.2	1.0.2		1.0.2
tibble	CRAN	1.2	1.2	1.2	1.2
tidyverse	GitHub hadley	1.1.1			
timeline	CRAN		0.9		
tools	base	3.3.2	3.3.2	3.3.1	3.3.0
tpr	CRAN		0.3-1		
trimcluster	CRAN	0.1-2	0.1-2	0.1-2	
Unicode	CRAN	9.0.0-1	9.0.0-1	9.0.0-1	
utils	base	3.3.2	3.3.2	3.3.1	3.3.0
vioplot	CRAN		0.2		
vioplotx	GitHub TomKellyGenetics	0.0.0.9000	0.0.0.9000		
viridis	CRAN	0.3.4	0.3.4	0.3.4	
visNetwork	CRAN		1.0.3	1.0.3	
whisker	CRAN	0.3-2	0.3-2	0.3-2	0.3-2
withr	CRAN	1.0.2	1.0.2	1.0.2	1.0.2
XML	base	3.98-1.3	3.98-1.1	3.98-1.5	3.98-1.4

xml2	CRAN	1.1.1	1.1.1	1.0.0
xtable	CRAN	1.8-2	1.8-2	1.8-2
XVector	Bioconductor		0.14.0	0.14.0
yaml	CRAN		2.1.14	2.1.14
zlibbioc	CRAN		1.20.0	1.20.0
zoo	CRAN	1.7-13	1.7-14	1.7-13

Appendix C

Mutation Analysis in Breast Cancer

C.1 Synthetic Lethal Genes and Pathways

SLIPT expression analysis (described in Section 3.1) on TCGA breast cancer data ($n = 969$) found the following genes and pathways, described in sections 4.1 and 4.1.1.

Table C.1: Candidate synthetic lethal gene partners of *CDH1* from mtSLIPT

Gene	Observed*	Expected*	χ^2 value	p-value	p-value (False discovery rate (FDR))
<i>TFAP2B</i>	8	36.7	89.5	3.60×10^{-20}	8.37×10^{-17}
<i>ZNF423</i>	15	36.7	78.8	7.89×10^{-18}	1.22×10^{-14}
<i>CALCOCO1</i>	11	36.7	76.8	2.09×10^{-17}	2.59×10^{-14}
<i>RBM5</i>	13	36.7	75.7	3.65×10^{-17}	4.00×10^{-14}
<i>BTG2</i>	7	36.7	71.7	2.72×10^{-16}	1.81×10^{-13}
<i>RXRA</i>	6	36.7	70.5	5.00×10^{-16}	2.97×10^{-13}
<i>SLC27A1</i>	11	36.7	70.3	5.42×10^{-16}	2.97×10^{-13}
<i>MEF2D</i>	12	36.7	69.6	7.86×10^{-16}	3.95×10^{-13}
<i>NISCH</i>	12	36.7	69.6	7.86×10^{-16}	3.95×10^{-13}
<i>AVPR2</i>	9	36.7	69.2	9.36×10^{-16}	4.58×10^{-13}
<i>CRY2</i>	13	36.7	68.9	1.07×10^{-15}	4.98×10^{-13}
<i>RAPGEF3</i>	13	36.7	68.9	1.07×10^{-15}	4.98×10^{-13}
<i>NRIP2</i>	10	36.7	68.2	1.58×10^{-15}	7.18×10^{-13}
<i>DARC</i>	12	36.7	66.4	3.76×10^{-15}	1.54×10^{-12}
<i>SFRS5</i>	12	36.7	66.4	3.76×10^{-15}	1.54×10^{-12}
<i>NOSTRIN</i>	5	36.7	65.1	7.40×10^{-15}	2.70×10^{-12}
<i>KIF13B</i>	12	36.7	63.4	1.69×10^{-14}	5.16×10^{-12}
<i>TENC1</i>	10	36.7	62.5	2.67×10^{-14}	7.40×10^{-12}
<i>MFAP4</i>	12	36.7	60.5	7.17×10^{-14}	1.67×10^{-11}
<i>ELN</i>	13	36.7	59.7	1.07×10^{-13}	2.32×10^{-11}
<i>SGK223</i>	14	36.7	59	1.51×10^{-13}	3.05×10^{-11}
<i>KIF12</i>	11	36.7	58.8	1.74×10^{-13}	3.34×10^{-11}
<i>SELP</i>	11	36.7	58.8	1.74×10^{-13}	3.34×10^{-11}
<i>CIRBP</i>	9	36.7	58.7	1.83×10^{-13}	3.41×10^{-11}
<i>CTDSP1</i>	9	36.7	58.7	1.83×10^{-13}	3.41×10^{-11}

Strongest candidate synthetic lethal partners for *CDH1* by mtSLIPT in TCGA in breast cancer expression and mutation data

* Observed and expected numbers of *CDH1* mutant TCGA breast tumours with low expression of partner genes

Table C.2: Pathways for *CDH1* partners from mtSLIPT

Pathways Over-represented	Pathway Size	SL Genes	p-value (FDR)
Eukaryotic Translation Elongation	86	60	2.0×10^{-128}
Peptide chain elongation	83	59	2.0×10^{-128}
Eukaryotic Translation Termination	83	58	2.3×10^{-125}
Viral mRNA Translation	81	57	2.5×10^{-124}
Nonsense Mediated Decay independent of the Exon Junction Complex	88	59	8.6×10^{-124}
Nonsense-Mediated Decay	103	61	5.2×10^{-117}
Nonsense Mediated Decay enhanced by the Exon Junction Complex	103	61	5.2×10^{-117}
Formation of a pool of free 40S subunits	93	58	1.6×10^{-116}
L13a-mediated translational silencing of Ceruloplasmin expression	103	59	1.3×10^{-111}
3' -UTR-mediated translational regulation	103	59	1.3×10^{-111}
GTP hydrolysis and joining of the 60S ribosomal subunit	104	59	6.2×10^{-111}
SRP-dependent cotranslational protein targeting to membrane	104	58	2.9×10^{-108}
Eukaryotic Translation Initiation	111	59	3.0×10^{-106}
Cap-dependent Translation Initiation	111	59	3.0×10^{-106}
Influenza Viral RNA Transcription and Replication	108	57	5.1×10^{-103}
Influenza Infection	117	59	1.5×10^{-102}
Translation	141	64	3.7×10^{-101}
Influenza Life Cycle	112	57	1.4×10^{-100}
GPCR downstream signalling	472	116	1.0×10^{-80}
Hemostasis	422	105	1.4×10^{-78}

Gene set over-representation analysis (hypergeometric test) for Reactome pathways in mtSLIPT partners for *CDH1*.

The genes and pathways identified in Tables C.1 and C.2 were derived from comparing the expression profiles of potential partners to the mutation status of *CDH1* (as shown in Figure 3.2). The following analysis was limited to the samples for which both expression and somatic mutation data were available from TCGA.

C.2 Synthetic Lethal Expression Profiles

Similar to the analysis of synthetic lethal partners against low *CDH1* expression in 4.1.2, the partners detected from *CDH1* mutation were also examined for their expression profiles and the pathway composition of gene clusters. Hierarchical clustering was performed on mtSLIPT partners for *CDH1* as showing in Figure C.1. Over-representation for Reactome pathways for each of the gene clusters identified is given in Table C.3.




Figure C.1: Synthetic lethal expression profiles of analysed samples. Gene expression profile heatmap (correlation distance) of all samples (separated by *CDH1* somatic mutation status) analysed in TCGA breast cancer dataset for gene expression of 3743 candidate partners of E-cadherin (*CDH1*) from mtSLIPT prediction (with significant FDR adjusted $p < 0.05$). Deeply clustered, inter-correlated genes form several main groups, each containing genes that were SL candidates or toxic in an siRNA screen Telford *et al.* (2015). Clusters had different sample groups highly expressing the synthetic lethal candidates in *CDH1* mutant samples and often lowly expressing *CDH1* wild-type samples (which were not tested for), although many of the *CDH1* mutant samples had among the lowest *CDH1* expression. In contrast to the expression analysis the (predominantly *CDH1* wild-type) basal subtype and ER negative samples have depleted expression among most candidate synthetic lethal partners.

Table C.3: Pathways for clusters of *CDH1* partners from mtSLIPT

Pathways Over-represented in Cluster 1	Pathway Size	Cluster Genes	p-value (FDR)
Olfactory Signalling Pathway	57	8	7.1×10^{-9}
Assembly of the primary cilium	149	14	8.0×10^{-9}
Sphingolipid metabolism	62	8	9.6×10^{-9}
Signalling by ERBB4	133	12	5.1×10^{-8}
PI3K Cascade	65	7	4.9×10^{-7}
Circadian Clock	33	5	4.9×10^{-7}
Nuclear signalling by ERBB4	34	5	4.9×10^{-7}
Intraflagellar transport	35	5	4.9×10^{-7}
PI3K events in ERBB4 signalling	87	8	4.9×10^{-7}
PIP3 activates AKT signalling	87	8	4.9×10^{-7}
PI3K events in ERBB2 signalling	87	8	4.9×10^{-7}
PI-3K cascade:FGFR1	87	8	4.9×10^{-7}
PI-3K cascade:FGFR2	87	8	4.9×10^{-7}
PI-3K cascade:FGFR3	87	8	4.9×10^{-7}
PI-3K cascade:FGFR4	87	8	4.9×10^{-7}
Deadenylation of mRNA	22	4	5.6×10^{-7}
PI3K/AKT activation	90	8	5.6×10^{-7}
Cargo trafficking to the periciliary membrane	38	5	5.6×10^{-7}
Pathways Over-represented in Cluster 2	Pathway Size	Cluster Genes	p-value (FDR)
G _{αs} signalling events	83	19	5.1×10^{-25}
Extracellular matrix organization	238	30	1.4×10^{-18}
Hemostasis	422	46	2.7×10^{-16}
Aquaporin-mediated transport	32	9	2.7×10^{-16}
Transcriptional regulation of white adipocyte differentiation	56	11	1.7×10^{-15}
Degradation of the extracellular matrix	102	15	1.7×10^{-15}
Integration of energy metabolism	84	13	8.8×10^{-15}
GPCR downstream signalling	472	48	2.8×10^{-14}
G _{αx} signalling events	15	6	5.0×10^{-14}
Molecules associated with elastic fibres	33	8	5.4×10^{-14}
Phase 1 - Functionalization of compounds	67	11	5.6×10^{-14}
Platelet activation, signalling and aggregation	179	20	5.6×10^{-14}
Vasopressin regulates renal water homeostasis via Aquaporins	24	7	6.1×10^{-14}
Elastic fibre formation	37	8	$.03 \times 10^{-13}$
Calmodulin induced events	27	7	3.3×10^{-13}
CaM pathway	27	7	3.3×10^{-13}
cGMP effects	18	6	3.6×10^{-13}
G _{αi} signalling events	167	18	6.3×10^{-13}
Pathways Over-represented in Cluster 3	Pathway Size	Cluster Genes	p-value (FDR)
Eukaryotic Translation Elongation	86	55	1.1×10^{-112}
Peptide chain elongation	83	54	1.3×10^{-112}
Viral mRNA Translation	81	53	1.6×10^{-111}
Eukaryotic Translation Termination	83	53	7.1×10^{-110}
Nonsense Mediated Decay independent of the Exon Junction Complex	88	54	1.0×10^{-108}
Formation of a pool of free 40S subunits	93	53	4.1×10^{-102}
Nonsense-Mediated Decay	103	54	3.9×10^{-98}
Nonsense Mediated Decay enhanced by the Exon Junction Complex	103	54	3.9×10^{-98}
L13a-mediated translational silencing of Ceruloplasmin expression	103	53	1.2×10^{-95}
3' -UTR-mediated translational regulation	103	53	1.2×10^{-95}
SRP-dependent cotranslational protein targeting to membrane	104	53	4.3×10^{-95}
GTP hydrolysis and joining of the 60S ribosomal subunit	104	53	4.3×10^{-95}
Influenza Viral RNA Transcription and Replication	108	53	9.6×10^{-93}
Eukaryotic Translation Initiation	111	53	4.2×10^{-91}
Cap-dependent Translation Initiation	111	53	4.2×10^{-91}
Influenza Life Cycle	112	53	1.4×10^{-90}
Influenza Infection	117	53	6.2×10^{-88}
Translation	141	55	3×10^{-81}
Pathways Over-represented in Cluster 4	Pathway Size	Cluster Genes	p-value (FDR)
ECM proteoglycans	66	10	2.9×10^{-11}
deactivation of the beta-catenin transactivating complex	38	7	5.1×10^{-10}
Arachidonic acid metabolism	41	7	1.1×10^{-9}
G _{αq} signalling events	149	14	4.0×10^{-9}
HS-GAG degradation	21	5	4.5×10^{-9}
Uptake and actions of bacterial toxins	22	5	6.1×10^{-9}
Gastrin-CREB signalling pathway via PKC and MAPK	170	15	6.1×10^{-9}
RNA Polymerase I, RNA Polymerase III, and Mitochondrial Transcription	64	8	6.1×10^{-9}
Non-integrin membrane-ECM interactions	53	7	1.5×10^{-8}
Syndecan interactions	25	5	1.5×10^{-8}
NOTCH1 Intracellular Domain Regulates Transcription	40	6	2.3×10^{-8}
Synthesis of Leukotrienes and Eoxins	15	4	3.2×10^{-8}
Signalling by NOTCH1	59	7	5.3×10^{-8}
Regulation of insulin secretion	44	6	6.0×10^{-8}
Metabolism of lipids and lipoproteins	471	37	8.2×10^{-8}
Signalling by NOTCH1	80	8	1.2×10^{-7}
Platelet activation, signalling and aggregation	179	14	1.2×10^{-7}
Recruitment of mitotic centrosome proteins and complexes	64	7	1.2×10^{-7}

Pathway over-representation analysis for Reactome pathways with the number of genes in each pathway (Pathway Size), number of genes within the pathway identified (Cluster Genes), and the pathway over-representation p-value (adjusted by FDR) from the hypergeometric test.

C.3 Comparison to Primary Screen

The mutation synthetic lethal partners with *CDH1* were also compared to siRNA primary screen data (Telford *et al.*, 2015), as performed in Section 4.2.1. These were expected to be more concordant with the experimental results performed on a null mutant, however this was not the case at the gene level: less genes overlapped with experimental candidates in Figure C.2. This discrepancy was may be due to lower sample size for mutations in TCGA data or lower frequency (expected value) of *CDH1* mutations compared to low expression.




Figure C.2: **Comparison of mtSLIPT to siRNA.** Testing the overlap of gene candidates for E-cadherin synthetic lethal partners between computational (SLIPT) and experimental screening (siRNA) approaches. The χ^2 test suggests that the overlap is no more than would be expected by chance ($p = 0.281$).

Despite a lower sample size (and low number of a predicted partners) for mutation analysis, the pathway composition (Tables C.2 and C.4) was similar to expression analysis, as described in Section 4.2.5. In particular, the resampling analysis (Section C.3.1) supported many of the results of expression analysis (Section 4.2.5.1). Tables C.5 and C.6 detected many of the same or functionally-related pathways.

Table C.4: Pathways for *CDH1* partners from mtSLIPT and siRNA

Predicted only by SLIPT (2901 genes)	Pathway Size	Genes Identified	p-value (FDR)
Eukaryotic Translation Elongation	87	57	2.8×10^{-120}
Peptide chain elongation	84	56	3.1×10^{-120}
Eukaryotic Translation Termination	84	55	2.8×10^{-117}
Viral mRNA Translation	82	54	4.1×10^{-116}
Nonsense Mediated Decay independent of the Exon Junction Complex	89	55	3.7×10^{-113}
Formation of a pool of free 40S subunits	94	55	2.8×10^{-109}
Nonsense-Mediated Decay	104	57	8.4×10^{-108}
Nonsense Mediated Decay enhanced by the Exon Junction Complex	104	57	8.4×10^{-108}
L13a-mediated translational silencing of Ceruloplasmin expression	104	56	3.4×10^{-105}
3' -UTR-mediated translational regulation	104	56	3.4×10^{-105}
GTP hydrolysis and joining of the 60S ribosomal subunit	105	56	1.4×10^{-104}
Eukaryotic Translation Initiation	112	56	2.8×10^{-100}
Cap-dependent Translation Initiation	112	56	2.8×10^{-100}
SRP-dependent cotranslational protein targeting to membrane	105	54	2.2×10^{-99}
Influenza Viral RNA Transcription and Replication	109	54	5.3×10^{-97}
Influenza Life Cycle	113	54	9.6×10^{-95}
Influenza Infection	118	55	1.7×10^{-94}
Translation	142	60	3.5×10^{-94}
Infectious disease	349	77	5.9×10^{-62}
Extracellular matrix organization	241	54	3.0×10^{-52}

Detected only by siRNA screen (1752 genes)	Pathway Size	Genes Identified	p-value (FDR)
Class A/1 (Rhodopsin-like receptors)	282	69	1.9×10^{-59}
GPCR ligand binding	363	78	2.7×10^{-54}
Peptide ligand-binding receptors	175	41	1.5×10^{-42}
$G_{\alpha i}$ signalling events	184	41	1.1×10^{-40}
Gastrin-CREB signalling pathway via PKC and MAPK	180	37	1.5×10^{-35}
$G_{\alpha q}$ signalling events	159	34	3.7×10^{-35}
DAP12 interactions	159	27	1.1×10^{-24}
VEGFA-VEGFR2 Pathway	91	19	1.0×10^{-23}
Downstream signal transduction	146	24	1.9×10^{-22}
Signalling by VEGF	99	19	2.6×10^{-22}
DAP12 signalling	149	24	4.2×10^{-22}
Organelle biogenesis and maintenance	264	34	4.3×10^{-20}
Downstream signalling of activated FGFR1	134	21	4.3×10^{-20}
Downstream signalling of activated FGFR2	134	21	4.3×10^{-20}
Downstream signalling of activated FGFR3	134	21	4.3×10^{-20}
Downstream signalling of activated FGFR4	134	21	4.3×10^{-20}
Signalling by ERBB2	146	22	5.3×10^{-20}
Signalling by FGFR	146	22	5.3×10^{-20}
Signalling by FGFR1	146	22	5.3×10^{-20}
Signalling by FGFR2	146	22	5.3×10^{-20}

Intersection of SLIPT and siRNA screen (450 genes)	Pathway Size	Genes Identified	p-value (FDR)
HS-GAG degradation	21	4	4.9×10^{-6}
Retinoid metabolism and transport	39	5	4.9×10^{-6}
Platelet activation, signalling and aggregation	186	13	4.9×10^{-6}
Signalling by NOTCH4	11	3	4.9×10^{-6}
$G_{\alpha s}$ signalling events	100	8	5.0×10^{-6}
Defective EXT2 causes exostoses 2	12	3	5.0×10^{-6}
Defective EXT1 causes exostoses 1, TRPS2 and CHDS	12	3	5.0×10^{-6}
Class A/1 (Rhodopsin-like receptors)	289	18	2.2×10^{-5}
Signalling by PDGF	173	11	2.9×10^{-5}
Circadian Clock	34	4	2.9×10^{-5}
Signalling by ERBB4	139	9	4.3×10^{-5}
Role of LAT2/NTAL/LAB on calcium mobilization	99	7	4.4×10^{-5}
Peptide ligand-binding receptors	181	11	4.5×10^{-5}
Defective B4GALT7 causes EDS, progeroid type	19	3	4.5×10^{-5}
Defective B3GAT3 causes JDSSDHD	19	3	4.5×10^{-5}
Signalling by NOTCH	80	6	4.5×10^{-5}
$G_{\alpha q}$ signalling events	164	10	5.1×10^{-5}
Response to elevated platelet cytosolic Ca^{2+}	84	6	7.1×10^{-5}
Signalling by ERBB2	148	9	7.1×10^{-5}
Signalling by SCF-KIT	129	8	8.3×10^{-5}

C.3.1 Resampling Analysis

Table C.5: Pathways for *CDH1* partners from mtSLIPT

Reactome Pathway	Over-representation	Permutation
Eukaryotic Translation Elongation	3.2×10^{-128}	$< 7.035 \times 10^{-4}$
Peptide chain elongation	3.2×10^{-128}	$< 7.035 \times 10^{-4}$
Eukaryotic Translation Termination	3.7×10^{-125}	$< 7.035 \times 10^{-4}$
Viral mRNA Translation	4.1×10^{-124}	$< 7.035 \times 10^{-4}$
Nonsense Mediated Decay independent of the Exon Junction Complex	1.4×10^{-123}	$< 7.035 \times 10^{-4}$
Nonsense-Mediated Decay	8.4×10^{-117}	$< 7.035 \times 10^{-4}$
Nonsense Mediated Decay enhanced by the Exon Junction Complex	8.4×10^{-117}	$< 7.035 \times 10^{-4}$
Formation of a pool of free 40S subunits	2.6×10^{-116}	$< 7.035 \times 10^{-4}$
L13a-mediated translational silencing of Ceruloplasmin expression	2.0×10^{-111}	$< 7.035 \times 10^{-4}$
3' -UTR-mediated translational regulation	2.0×10^{-111}	$< 7.035 \times 10^{-4}$
GTP hydrolysis and joining of the 60S ribosomal subunit	9.9×10^{-111}	$< 7.035 \times 10^{-4}$
SRP-dependent cotranslational protein targeting to membrane	4.7×10^{-108}	$< 7.035 \times 10^{-4}$
Eukaryotic Translation Initiation	4.8×10^{-106}	$< 7.035 \times 10^{-4}$
Cap-dependent Translation Initiation	4.8×10^{-106}	$< 7.035 \times 10^{-4}$
Influenza Viral RNA Transcription and Replication	8.1×10^{-103}	$< 7.035 \times 10^{-4}$
Influenza Infection	2.4×10^{-102}	$< 7.035 \times 10^{-4}$
Translation	6.0×10^{-101}	$< 7.035 \times 10^{-4}$
Influenza Life Cycle	2.2×10^{-100}	$< 7.035 \times 10^{-4}$
Disease	2.1×10^{-90}	0.013347
GPCR downstream signalling	1.6×10^{-80}	0.095478
Hemostasis	2.1×10^{-78}	0.2671
Signalling by GPCR	1.2×10^{-73}	0.44939
<i>Extracellular matrix organization</i>	2.2×10^{-67}	0.054008
Metabolism of proteins	1.4×10^{-66}	0.9607
Signal Transduction	2.1×10^{-66}	0.48184
Developmental Biology	2.5×10^{-66}	0.54075
Innate Immune System	5.3×10^{-66}	0.9589
Infectious disease	9.6×10^{-66}	0.21075
Signalling by NGF	1.1×10^{-62}	0.43356
Immune System	2.8×10^{-62}	0.23052

Over-representation (hypergeometric test) and Permutation p-values adjusted for multiple tests across pathways (FDR). Significant pathways were marked in bold (FDR < 0.05) and italics (FDR < 0.1).

Table C.6: Pathways for *CDH1* partners from mtSLIPT and siRNA primary screen

Reactome Pathway	Over-representation	Permutation
Visual phototransduction	1.2×10^{-9}	0.86279
G_{as} signalling events	2.9×10^{-7}	0.023066
Retinoid metabolism and transport	2.9×10^{-7}	0.299
Acylic chain remodelling of PS	1.1×10^{-5}	0.42584
Transcriptional regulation of white adipocyte differentiation	1.1×10^{-5}	0.53928
Chemokine receptors bind chemokines	1.1×10^{-5}	0.95259
<i>Signalling by NOTCH4</i>	1.2×10^{-5}	0.079229
Defective EXT2 causes exostoses 2	1.2×10^{-5}	0.22292
Defective EXT1 causes exostoses 1, TRPS2 and CHDS	1.2×10^{-5}	0.22292
Platelet activation, signalling and aggregation	1.2×10^{-5}	0.48853
Serotonin receptors	1.4×10^{-5}	0.34596
Nicotinamide salvaging	1.4×10^{-5}	0.70881
Phase 1 - Functionalization of compounds	2×10^{-5}	0.31142
Amine ligand-binding receptors	2.5×10^{-5}	0.34934
Acylic chain remodelling of PE	3.8×10^{-5}	0.42615
Signalling by GPCR	3.8×10^{-5}	0.93888
Molecules associated with elastic fibres	3.9×10^{-5}	0.017982
DAP12 interactions	3.9×10^{-5}	0.71983
Beta defensins	3.9×10^{-5}	0.91458
Cytochrome P ₄₅₀ - arranged by substrate type	4.7×10^{-5}	0.83493
GPCR ligand binding	5.7×10^{-5}	0.95258
Acylic chain remodelling of PC	6.1×10^{-5}	0.42584
Response to elevated platelet cytosolic Ca ²⁺	6.4×10^{-5}	0.54046
Arachidonic acid metabolism	6.7×10^{-5}	0.026696
Defective B4GALT7 causes EDS, progeroid type	7.3×10^{-5}	0.24921
Defective B3GAT3 causes JDSSDHD	7.3×10^{-5}	0.24921
Hydrolysis of LPC	7.3×10^{-5}	0.80663
Elastic fibre formation	7.4×10^{-5}	0.0058768
HS-GAG degradation	9.4×10^{-5}	0.0083179
<i>Bile acid and bile salt metabolism</i>	9.4×10^{-5}	0.079905
Netrin-1 signalling	0.00011	0.92216
Integration of energy metabolism	0.00011	0.011152
Dectin-2 family	0.00012	0.10385
Platelet sensitization by LDL	0.00012	0.34596
DAP12 signalling	0.00012	0.62787
Defensins	0.00012	0.77542
GPCR downstream signalling	0.00012	0.79454
<i>Diseases associated with glycosaminoglycan metabolism</i>	0.00013	0.065927
<i>Diseases of glycosylation</i>	0.00013	0.065927
Signalling by Retinoic Acid	0.00013	0.22292
Signalling by Leptin	0.00013	0.34596
Signalling by SCF-KIT	0.00013	0.70881
Opioid Signalling	0.00013	0.96053
Signalling by NOTCH	0.00015	0.26884
Platelet homeostasis	0.00015	0.4878
Signalling by NOTCH1	0.00016	0.13043
Class B/2 (Secretin family receptors)	0.00016	0.13994
<i>Diseases of Immune System</i>	0.0002	0.0795
<i>Diseases associated with the TLR signalling cascade</i>	0.0002	0.0795
A tetrasaccharide linker sequence is required for GAG synthesis	0.0002	0.42615

Over-representation (hypergeometric test) and Permutation p-values adjusted for multiple tests across pathways (FDR). Significant pathways were marked in bold (FDR < 0.05) and italics (FDR < 0.1).

C.4 Compare SLIPT genes

The mutation synthetic lethal partners with *CDH1* were also compared to siRNA primary screen data (Telford *et al.*, 2015), by correlation and siRNA viability as described in sections 4.2.2 and 4.2.3.




Figure C.3: Compare mtSLIPT and siRNA genes with correlation. The mtSLIPT p-values were compared against Pearson correlation of expression with *CDH1*. Genes detected by SLIPT or siRNA were coloured according to the legend.




Figure C.4: Compare mtSLIPT and siRNA genes with correlation. Genes detected by mtSLIPT against *CDH1* mutation and siRNA screening were compared against Pearson correlation of expression with *CDH1*. There were no differences in correlation between the gene groups.




Figure C.5: Compare mtSLIPT and siRNA genes with siRNA viability. Genes detected as candidate synthetic lethal partners by mtSLIPT (in TCGA breast cancer) expression analysis against *CDH1* mutation and experimental screening (with siRNA) were compared against the viability ratio of *CDH1* mutant and wild-type cells in the primary siRNA screen. There were clear no differences in viability between genes detected by mtSLIPT and those not with the differences being primarily due to viability thresholds that were used to detect synthetic lethality by Telford *et al.* (2015).

Appendix D

Metagene Analysis

The gene signatures (Gatza *et al.*, 2011, 2014) were used to demonstrate the utility of the metagene approach for use on a wider range of pathways as was performed with the Reactome (Croft *et al.*, 2014) pathways as an alternative approach to identification of synthetic lethal pathways. The direction of metagenes is arbitrary but they have been corrected to ensure the metagene increases in a direction which reflects overall activation of the pathways (as described in Section 2.2.3) which was verified by examining in the following pathway signatures in breast cancer.

Metagenes were derived for well characterised gene signatures in breast cancer (Gatza *et al.*, 2011, 2014). These pathways signatures have expected molecular properties in clinical and molecular subtypes (Parker *et al.*, 2009; Perou *et al.*, 2000). This was performed by examining the pathways expression of these breast cancer gene signatures in TCGA expression data. These metagenes were also compared to somatic mutation to evaluate mutation as a measure of gene activity in comparison to gene and protein expression.

Having established that metagenes generated with this procedure reflect gene activity, the metagene procedure (in Section 2.2.3) was then applied to the Reactome pathways (Croft *et al.*, 2014). These Reactome metagenes were used for synthetic lethal analysis of pathways with SLIPT, directly using pathways activity for identifying synthetic lethal pathways with *CDH1*.

D.1 Pathway Signature Expression

Pathway metagenes (generated as described in Section 2.2.3) for gene signatures of key processes in breast cancer (Gatza *et al.*, 2011) were used to check that metagenes were generated in the correct direction to indicate pathways activation. Some of these

gene signatures are plotted in Figure D.1 for comparison with clinical factors and somatic mutations. The “intrinsic subtypes” was computed by performing the Prediction Analysis of Microarray 50 (PAM50) procedure Parker *et al.* (2009) for RNA-Seq data which was highly concordant ($\chi^2 = 1305.9$, $p = 2.73 \times 10^{-268}$) with the subtypes provided by University of California, Santa Cruz (UCSC) (UCSC, 2012) for TCGA samples (Koboldt *et al.*, 2012) previously analysed by microarrays (as shown in Appendix E). Somatic mutations were reported for gislinkrecurrent mutationrecurrently mutated genes in breast cancer, as reported by TCGA (Koboldt *et al.*, 2012), related genes, and those previously discussed to be important in hereditary breast cancers (*BRCA1*, *BRCA2*, and *CDH1*).

These gene signatures reflect intrinsic subtypes as expected. In particular, the estrogen and progesterone receptor signatures are low in the predominantly ER⁻ and Progesterone receptor (PR)⁻ basal-like subtype tumours. These tumours also had the highest frequency of *TP53* mutations and a corresponding reduction of p53 metagene activity, as expected for loss of a tumour suppressor. The luminal A and luminal B tumour subtypes are the most similar, which is reflected in these metagenes signatures, although they are distinguishable molecular subtypes as shown by elevated phosphoinositide 3-kinase (PI3K), AKT, RAS, and β -catenin signalling in luminal B tumours. However, these pathways were also elevated in basal-like and HER2-enriched subtypes and lowly expressed in the “normal-like” subtype (which contained the normal samples). These intrinsic subtypes specific gene signature profiles were further supported with metagenes for an extended set of signatures (Gatza *et al.*, 2014), as shown in Figure D.4.

TP53 mutations were the most frequent and more common in the basal-like subtype. Similarly, *GATA3* mutations were more common in luminal subtype tumours. PI3K mutations were more frequent across breast tumours, although these were less common in the basal-like subtype despite an elevated metagene (this discrepancy will be discussed further in Section D.2). *CDH1* mutations similarly occurred across molecular subtypes with the exception of the basal-like subtype (as observed in gene expression with Figure 4.1). *CDH1* low samples occurred in all subtypes but were predominantly of the lobular histological subtype. Apart from these genes, mutations did not show clear specificity to a particular subtype and the variation between samples reflects the range of molecular cascades that can result in tumours with similar molecular profiles, supporting the use of gene expression data for cancer diagnostics and identification of molecular targets.




Figure D.1: Pathway metagene expression profiles. Expression profiles for metagene signatures from Gatza *et al.* (2011) in TCGA breast data, annotated for clinical factors (with sample types and histological results coloured according to the legend) and cancer gene mutations (Negative values for mutation are light grey with missing data in white). Intrinsic subtypes are shown as derived from microarray (UCSC) and RNA-Seq (PAM50) data (Koboldt *et al.*, 2012; Parker *et al.*, 2009). Samples were clustered independently for each intrinsic subtypes and by *CDH1* expression status. Pathway expression signatures are consistent with mutations and clinical subgroups.

The direction of each metagene was consistent with the clinical characteristics, which formed a consensus of gene activity as shown for the PI3K and ER signatures (Gatza *et al.*, 2011) in Figures D.2 and D.3, respectively. Supporting data for p53 and BRCA metagenes (Gatza *et al.*, 2011, 2014) are given in the Appendix (Figures D.5 and D.6). In each of the examples for gene signatures, the expression of the majority of the genes were highly concordant with the metagene, being either positively or negatively correlated. These were generally consistent with established clinical and molecular subtypes of breast cancer and the recurrent mutations shown. However, the *PIK3CA* and *PIK3R1* mutant samples did not necessarily have elevated PI3K pathways metagene activity (as shown in Figure D.2).




Figure D.2: **Expression profiles for constituent genes of PI3K.** Expression profiles the genes contained in the PI3K gene signature from Gatza *et al.* (2011) in TCGA breast data, annotated for clinical factors and cancer gene mutations. Samples are separated by *CDH1* expression status and sorted by the metagene. In both cases, the majority of genes were consistent with the direction of the PI3K metagene, although considerable proportion were inversely correlated with the metagene. Normal samples had low PI3K metagene expression and *TP53* mutant samples had high PI3K expression. Although, oncogenic *PIK3CA* and tumour suppressor *PIK3R1* mutations across samples including those with low metagene response.




Figure D.3: **Expression profiles for estrogen receptor related genes.** Expression profiles the genes contained in the estrogen receptor (ER) gene signature from Gatza *et al.* (2011) in TCGA breast data, annotated for clinical factors and cancer gene mutations. Samples are separated by *CDH1* expression status and sorted by the metagene. In both cases, the majority of genes were consistent with the direction of the metagene, with very few exceptions being inversely correlated. Estrogen receptor (by antibody staining) negative samples had low metagene expression, as expected. These were more likely to be ductal and basal subtypes, lacking *CDH1* or *PIK3CA* mutations.




Figure D.4: **Pathway metagene expression profiles.** Expression profiles for metagene signatures from Gatza *et al.* (2014) in TCGA breast data, annotated for clinical factors and cancer gene mutations.




Figure D.5: Expression profiles for p53 related genes. Expression profiles the genes contained in the *TP53* gene signature from Gatza *et al.* (2011) in TCGA breast data, annotated for clinical factors and cancer gene mutations. Samples were separated by *CDH1* expression status and sorted by the metagene. In both cases, the majority of genes were consistent with the direction of the metagene, with few very exceptions. *TP53* mutant samples had low metagene expression, consistent with loss of tumour suppressor functions, and were less likely to have *CDH1* or *PIK3CA* mutations.




Figure D.6: **Expression profiles for BRCA related genes.** Expression profiles the genes contained in the gene signature related to *BRCA1* and *BRCA2* functions from Gatza *et al.* (2014) in TCGA breast data, annotated for clinical factors and cancer gene mutations. Samples were separated by *CDH1* expression status and sorted by the metagene. In both cases, the majority of genes were consistent with the direction of the metagene, with few very exceptions. *BRCA1* and *BRCA2* mutant samples had higher metagene expression than most samples for the ductal subtype, although this was not the case (for the lobular samples for which the metagene was lower). However, the metagene was higher for basal subtype and ER negative samples.

D.2 Somatic Mutation




Figure D.7: Somatic mutation against the PI3K metagene. Mutations in *PIK3CA*, *PIK3R1*, *CDH1*, and *TP53* were examined in TCGA breast cancer for their association with the PI3K (Gatza *et al.*, 2011) pathways metagene. The tumour suppressors *CDH1* and *TP53* showed an increase and decrease in the metagene respectively, whereas *PIK3CA* and *PIK3R1* mutations had little effect on the metagene levels.

It should be noted that metagenes, while consistent with the consensus of constituent expressed genes, were not necessarily reflecting the somatic mutation status. The PI3K (Gatza *et al.*, 2011) metagene levels in particular, were not statistically significantly varying between mutant and wild-type *PIK3CA* samples (shown in Figure D.7). However, the PI3K metagene differed across *CDH1* and *TP53* mutations, remarkably in opposite directions considering that PI3K is an oncogenic growth pathways and these are both most frequently tumour suppressors inactivated in cancers. This shows that

CDH1 and *TP53* deficient tumours have distinct molecular growth pathways and that synthetic lethal interventions against loss of *CDH1* function may not be applicable to other cancers with driver mutations such as *TP53*, although these were kept in the analysis for comparison. These differences may be related to these mutations being more frequent in tumours with difference clinical characteristics (as observed in Section D.1). Thus mutations do not necessarily have corresponding changes in pathways expression, particularly for oncogenes which may change in function rather than being upregulated.

While the more specific *PIK3CA* (Gatza *et al.*, 2014) metagene showed significant differences with *PIK3CA* and *PIK3R1* mutations (as shown in Figure D.8), this metagene replicated stronger differences for *CDH1* and *TP53*. These differences were less pronounced in the protein levels of p110 α (enocded by *PIK3CA*) and the downstream AKT gene (shown in Figures D.9 and D.10 respectively). However, this may be due to this regulatory cascade (kinases) being transmitted as a change in protein state (phosphorylation) rather than changes in expression levels. Another consideration is that mutations at different loci have different effects on protein function, particularly for oncogenes.

D.3 Synthetic Lethal Reactome Metagenes

Metagene analysis was performed for synthetic lethal pathways against *CDH1* mutation. These were described and compared to expression analysis in Section 4.3.

Table D.1: Candidate synthetic lethal metagenes against *CDH1* from mtSLIPT

Pathway	ID	Observed	Expected	χ^2 value	p-value	p-value (FDR)
Neurotoxicity of clostridium toxins	168799	8	36.7	79.4	5.71×10^{-18}	3.14×10^{-15}
Aquaporin-mediated transport	445717	8	36.7	76.3	2.73×10^{-17}	9.01×10^{-15}
Toxicity of botulinum toxin type G (BoNT/G)	5250989	8	36.7	76.3	2.73×10^{-17}	9.01×10^{-15}
ABC-family proteins mediated transport	382556	10	36.7	68.2	1.58×10^{-15}	1.86×10^{-13}
G _{αz} signalling events	418597	10	36.7	59.9	9.97×10^{-14}	5.48×10^{-12}
Regulation of IGF transport and uptake by IGFBPs	381426	9	36.7	56.3	5.88×10^{-13}	2.11×10^{-11}
GP1b-IX-V activation signalling	430116	8	36.7	55.7	8.20×10^{-13}	2.76×10^{-11}
GABA receptor activation	977443	12	36.7	55.1	1.07×10^{-12}	3.26×10^{-11}
Vasopressin regulates renal water homeostasis via Aquaporins	432040	9	36.7	54.1	1.77×10^{-12}	4.88×10^{-11}
Toxicity of botulinum toxin type D (BoNT/D)	5250955	14	36.7	53.4	2.54×10^{-12}	6.64×10^{-11}
Toxicity of botulinum toxin type F (BoNT/F)	5250981	14	36.7	53.4	2.54×10^{-12}	6.64×10^{-11}
STAT6-mediated induction of chemokines	3249367	16	36.7	52.2	4.72×10^{-12}	1.13×10^{-10}
Toxicity of botulinum toxin type B (BoNT/B)	5250958	14	36.7	50.8	9.5×10^{-12}	1.98×10^{-10}
S6K1 signalling	165720	12	36.7	50.2	1.24×10^{-11}	2.5×10^{-10}
G _{αs} signalling events	418555	11	36.7	49.2	2.08×10^{-11}	3.85×10^{-10}
RHO GTPases activate CIT	5625900	14	36.7	48.2	3.34×10^{-11}	5.9×10^{-10}
NADE modulates death signalling	205025	15	36.7	47.4	5.00×10^{-11}	8.32×10^{-10}
Keratan sulfate degradation	2022857	10	36.7	46.6	7.5×10^{-11}	1.15×10^{-9}
Signalling by Retinoic Acid	5362517	10	36.7	46.6	7.5×10^{-11}	1.15×10^{-9}
Adenylate cyclase inhibitory pathway	170670	14	36.7	45.9	1.11×10^{-10}	1.59×10^{-9}
Inhibition of adenylate cyclase pathway	997269	14	36.7	45.9	1.11×10^{-10}	1.59×10^{-9}
Fatty acids	211935	6	36.7	45.7	1.21×10^{-10}	1.72×10^{-9}
Ionotropic activity of Kainate Receptors	451306	13	36.7	44.6	2.03×10^{-10}	2.58×10^{-9}
Activation of Ca-permeable Kainate Receptor	451308	13	36.7	44.6	2.03×10^{-10}	2.58×10^{-9}
RA biosynthesis pathway	5365859	13	36.7	44.6	2.03×10^{-10}	2.58×10^{-9}

Strongest candidate synthetic lethal partners for *CDH1* by mtSLIPT with observed and expected numbers of mutant *CDH1* TCGA breast cancer tumours with low expression of partner metagenes.

D.4 Expression of Somatic Mutations




Figure D.8: **Somatic mutation against PIK3CA metagene.** Mutations in *PIK3CA*, *PIK3R1*, *CDH1*, and *TP53* were examined in TCGA breast cancer for their effect on the PIK3CA (Gatza *et al.*, 2014) pathway metagene. The tumour suppressors *CDH1* and *TP53* showed an increase and decrease in the metagene respectively, whereas *PIK3CA* and *PIK3R1* mutations weaker evidence of decrease in metagene levels.




Figure D.9: Somatic mutation against PI3K protein. Mutations in *PIK3CA*, *PIK3R1*, *CDH1*, and *TP53* were examined in TCGA breast cancer for their effect on the expression of the p110 α protein (encoded by *PIK3CA*). Protein levels were significantly elevated in samples with *PIK3CA* or *PIK3R1* mutations and lower in samples with *TP53* mutations.




Figure D.10: Somatic mutation against AKT protein. Mutations in *PIK3CA*, *PIK3R1*, *CDH1*, and *TP53* were examined in TCGA breast cancer for their effect on the expression of the AKT protein (a downstream target of *PIK3CA*). Protein levels were not significantly different in samples with mutations in any of these cancer genes.

Appendix E

Intrinsic Subtyping

The intrinsic subtypes for TCGA breast cancer samples provided by UCSC (Koboldt *et al.*, 2012; UCSC, 2012) that were derived from microarray analysis have been compared to the PAM50 results for performing subtyping from RNA-Seq data (Parker *et al.*, 2009). As shown in Table E.1, these subtypes were highly concordant for samples which had both procedures performed upon them ($\chi^2 = 1305.9$, $p = 2.73 \times 10^{-268}$). The main exception were the luminal A samples some of which were reclassified as luminal B or “normal-like”.

Table E.1: Comparison of intrinsic subtypes

UCSC Subtype					
	Basal-like	HER2-enriched	Luminal A	Luminal B	Normal-like
	100	58	232	128	30
PAM50 Subtype					
	Basal-like	HER2-enriched	Luminal A	Luminal B	Normal-like
	208	94	314	334	227
UCSC Subtype					
PAM50 Subtype	Basal-like	HER2-enriched	Luminal A	Luminal B	Normal-like
Basal-like	96	4	2	2	1
HER2-enriched	0	47	5	3	0
Luminal A	1	0	141	1	0
Luminal B	2	7	49	121	0
Normal-like	1	0	35	1	29

The intrinsic subtypes of TCGA breast samples were compared between those provided by UCSC (Koboldt *et al.*, 2012) from microarray expression to those derived from RNA-Seq data (Parker *et al.*, 2009). Comparisons between these were limited to samples for which both data types were available.

The PAM50 subtypes could be more accurate given similarity of these subtypes and that the remainder of the subtypes were accurately recapitulated with RNA-Seq data. Furthermore, UCSC subtypes correctly identified 22/22 normal samples as “normal-like” and PAM50 subtyping in RNA-Seq data had a success rate of 112/113 (including all of those identified from microarrays). Therefore the PAM50 subtypes (performed on a larger cohort of samples) are appropriate to use for further interpretation, superseding the UCSC subtypes available for a limited set of samples.

Appendix F

Stomach Expression Analysis

The following results are a replication of the TCGA results (in Chapter 4) with stomach cancer data, using synthetic lethality (SLIPT) against *CDH1*.

F.1 Synthetic Lethal Genes and Pathways

Table F.1: Synthetic lethal gene partners of *CDH1* from SLIPT in stomach cancer

Gene	Observed*	Expected*	χ^2 value	p-value	p-value (FDR)
<i>PRAF2</i>	17	50.4	121	3.54×10^{-25}	1.45×10^{-21}
<i>EMP3</i>	17	50.4	115	5.06×10^{-24}	1.48×10^{-20}
<i>PLEKHO1</i>	22	50.4	112	2.14×10^{-23}	4.75×10^{-20}
<i>SELM</i>	20	50.4	111	5.13×10^{-23}	8.09×10^{-20}
<i>GYPC</i>	20	50.4	110	5.77×10^{-23}	8.45×10^{-20}
<i>COX7A1</i>	18	50.4	109	1.15×10^{-22}	1.39×10^{-19}
<i>TNFSF12</i>	20	50.4	106	4.06×10^{-22}	4.38×10^{-19}
<i>SEPT4</i>	17	50.4	106	6.58×10^{-22}	5.91×10^{-19}
<i>LGALS1</i>	19	50.4	105	6.64×10^{-22}	5.91×10^{-19}
<i>RARRES2</i>	27	50.4	105	8.02×10^{-22}	6.85×10^{-19}
<i>VEGFB</i>	16	50.4	104	1.19×10^{-21}	9.74×10^{-19}
<i>PRR24</i>	22	50.4	102	2.96×10^{-21}	2.02×10^{-18}
<i>SYNC</i>	19	50.4	102	3.73×10^{-21}	2.39×10^{-18}
<i>MAGEH1</i>	17	50.4	100	9.52×10^{-21}	5.01×10^{-18}
<i>HSPB2</i>	23	50.4	99.6	1.19×10^{-20}	5.82×10^{-18}
<i>SMARCD3</i>	19	50.4	99	1.59×10^{-20}	7.57×10^{-18}
<i>CREM</i>	13	50.4	98.1	2.48×10^{-20}	1.13×10^{-17}
<i>GNG11</i>	20	50.4	97.3	3.68×10^{-20}	1.59×10^{-17}
<i>GNAI2</i>	17	50.4	96.4	5.75×10^{-20}	2.36×10^{-17}
<i>FUNDC2</i>	22	50.4	95.9	7.39×10^{-20}	2.91×10^{-17}
<i>CNRIP1</i>	21	50.4	95.3	1.0×10^{-19}	3.66×10^{-17}
<i>CALHM2</i>	22	50.4	93.1	2.94×10^{-19}	1.06×10^{-16}
<i>ARID5A</i>	18	50.4	92.7	3.47×10^{-19}	1.22×10^{-16}
<i>ST3GAL3</i>	27	50.4	92.2	4.49×10^{-19}	1.56×10^{-16}
<i>LOC339524</i>	21	50.4	92.1	4.8×10^{-19}	1.59×10^{-16}

Strongest candidate synthetic lethal partners for *CDH1* by SLIPT in TCGA stomach cancer expression data

* Observed and expected numbers of samples which had low expression of both genes

Table F.2: Pathways for *CDH1* partners from SLIPT in stomach cancer

Pathways Over-represented	Pathway Size	SL Genes	p-value (FDR)
Extracellular matrix organization	241	104	7.5×10^{-140}
Hemostasis	445	138	1.8×10^{-121}
Developmental Biology	432	125	9.2×10^{-107}
Axon guidance	289	94	1.5×10^{-102}
Eukaryotic Translation Termination	84	49	1.9×10^{-99}
GPCR ligand binding	373	108	3.8×10^{-99}
Viral mRNA Translation	82	48	3.3×10^{-98}
Formation of a pool of free 40S subunits	94	51	3.3×10^{-98}
Eukaryotic Translation Elongation	87	49	1.6×10^{-97}
Peptide chain elongation	84	48	7.2×10^{-97}
Class A/1 (Rhodopsin-like receptors)	289	90	2.7×10^{-96}
Nonsense Mediated Decay independent of the Exon Junction Complex	89	49	3.0×10^{-96}
Infectious disease	349	100	2.6×10^{-94}
GTP hydrolysis and joining of the 60S ribosomal subunit	105	52	3.4×10^{-94}
L13a-mediated translational silencing of Ceruloplasmin expression	104	51	2.8×10^{-92}
3' -UTR-mediated translational regulation	104	51	2.8×10^{-92}
Neuronal System	272	84	8.4×10^{-92}
SRP-dependent cotranslational protein targeting to membrane	105	51	9.5×10^{-92}
Eukaryotic Translation Initiation	112	52	2.0×10^{-90}
Cap-dependent Translation Initiation	112	52	2.0×10^{-90}

Gene set over-representation analysis (hypergeometric test) for Reactome pathways for *CDH1*.




Figure F.1: Synthetic lethal expression profiles of analysed samples. Gene expression profile heatmap (correlation distance) of all samples (separated by the $1/3$ quantile of *CDH1* expression) analysed in TCGA stomach cancer dataset for gene expression of 4365 candidate partners of E-cadherin (*CDH1*) from SLIPT prediction (with significant FDR adjusted $p < 0.05$). Deeply clustered, inter-correlated genes form several main groups, each containing genes that were SL candidates or toxic in an siRNA screen Telford *et al.* (2015). Clusters had different sample groups highly expressing the synthetic lethal candidates in *CDH1* low samples. Notably, diffuse and *CDH1* mutant samples had elevated expression in one or more distinct clusters, although there was less complexity and variation among candidate synthetic lethal partners than in breast data. *CDH1* low samples also contained most of samples with *CDH1* mutations.

Table F.3: Pathways for clusters of *CDH1* partners in stomach SLIPT

Pathways Over-represented in Cluster 1	Pathway Size	Cluster Genes	p-value (FDR)
Viral mRNA Translation	82	48	1.3×10^{-97}
Formation of a pool of free 40S subunits	94	51	1.3×10^{-97}
Eukaryotic Translation Elongation	87	49	4.8×10^{-97}
Peptide chain elongation	84	48	1.4×10^{-96}
Eukaryotic Translation Termination	84	48	1.4×10^{-96}
GTP hydrolysis and joining of the 60S ribosomal subunit	105	52	7.9×10^{-94}
Nonsense Mediated Decay independent of the Exon Junction Complex	89	48	3.1×10^{-93}
L13a-mediated translational silencing of Ceruloplasmin expression	104	51	5.1×10^{-92}
3'-UTR-mediated translational regulation	104	51	5.1×10^{-92}
SRP-dependent cotranslational protein targeting to membrane	105	51	1.7×10^{-91}
Eukaryotic Translation Initiation	112	52	3.3×10^{-90}
Cap-dependent Translation Initiation	112	52	3.3×10^{-90}
Translation	142	56	3.6×10^{-85}
Nonsense-Mediated Decay	104	48	1.2×10^{-84}
Nonsense Mediated Decay enhanced by the Exon Junction Complex	104	48	1.2×10^{-84}
Influenza Viral RNA Transcription and Replication	109	48	4.1×10^{-82}
Influenza Life Cycle	113	48	3.4×10^{-80}
Influenza Infection	118	48	6.4×10^{-78}
Pathways Over-represented in Cluster 2	Pathway Size	Cluster Genes	p-value (FDR)
Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell	65	12	1.3×10^{-15}
Phosphorylation of CD3 and TCR zeta chains	18	6	1.7×10^{-12}
Generation of second messenger molecules	29	7	2.7×10^{-12}
PD-1 signalling	21	6	7.4×10^{-12}
TCR signalling	62	9	4.3×10^{-11}
Translocation of ZAP-70 to Immunological synapse	16	5	1.1×10^{-10}
Interferon alpha/beta signalling	68	9	1.6×10^{-10}
Initial triggering of complement	17	5	1.6×10^{-10}
IKK complex recruitment mediated by RIP1	19	5	5.1×10^{-10}
TRIF-mediated programmed cell death	10	4	6.2×10^{-10}
Creation of C4 and C2 activators	11	4	1.3×10^{-9}
RHO GTPases Activate NADPH Oxidases	11	4	1.3×10^{-9}
Interferon Signalling	175	15	2.3×10^{-9}
Chemokine receptors bind chemokines	52	7	4.0×10^{-9}
Interferon gamma signalling	74	8	1.6×10^{-8}
TRAF6 mediated induction of TAK1 complex	15	4	1.6×10^{-8}
Activation of IRF3/IRF7 mediated by TBK1/IKK epsilon	16	4	2.7×10^{-8}
Downstream TCR signalling	45	6	3.5×10^{-8}
Pathways Over-represented in Cluster 3	Pathway Size	Cluster Genes	p-value (FDR)
Uptake and actions of bacterial toxins	22	4	3.5×10^{-6}
Neurotoxicity of clostridium toxins	10	3	3.5×10^{-6}
Activation of PPARGC1A (PGC-1alpha) by phosphorylation	10	3	3.5×10^{-6}
SMAD2/SMAD3:SMAD4 heterotrimer regulates transcription	28	4	1.4×10^{-5}
Assembly of the primary cilium	149	10	2.5×10^{-5}
Serotonin Neurotransmitter Release Cycle	15	3	2.5×10^{-5}
Glycosaminoglycan metabolism	114	8	3.3×10^{-5}
Platelet homeostasis	54	5	3.3×10^{-5}
Norepinephrine Neurotransmitter Release Cycle	17	3	3.3×10^{-5}
Acetylcholine Neurotransmitter Release Cycle	17	3	3.3×10^{-5}
G _{αs} signalling events	100	7	5.5×10^{-5}
GABA synthesis, release, reuptake and degradation	19	3	5.6×10^{-5}
deactivation of the beta-catenin transactivating complex	39	4	6.7×10^{-5}
Dopamine Neurotransmitter Release Cycle	20	3	6.7×10^{-5}
IRS-related events triggered by IGF1R	83	6	7.1×10^{-5}
Generic Transcription Pathway	186	11	7.1×10^{-5}
Termination of O-glycan biosynthesis	21	3	7.4×10^{-5}
Kinesins	22	3	8.5×10^{-5}
Pathways Over-represented in Cluster 4	Pathway Size	Cluster Genes	p-value (FDR)
Extracellular matrix organization	241	97	8.8×10^{-126}
Axon guidance	289	75	8.3×10^{-72}
Hemostasis	445	101	8.3×10^{-72}
Developmental Biology	432	95	3.0×10^{-67}
Response to elevated platelet cytosolic Ca ²⁺	84	37	5.8×10^{-67}
Platelet degranulation	79	36	5.8×10^{-67}
Degradation of the extracellular matrix	104	39	6.7×10^{-63}
Platelet activation, signalling and aggregation	186	52	6.6×10^{-62}
ECM proteoglycans	66	31	8.1×10^{-61}
Neuronal System	272	64	5.1×10^{-60}
Signalling by PDGF	173	47	9.7×10^{-57}
Integrin cell surface interactions	82	31	1.9×10^{-53}
Collagen biosynthesis and modifying enzymes	56	26	1.1×10^{-52}
Collagen formation	67	28	1.4×10^{-52}
Class A/1 (Rhodopsin-like receptors)	289	61	2.3×10^{-52}
GPCR ligand binding	373	73	2.8×10^{-52}
Elastic fibre formation	38	22	4.7×10^{-52}
Non-integrin membrane-ECM interactions	53	24	7.0×10^{-49}

Pathway over-representation analysis for Reactome pathways with the number of genes in each pathway (Pathway Size), number of genes within the pathway identified (Cluster Genes), and the pathway over-representation p-value (adjusted by FDR) from the hypergeometric test.

F.2 Comparison to Primary Screen

The synthetic lethal partners with *CDH1* expression in stomach cancers were also compared to siRNA primary screen data (Telford *et al.*, 2015), as performed in Section 4.2.1. These were expected to be more concordant with the experimental results performed on a null mutant, however this was not the case at the gene level: less genes overlapped with experimental candidates in Figure F.2. This may be due to lower sample size for mutations in TCGA data or lower frequency (expected value) of *CDH1* mutations compared to low expression.




Figure F.2: **Comparison of SLIPT in stomach to siRNA.** The overlap of gene candidates for E-cadherin synthetic lethal partners between computational (SLIPT) and experimental screening (siRNA) approaches. The χ^2 test suggests that the overlap is no more than would be expected by chance ($p = 0.281$).

Table F.4: Pathways for *CDH1* partners from SLIPT and siRNA

Predicted only by SLIPT (3392 genes)	Pathway	Size	Genes Identified	p-value (FDR)
Extracellular matrix organization		238	90	3.4×10^{-107}
Eukaryotic Translation Termination		79	46	7.6×10^{-91}
Viral mRNA Translation		77	45	1.2×10^{-89}
Eukaryotic Translation Elongation		82	46	5.8×10^{-89}
Peptide chain elongation		79	45	2.1×10^{-88}
Nonsense Mediated Decay independent of the Exon Junction Complex		84	46	9.4×10^{-88}
Formation of a pool of free 40S subunits		89	47	3.3×10^{-87}
GTP hydrolysis and joining of the 60S ribosomal subunit		100	48	3.2×10^{-83}
Axon guidance		284	84	3.9×10^{-82}
Developmental Biology		426	111	4.2×10^{-82}
L13a-mediated translational silencing of Ceruloplasmin expression		99	47	1.4×10^{-81}
3' -UTR-mediated translational regulation		99	47	1.4×10^{-81}
SRP-dependent cotranslational protein targeting to membrane		99	47	1.4×10^{-81}
Nonsense-Mediated Decay		99	47	1.4×10^{-81}
Nonsense Mediated Decay enhanced by the Exon Junction Complex		99	47	1.4×10^{-81}
Hemostasis		438	112	1.2×10^{-80}
Eukaryotic Translation Initiation		107	48	8.0×10^{-80}
Cap-dependent Translation Initiation		107	48	8.0×10^{-80}
Infectious disease		338	90	1.6×10^{-76}
Neuronal System		267	77	1.6×10^{-76}
Detected only by siRNA screen (1803 genes)	Pathway	Size	Genes Identified	p-value (FDR)
Class A/1 (Rhodopsin-like receptors)		282	62	8.1×10^{-50}
GPCR ligand binding		363	71	4.9×10^{-46}
Peptide ligand-binding receptors		175	38	7.9×10^{-38}
G _{αi} signalling events		184	37	1.1×10^{-34}
Gastrin-CREB signalling pathway via PKC and MAPK		180	35	1.4×10^{-32}
G _{αq} signalling events		159	32	4.8×10^{-32}
DAP12 interactions		159	29	1.4×10^{-27}
Downstream signal transduction		146	26	2.4×10^{-25}
DAP12 signalling		149	26	6.4×10^{-25}
VEGFA-VEGFR2 Pathway		91	19	8.1×10^{-24}
Signalling by PDGF		172	27	5.7×10^{-23}
Signalling by ERBB2		146	24	1.4×10^{-22}
Signalling by VEGF		99	19	2.0×10^{-22}
Visual phototransduction		85	17	1.3×10^{-21}
Downstream signalling of activated FGFR1		134	22	1.3×10^{-21}
Downstream signalling of activated FGFR2		134	22	1.3×10^{-21}
Downstream signalling of activated FGFR3		134	22	1.3×10^{-21}
Downstream signalling of activated FGFR4		134	22	1.3×10^{-21}
Signalling by FGFR		146	23	2.0×10^{-21}
Signalling by FGFR1		146	23	2.0×10^{-21}
Intersection of SLIPT and siRNA screen (547 genes)	Pathway	Size	Genes Identified	p-value (FDR)
Class A/1 (Rhodopsin-like receptors)		282	25	3.9×10^{-9}
Platelet activation, signalling and aggregation		182	17	3.9×10^{-9}
Response to elevated platelet cytosolic Ca ²⁺		82	9	5.5×10^{-8}
Platelet homeostasis		53	7	5.7×10^{-8}
Nucleotide-like (purinergic) receptors		16	4	1.8×10^{-7}
Platelet degranulation		77	8	2.8×10^{-7}
Peptide ligand-binding receptors		175	14	3.8×10^{-7}
Molecules associated with elastic fibres		34	5	7.1×10^{-7}
Amine ligand-binding receptors		35	5	8.6×10^{-7}
G _{αi} signalling events		184	14	9.8×10^{-7}
GPCR ligand binding		363	27	1.1×10^{-6}
Elastic fibre formation		38	5	1.5×10^{-6}
G _{αq} signalling events		159	12	1.9×10^{-6}
Serotonin receptors		12	3	3.8×10^{-6}
P2Y receptors		12	3	3.8×10^{-6}
Signal amplification		16	3	2.3×10^{-5}
Gastrin-CREB signalling pathway via PKC and MAPK		180	12	2.3×10^{-5}
Complement cascade		33	4	2.4×10^{-5}
Glycosaminoglycan metabolism		110	8	2.5×10^{-5}
Glycogen breakdown (glycogenolysis)		17	3	2.7×10^{-5}

F.2.1 Resampling Analysis

Table F.5: Pathways for *CDH1* partners from SLIPT in stomach cancer

Reactome Pathway	Over-representation	Permutation
<i>Extracellular matrix organization</i>	7.5×10^{-140}	0.070215
Hemostasis	1.8×10^{-121}	0.25804
Developmental Biology	9.2×10^{-107}	0.53032
Axon guidance	1.5×10^{-102}	0.6704
Eukaryotic Translation Termination	1.9×10^{-99}	$> 1.031 \times 10^{-5}$
GPCR ligand binding	3.8×10^{-99}	0.54914
Viral mRNA Translation	3.3×10^{-98}	$> 1.031 \times 10^{-5}$
Formation of a pool of free 40S subunits	3.3×10^{-98}	$> 1.031 \times 10^{-5}$
Eukaryotic Translation Elongation	1.6×10^{-97}	$> 1.031 \times 10^{-5}$
Peptide chain elongation	7.2×10^{-97}	$> 1.031 \times 10^{-5}$
Class A/1 (Rhodopsin-like receptors)	2.7×10^{-96}	0.58174
Nonsense Mediated Decay independent of the Exon Junction Complex	3×10^{-96}	$> 1.031 \times 10^{-5}$
Infectious disease	2.6×10^{-94}	0.25484
GTP hydrolysis and joining of the 60S ribosomal subunit	3.4×10^{-94}	$> 1.031 \times 10^{-5}$
L13a-mediated translational silencing of Ceruloplasmin expression	2.8×10^{-92}	$> 1.031 \times 10^{-5}$
3' -UTR-mediated translational regulation	2.8×10^{-92}	$> 1.031 \times 10^{-5}$
Neuronal System	8.4×10^{-92}	0.53433
SRP-dependent cotranslational protein targeting to membrane	9.5×10^{-92}	$> 1.031 \times 10^{-5}$
Eukaryotic Translation Initiation	2.0×10^{-90}	$> 1.031 \times 10^{-5}$
Cap-dependent Translation Initiation	2.0×10^{-90}	$> 1.031 \times 10^{-5}$
Nonsense-Mediated Decay	7.4×10^{-90}	$> 1.031 \times 10^{-5}$
Nonsense Mediated Decay enhanced by the Exon Junction Complex	7.4×10^{-90}	$> 1.031 \times 10^{-5}$
Adaptive Immune System	8.1×10^{-88}	0.14116
Translation	1.3×10^{-87}	$> 1.031 \times 10^{-5}$
Platelet activation, signalling and aggregation	1.3×10^{-86}	0.28959
Influenza Infection	1×10^{-82}	$> 1.031 \times 10^{-5}$
Influenza Viral RNA Transcription and Replication	2.4×10^{-82}	$> 1.031 \times 10^{-5}$
Influenza Life Cycle	2×10^{-80}	$> 1.031 \times 10^{-5}$
Response to elevated platelet cytosolic Ca ²⁺	4.9×10^{-78}	0.50817
Signalling by NGF	1.6×10^{-75}	0.38518
Rho GTPase cycle	5.1×10^{-75}	0.14864
Signalling by PDGF	7.4×10^{-74}	0.40493
<i>Signalling by Rho GTPases</i>	5.1×10^{-73}	0.077217
Glycosaminoglycan metabolism	1.4×10^{-68}	0.52984
<i>G_{ai} signalling events</i>	1.8×10^{-66}	0.9254
Metabolism of carbohydrates	1.1×10^{-65}	0.39501
G_{as} signalling events	2.7×10^{-65}	0.0050293
Potassium Channels	2.7×10^{-65}	0.53359
Transmission across Chemical Synapses	1.8×10^{-64}	0.81833
ECM proteoglycans	3.4×10^{-64}	0.083482
Peptide ligand-binding receptors	4.8×10^{-64}	0.62817
Degradation of the extracellular matrix	1.1×10^{-63}	0.80879
Platelet homeostasis	5.3×10^{-63}	0.53134
NGF signalling via TRKA from the plasma membrane	6.1×10^{-63}	0.57117
Integration of energy metabolism	4.5×10^{-61}	0.10889
Collagen formation	5.4×10^{-61}	0.29896
Integrin cell surface interactions	7×10^{-59}	0.18167
Collagen biosynthesis and modifying enzymes	7×10^{-59}	0.30208
Neurotransmitter Receptor Binding And Downstream Transmission	8.7×10^{-57}	0.82522
In The Postsynaptic Cell		
Signalling by Wnt	8.7×10^{-57}	0.25468

Over-representation (hypergeometric test) and Permutation p-values adjusted for multiple tests across pathways (FDR). Significant pathways were marked in bold (FDR < 0.05) and italics (FDR < 0.1).

Table F.6: Pathways for *CDH1* partners from SLIPT in stomach and siRNA

Reactome Pathway	Over-representation	Permutation
Platelet activation, signalling and aggregation	3.9×10^{-9}	0.49557
Class A/1 (Rhodopsin-like receptors)	3.9×10^{-9}	0.98432
Response to elevated platelet cytosolic Ca ²⁺	5.5×10^{-8}	0.54349
Platelet homeostasis	5.7×10^{-8}	0.45017
Nucleotide-like (purinergic) receptors	1.8×10^{-7}	0.36966
Peptide ligand-binding receptors	3.8×10^{-7}	0.91294
Molecules associated with elastic fibres	7.1×10^{-7}	0.0025868
Amine ligand-binding receptors	8.6×10^{-7}	0.43303
G _{ai} signalling events	9.8×10^{-7}	0.99626
GPCR ligand binding	1.1×10^{-6}	0.97733
Elastic fibre formation	1.5×10^{-6}	0.0025868
G _{aq} signalling events	1.9×10^{-6}	0.86089
P2Y receptors	3.8×10^{-6}	0.18795
Serotonin receptors	3.8×10^{-6}	0.37853
Signal amplification	2.3×10^{-5}	0.47856
Gastrin-CREB signalling pathway via PKC and MAPK	2.3×10^{-5}	0.98567
Complement cascade	2.4×10^{-5}	$> 3.4628 \times 10^{-6}$
Glycosaminoglycan metabolism	2.5×10^{-5}	0.38953
Glycogen breakdown (glycogenolysis)	2.7×10^{-5}	0.83772
Defective B4GALT7 causes EDS, progeroid type	4.9×10^{-5}	0.10792
Defective B3GAT3 causes JDSSDHD	4.9×10^{-5}	0.10792
Role of LAT2/NTAL/LAB on calcium mobilization	5.6×10^{-5}	0.35373
Cell surface interactions at the vascular wall	5.6×10^{-5}	0.47642
G_{as} signalling events	6×10^{-5}	0.019858
Signalling by NOTCH	6×10^{-5}	0.19008
A tetrasaccharide linker sequence is required for GAG synthesis	0.00017	0.47642
Extracellular matrix organization	0.00018	0.0047308
Collagen formation	0.00018	0.19245
Effects of PIP2 hydrolysis	0.0002	0.37779
Syndecan interactions	0.0002	0.37779
Diseases associated with glycosaminoglycan metabolism	0.00023	0.01028
Diseases of glycosylation	0.00023	0.01028
<i>Chondroitin sulfate/dermatan sulfate metabolism</i>	0.00023	0.085541
Integrin alphaIIb beta3 signalling	0.00028	0.76936
Keratan sulfate biosynthesis	0.00034	0.68744
Rho GTPase cycle	0.00034	0.15675
Creation of C4 and C2 activators	0.00035	0.12275
Abacavir transport and metabolism	0.00035	0.12443
Amine compound SLC transporters	0.00037	0.69773
FCER1 mediated NF-kB activation	0.00037	0.69846
Fc epsilon receptor (FCER1) signalling	0.00056	0.43303
Defective EXT2 causes exostoses 2	0.00067	0.16053
Defective EXT1 causes exostoses 1, TRPS2 and CHDS	0.00067	0.16053
<i>Collagen biosynthesis and modifying enzymes</i>	0.00071	0.052911
Keratan sulfate/keratin metabolism	0.00073	0.46533
G alpha (12/13) signalling events	0.00078	0.59164
SEMA3A-Plexin repulsion signalling by inhibiting Integrin adhesion	0.00084	0.038504
Signal attenuation	0.00084	0.37779
Eicosanoid ligand-binding receptors	0.0011	0.11117
SOS-mediated signalling	0.0011	0.25387

Over-representation (hypergeometric test) and Permutation p-values adjusted for multiple tests across pathways (FDR). Significant pathways were marked in bold (FDR < 0.05) and italics (FDR < 0.1).

F.3 Metagene Analysis

Metagenes used to detect synthetic lethal pathways with *CDH1* in stomach cancer.

Table F.7: Synthetic lethal metagenes against *CDH1* in stomach cancer

Pathway	ID	Observed	Expected	χ^2 value	p-value	p-value (FDR)
Cell-Cell communication	1500931	18	50.4	110	7.43×10^{-23}	1.53×10^{-20}
VEGFR2 mediated vascular permeability	5218920	19	50.4	109	1.36×10^{-22}	2.49×10^{-20}
Sema4D in semaphorin signalling	400685	20	50.4	104	1.62×10^{-21}	2.12×10^{-19}
Ion transport by P-type ATPases	936837	17	50.4	100	8.29×10^{-21}	8.06×10^{-19}
Sialic acid metabolism	4085001	19	50.4	95.3	9.95×10^{-20}	7.82×10^{-18}
Synthesis of pyrophosphates in the cytosol	1855167	26	50.4	94	1.86×10^{-19}	1.23×10^{-17}
Keratan sulfate/keratin metabolism	1638074	25	50.4	93.5	2.36×10^{-19}	1.44×10^{-17}
Ion channel transport	983712	19	50.4	92.8	3.37×10^{-19}	1.99×10^{-17}
Keratan sulfate biosynthesis	2022854	26	50.4	91.4	6.79×10^{-19}	3.62×10^{-17}
Arachidonic acid metabolism	2142753	22	50.4	90.6	9.81×10^{-19}	5.07×10^{-17}
RHO GTPases activate CIT	5625900	22	50.4	87	5.80×10^{-18}	2.66×10^{-16}
Stimuli-sensing channels	2672351	25	50.4	85.8	1.03×10^{-17}	4.58×10^{-16}
Synthesis of PI	1483226	19	50.4	85.6	1.15×10^{-17}	4.89×10^{-16}
G-protein activation	202040	19	50.4	85.3	1.34×10^{-17}	5.53×10^{-16}
NrCAM interactions	447038	22	50.4	84.3	2.1×10^{-17}	8.27×10^{-16}
Inwardly rectifying K^+ channels	1296065	24	50.4	83.5	3.19×10^{-17}	1.22×10^{-15}
Calcitonin-like ligand receptors	419812	20	50.4	82.2	6.07×10^{-17}	2.13×10^{-15}
Prostacyclin signalling through prostacyclin receptor	392851	24	50.4	81.8	7.27×10^{-17}	2.5×10^{-15}
Presynaptic function of Kainate receptors	500657	26	50.4	79.7	2.00×10^{-16}	6.34×10^{-15}
ADP signalling through P2Y purinoceptor 12	392170	23	50.4	79.2	2.57×10^{-16}	7.71×10^{-15}
regulation of FZD by ubiquitination	4641263	22	50.4	78.8	3.15×10^{-16}	9.3×10^{-15}
Toxicity of tetanus toxin (TeNT)	5250982	27	50.4	78.7	3.36×10^{-16}	9.75×10^{-15}
Gap junction degradation	190873	21	50.4	78.5	3.66×10^{-16}	1.04×10^{-14}
Nephrin interactions	373753	25	50.4	78.2	4.21×10^{-16}	1.14×10^{-14}
GABA synthesis, release, reuptake and degradation	888590	26	50.4	77	7.69×10^{-16}	1.95×10^{-14}

Strongest candidate synthetic lethal partners for *CDH1* by SLIPT with observed and expected numbers of TCGA stomach cancer samples with low expression of both genes.

Appendix G

Synthetic Lethal Genes in Pathways




Figure G.1: Synthetic lethality in the PI3K/AKT pathway. The Reactome PI3K/AKT pathway with synthetic lethal candidates, coloured as shown in the legend.




Figure G.2: Synthetic lethality in the PI3K/AKT pathway in cancer. The Reactome PI3K/AKT in cancer pathway with synthetic lethal candidates, coloured as shown in the legend.




Figure G.3: Synthetic lethality in the Extracellular Matrix. The Reactome Extracellular Matrix pathway with synthetic lethal candidates, coloured as shown in the legend.




Figure G.4: Synthetic lethality in the GPCRs. The Reactome $G_{\alpha i}$ pathway with synthetic lethal candidates, coloured as shown in the legend.




Figure G.5: Synthetic lethality in the GPCR Downstream. The Reactome G protein coupled receptor (GPCR) Downstream pathway with synthetic lethal candidates, coloured as shown in the legend.




Figure G.6: **Synthetic lethality in the Translation Elongation.** The Reactome Translation Elongation pathway with synthetic lethal candidates, coloured as shown in the legend.




Figure G.7: Synthetic lethality in the Nonsense-mediated Decay. The Reactome nonsense-mediated decay (NMD) pathway with synthetic lethal candidates, coloured as shown in the legend.




Figure G.8: **Synthetic lethality in the 3' UTR.** The Reactome 3' untranslated region (UTR) pathway with synthetic lethal candidates, coloured as shown in the legend.

Appendix H

Pathway Connectivity for Mutation SLIPT




Figure H.1: **Synthetic lethality and vertex degree.** The number of connected genes (vertex degree) was compared (on a log-scale across genes detected by mtSLIPT and siRNA screening in the Reactome PI3K cascade pathway. There were very few differences in vertex degree between the groups, although genes detected by siRNA included those with the fewest connections.




Figure H.2: Synthetic lethality and centrality. The information centrality was compared (on a log-scale across genes detected by mtSLIPT and siRNA screening in the Reactome PI3K cascade pathway. Genes detected by mtSLIPT or siRNA did not have higher connectivity than genes not detected by either approach. The gene with the highest centrality was detected by siRNA.




Figure H.3: Synthetic lethality and PageRank. The PageRank centrality was compared (on a log-scale across genes detected by mtSLIPT and siRNA screening in the Reactome PI3K cascade pathway. Genes detected by siRNA had a more restricted range of centrality values than other genes not detected by either approach, although these groups also had fewer genes.

Table H.1: ANOVA for synthetic lethality and vertex degree

	DF	Sum Squares	Mean Squares	F-value	p-value
siRNA	1	15	15.50	0.0134	0.9084
mtSLIPT	1	196	195.94	0.1689	0.6825
siRNA×mtSLIPT	1	9	9.17	0.0079	0.9294

Analysis of variance for vertex degree against synthetic lethal detection approaches (with an interaction term)

Table H.2: ANOVA for synthetic lethality and information centrality

	DF	Sum Squares	Mean Squares	F-value	p-value
siRNA	1	0.000256	0.0002561	0.1851	0.6685
mtSLIPT	1	0.003225	0.0032247	2.3308	0.1318
siRNA×mtSLIPT	1	0.001238	0.0012385	0.8952	0.3476

Analysis of variance for information centrality against synthetic lethal detection approaches (with an interaction term)

Table H.3: ANOVA for synthetic lethality and PageRank centrality

	DF	Sum Squares	Mean Squares	F-value	p-value
siRNA	1	0.0002038	2.0385×10^{-4}	1.1423	0.2892
mtSLIPT	1	0.0000208	2.0752×10^{-5}	0.1163	0.7342
siRNA×mtSLIPT	1	0.0000137	1.3743×10^{-5}	0.0770	0.7823

Analysis of variance for PageRank centrality against synthetic lethal detection approaches (with an interaction term)

Appendix I

Information Centrality for Gene Essentiality

Network structure could be used to analyse gene function. This has been performed to investigate network properties of a network constructed from Reactome pathways (Croft *et al.*, 2014) imported via Pathway Commons with Paxtools (Cerami *et al.*, 2011; Demir *et al.*, 2013). Information centrality, which has been proposed as a measure of gene essentiality, was calculated as performed by Kranthi *et al.* (2013) using the efficiency and shortest path between each pair of nodes in the network before and after a node of interest is removed to test the importance of a node to network connectivity. Reactome contains substrates and cofactors in addition to genes or proteins. In support of centrality as a measure of essentiality, a number of nodes with the highest centrality (shown in Table I.1) were essential nutrients including Mg²⁺, Ca²⁺, Zn²⁺, and Fe. In addition, there were genes important in development of epithelial tissues and breast cancer such as *IL8*, *GATA3*, and *CTNNB1* detected with relatively high information centrality.

Table I.1: Information centrality for genes and molecules in the Reactome network

Node	Centrality
<i>ZNF473</i>	0.0510
Magnesium (Mg^{2+})	0.0082
<i>XBP1</i>	0.0053
Calcium (Ca^{2+})	0.0050
Zinc (Zn^{2+})	0.0048
Iron atom (Fe)	0.0041
<i>FMN</i>	0.0040
<i>AGT</i>	0.0037
<i>HSP90AA1</i>	0.0029
Phosphatidyl-L-serine	0.0029
<i>P2RX7</i>	0.0026
<i>PANX1</i>	0.0024
<i>NCAM1</i>	0.0022
<i>NUDT1</i>	0.0021
<i>PLAUR</i>	0.0020
<i>IL8</i>	0.0020
<i>HSPA8</i>	0.0019
<i>TYROBP</i>	0.0019
<i>CASP3</i>	0.0017
<i>GNAL</i>	0.0015
<i>CBLB</i>	0.0015
<i>HBB</i>	0.0014
<i>GATA4</i>	0.0013
<i>TGS1</i>	0.0013
<i>CTNNB1</i>	0.0012

Highest information centrality for genes (proteins), cofactors, and minerals in the Reactome network




Figure I.1: **Information centrality distribution.** Information centrality in the Reactome network for nodes, including genes/proteins and other biomolecules.

Appendix J

Pathway Structure for Mutation SLIPT




Figure J.1: **Synthetic lethality and heirarchy score in PI3K.** The hierarchical distance scores were similarly distributed across mtSLIPT and siRNA genes. Genes detected by both methods had a higher (downstream) median than either group.

Table J.1: ANOVA for synthetic lethality and PI3K hierarchy

	DF	Sum Squares	Mean Squares	F-value	p-value
siRNA	1	0.001	0.00070	0.0004	0.9841
mtSLIPT	1	0.007	0.0066	0.0040	0.9496
siRNA×mtSLIPT	1	3.906	3.9056	2.3829	0.1250

Analysis of variance for PI3K hierarchy score against synthetic lethal detection approaches (with an interaction term)




Figure J.2: **Hierarchy score in PI3K against synthetic lethality in PI3K.** The number of mtSLIPT and siRNA genes against the hierarchical distance scores showing no significant tendency for either method to either of the pathway upstream or downstream extremities.




Figure J.3: **Structure of synthetic lethality in PI3K.** The number of mtSLIPT and siRNA genes against the hierarchical distance scores showing no significant tendency for either method to either of the pathway upstream or downstream extremities. The number of mtSLIPT and siRNA genes upstream or downstream of each gene in the Reactome PI3K pathway were tested (by the χ^2 -test). These were plotted as a split jitter stripchart against the hierarchical distance scores showing no significant tendency for either method to either of the pathway upstream or downstream extremities.




Figure J.4: Structure of synthetic lethality resampling. A null distribution (10,000 iterations) of the siRNA genes upstream or downstream of mtSLIPT genes (shown by the difference) in the PI3K pathway. The observed events (red) were compared to the the distribution (violet) and were not significant. Genes detected by both methods were fixed for the distribution (blue). The genes detected by both approaches were used.

Table J.2: Resampling for pathway structure of synthetic lethal detection methods

Pathway	Graph		States		Observed				Permutation p-value	
	Nodes	Edges	mtSL	siRNA	Up	Down	Up-Down	Up/Down	Up-Down	Down-Up
PI3K Cascade	138	1495	42	25	131	123	8	1.065	0.4473	0.5466
PI3K/AKT Signalling in Cancer	275	12882	56	44	478	440	38	1.086	0.4163	0.5810
G_{αi} Signalling	292	22003	57	58	543	866	-323	0.627	0.9507	0.0488
GPCR downstream	1270	142071	218	160	7632	6500	1132	1.174	0.1707	0.8291
Elastic fibre formation	42	175	16	7	6	7	-1	0.857	0.5512	0.3681
Extracellular matrix	299	3677	81	29	313	347	-34	0.902	0.5762	0.4215
Formation of Fibrin	52	243	11	5	8	19	-11	0.421	0.7993	0.1800
Nonsense-Mediated Decay	103	102	56	2	0	0	0		0.197	0.1373
3'-UTR-mediated translational regulation	107	2860	56	1	52	1	51	52	0.1210	0.8751
Eukaryotic Translation Elongation	92	3746	57	0	0	0	0		0.4952	0.4892

Pathways in the Reactome network tested for structural relationships between mtSLIPT and siRNA genes by resampling. The raw p-value (computed without adjusting for multiple comparisons over pathways) is given for the difference in upstream and downstream paths from mtSLIPT to siRNA gene candidate partners of CDH1 with significant pathways highlighted in bold. Sampling was performed only in the target pathway and shortest paths were computed within it. Loops or paths in either direction that could not be resolved were excluded from the analysis. The gene detected by both mtSLIPT and siRNA (or resampling for them) were included in the analysis and the number of these were fixed to the number observed.

Appendix K

Performance of SLIPT and χ^2




Figure K.1: **Performance of χ^2 and SLIPT across quantiles.** (continued on next page)




Figure K.1: Performance of χ^2 and SLIPT across quantiles. Synthetic lethal detection with quantiles as in axis labels. The barplots have the same hue for each quantile (grey for correlation) and darker for χ^2 (and positive correlation). SLIPT and χ^2 performed similarly, peaking at $1/3$ -quantiles and converging to random (0.5). Negative correlation was higher than positive but not optimal quantiles for SLIPT or χ^2 . These findings were robust across different numbers of underlying synthetic lethal genes in 10,000 simulations of 100 genes and 1000 samples. SLIPT performed better than χ^2 for higher numbers of synthetic lethal genes and finer quantiles.




Figure K.2: **Performance of χ^2 and SLIPT across quantiles.** (continued on next page)




Figure K.2: Performance of χ^2 and SLIPT across quantiles. Synthetic lethal detection with quantiles as in axis labels. The line plots are coloured for SLIPT (red), χ^2 (blue) and correlation (grey), according to the legend. SLIPT and χ^2 performed similarly, peaking at $1/3$ -quantiles and converging to random (0.5). Negative correlation was higher than positive but not optimal quantiles for SLIPT or χ^2 . These findings were robust across different numbers of underlying synthetic lethal genes in 10,000 simulations of 100 genes and 1000 samples. SLIPT performed better than χ^2 for higher numbers of synthetic lethal genes and finer quantiles.




Figure K.3: **Performance of χ^2 and SLIPT across quantiles with more genes.**
 (continued on next page)




Figure K.3: Performance of χ^2 and SLIPT across quantiles with more genes.
 Synthetic lethal detection with quantiles as in axis labels. The line plots are coloured for SLIPT (red), χ^2 (blue) and correlation (grey), according to the legend. SLIPT and χ^2 performed similarly, peaking at $1/3$ -quantiles and converging to random (0.5). Negative correlation was higher than positive but not optimal quantiles for SLIPT or χ^2 . These findings were robust across different numbers of underlying synthetic lethal genes in 1000 simulations of 20,000 genes and 1000 samples. SLIPT performed better than χ^2 for higher numbers of synthetic lethal genes and finer quantiles.

K.1 Correlated Query Genes affects Specificity




Figure K.4: Performance of χ^2 and SLIPT across quantiles with query correlation. (continued on next page)




Figure K.4: Performance of χ^2 and SLIPT across quantiles with query correlation. Synthetic lethal detection with quantiles as in axis labels. The barplots have the same hue for each quantile (grey for correlation) and darker for χ^2 (and positive correlation). SLIPT and χ^2 performed similarly, peaking at $1/3$ -quantiles and converging to random (0.5). Negative correlation was higher than positive but not optimal quantiles for SLIPT or χ^2 . These findings were robust across different numbers of underlying synthetic lethal genes in 10,000 simulations of 100 genes (including 10 correlated with the query) and 1000 samples. SLIPT performed consistently better than χ^2 with positively correlated genes.




Figure K.5: **Performance of χ^2 and SLIPT across quantiles with query correlation.** (continued on next page)




Figure K.5: Performance of χ^2 and SLIPT across quantiles with query correlation. Synthetic lethal detection with quantiles as in axis labels. The line plots are coloured for SLIPT (red), χ^2 (blue) and correlation (grey), according to the legend. SLIPT and χ^2 performed similarly, peaking at $1/3$ -quantiles and converging to random (0.5). Negative correlation was higher than positive but not optimal quantiles for SLIPT or χ^2 . These findings were robust across different numbers of underlying synthetic lethal genes in 10,000 simulations of 100 genes (including 10 correlated with the query) and 1000 samples. SLIPT performed consistently better than χ^2 with positively correlated genes.




Figure K.6: **Performance of χ^2 and SLIPT across quantiles with query correlation and more genes.** (continued on next page)




Figure K.6: Performance of χ^2 and SLIPT across quantiles with query correlation and more genes. Synthetic lethal detection with quantiles as in axis labels. The line plots are coloured for SLIPT (red), χ^2 (blue) and correlation (grey), according to the legend. SLIPT and χ^2 performed similarly, peaking at $1/3$ -quantiles and converging to random (0.5). Negative correlation was higher than positive but not optimal quantiles for SLIPT or χ^2 . These findings were robust across different numbers of underlying synthetic lethal genes in 1000 simulations of 20,000 genes (including 1000 correlated with the query) and 1000 samples. SLIPT performed consistently better than χ^2 with positively correlated genes.

Appendix L

Simulations on Graph Structures




Figure L.1: Performance of simulations on a simple graph. Simulation of synthetic lethality was performed using a multivariate normal distribution from a converging graph. For each parameter, 10,000 simulations were used. Colours in Figure L.1b match Figure L.1d.

L.0.1 Simulations from Inhibiting Graph Structures






Figure L.2: Performance of simulations on an inhibiting graph. Simulation of synthetic lethality used a multivariate normal distribution from a converging graph. For each parameter, 10,000 simulations were used. Colours in Figure L.2b match Figure L.2d.



(a) Statistical evaluation


(b) ROC



(c) Graph Structure


(d) Statistical performance

Figure L.3: Performance of simulations on a constructed graph with inhibition.
 Simulation of synthetic lethality used a multivariate normal distribution from a pathway with only inhibitions. For each parameter, 10,000 simulations were used. Colours in Figure L.3b match Figure L.3d.



(a) Statistical evaluation

(b) ROC




(c) Graph Structure


(d) Statistical performance

Figure L.4: Performance of simulations on a constructed graph with inhibition.
 Simulation of synthetic lethality used a multivariate normal distribution from a pathway with a combination of inhibitions. For each parameter, 10,000 simulations were used. Colours in Figure L.4b match Figure L.4d.


L.1 Simulation across Graph Structures



(a) Activating Graph Structure (b) χ^2 distribution for “A” SL



(c) Gene “B” SL (d) Gene “C” SL



(e) Gene “D” SL (f) Gene “E” SL

Figure L.5: **Detection of synthetic lethality within a graph Structure.** (continued on next page)







Figure L.5: Detection of synthetic lethality within a graph structure. Each gene was designated to be synthetic lethal separately and the χ^2 value from SLIPT was computed for each gene across the graph. For each synthetic lethal gene (highlighted in the respective colours), the χ^2 values were computed in 100 simulations of datasets of 20,000 genes including the graph structure and 1000 samples. For each synthetic lethal gene, the adjacent genes in the network also had elevated test statistics.




(a) Inhibiting Graph Structure




(b) χ^2 distribution for "A" SL




(c) Gene "B" SL




(d) Gene "C" SL




(e) Gene "D" SL




(f) Gene "E" SL




(g) Gene "F" SL



(h) Gene "G" SL




(i) Gene "H" SL




(j) Gene "I" SL


Figure L.6: Detection of synthetic lethality within an inhibiting graph. Each gene was designated to be synthetic lethal separately and the χ^2 value from SLIPT was computed for each gene across the graph structure with inhibiting relationships. For each synthetic lethal gene (highlighted in the respective colours), the χ^2 values were computed in 100 simulations of datasets of 20,000 genes including the graph structure and 1000 samples. For each synthetic lethal gene, the adjacent genes exhibited lower χ^2 values with inhibiting relationships.




(a) Inhibiting Graph Structure




(b) χ^2 distribution for "A" SL




(c) Gene "B" SL




(d) Gene "C" SL




(e) Gene "D" SL




(f) Gene "E" SL




(g) Gene "F" SL



(h) Gene "G" SL



(i) Gene "H" SL



(j) Gene "I" SL

Figure L.7: Detection of synthetic lethality within an inhibiting graph. Each gene was designated to be synthetic lethal separately and the χ^2 value from SLIPT was computed for each gene across the graph structure with inhibiting and relationships. For each synthetic lethal gene (highlighted in the respective colours), the χ^2 values were computed in 100 simulations of datasets of 20,000 genes including the graph structure and 1000 samples.

L.2 Simulations from Complex Graph Structures






Figure L.8: **Performance of simulations on a branching graph.** Simulation of synthetic lethality used a multivariate normal distribution from a branching graph. For each parameter, 10,000 simulations were used. Colours in Figure L.8b match Figure L.8d.




(a) Statistical evaluation

(b) ROC




(c) Graph Structure




(d) Statistical performance

Figure L.9: Performance of simulations on a complex graph. Simulation of synthetic lethality used a multivariate normal distribution from a complex graph. For each parameter, 10,000 simulations were used. Colours in Figure L.9b match Figure L.9d.




(a) Statistical evaluation

(b) ROC



(c) Graph Structure



(d) Statistical performance

Figure L.10: Performance of simulations on a large graph. Simulation of synthetic lethality used a multivariate normal distribution from a large graph. For each parameter, 10,000 simulations were used. Colours in Figure L.10b match Figure L.10d.

L.2.1 Simulations from Complex Inhibiting Graphs






Figure L.11: Performance of simulations on a branching graph with inhibition.
 Simulation of synthetic lethality used a multivariate normal distribution from a branching graph with only inhibitions. For each parameter, 10,000 simulations were used. Colours in Figure L.11b match Figure L.11d.




(a) Statistical evaluation

(b) ROC




(c) Graph Structure




(d) Statistical performance

Figure L.12: Performance of simulations on a branching graph with inhibition.
 Simulation of synthetic lethality used a multivariate normal distribution from a branching graph with alternating inhibitions. For each parameter, 10,000 simulations were used. Colours in Figure L.12b match Figure L.12d.




(a) Statistical evaluation

(b) ROC




(c) Graph Structure




(d) Statistical performance

Figure L.13: Performance of simulations on a complex graph with inhibition.
 Simulation of synthetic lethality used a multivariate normal distribution from a complex graph with only inhibitions. For each parameter, 10,000 simulations were used. Colours in Figure L.13b match Figure L.13d.




(a) Statistical evaluation

(b) ROC




(c) Graph Structure




(d) Statistical performance

Figure L.14: Performance of simulations on a complex graph with inhibition.
 Simulation of synthetic lethality used a multivariate normal distribution from a complex graph with a combination of relationships. For each parameter, 10,000 simulations were used. Colours in Figure L.14b match Figure L.14d.




(a) Statistical evaluation

(b) ROC




(c) Graph Structure



(d) Statistical performance

Figure L.15: Performance of simulations on a large constructed graph with inhibition. Simulation of synthetic lethality used a multivariate normal distribution from a large graph with only inhibitions. For each parameter, 10,000 simulations were used. Colours in Figure L.15b match Figure L.15d.




(a) Statistical evaluation

(b) ROC



(c) Graph Structure



(d) Statistical performance

Figure L.16: Performance of simulations on a large constructed graph with inhibition. Simulation of synthetic lethality used a multivariate normal distribution from a large graph with alternating inhibitions. For each parameter, 10,000 simulations were used. Colours in Figure L.16b match Figure L.16d.

L.3 Simulations from Pathway Graph Structures




Figure L.17: Performance of simulations on the $G_{\alpha i}$ signalling pathway. Simulation of synthetic lethality used a multivariate normal distribution based on the Reactome $G_{\alpha i}$ signalling pathway. Performance of SLIPT was high across parameters for detecting synthetic lethality in the graph structure within a larger dataset. The performance decreased for a greater number of true positives to detect but the accuracy increased with a low false positive rate.




Figure L.18: Performance of simulations including the $G_{\alpha i}$ signalling pathway. Simulation of synthetic lethality used a multivariate normal distribution (without correlation structure apart from the Reactome $G_{\alpha i}$ signalling pathway. Performance of SLIPT was high across parameters for detecting synthetic lethality in the graph structure within a larger dataset. The sensitivity decreased for a greater number of true positives to detect but the specificity remained high with a low false positive rate.