

Contents

Glossary	xi
Acronyms	xiii
1 Introduction and Literature Review	1
1.1 Cancer Research in the Post-Genomic Era	1
1.1.1 Cancer is a Global Health Issue	2
1.1.1.1 The Genetics and Molecular Biology of Cancers	3
1.1.2 The Genomics Revolution in Cancer Research	3
1.1.2.1 High-Throughput Technologies	4
1.1.2.2 Bioinformatics and Genomic Data	5
1.1.3 Genomics Projects	5
1.1.3.1 The Cancer Genome Project	6
1.1.3.2 The Cancer Genome Atlas Project	6
1.1.4 Genomic Cancer Medicine	8
1.1.4.1 Cancer Genes and Driver Mutations	8
1.1.4.2 Precision Cancer Medicine	9
1.1.4.3 Molecular Diagnostics and Pan-Cancer Medicine	9
1.1.4.4 Targeted Therapeutics and Pharmacogenomics	10
1.1.5 Systems and Network Biology	11
1.2 Synthetic Lethal Cancer Medicine	12
1.2.1 Synthetic Lethal Genetic Interactions	12
1.2.2 Synthetic Lethal Concepts in Genetics	14
1.2.3 Synthetic Lethality in Model Systems	14
1.2.3.1 Synthetic Lethal Pathways and Networks	15
1.2.3.2 Evolution of Synthetic Lethality	15
1.2.4 Synthetic Lethality in Cancer	16
1.2.5 Clinical Impact of Synthetic Lethality in Cancer	18
1.2.6 High-throughput Screening for Synthetic Lethality	19
1.2.6.1 Synthetic Lethal Screens	21
1.2.7 Computational Prediction of Synthetic Lethality	22
1.2.7.1 Bioinformatics Approaches to Genetic Interactions	22
1.2.7.2 Comparative Genomics	23
1.2.7.3 Analysis and Modelling of Protein Data	26
1.2.7.4 Differential Gene Expression	28
1.2.7.5 Data Mining and Machine Learning	29

1.2.7.6	Mutual Exclusivity and Bimodality	31
1.2.7.7	Rationale for Further Development	33
1.3	E-cadherin as a Synthetic Lethal Target	33
1.3.1	The <i>CDH1</i> gene and its Biological Functions	33
1.3.2	Hereditary Diffuse Gastric (and Lobular Breast) Cancer	34
1.3.3	Cell Line Models of <i>CDH1</i> Null Mutations	35
1.4	Summary and Research Direction of Thesis	36
1.4.1	Thesis Aims	37
2	Methods and Resources	38
2.1	Bioinformatics Resources for Genomics Research	38
2.1.1	Public Data and Software Packages	38
2.1.1.1	Cancer Genome Atlas Data	39
2.1.1.2	Reactome and Annotation Data	40
2.2	Data Handling	40
2.2.1	Normalisation	40
2.2.2	Sample Triage	40
2.2.3	Metagenes and the Singular Value Decomposition	41
2.2.4	Candidate Triage and Integration with Screen Data	43
2.3	Techniques	43
2.3.1	Statistical Procedures and Tests	44
2.3.2	Gene Set Over-representation Analysis	45
2.3.3	Clustering	45
2.3.4	Heatmap	45
2.3.5	Modelling and Simulations	46
2.3.5.1	Receiver Operating Characteristic Curves	47
2.3.6	Resampling Analysis	47
2.4	Pathway Structure Methods	48
2.4.1	Network and Graph Analysis	48
2.4.2	Sourcing Graph Structure Data	49
2.4.3	Constructing Pathway Subgraphs	49
2.4.4	Network Analysis Metrics	50
2.5	Implementation	51
2.5.1	Computational Resources and Linux Utilities	51
2.5.2	R Language and Packages	52
2.5.3	High Performance and Parallel Computing	55
3	Methods Developed During Thesis	57
3.1	A Synthetic Lethal Detection Methodology	57
3.2	Synthetic Lethal Simulation and Modelling	59
3.2.1	A Model of Synthetic Lethality in Expression Data	60
3.2.2	Simulation Procedure	64
3.3	Detecting Simulated Synthetic Lethal Partners	67
3.3.1	Binomial Simulation of Synthetic Lethality	67
3.3.2	Multivariate Normal Simulation of Synthetic Lethality	69
3.3.2.1	Multivariate Normal Simulation with Correlated Genes	71

3.3.2.2	Specificity with Query-Correlated Pathways	79
3.4	Graph Structure Methods	81
3.4.1	Upstream and Downstream Gene Detection	81
3.4.1.1	Permutation Analysis for Statistical Significance	82
3.4.2	Simulating Gene Expression from Graph Structures	83
3.5	Customised Functions and Packages Developed	87
3.5.1	Synthetic Lethal Interaction Prediction Tool	87
3.5.2	Data Visualisation	88
3.5.3	Extensions to the iGraph Package	89
3.5.3.1	Sampling Simulated Data from Graph Structures	89
3.5.3.2	Plotting Directed Graph Structures	89
3.5.3.3	Computing Information Centrality	91
3.5.3.4	Testing Pathway Structure with Permutation Testing	91
3.5.3.5	Metapackage to Install iGraph Functions	92
4	Synthetic Lethal Analysis of Gene Expression Data	93
4.1	Synthetic Lethal Genes in Breast Cancer	94
4.1.1	Synthetic Lethal Pathways in Breast Cancer	95
4.1.2	Expression Profiles of Synthetic Lethal Partners	97
4.1.2.1	Subgroup Pathway Analysis	100
4.2	Comparing Synthetic Lethal Gene Candidates	102
4.2.1	Primary siRNA Screen Candidates	102
4.2.2	Comparison with Correlation	102
4.2.3	Comparison with Primary Screen Viability	105
4.2.4	Comparison with Secondary siRNA Screen Validation	107
4.2.5	Comparison to Primary Screen at Pathway Level	108
4.2.5.1	Resampling Genes for Pathway Enrichment	110
4.2.6	Integrating Synthetic Lethal Pathways and Screens	115
4.3	Synthetic Lethal Pathway Metagenes	116
4.4	Replication in Stomach Cancer	118
4.5	Discussion	119
4.5.1	Strengths of the SLIPT Methodology	119
4.5.2	Synthetic Lethal Pathways for E-cadherin	120
4.5.3	Replication and Validation	122
4.5.3.1	Integration with siRNA Screening	122
4.5.3.2	Replication across Tissues	123
4.6	Summary	123
5	Synthetic Lethal Pathway Structure	125
5.1	Synthetic Lethal Genes in Reactome Pathways	125
5.1.1	The PI3K/AKT Pathway	126
5.1.2	The Extracellular Matrix	128
5.1.3	G Protein Coupled Receptors	131
5.1.4	Gene Regulation and Translation	131
5.2	Network Analysis of Synthetic Lethal Genes	133
5.2.1	Gene Connectivity and Vertex Degree	134

5.2.2	Gene Importance and Centrality	135
5.2.2.1	Information Centrality	135
5.2.2.2	PageRank Centrality	137
5.3	Relationships between Synthetic Lethal Genes	138
5.3.1	Detecting Upstream or Downstream Synthetic Lethality	139
5.3.2	Resampling for Synthetic Lethal Pathway Structure	141
5.4	Discussion	143
5.5	Summary	145
6	Simulation and Modelling of Synthetic Lethal Pathways	147
6.1	Synthetic Lethal Detection Methods	148
6.1.1	Performance of SLIPT and χ^2 across Quantiles	148
6.1.1.1	Correlated Query Genes affects Specificity	152
6.1.2	Alternative Synthetic Lethal Detection Strategies	154
6.1.2.1	Correlation for Synthetic Lethal Detection	154
6.1.2.2	Testing for Bimodality with BiSEp	156
6.2	Simulations with Graph Structures	157
6.2.1	Performance over Graph Structures	158
6.2.1.1	Simple Graph Structures	158
6.2.1.2	Constructed Graph Structures	160
6.2.2	Performance with Inhibitions	163
6.2.3	Synthetic Lethality across Graph Structures	168
6.2.4	Performance within a Large Simulated Datasets	171
6.3	Simulations in More Complex Graph Structures	175
6.3.1	Simulations over Pathway-based Graphs	176
6.3.2	Pathway Structures in a Large Simulated Datasets	179
6.4	Discussion	182
6.4.1	Simulation Procedure	182
6.4.2	Comparing Methods with Simulated Data	183
6.4.3	Design and Performance of SLIPT	184
6.4.4	Simulations from Graph Structures	186
6.5	Summary	187
7	Discussion	188
7.1	Synthetic Lethality and <i>CDH1</i> Biology	188
7.1.1	Established Functions of <i>CDH1</i>	189
7.1.2	The Molecular Role of <i>CDH1</i> in Cancer	189
7.2	Significance	190
7.2.1	Synthetic Lethality in the Genomic Era	190
7.2.2	Clinical Interventions based on Synthetic Lethality	192
7.3	Future Directions	193
7.4	Conclusions	195
	Bibliography	197

A	Sample Quality	222
A.1	Sample Correlation	222
A.2	Replicate Samples in TCGA Breast Cancer Data	225
B	Software Used for Thesis	229
C	Mutation Analysis in Breast Cancer	238
C.1	Synthetic Lethal Genes and Pathways	238
C.2	Synthetic Lethal Expression Profiles	239
C.3	Comparison to Primary Screen	242
C.3.1	Resampling Analysis	244
C.4	Compare SLIPT genes	246
D	Metagene Analysis	248
D.1	Pathway Signature Expression	248
D.2	Synthetic Lethal Reactome Metagenes	252
E	Intrinsic Subtyping	253
F	Stomach Expression Analysis	255
F.1	Synthetic Lethal Genes and Pathways	255
F.2	Comparison to Primary Screen	259
F.2.1	Resampling Analysis	261
F.3	Metagene Analysis	263
G	Synthetic Lethal Genes in Pathways	264
H	Network Analysis for Mutation SLIPT	271
I	Pathway Structure for Mutation SLIPT	274
J	Performance of SLIPT and χ^2	276
J.1	Correlated Query Genes affects Specificity	282
K	Simulations on Graph Structures	288
K.0.1	Simulations from Inhibiting Graph Structures	289
K.1	Simulation across Graph Structures	292
K.2	Simulations from Complex Graph Structures	296
K.2.1	Simulations from Complex Inhibiting Graphs	299
K.3	Simulations from Pathway Graph Structures	305

List of Figures

1.1	Synthetic genetic interactions	13
1.2	Synthetic lethality in cancer	17
2.1	Read count density	42
2.2	Read count sample mean	42
3.1	Framework for synthetic lethal prediction	58
3.2	Synthetic lethal prediction adapted for mutation	59
3.3	A model of synthetic lethal gene expression	61
3.4	Modelling synthetic lethal gene expression	62
3.5	Synthetic lethality with multiple genes	63
3.6	Simulating gene function	65
3.7	Simulating synthetic lethal gene function	65
3.8	Simulating synthetic lethal gene expression	66
3.9	Performance of binomial simulations	68
3.10	Comparison of statistical performance	68
3.11	Performance of multivariate normal simulations	70
3.12	Simulating expression with correlated gene blocks	72
3.13	Simulating expression with correlated gene blocks	73
3.14	Synthetic lethal prediction across simulations	75
3.15	Performance with correlations	76
3.16	Comparison of statistical performance with correlation structure	77
3.17	Performance with query correlations	78
3.18	Statistical evaluation of directional criteria	79
3.19	Performance of directional criteria	80
3.20	Simulated graph structures	84
3.21	Simulating expression from a graph structure	85
3.22	Simulating expression from graph structure with inhibitions	86
3.23	Demonstration of violin plots with custom features	90
3.24	Demonstration of annotated heatmap	90
3.25	Simulating graph structures	91
4.1	Synthetic lethal expression profiles of analysed samples	98
4.2	Comparison of SLIPT with siRNA	103
4.3	Comparison of SLIPT and siRNA genes with correlation	103
4.4	Comparison of SLIPT and siRNA genes with correlation	105
4.5	Comparison of SLIPT and siRNA genes with screen viability	106

4.6	Comparison of SLIPT genes with siRNA screen viability	106
4.7	Resampled intersection of SLIPT and siRNA candidate genes	111
5.1	Synthetic lethality in the PI3K cascade	127
5.2	Synthetic lethality in Elastic Fibre Formation	129
5.3	Synthetic lethality in Fibrin Clot Formation	130
5.4	Synthetic lethality in the GPCRs	132
5.5	Synthetic lethality and vertex degree	134
5.6	Synthetic lethality and centrality	136
5.7	Synthetic lethality and PageRank	138
5.8	Structure of synthetic lethality resampling	140
6.1	Performance of χ^2 and SLIPT across quantiles	150
6.2	Performance of χ^2 and SLIPT across quantiles with more genes	151
6.3	Performance of χ^2 and SLIPT across quantiles with query correlation	152
6.4	Performance of χ^2 and SLIPT across quantiles with query correlation and more genes	153
6.5	Performance of negative correlation and SLIPT	155
6.6	Simple graph structures	158
6.7	Performance of simulations on a simple graph	159
6.8	Performance of simulations is similar in simple graphs	161
6.9	Performance of simulations on a pathway	162
6.10	Performance of simulations on a simple graph with inhibition	164
6.11	Performance is higher on a simple inhibiting graph	165
6.12	Performance of simulations on a constructed graph with inhibition	166
6.13	Performance is affected by inhibition in graphs	168
6.14	Detection of synthetic lethality within a graph structure	170
6.15	Performance of simulations including a simple graph	172
6.16	Performance on a simple graph improves with more genes	174
6.17	Performance on an inhibiting graph improves with more genes	175
6.18	Performance of simulations on the PI3K cascade	178
6.19	Performance of simulations including the PI3K cascade	180
6.20	Performance on pathways improves with more genes	181
A.1	Correlation profiles of removed samples	223
A.2	Correlation analysis and sample removal	224
A.3	Replicate excluded samples	225
A.4	Replicate samples with all remaining	226
A.5	Replicate samples with some excluded	227
C.1	Synthetic lethal expression profiles of analysed samples	240
C.2	Comparison of mtSLIPT to short interfering RNA (siRNA)	242
C.3	Compare mtSLIPT and siRNA genes with correlation	246
C.4	Compare mtSLIPT and siRNA genes with correlation	246
C.5	Compare mtSLIPT and siRNA genes with siRNA viability	247
D.1	Pathway metagene expression profiles	250

D.2	Expression profiles for estrogen receptor related genes	251
F.1	Synthetic lethal expression profiles of stomach samples	257
F.2	Comparison of SLIPT in stomach to siRNA	259
G.1	Synthetic lethality in the PI3K/AKT pathway	264
G.2	Synthetic lethality in the PI3K/AKT pathway in cancer	265
G.3	Synthetic lethality in the Extracellular Matrix	266
G.4	Synthetic lethality in the GPCR Downstream	267
G.5	Synthetic lethality in the Translation Elongation	268
G.6	Synthetic lethality in the Nonsense-mediated Decay	269
G.7	Synthetic lethality in the 3' UTR	270
H.1	Synthetic lethality and vertex degree	271
H.2	Synthetic lethality and centrality	272
H.3	Synthetic lethality and PageRank	272
I.1	Structure of synthetic lethality resampling	274
J.1	Performance of χ^2 and SLIPT across quantiles	276
J.2	Performance of χ^2 and SLIPT across quantiles	278
J.3	Performance of χ^2 and SLIPT across quantiles with more genes	280
J.4	Performance of χ^2 and SLIPT across quantiles with query correlation	282
J.5	Performance of χ^2 and SLIPT across quantiles with query correlation	284
J.6	Performance of χ^2 and SLIPT across quantiles with query correlation and more genes	286
K.1	Performance of simulations on a simple graph	288
K.2	Performance of simulations on an inhibiting graph	289
K.3	Performance of simulations on a constructed graph with inhibition	290
K.4	Performance of simulations on a constructed graph with inhibition	291
K.5	Detection of synthetic lethality within a graph structure	292
K.6	Detection of synthetic lethality within an inhibiting graph	294
K.7	Detection of synthetic lethality within an inhibiting graph	295
K.8	Performance of simulations on a branching graph	296
K.9	Performance of simulations on a complex graph	297
K.10	Performance of simulations on a large graph	298
K.11	Performance of simulations on a branching graph with inhibition	299
K.12	Performance of simulations on a branching graph with inhibition	300
K.13	Performance of simulations on a complex graph with inhibition	301
K.14	Performance of simulations on a complex graph with inhibition	302
K.15	Performance of simulations on a large constructed graph with inhibition	303
K.16	Performance of simulations on a large constructed graph with inhibition	304
K.17	Performance of simulations on the $G_{\alpha i}$ signalling pathway	305
K.18	Performance of simulations including the $G_{\alpha i}$ signalling pathway	306

List of Tables

1.1	Methods for predicting genetic interactions	23
1.2	Methods for predicting synthetic lethality in cancer	23
1.3	Methods used by Wu <i>et al.</i> (2014)	25
2.1	Excluded samples by batch and clinical characteristics.	41
2.2	Computers used during thesis	51
2.3	Linux utilities and applications used during thesis	52
2.4	R installations used during thesis	53
2.5	R Packages used during thesis	53
2.6	R packages developed during thesis	55
4.1	Candidate synthetic lethal gene partners of <i>CDH1</i> from SLIPT	95
4.2	Pathways for <i>CDH1</i> partners from SLIPT	96
4.3	Pathways for clusters of <i>CDH1</i> partners from SLIPT	101
4.4	ANOVA for synthetic lethality and correlation with <i>CDH1</i>	104
4.5	Comparison of Synthetic Lethal Interaction Prediction Tool (SLIPT) genes against secondary siRNA screen	108
4.6	Pathways for <i>CDH1</i> partners from SLIPT and siRNA	109
4.7	Pathways for <i>CDH1</i> partners from SLIPT	112
4.8	Pathways for <i>CDH1</i> partners from SLIPT and siRNA primary screen .	113
4.9	Examples of candidate metagenes synthetic lethal for <i>CDH1</i> from SLIPT	117
5.1	ANOVA for synthetic lethality and vertex degree	135
5.2	ANOVA for synthetic lethality and information centrality	136
5.3	ANOVA for synthetic lethality and PageRank centrality	137
5.4	Resampling for pathway structure of synthetic lethal detection methods	142
B.1	Complete list of R packages used during this thesis	229
C.1	Candidate synthetic lethal gene partners of <i>CDH1</i> from mtSLIPT . . .	238
C.2	Pathways for <i>CDH1</i> partners from mtSLIPT	239
C.3	Pathways for clusters of <i>CDH1</i> partners from mtSLIPT	241
C.4	Pathways for <i>CDH1</i> partners from mtSLIPT and siRNA	243
C.5	Pathways for <i>CDH1</i> partners from mtSLIPT	244
C.6	Pathways for <i>CDH1</i> partners from mtSLIPT and siRNA primary screen	245
D.1	Candidate synthetic lethal metagenes against <i>CDH1</i> from mtSLIPT . .	252

E.1	Comparison of intrinsic subtypes	253
F.1	Synthetic lethal gene partners of <i>CDH1</i> from SLIPT in stomach cancer	255
F.2	Pathways for <i>CDH1</i> partners from SLIPT in stomach cancer	256
F.3	Pathways for clusters of <i>CDH1</i> partners in stomach SLIPT	258
F.4	Pathways for <i>CDH1</i> partners from SLIPT and siRNA	260
F.5	Pathways for <i>CDH1</i> partners from SLIPT in stomach cancer	261
F.6	Pathways for <i>CDH1</i> partners from SLIPT in stomach and siRNA	262
F.7	Synthetic lethal metagenes against <i>CDH1</i> in stomach cancer	263
H.1	ANOVA for synthetic lethality and vertex degree	273
H.2	ANOVA for synthetic lethality and information centrality	273
H.3	ANOVA for synthetic lethality and PageRank centrality	273
I.1	Resampling for pathway structure of synthetic lethal detection methods	275

Glossary

bioinformatics	Statistical or computational approaches to biological data or research tools.
centrality	A network metric which identifies important vertices .
edge or link	A relationship connecting a pair of elements of a graph structure or network, may be weighted or directional.
essential	A gene which is required to be functional or expressed for a cell or organism to be viable, grow or develop.
gene expression	A measure of the relative expression of each gene from the mRNA extracted from (pooled) cells.
graph or network	A mathematical structure modelling or depicting the relationships between elements.
hub	A central or highly connected component of a network.
induced essentiality	A gene becoming essential to viability under certain conditions, including inactivation of a synthetic lethal partner.
information centrality	A network centrality metric which uses the impact of removing a vertex or node on connections in the network.
metagene	A consistent signal of expression for a collection of genes such as a biological pathway, derived from singular value decomposition.
mutation	A change in DNA sequence that disrupts gene function.

non-oncogene addiction	The dependence of a cancer cell on functioning non-mutant genes.
oncogene	A gene that potentially causes cancer, typically by over-expression or mutant gene variants.
oncogene addiction	The dependence of a cancer cell on a specific oncogenic pathway.
PageRank centrality	A network centrality metric which uses eigenvectors with a scaling factor (Brin and Page, 1998).
pathway	A series of biomolecules that produces a particular product or biological function.
scale-free	A property of a network which has a power law vertex degree distribution, that is several highly connected hub genes and many with very few connections.
shortest path	A path with the fewest possible edges which connects two particular vertices .
synthetic lethal	Genetic interactions where inactivation of multiple genes is inviable (or deleterious) which are viable if inactivated separately.
targeted therapy	Cancer treatment that specifically acts against a molecular target, in contrast to standard chemotherapy.
tumour suppressor	A gene potentially causes cancer, typically by disruption of functions which protect the cell from cancer.
vertex degree	A network metric of connectivity of vertices which uses the number of edges connected to each vertex or node .
vertex or node	An element of a graph structure or network.

Acronyms

AMP	Adenosine Monophosphate.
AMPK	AMP -activated Protein Kinase.
ANOVA	Analysis of Variance.
BioPAX	Biological Pathway Exchange.
BMP	Bone Morphogenic Protein.
CXCR	Chemokine Receptor.
EMT	Epithelial-Mesenchymal Transition.
GPCR	G Crotein Coupled Receptor.
JAK	Janus Kinase.
mtSLIPT	Synthetic Lethal Interaction Prediction Tool (against mutation).
NMD	Nonsense-Mediated Decay.
PDE	Phosphodiesterase.
PI3K	Phosphoinositide 3-kinase.
RGS	G-protein Signalling.
RHO	Ras Homolog Family.
RNA	Ribonucleic Acid.
siRNA	Short Interfering RNA.
SLIPT	Synthetic Lethal Interaction Prediction Tool.
TCGA	The Cancer Genome Atlas (genomics project).
TGF β	Transforming Growth Factor β .
UTR	Untranslated Region (of mRNA).

Chapter 5

Synthetic Lethal Pathway Structure

Having identified key [pathways](#) implicated in [synthetic lethal](#) genetic interactions with *CDH1* (in Chapter 4), these were investigated in this Chapter for the [synthetic lethal](#) genes within them, and for their relationships to [pathway](#) structure. This chapter will focus on the Reactome biological [pathways](#) detected across analyses in Chapter 4. Specifically, investigations were performed to determine whether [synthetic lethal](#) candidates, detected by [SLIPT](#) or [siRNA](#), exhibited differences with respect to metrics of [pathway](#) structure of network connectivity and importance (as described in Sections 2.4.4 and 3.5.3). The relationships between [synthetic lethal](#) candidates, detected by either approach, were also examined to determine whether [SLIPT](#) candidate genes were upstream or downstream [siRNA](#) candidate genes. These directional relationships were tested by resampling (as described in Sections 3.4.1 and 3.4.1.1) and comparisons to the [pathway](#) hierarchical score based on biological context (as derived in Section ??). Together these investigations into structural relationships demonstrate how a combination of network biology and statistical techniques can be performed with genes identified by a [bioinformatics](#) analysis.

5.1 Synthetic Lethal Genes in Reactome Pathways

The [graph](#) structure for Reactome [pathways](#) was obtained from Pathway Commons via [Biological Pathway eXchange \(BioPAX\)](#) (as described in Section 2.4.2). The [pathways](#) describe the (directional) relationships between biomolecules, including genes that encode proteins in biological [pathways](#). These relationships include cell signalling (e.g., kinase phosphorylation cascades), gene regulation (e.g., transcription factors, chromatin modifiers, [RNA](#) binding proteins), and metabolism (e.g., the product of an enzyme being the substrate of another). Together these relationships describe the

known functional [pathways](#) in a human cell with a reasonable resolution, from a curated database supported by publications documenting [pathway](#) relationships.

Pathway structures from the Reactome network (as described in Section 2.4.3) were used to derive the [graph](#) structure of each biological [pathway](#). The [synthetic lethal](#) candidate genes for notable [pathways](#) discussed in Chapter 4, including candidate [synthetic lethal pathways](#) of *CDH1*, were examined to show the [SLIPT](#) and [siRNA](#) candidates within these [pathways](#). The [synthetic lethal](#) genes considered here are those candidates detected by [SLIPT](#) (as described in Section 3.1) in The Cancer Genome Atlas (TCGA) breast cancer [expression](#) and [mutation](#) data (Koboldt *et al.*, 2012) in comparison to the candidate gene partners from the [siRNA](#) screening in breast cell lines (Telford *et al.*, 2015).

5.1.1 The PI3K/AKT Pathway

The [phosphoinositide 3-kinase \(PI3K\)](#) cascade signalling [pathway](#) is important in cancer because it is involved in mediating signals between the [G protein coupled receptors](#) and regulation of protein translation, which have both been strongly implicated to be [synthetic lethal pathways](#) with loss of *CDH1* function (Chapter 4). These [pathways](#) are all subject to dysregulation in cancer (Courtney *et al.*, 2010; Dorsam and Gutkind, 2007; Gao and Roux, 2015). Thus the PI3K cascade will be examined along with the most supported [synthetic lethal pathways](#) (as identified in Chapter 4).

The [phosphoinositide 3-kinase \(PI3K\)](#) [pathway](#) is well characterised and has an established direction of signal transduction from extracellular stimuli (and membrane bound receptors) to the inner mechanisms of the cell, namely, the regulation of protein translation. The production of proteins is necessary for the growth of the cell so it is reasonable to suggest that these processes may be subject to (non-[oncogene](#)) addiction in some cancer cells which rely upon them for sustained protein production and cell growth. This is also supported by the [oncogenes](#) *PIK3CA* and *AKT1* being involved with the PI3K cascade and the related PI3K/AKT [pathway](#), which may be subject to [oncogene addiction](#) when these proto-oncogenes are activated.

The [PI3K](#) cascade was not supported across [SLIPT](#) in TCGA breast [expression](#) data and the [siRNA](#) primary screen by over-representation (in Section 4.2.5) or resampling (in Section 4.2.5.1) but genes within this [pathway](#) were detectable by either approach (as shown in Figure 5.1). While few genes were identified by both approaches, they include genes that are highly connected in the PI3K cascade and are hubs to information transmission such as *FGF9*, *PDE3B*, and *PDE4A*. The key upstream genes *PIK3CA* and




Figure 5.1: **Synthetic lethality in the PI3K cascade.** The Reactome [PI3K](#) Cascade pathway with [synthetic lethal](#) candidates coloured as shown in the legend.

PIK3CG were detected by [siRNA](#) whereas the downstream *PIK3R1* and *AKT2* genes were detected by [SLIPT](#). Genes detected by either method were also prevalent in the [PI3K](#), [phosphodiesterase \(PDE\)](#), and [AMP-activated protein kinase \(AMPK\)](#) modules, in addition to the downstream translation factors and ribosomal genes (*EIF4B*, *EEF2K*, and *RPS6*). Together these suggest that there may be further structure between the [SLIPT](#) and [siRNA](#) candidate partners of *CDH1* in [pathways](#) as illustrated by [PI3K](#). As such, [pathway](#) structure will be investigated to detect differences in the upstream and downstream gene candidates detected by either method. Pathway structure may

account for the disparity between SLIPT and siRNA genes, even in pathways such as PI3K where they did not significantly intersect. For instance, SLIPT gene partners may be downstream of siRNA candidates rather than replicating them directly.

This disparity between SLIPT and siRNA gene candidates synthetic lethal partners of CDH1 (i.e., a high number of genes detected by either approach with few detected by both) was replicated in the related PI3K/AKT pathway and the “PI3K/AKT in cancer” pathway (shown in Appendix Figures G.1 and G.2). Many synthetic lethal candidates were at the upstream core of these pathway networks and the downstream extremities. It is particularly notable that many genes important in cell signalling and gene regulation were detected by either synthetic lethal detection approach. These include *AKT1*, *AKT2*, and *AKT3*, the Calmodulin signalling genes *CALM1* and *CAMK4*, and the forkhead family transcription factors *FOXO1* (a tumour suppressor) and *FOXO4* (an inhibitor of EMT).

5.1.2 The Extracellular Matrix

The extracellular pathways “elastic fibre formation” and “fibrin clot formation” (shown in Figures 5.2 and 5.3 respectively) were both supported across analyses (in Chapter 4). These pathways were identified by both SLIPT (for TCGA breast cancer) and siRNA gene candidates as they had significant over-representation and resampling analyses.

Particularly for elastic fibres (Figure 5.2), the vast majority of genes were detected by either approach, in addition to a significant proportion of genes detected by both approaches (as determined in Section 4.2.5). The genes detected by both approaches also appeared to have a non-random distribution in the network, with *TFGB1*, *ITGB8*, and *MFAP2* exhibiting high connectivity, and having a central role in their respective pathway modules. In addition to a structural role in the extracellular matrix and connective tissue (including the tumour microenvironment), these proteins including Furin, transforming growth factor β (TGF β), and the bone morphogenic proteins (BMPs), are also involved in responses to endocrine signals and interact with the cellular receptors for signalling pathways. Therefore it is plausible that *CDH1* deficient tumours will be subject to non-oncogene addiction to the extracellular environment and growth signals arising from this pathway. The pathway structure also indicated that the genes detected by siRNA (or by both approaches) may be downstream of those detected by SLIPT, in addition to whether connectivity or centrality is higher for synthetic lethal candidates than other genes in the pathway.




Figure 5.2: **Synthetic lethality in Elastic Fibre Formation.** The Reactome Elastic Fibre Formation [pathway](#) with [synthetic lethal](#) candidates coloured as shown in the legend.

Genes detected as [synthetic lethal](#) partners of *CDH1* by [SLIPT](#) or [siRNA](#) screening were also common in the Fibrin clot formation [pathway](#) (shown in Figure 5.3). This is consistent with the established pleiotropic role of *CDH1* in regulating fibrin clotting. It is also notable that the genes detected by either method appear to be highly connected such as *C1QBP*, *KNG1*, *F8*, *F10*, *F12*, *F13A*, and *PROC* (including many of the coagulation factors). [Synthetic lethal](#) candidates also include *SERPINE2* and *PRCP*, which only affect downstream genes, in addition to *PROCR* and *VWF*, which are only affected by upstream genes.




Figure 5.3: **Synthetic lethality in Fibrin Clot Formation.** The Reactome Fibrin Clot Formation [pathway](#) with [synthetic lethal](#) candidates coloured as shown in the legend.

Many of these genes are involved in the larger Extracellular Matrix [pathway](#) (shown in Appendix Figure [G.3](#)), including many of the [synthetic lethal](#) candidates discussed for elastic fibres. The number of [SLIPT](#) candidate genes outnumbers those identified by [siRNA](#), as expected from an isolated cell model. However, the endocrine response genes (e.g., *TGFB1* and *LTBP4*) which are potentially artifacts of the cell line growth process were replicated with [SLIPT](#) analysis in patient tumours (TCGA breast cancer data). There is also additional support for [synthetic lethal](#) genes (e.g., *ITGB2*, *MFAP2*, and *SPARC*) being highly connected networks hubs of the [pathway](#). The complexity

of the extracellular matrix [pathway](#) lends credence to the need for formal network analysis approaches to interpret the [pathway](#) structure of [synthetic lethal](#) candidates. Furthermore, statistical approaches are needed to determine whether the apparent structural relationships between [synthetic lethal](#) candidates could have occurred by chance

5.1.3 G Protein Coupled Receptors

G protein coupled receptor (GPCR) [pathways](#) are highly complex (as shown in Figure 5.4 and Appendix Figure G.4). Many of genes in these [pathways](#) were [synthetic lethal](#) candidates, detected by either SLIPT or siRNA screening, including genes frequently detected by both approaches, consistent with these [pathways](#) being supported by prior analyses (in Sections 4.2.5 and 4.2.5.1). [Synthetic lethal](#) candidates include the PDE and Calmodulin genes (as discussed in Section 5.1.3) in addition to others such as the regulators of G-protein signalling (RGS), chemokine receptors (CXCR), Janus kinase (JAK), and the Ras homolog family (RHO) genes. These are important regulatory signalling [pathways](#) necessary for cellular growth and cancer proliferation. Thus the GPCR [pathways](#) (and downstream PI3K/AKT signals) are a potentially actionable vulnerability against *CDH1* deficient cancers, particularly since many existing drug targets are in these signalling [pathways](#), some of which have been experimentally validated (Telford *et al.*, 2015). While statistically significant numbers of genes in GPCR [pathways](#) were detected by both approaches (in Sections 4.2.5 and 4.2.5.1), the complexity of GPCR networks (containing hundreds of genes) further support the needs for a rational network-based approach to the relationships between SLIPT and experimental candidates.

5.1.4 Gene Regulation and Translation

While very few [synthetic lethal](#) genes were detected in translational [pathways](#) in an experimental screen against *CDH1* (Telford *et al.*, 2015), these were highly over-represented in translational elongation (as shown in Appendix Figure G.5). These SLIPT genes include many ribosomal proteins and the regulatory “elongation factors” which may be subject to responses in the upstream signalling [pathways](#). This observation further indicates that [pathway](#) structure may be used to identify relationships between [synthetic lethal](#) candidates detected by SLIPT and siRNA. The computational approach with SLIPT may exhibit the ability to detect downstream genes in the core translational processes, which experimental screening did not identify. The experimental screening may similarly detect upstream regulatory genes less sensitive




Figure 5.4: **Synthetic lethality in the GPCRs.** The Reactome $G_{\alpha i}$ pathway with **synthetic lethal** candidates, coloured as shown in the legend.

to inactivation, that is, genes that are less likely to be indiscriminately lethal to both genotypes at high doses of inactivation.

Many of these SLIPT candidate genes are also among the nonsense-mediated decay (NMD) pathway (shown in Appendix Figure G.6) or 3' untranslated region (UTR) mediated translational regulation (shown in Appendix Figure G.7). While genes in these pathways were also supported by experimental screening with siRNA, there were differences in which genes were detected within the pathway structures. In particular, *UPF1* was detected in the siRNA screen and is the focal downstream gene for the entire NMD pathway showing that (in this case) siRNA genes are downstream effectors of those detected by SLIPT. 3' UTR mediated translational regulation has a similar structure with two modules connected solely by *RPL13A*, giving an example of SLIPT candidate genes with high connectivity, although there were many ribosomal proteins detected by SLIPT. However, the detection of *EIF3K*, a regulatory elongation factor (not essential to ribosomal function) was replicated across SLIPT and siRNA screening, while the majority of the elongation factors were not detected by either approach. Regulatory genes, being more amenable to experimental validation, also support further investigation into pathway structure. The SLIPT candidates may support experimental candidates in biological pathways by detecting downstream genes, which may not be detectable by experimental screening with high dose inhibitors. This difference between the approaches may explain the greater number of SLIPT candidate partners of *CDH1* than those experimentally identified.

5.2 Network Analysis of Synthetic Lethal Genes

To demonstrate the network properties of synthetic lethal candidates in a pathway, a network analysis was performed on the genes detected as synthetic lethal partners of *CDH1* with the SLIPT computational approach and the siRNA screen (Telford *et al.*, 2015) in $G_{\alpha i}$ signalling, a GPCR pathway. This pathway was used to demonstrate deeper network analysis approaches to synthetic lethal candidates within complex pathways, as it was supported across analyses (in Chapter 4), with significant over-representation in both SLIPT and siRNA screening, and the genes differed considerably between synthetic lethal detection methods (shown in Appendix Figures 5.4). These network metrics were used to measure whether the network properties differed between groups of genes detected by either or both approaches. These analyses serve to test both whether synthetic lethal gene candidates had higher connectivity or importance in

a network and whether either detection approach is biased towards genes with different network properties.

5.2.1 Gene Connectivity and Vertex Degree

Vertex degree (the number of connections) for each gene is a fundamental property of a network. The vast majority of genes had a relatively modest number of connections, each with only a few genes in the $G_{\alpha i}$ pathway (shown in Figure 5.5) having pathway relationships with a high number of genes, consistent with the *scale-free* property of biological networks (Barabási and Oltvai, 2004). The number of connections was similar between gene groups (by *synthetic lethal* detection). Genes detected by *siRNA* included those with the fewest connections, despite there being fewer genes that were detected by either approach. There was no statistically significant effect of either computational or experimental *synthetic lethal* detection method on *vertex* degree, as determined by *analysis of variance* (ANOVA) (shown by Table 5.1).




Figure 5.5: **Synthetic lethality and vertex degree.** The number of connected genes (*vertex degree*) was compared (on a log-scale) across genes detected by *SLIPT* and *siRNA* screening in the Reactome $G_{\alpha i}$ cascade pathway. There were no differences in *vertex* degree between the groups (shown in Table 5.1), although genes detected by *siRNA* included those with the fewest connections.

Table 5.1: ANOVA for synthetic lethality and vertex degree

	DF	Sum Squares	Mean Squares	F-value	p-value
siRNA	1	21	20.8	0.0030	0.9561
SLIPT	1	16215	16215	2.3722	0.1246
siRNA×SLIPT	1	17	17	0.0025	0.9603

Analysis of variance for **vertex** degree against **synthetic lethal** detection approaches (with an interaction term)

The results for the $G_{\alpha i}$ **pathway** were very similar when testing **synthetic lethality** against *CDH1* **mutation** (mtSLIPT). In either case, there was no significant evidence that SLIPT or mtSLIPT-specific genes had higher connectivity than those detected by siRNA screening (shown in Appendix Figure H.1 and Appendix Table H.1). Thus **synthetic lethal** detection does not discriminate among genes by their connectivity in this **pathway** network, nor is either approach constrained to detecting highly connected genes. Both approaches have been demonstrated to detect genes with many and very few connections in the $G_{\alpha i}$ signalling **pathway**.

5.2.2 Gene Importance and Centrality

5.2.2.1 Information Centrality

Information centrality is a measure of the importance of **nodes** in a network in terms of how vital they are to the transmission of information throughout the network. This applies well to biological **pathways**, particularly gene regulation and cell signalling. The **nodes** with the highest **information centrality** are not necessarily the most connected, as they may also include **nodes** that pass signals between highly connected network hubs. **Information centrality** therefore provides a distinct metric for the connectivity of a gene in a **pathway**, which has the added benefit of being directly related to the disruption of **pathway** function were it to be inactivated or removed. **Information centrality** has also been suggested to be indicative of the **essentiality** of genes or proteins (Kranthi *et al.*, 2013).

Within the $G_{\alpha i}$ **pathway** (shown in Figure 5.6), the **information centrality** across gene groups detected by either **synthetic lethal** approach did not differ significantly (shown by Table 5.2). Genes detected by SLIPT span the complete range of **PageRank centrality** values for this **pathway**. These findings were replicated (shown in Appendix Figure H.2 and Appendix Table H.2). Thus neither method was unable to detect

synthetic lethal genes in the $G_{\alpha i}$ pathway with particular centrality constraints but they were also not detecting genes with higher centrality than expected by chance.




Figure 5.6: **Synthetic lethality and centrality.** The information centrality was compared (on a log-scale) across genes detected by SLIPT and siRNA screening in the Reactome $G_{\alpha i}$ pathway. Genes detected by SLIPT or siRNA did not have higher centrality than other genes (shown in Table 5.2). Genes detected by SLIPT spanned the range of centrality values.

Table 5.2: ANOVA for synthetic lethality and information centrality

	DF	Sum Squares	Mean Squares	F-value	p-value
siRNA	1	0.00000000	2.7000×10^{-9}	0.0016	0.96783
SLIPT	1	0.00000548	5.4831×10^{-6}	3.3253	0.06926
siRNA×SLIPT	1	0.00000002	1.8800×10^{-8}	0.0114	0.91511

Analysis of variance for information centrality against synthetic lethal detection approaches (with an interaction term)

5.2.2.2 PageRank Centrality

PageRank centrality is another network analysis procedure to infer a hierarchy of gene importance from a network using connections and structure (Brin and Page, 1998). In contrast to the information centrality approach of removing nodes, PageRank uses the eigenvalue properties of the adjacency matrix to rank genes according to the number of connections and paths they are involved in.

This distinction is immediately clear within the $G_{\alpha i}$ pathway (shown in Figure 5.7), which differs considerably from the information centrality scores (as shown in Figure 5.6). Genes detected by either synthetic lethal approach did not include those with the highest PageRank centrality. There was a significant association between genes detected by SLIPT (which had a lower median) with PageRank centrality (shown by Table 5.3).

The genes detected by SLIPT span the range of centrality values of siRNA showing that both approaches were capable of detecting genes of moderately high centrality (as shown for information centrality) and that the lower centrality of SLIPT candidates in the $G_{\alpha i}$ pathway may be due to synthetic lethal partners being less critical to the pathway, rather than a limitation of the methodology. While it is expected that some synthetic lethal genes will be important to the function of the pathway, it is possible that genes with high centrality were avoided if they are essential to cellular viability.

There was not a significant association between siRNA candidates and PageRank centrality. The significant result for SLIPT was not replicated when testing synthetic lethality against *CDH1* mutation (shown in Appendix Figure H.3 and Appendix Table H.3). However, this may be due to fewer genes being detected by mtSLIPT and siRNA.

Table 5.3: ANOVA for synthetic lethality and PageRank centrality

	DF	Sum Squares	Mean Squares	F-value	p-value
siRNA	1	0.0001059	1.0589×10^{-4}	2.1021	0.14818
SLIPT	1	0.0002881	2.8808×10^{-4}	5.7188	0.01743
siRNA×SLIPT	1	0.0000477	4.7704×10^{-5}	0.9470	0.33131

Analysis of variance for PageRank centrality against synthetic lethal detection approaches (with an interaction term)




Figure 5.7: **Synthetic lethality and PageRank.** The [PageRank centrality](#) was compared (on a log-scale) across genes detected by [mtSLIPT](#) and [siRNA](#) screening in the Reactome $G_{\alpha i}$ [pathway](#). Genes detected by with either [synthetic lethal](#) detection approach had a more restricted range of [centrality](#) values but only [SLIPT](#) genes had a significant association with [centrality](#) (shown in Table 5.3).

5.3 Relationships between Synthetic Lethal Genes

The network analyses so far have tested whether [synthetic lethal](#) candidate genes were more connected or important within a [pathway](#) structure, such as the $G_{\alpha i}$ [pathway](#). However these metrics do not ascertain whether there were relationships between [SLIPT](#) and [siRNA](#) candidate partners of *CDH1*. In particular, it is plausible that they may be upstream or downstream of one and other within a [pathway](#).

The direction of a biological [pathway](#) is important, particularly those involved in cell signalling which respond to extracellular stimuli and transmit these signals via intermediary proteins to regulate core functions and responses of the cell. These [pathways](#) regulate process such as [gene expression](#) and protein translation, which are important in the proliferation of cancers (Gao and Roux, 2015). Therefore it is important to determine which [synthetic lethal](#) candidates were upstream or downstream in the con-

text of a biological [pathway](#). In particular, [pathway](#) structure may be used to identify relationships between [SLIPT](#) and [siRNA](#) gene candidates.

A [pathway](#) structure method was devised to use [network](#) structures to identify directional relationships between individual [SLIPT](#) and [siRNA](#) genes. This [pathway](#) structure methodology was applied (as described in Section 3.4.1) to detect the direction of [shortest paths](#) between [SLIPT](#) and [siRNA](#) gene candidates. This is used to demonstrate the methodology on the [PI3K](#) and $G_{\alpha i}$ [pathways](#), to develop a statistical test for [pathway](#) structure between [SLIPT](#) and [siRNA](#) gene candidates using resampling (as described in Section 3.4.1.1), and to apply this test for [pathway](#) structure among [synthetic lethal](#) gene candidates to the [pathways](#) identified in Chapter 4 and discussed in Section 5.1.

5.3.1 Detecting Upstream or Downstream Synthetic Lethality

Shortest paths in a [pathway](#) network were used to devise a strategy to detect [pathway](#) structure between [SLIPT](#) and [siRNA](#) gene candidates partners of *CDH1* (as described in Section 3.4.1). Thus we can determine whether individual [SLIPT](#) genes have upstream or downstream [siRNA](#) candidates (scored as “up” or “down” events respectively). This procedure enables the detection of directional relationships between [SLIPT](#) and [siRNA](#) gene candidates (e.g., if genes detected by [siRNA](#) are more likely to be downstream of genes detected by [SLIPT](#) in the same pathway).

The total number of gene candidate pairs in either direction can be compared within a [pathway](#) network to assess the overall directional relationships in a [pathway](#). This directionality is detectable by the difference between the number of [SLIPT](#) candidate genes with upstream and downstream [siRNA](#) gene partners. However, this measure alone is not sufficient to determine whether there is evidence of [pathway](#) structure between [SLIPT](#) and [siRNA](#) gene candidates partners of *CDH1* in a [pathway](#) network. Nevertheless, it does serve to measure the magnitude (and direction) of the consensus of directional relationships (upstream and downstream) between [SLIPT](#) and [siRNA](#) gene candidates partners. This measure of [pathway](#) structure can be used for testing for statistical significance of [pathway](#) structure by resampling, using a permutation procedure to test whether these relationships are detectable among randomly selected gene groups rather than the detected [SLIPT](#) and [siRNA](#) gene candidates partners (as described in Sections 2.3.6 and 3.4.1.1).

This resampling procedure was performed for the $G_{\alpha i}$ and [PI3K pathways](#) to generate a null distribution for the difference in the number of “up events” and “down




Figure 5.8: **Structure of synthetic lethality resampling.** A null distribution with 10,000 iterations of the number of **siRNA** genes upstream or downstream of **SLIPT** genes (depicted as the difference of these) in each **pathway**. To assess significance, the observed events (with **shortest paths**) were compared to the 90% and 95% intervals for the null distribution (shown in blue). Genes detected by both methods were not fixed to the same number as observed for the alternative null distribution (shown in red), although the significance of the observed number of events (red) was changed in either case. The genes detected by both approaches were included in computing the number of **shortest paths** (in either direction) between **SLIPT** and **siRNA** genes. The permutations show (a) a significant **pathway** relationship for $G_{\alpha i}$ signalling and (b) and non-significant relationship for the **PI3K** cascade.

events” for these **pathway** structures (as shown in Figures 5.1 and 5.4). The resulting null distributions (as shown in Figure 5.8) were used to detect whether genes detected by **SLIPT** had significantly more upstream or downstream **siRNA** candidates in either **pathway**. It can be seen that **siRNA** genes were significantly downstream of **SLIPT** candidate genes by resampling for the $G_{\alpha i}$ signalling **pathway** (Figure 5.8a). This demonstrates that **pathway** relationships can be detected between **synthetic lethal** candidates by this procedure and that **siRNA** genes were downstream of gene detected by **SLIPT** in an example of GPCR signalling expanding on support for **synthetic lethality** in this **pathway** (as shown in Chapter 4). These structural relationships may also account for why each the computational and experimental approaches did not detect many of the same specific genes because they are detecting different parts of the **pathway**.

In contrast, there was not significant evidence of such **pathway** structure between **siRNA** and **SLIPT** candidate genes when resampling within the **PI3K** cascade **pathway** (as shown in Figure 5.8b). This indicates that such relationships may be **pathway** spe-

cific rather than a general property of these [synthetic lethal](#) detection methods. These results were robustly reproducible, with similar findings (as shown in Appendix Figure I.1) for each [pathway](#) when testing for [synthetic lethality](#) against *CDH1* mutation (mtSLIPT).

The number of genes detected by both approaches was fixed (to the number observed) for deriving p-values for pathway relationships (as described in Section 3.4.1.1). These genes were included in the analysis because they can be disproportionately upstream (or downstream) of more [siRNA](#) genes than [SLIPT](#) genes, which may lead to them having different proportions of genes detected by either approach upstream (or downstream) of them. However, allowing the number of jointly detected genes to vary during resampling (as shown in Figure 5.8 and Appendix Figure I.1) or excluding these jointly detected genes did not alter the findings of this approach. Furthermore, expanding the range of [shortest paths](#) to consider [links](#) in related [pathways](#) (using the “meta-[pathways](#)” constructed in Section 2.4.3) also had little effect on the null distribution generated, despite increasing the computational complexity of the procedure.

5.3.2 Resampling for Synthetic Lethal Pathway Structure

The permutation procedure (as described in Section 3.4.1.1) that was performed in Section 5.3.1 for the $G_{\alpha i}$ and [PI3K pathways](#) was also applied to other [pathways](#) identified in Chapter 4 and discussed in Section 5.1. In addition to the cell signalling [pathways](#) ([PI3K/AKT](#) and [GCPRs](#) demonstrated in Section 5.3.1), the [pathways](#) tested include extracellular matrix (with constituent elastic fibre and fibrin [pathways](#)), and translational [pathways](#) (with [NMD](#) and 3'UTR regulation).

The resampling results across these [pathways](#) (as shown in Table 5.4) had limited support for association between [pathway](#) structure and detection of [synthetic lethal](#) genes, with the majority of these being non-significant (as shown for [PI3K](#) in Figure 5.8b), with the exception of $G_{\alpha i}$ signalling (as shown in Figure 5.8a). However, the exact distribution for these [pathways](#) will differ depending on their structure, the number of genes they contain, and the proportion of [synthetic lethal](#) candidates among them (including a higher frequency of genes detected by both methods for the [pathways](#) identified in Section 4.2.5.1). This resampling is therefore an appropriate procedure to use to detect structural relationships across [pathways](#) as it does not assume an underlying test statistic distribution.

Pathway structure was supported for the [NMD pathway](#) (which is consistent with [siRNA](#) being downstream in Appendix Figure G.6). However, this observation rests

upon a single gene and was not replicated when testing **synthetic lethality** (**mtSLIPT**) against *CDH1* **mutation** (as shown in Appendix Table I.1) nor was it supported by the related 3'UTR regulation and translational elongation **pathways**.

Table 5.4: Resampling for **pathway** structure of **synthetic lethal** detection methods

Pathway	Graph		Candidates		Observed				Permutation p-value		p-value (FDR)
	Nodes	Edges	SLIPT	siRNA	Up ¹	Down ²	Up-Down	Up/Down	Up-Down	Down-Up	Down-Up
PI3K Cascade	138	1495	38	25	122	128	-6	0.953	0.5326	0.4606	0.6734
PI3K/AKT Signalling in Cancer	275	12882	98	44	779	679	100	1.147	0.3255	0.6734	0.6734
G_{αi} Signalling	292	22003	95	58	836	1546	-710	0.541	0.9971	0.0029	0.0145
GPCR downstream	1270	142071	312	160	9755	9261	494	1.053	0.3692	0.6305	0.6734
Elastic fibre formation	42	175	24	7	1	2	-1	0.500	0.5461	0.3865	0.6734
Extracellular matrix	299	3677	127	29	547	455	92	1.202	0.3351	0.6636	0.6734
Formation of Fibrin	52	243	18	5	12	17	-5	0.706	0.6198	0.3564	0.6734
Nonsense-Mediated Decay	103	102	74	2	0	74	-74	0	1.0000	< 0.0001	< 0.0010
3' -UTR-mediated translational regulation	107	2860	77	1	0	0	0		0.4902	0.5027	0.6734
Eukaryotic Translation Elongation	92	3746	76	0	0	0	0		0.4943	0.4933	0.6734

Pathways in the Reactome network tested for structural relationships between **SLIPT** and **siRNA** genes by resampling. The raw p-value (computed without adjusting for multiple comparisons over **pathways**) is given for the difference in upstream and downstream paths from **SLIPT** to **siRNA** gene candidate partners of *CDH1* with significant **pathways** highlighted in bold. Sampling was performed only in the target **pathway** and **shortest paths** were computed within it. Loops or paths in either direction that could not be resolved were excluded from the analysis. The genes detected by both **SLIPT** and **siRNA** (or resampling for them) were included in the analysis and the number of these were fixed to the number observed.

¹ The number of paths where the **siRNA** candidate was upstream of a **SLIPT** candidate

² The number of paths where the **siRNA** candidate was downstream of a **SLIPT** candidate

There does not appear to be a consensus on the directionality of **SLIPT** and **siRNA** candidates across **pathways** as distinct **pathways** showed stronger tendency for **siRNA** genes to be either upstream or downstream. Even related **pathways** such as **PI3K** and **PI3K/AKT** signalling showed directional events in opposite directions. The strongest **pathway** (among those tested) with support for directional **pathways** structure is **G_{αi} signalling** (as shown in Figure 5.8a). In contrast to the other **pathways** **G_{αi} signalling** showed significant downstream **siRNA** genes for **SLIPT** from a large number of **shortest paths** (in Table 5.4). This would indicate that **SLIPT** detects upstream regulators of genes experimentally validated by **siRNA** in this **pathway**. This result was **pathway** was also the strongest result in **mtSLIPT** results (Appendix Table I.1), although it was not significant after adjusting for multiple testing in this case.

There is insufficient evidence to determine whether there is **pathway** structure, that genes were detected upstream or downstream by either method, between the **SLIPT** and **siRNA** candidates in many of the **synthetic lethal pathways** (identified in Chapter 4). In particular, directional structure among **synthetic lethal** candidates for *CDH1* was not strongly supported in most of the signalling **pathways** (with the exception of **G_{αi} signalling**) upon which the rationale for **pathway** structure hypotheses were based. While there is statistically significant over-representation of many of these **pathways** in genes detected by both **SLIPT** and **siRNA** (as described in Chapter 4),

many of these did not show relationships with respect to [pathway](#) structure. Despite the design of a robust resampling approach to test relationships between gene groups, the detection of structural relationships between [SLIPT](#) and [siRNA](#) gene candidates did not generalise across [pathways](#) (and was specific to a few). Such structural relationships may apply more broadly to gene networks as different biological [pathways](#) were more over-represented among [SLIPT](#) and [siRNA](#) gene candidates. Furthermore, [pathway](#) structure did not account for the discrepancy between [SLIPT](#) and [siRNA](#) gene candidates which did not significantly intersect, such as the [PI3K](#) cascade.

5.4 Discussion

These investigations used a functional [pathway](#) network that encapsulates protein complexes and functional modules. The Reactome network uses curated, experimentally identified [pathways](#) to determine relationships between genes and does not have the limitation of relying solely on protein binding or text-mining which are prone to false positives (Croft *et al.*, 2014). While it is not documented whether these relationships are activating or inhibitory, the Reactome network (Croft *et al.*, 2014) is sufficient to test [pathway](#) relationships with directional information.

Synthetic lethal genes and [pathways](#) (for *CDH1* loss in cancer) were identified across [gene expression](#) and [mutation](#) datasets in Chapter 4. The investigations presented here extend those findings to consider [synthetic lethal](#) gene candidates within [pathway](#) structures, including exploring whether the discrepancy between individual [SLIPT](#) and [siRNA](#) candidate genes can be accounted for within a [synthetic lethal pathway](#). Pathways with replicated [synthetic lethal](#) genes across these detection methods, breast and stomach cancer data, were investigated, including [pathways](#) from the extracellular microenvironment to core translational [pathways](#) and the signalling [pathways](#) between them.

Examining [synthetic lethal](#) gene candidates in the context of [pathway](#) structures may also provide additional mechanisms by which the function of particular genes is subject to [induced essentiality](#) and support for them belonging to a [synthetic lethal pathway](#). Gene candidates with characterised functions important to cellular viability are ideal for triage of targets specific to *CDH1* deficient tumours and for further experimental validation in preclinical models. This chapter presents computational methods to use [pathway](#) structure in an attempt to detect genes with importance in a [pathway](#) and reconcile the differences between [SLIPT](#) and [siRNA](#) candidate genes with [pathway](#) relationships (e.g., one group being downstream of the other).

Many genes were detected by either **SLIPT** or **siRNA**. The differences between these computational and experimental screening approaches could feasibly lead to differences in which genes within a **synthetic lethal pathway** are identified. Genes detected by **synthetic lethal** detection strategies included those of biological importance within **synthetic lethal pathways**, those which are actionable drug targets, and those with functional implications for the biological growth mechanisms or vulnerabilities of *CDH1* deficient tumours. It appeared that genes detected by both approaches were highly connected (or of importance) in the **network** structure of some **pathways**, and that there may be some structure with **SLIPT** and **siRNA** candidates tending to appear upstream or downstream of each other.

The complexity of biological **pathways** meant that relationships between gene candidates were difficult to discern without formal mathematical and computational approaches, and thus these were used to analyse large biological networks. Network analysis techniques were applied to formalise and quantify the connectivity and importance (centrality) of genes within **pathways** (using $G_{\alpha i}$ signalling as an example). However, these network techniques were unable to identify distinct differences in many of the network properties of genes between those detected as **synthetic lethal** candidates by computational or experimental methods. These network metrics support the application of **synthetic lethal** detection across **pathways** (and the findings using **pathways** as gene sets in Chapter 4) as neither **synthetic lethal** detection approach was pre-disposed towards genes of higher importance or connectivity and neither approach was insensitive to genes of lower importance or connectivity. **SLIPT** did not detect genes with a significantly more crucial role in the $G_{\alpha i}$ **pathway**, as inferred by **pathway** connectivity and **centrality** measures. However, **SLIPT** genes had significantly lower centrality in the $G_{\alpha i}$ **pathway** by **PageRank centrality** (as shown in Section 5.2.2.2), and so the highest scoring genes may be too **essential** to cellular viability to be **synthetic lethal**.

A measure of **pathway** structure between individual **SLIPT** and **siRNA** candidate genes within a **pathway** was devised using the direction of **shortest paths** in a directed **graph** structure. This is amenable to detecting the consensus directionality of the **pathway** across pairs of genes detected by either method. The **pathway** structure methodology developed here is generally applicable to comparison of **node** groups (which may intersect), including genes in biological **pathways** and their detection by different methodologies. While the **pathway** structure measure alone is not able to detect structural relationships between gene groups (e.g., **SLIPT** and **siRNA** gene candidates), it is

amenable to resampling to determine whether these relationships are statistically significant. This approach successfully detected a statistically robust relationship between SLIPT and siRNA candidate genes on the $G_{\alpha i}$ signalling pathway, despite there being few differences between these genes with respect to network metrics of connectivity or centrality.

5.5 Summary

Together these analyses of biological pathways, network metrics, and statistical procedures devised specifically for this purpose were applied to Reactome pathway structures to test whether structural relationships existed between synthetic lethal candidates. Of particular interest was whether these relationships related to the differences between the computational (SLIPT) and experimental (siRNA) synthetic lethal candidate partners of *CDH1* (in the pathways discussed in Chapter 4).

While biologically relevant relationships were observed in specific pathways, there were few detectable structural relationships between SLIPT and siRNA gene candidates, apart from structural relationships specific to $G_{\alpha i}$ signalling. In this pathway, synthetic lethal candidates did not exhibit significant differences in network connectivity or centrality measures. These network analyses were also unable to ascertain whether the candidates detected by either method stratified into upstream and downstream genes on the pathway.

A statistical resampling procedure was applied to shortest path analysis to test whether pairs of SLIPT and siRNA gene candidates were more likely to be upstream or downstream of each other. This approach did not detect many structural relationships in the synthetic lethal pathways identified in Chapter 4. Overall, support for pathway structure between SLIPT and siRNA gene candidates was weak and the direction was inconsistent across pathways. Therefore pathway structure does not appear to generally account for the differences between the SLIPT and siRNA gene candidates, although it may apply in specific pathways as demonstrated with $G_{\alpha i}$ signalling. It was possible to detect some pathway relationships between candidate genes in synthetic lethal pathways, in addition to the significantly over-represented genes shared between SLIPT and siRNA (as identified in Chapter 4).

Furthermore, the resampling procedure demonstrated in this chapter is more widely applicable to gene states in network structures and may be of further utility in the analysis of biological pathways or networks. This approach was able to quantify structural relationships that were otherwise difficult to interpret and to conclusively exclude many

potential relationships. In this respect, the network resampling methodology may also be applicable to triage of [therapeutic targets](#) for experimental validation.

Bibliography

- Aarts, M., Bajrami, I., Herrera-Abreu, M.T., Elliott, R., Brough, R., Ashworth, A., Lord, C.J., and Turner, N.C. (2015) Functional genetic screen identifies increased sensitivity to weel inhibition in cells with defects in fanconi anemia and hr pathways. *Mol Cancer Ther*, **14**(4): 865–76.
- Abeshouse, A., Ahn, J., Akbani, R., Ally, A., Amin, S., Andry, C.D., Annala, M., Aprikian, A., Armenia, J., Arora, A., *et al.* (2015) The Molecular Taxonomy of Primary Prostate Cancer. *Cell*, **163**(4): 1011–1025.
- Adler, D. (2005) *vioplot: Violin plot*. R package version 0.2.
- Akbani, R., Akdemir, K.C., Aksoy, B.A., Albert, M., Ally, A., Amin, S.B., Arachchi, H., Arora, A., Auman, J.T., Ayala, B., *et al.* (2015) Genomic Classification of Cutaneous Melanoma. *Cell*, **161**(7): 1681–1696.
- Akobeng, A.K. (2007) Understanding diagnostic tests 3: receiver operating characteristic curves. *Acta Pdiatrica*, **96**(5): 644–647.
- American Cancer Society (2017) Genetics and cancer. <https://www.cancer.org/cancer/cancer-causes/genetics.html>. Accessed: 22/03/2017.
- Anjomshoaa, A., Lin, Y.H., Black, M.A., McCall, J.L., Humar, B., Song, S., Fukuzawa, R., Yoon, H.S., Holzmann, B., Friederichs, J., *et al.* (2008) Reduced expression of a gene proliferation signature is associated with enhanced malignancy in colon cancer. *Br J Cancer*, **99**(6): 966–973.
- Araki, H., Knapp, C., Tsai, P., and Print, C. (2012) GeneSetDB: A comprehensive meta-database, statistical and visualisation framework for gene set analysis. *FEBS Open Bio*, **2**: 76–82.

- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., *et al.* (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet*, **25**(1): 25–29.
- Ashworth, A. (2008) A synthetic lethal therapeutic approach: poly(adp) ribose polymerase inhibitors for the treatment of cancers deficient in dna double-strand break repair. *J Clin Oncol*, **26**(22): 3785–90.
- Ashworth, A., Lord, C.J., and Reis-Filho, J.S. (2011) Genetic interactions in cancer progression and treatment. *Cell*, **145**(1): 30–38.
- Audeh, M.W., Carmichael, J., Penson, R.T., Friedlander, M., Powell, B., Bell-McGuinn, K.M., Scott, C., Weitzel, J.N., Oaknin, A., Loman, N., *et al.* (2010) Oral poly(adp-ribose) polymerase inhibitor olaparib in patients with *BRCA1* or *BRCA2* mutations and recurrent ovarian cancer: a proof-of-concept trial. *Lancet*, **376**(9737): 245–51.
- Babyak, M.A. (2004) What you see may not be what you get: a brief, nontechnical introduction to overfitting in regression-type models. *Psychosom Med*, **66**(3): 411–21.
- Bamford, S., Dawson, E., Forbes, S., Clements, J., Pettett, R., Dogan, A., Flanagan, A., Teague, J., Futreal, P.A., Stratton, M.R., *et al.* (2004) The COSMIC (Catalogue of Somatic Mutations in Cancer) database and website. *Br J Cancer*, **91**(2): 355–358.
- Barabási, A.L. and Albert, R. (1999) Emergence of scaling in random networks. *Science*, **286**(5439): 509–12.
- Barabási, A.L., Gulbahce, N., and Loscalzo, J. (2011) Network medicine: a network-based approach to human disease. *Nat Rev Genet*, **12**(1): 56–68.
- Barabási, A.L. and Oltvai, Z.N. (2004) Network biology: understanding the cell’s functional organization. *Nat Rev Genet*, **5**(2): 101–13.
- Barrat, A. and Weigt, M. (2000) On the properties of small-world network models. *The European Physical Journal B - Condensed Matter and Complex Systems*, **13**(3): 547–560.

- Barretina, J., Caponigro, G., Stransky, N., Venkatesan, K., Margolin, A.A., Kim, S., Wilson, C.J., Lehar, J., Kryukov, G.V., Sonkin, D., *et al.* (2012) The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. *Nature*, **483**(7391): 603–607.
- Barry, W.T. (2016) *safe: Significance Analysis of Function and Expression*. R package version 3.14.0.
- Baryshnikova, A., Costanzo, M., Dixon, S., Vizeacoumar, F.J., Myers, C.L., Andrews, B., and Boone, C. (2010a) Synthetic genetic array (sga) analysis in *saccharomyces cerevisiae* and *schizosaccharomyces pombe*. *Methods Enzymol*, **470**: 145–79.
- Baryshnikova, A., Costanzo, M., Kim, Y., Ding, H., Koh, J., Toufighi, K., Youn, J.Y., Ou, J., San Luis, B.J., Bandyopadhyay, S., *et al.* (2010b) Quantitative analysis of fitness and genetic interactions in yeast on a genome scale. *Nat Meth*, **7**(12): 1017–1024.
- Bass, A.J., Thorsson, V., Shmulevich, I., Reynolds, S.M., Miller, M., Bernard, B., Hinoue, T., Laird, P.W., Curtis, C., Shen, H., *et al.* (2014) Comprehensive molecular characterization of gastric adenocarcinoma. *Nature*, **513**(7517): 202–209.
- Bates, D. and Maechler, M. (2016) *Matrix: Sparse and Dense Matrix Classes and Methods*. R package version 1.2-7.1.
- Bateson, W. and Mendel, G. (1909) *Mendel's principles of heredity, by W. Bateson*. University Press, Cambridge [Eng.].
- Becker, K.F., Atkinson, M.J., Reich, U., Becker, I., Nekarda, H., Siewert, J.R., and Hfler, H. (1994) E-cadherin gene mutations provide clues to diffuse type gastric carcinomas. *Cancer Research*, **54**(14): 3845–3852.
- Bell, D., Berchuck, A., Birrer, M., Chien, J., Cramer, D., Dao, F., Dhir, R., DiSaia, P., Gabra, H., Glenn, P., *et al.* (2011) Integrated genomic analyses of ovarian carcinoma. *Nature*, **474**(7353): 609–615.
- Benjamini, Y. and Hochberg, Y. (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B (Methodological)*, **57**(1): 289–300.

- Berx, G., Cleton-Jansen, A.M., Nollet, F., de Leeuw, W.J., van de Vijver, M., Cornelisse, C., and van Roy, F. (1995) E-cadherin is a tumour/invasion suppressor gene mutated in human lobular breast cancers. *EMBO J*, **14**(24): 6107–15.
- Berx, G., Cleton-Jansen, A.M., Strumane, K., de Leeuw, W.J., Nollet, F., van Roy, F., and Cornelisse, C. (1996) E-cadherin is inactivated in a majority of invasive human lobular breast cancers by truncation mutations throughout its extracellular domain. *Oncogene*, **13**(9): 1919–25.
- Berx, G. and van Roy, F. (2009) Involvement of members of the cadherin superfamily in cancer. *Cold Spring Harb Perspect Biol*, **1**: a003129.
- Bitler, B.G., Aird, K.M., Garipov, A., Li, H., Amatangelo, M., Kossenkova, A.V., Schultz, D.C., Liu, Q., Shih, Ie, M., Conejo-Garcia, J.R., *et al.* (2015) Synthetic lethality by targeting ezh2 methyltransferase activity in arid1a-mutated cancers. *Nat Med*, **21**(3): 231–8.
- Blake, J.A., Christie, K.R., Dolan, M.E., Drabkin, H.J., Hill, D.P., Ni, L., Sitnikov, D., Burgess, S., Buza, T., Gresham, C., *et al.* (2015) Gene Ontology Consortium: going forward. *Nucleic Acids Res*, **43**(Database issue): D1049–1056.
- Boettcher, M., Lawson, A., Ladenburger, V., Fredebohm, J., Wolf, J., Hoheisel, J.D., Frezza, C., and Shlomi, T. (2014) High throughput synthetic lethality screen reveals a tumorigenic role of adenylate cyclase in fumarate hydratase-deficient cancer cells. *BMC Genomics*, **15**: 158.
- Boone, C., Bussey, H., and Andrews, B.J. (2007) Exploring genetic interactions and networks with yeast. *Nat Rev Genet*, **8**(6): 437–49.
- Borgatti, S.P. (2005) Centrality and network flow. *Social Networks*, **27**(1): 55 – 71.
- Boucher, B. and Jenna, S. (2013) Genetic interaction networks: better understand to better predict. *Front Genet*, **4**: 290.
- Bozovic-Spasojevic, I., Azambuja, E., McCaskill-Stevens, W., Dinh, P., and Cardoso, F. (2012) Chemoprevention for breast cancer. *Cancer treatment reviews*, **38**(5): 329–339.
- Breiman, L. (2001) Random forests. *Machine Learning*, **45**(1): 5–32.

- Brin, S. and Page, L. (1998) The anatomy of a large-scale hypertextual web search engine. *Computer Networks and ISDN Systems*, **30**(1): 107 – 117.
- Brückner, A., Polge, C., Lentze, N., Auerbach, D., and Schlattner, U. (2009) Yeast two-hybrid, a powerful tool for systems biology. *Int J Mol Sci*, **10**(6): 2763–2788.
- Bryant, H.E., Schultz, N., Thomas, H.D., Parker, K.M., Flower, D., Lopez, E., Kyle, S., Meuth, M., Curtin, N.J., and Helleday, T. (2005) Specific killing of *BRCA2*-deficient tumours with inhibitors of polyadprbose polymerase. *Nature*, **434**(7035): 913–7.
- Bussey, H., Andrews, B., and Boone, C. (2006) From worm genetic networks to complex human diseases. *Nat Genet*, **38**(8): 862–3.
- Butland, G., Babu, M., Diaz-Mejia, J.J., Bohdana, F., Phanse, S., Gold, B., Yang, W., Li, J., Gagarinova, A.G., Pogoutse, O., *et al.* (2008) esga: E. coli synthetic genetic array analysis. *Nat Methods*, **5**(9): 789–95.
- Cardiff, R.D., Couto, S., and Bolon, B. (2011) Three interrelated themes in current breast cancer research: gene addiction, phenotypic plasticity, and cancer stem cells. *Breast Cancer Res*, **13**(5): 216.
- cBioPortal for Cancer Genomics (cBioPortal) (2017) cBioPortal for Cancer Genomics. <http://www.cbioportal.org/>. Accessed: 26/03/2017.
- Cerami, E.G., Gross, B.E., Demir, E., Rodchenkov, I., Babur, O., Anwar, N., Schultz, N., Bader, G.D., and Sander, C. (2011) Pathway Commons, a web resource for biological pathway data. *Nucleic Acids Res*, **39**(Database issue): D685–690.
- Chen, A., Beetham, H., Black, M.A., Priya, R., Telford, B.J., Guest, J., Wiggins, G.A.R., Godwin, T.D., Yap, A.S., and Guilford, P.J. (2014) E-cadherin loss alters cytoskeletal organization and adhesion in non-malignant breast cells but is insufficient to induce an epithelial-mesenchymal transition. *BMC Cancer*, **14**(1): 552.
- Chen, S. and Parmigiani, G. (2007) Meta-analysis of BRCA1 and BRCA2 penetrance. *J Clin Oncol*, **25**(11): 1329–1333.
- Chipman, K. and Singh, A. (2009) Predicting genetic interactions with random walks on biological networks. *BMC Bioinformatics*, **10**(1): 17.

- Christofori, G. and Semb, H. (1999) The role of the cell-adhesion molecule E-cadherin as a tumour-suppressor gene. *Trends in Biochemical Sciences*, **24**(2): 73 – 76.
- Ciriello, G., Gatza, M.L., Beck, A.H., Wilkerson, M.D., Rhie, S.K., Pastore, A., Zhang, H., McLellan, M., Yau, C., Kandoth, C., *et al.* (2015) Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. *Cell*, **163**(2): 506–519.
- Clark, M.J. (2004) Endogenous Regulator of G Protein Signaling Proteins Suppress G o-Dependent μ -Opioid Agonist-Mediated Adenylyl Cyclase Supersensitization. *Journal of Pharmacology and Experimental Therapeutics*, **310**(1): 215–222.
- Collingridge, D.S. (2013) A primer on quantitized data analysis and permutation testing. *Journal of Mixed Methods Research*, **7**(1): 81–97.
- Collins, F.S. and Barker, A.D. (2007) Mapping the cancer genome. Pinpointing the genes involved in cancer will help chart a new course across the complex landscape of human malignancies. *Sci Am*, **296**(3): 50–57.
- Collisson, E., Campbell, J., Brooks, A., Berger, A., Lee, W., Chmielecki, J., Beer, D., Cope, L., Creighton, C., Danilova, L., *et al.* (2014) Comprehensive molecular profiling of lung adenocarcinoma. *Nature*, **511**(7511): 543–550.
- Costanzo, M., Baryshnikova, A., Bellay, J., Kim, Y., Spear, E.D., Sevier, C.S., Ding, H., Koh, J.L., Toufighi, K., Mostafavi, S., *et al.* (2010) The genetic landscape of a cell. *Science*, **327**(5964): 425–31.
- Costanzo, M., Baryshnikova, A., Myers, C.L., Andrews, B., and Boone, C. (2011) Charting the genetic interaction map of a cell. *Curr Opin Biotechnol*, **22**(1): 66–74.
- Courtney, K.D., Corcoran, R.B., and Engelman, J.A. (2010) The PI3K pathway as drug target in human cancer. *J Clin Oncol*, **28**(6): 1075–1083.
- Creighton, C.J., Morgan, M., Gunaratne, P.H., Wheeler, D.A., Gibbs, R.A., Robertson, A., Chu, A., Beroukhi, R., Cibulskis, K., Signoretti, S., *et al.* (2013) Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature*, **499**(7456): 43–49.
- Croft, D., Mundo, A.F., Haw, R., Milacic, M., Weiser, J., Wu, G., Caudy, M., Garapati, P., Gillespie, M., Kamdar, M.R., *et al.* (2014) The Reactome pathway knowledge-base. *Nucleic Acids Res*, **42**(database issue): D472D477.

- Crunkhorn, S. (2014) Cancer: Predicting synthetic lethal interactions. *Nat Rev Drug Discov*, **13**(11): 812.
- Csardi, G. and Nepusz, T. (2006) The igraph software package for complex network research. *InterJournal, Complex Systems*: 1695.
- Dai, X., Li, T., Bai, Z., Yang, Y., Liu, X., Zhan, J., and Shi, B. (2015) Breast cancer intrinsic subtype classification, clinical use and future trends. *Am J Cancer Res*, **5**(10): 2929–2943.
- Davierwala, A.P., Haynes, J., Li, Z., Brost, R.L., Robinson, M.D., Yu, L., Mnaimneh, S., Ding, H., Zhu, H., Chen, Y., *et al.* (2005) The synthetic genetic interaction spectrum of essential genes. *Nat Genet*, **37**(10): 1147–1152.
- De Leeuw, W.J., Berx, G., Vos, C.B., Peterse, J.L., Van de Vijver, M.J., Litvinov, S., Van Roy, F., Cornelisse, C.J., and Cleton-Jansen, A.M. (1997) Simultaneous loss of E-cadherin and catenins in invasive lobular breast cancer and lobular carcinoma in situ. *J Pathol*, **183**(4): 404–11.
- De Santis, G., Miotti, S., Mazzi, M., Canevari, S., and Tomassetti, A. (2009) E-cadherin directly contributes to PI3K/AKT activation by engaging the PI3K-p85 regulatory subunit to adherens junctions of ovarian carcinoma cells. *Oncogene*, **28**(9): 1206–1217.
- Demir, E., Babur, O., Rodchenkov, I., Aksoy, B.A., Fukuda, K.I., Gross, B., Sumer, O.S., Bader, G.D., and Sander, C. (2013) Using biological pathway data with Pax-tools. *PLoS Comput Biol*, **9**(9): e1003194.
- Deshpande, R., Asiedu, M.K., Klebig, M., Sutor, S., Kuzmin, E., Nelson, J., Piotrowski, J., Shin, S.H., Yoshida, M., Costanzo, M., *et al.* (2013) A comparative genomic approach for identifying synthetic lethal interactions in human cancer. *Cancer Res*, **73**(20): 6128–36.
- Dickson, D. (1999) Wellcome funds cancer database. *Nature*, **401**(6755): 729.
- Dijkstra, E.W. (1959) A note on two problems in connexion with graphs. *Numerische Mathematik*, **1**(1): 269–271.
- Dixon, S.J., Andrews, B.J., and Boone, C. (2009) Exploring the conservation of synthetic lethal genetic interaction networks. *Commun Integr Biol*, **2**(2): 78–81.

- Dixon, S.J., Fedyshyn, Y., Koh, J.L., Prasad, T.S., Chahwan, C., Chua, G., Toufighi, K., Baryshnikova, A., Hayles, J., Hoe, K.L., *et al.* (2008) Significant conservation of synthetic lethal genetic interaction networks between distantly related eukaryotes. *Proc Natl Acad Sci U S A*, **105**(43): 16653–8.
- Dorsam, R.T. and Gutkind, J.S. (2007) G-protein-coupled receptors and cancer. *Nat Rev Cancer*, **7**(2): 79–94.
- Erdős, P. and Rényi, A. (1959) On random graphs I. *Publ Math Debrecen*, **6**: 290–297.
- Erdős, P. and Rényi, A. (1960) On the evolution of random graphs. *Publ Math Inst Hung Acad Sci*, **5**(1): 17–61.
- Eroles, P., Bosch, A., Perez-Fidalgo, J.A., and Lluch, A. (2012) Molecular biology in breast cancer: intrinsic subtypes and signaling pathways. *Cancer Treat Rev*, **38**(6): 698–707.
- Farmer, H., McCabe, N., Lord, C.J., Tutt, A.N., Johnson, D.A., Richardson, T.B., Santarosa, M., Dillon, K.J., Hickson, I., Knights, C., *et al.* (2005) Targeting the dna repair defect in BRCA mutant cells as a therapeutic strategy. *Nature*, **434**(7035): 917–21.
- Fawcett, T. (2006) An introduction to ROC analysis. *Pattern Recognition Letters*, **27**(8): 861 – 874. {ROC} Analysis in Pattern Recognition.
- Fece de la Cruz, F., Gapp, B.V., and Nijman, S.M. (2015) Synthetic lethal vulnerabilities of cancer. *Annu Rev Pharmacol Toxicol*, **55**: 513–531.
- Ferlay, J., Soerjomataram, I., Dikshit, R., Eser, S., Mathers, C., Rebelo, M., Parkin, D.M., Forman, D., and Bray, F. (2015) Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*, **136**(5): E359–386.
- Fisher, R.A. (1919) Xv.the correlation between relatives on the supposition of mendelian inheritance. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, **52**(02): 399–433.
- Fong, P.C., Boss, D.S., Yap, T.A., Tutt, A., Wu, P., Mergui-Roelvink, M., Mortimer, P., Swaisland, H., Lau, A., O’Connor, M.J., *et al.* (2009) Inhibition of poly(adp-ribose) polymerase in tumors from BRCA mutation carriers. *N Engl J Med*, **361**(2): 123–34.

- Fong, P.C., Yap, T.A., Boss, D.S., Carden, C.P., Mergui-Roelvink, M., Gourley, C., De Greve, J., Lubinski, J., Shanley, S., Messiou, C., *et al.* (2010) Poly(adp)-ribose polymerase inhibition: frequent durable responses in BRCA carrier ovarian cancer correlating with platinum-free interval. *J Clin Oncol*, **28**(15): 2512–9.
- Forbes, S.A., Beare, D., Gunasekaran, P., Leung, K., Bindal, N., Boutselakis, H., Ding, M., Bamford, S., Cole, C., Ward, S., *et al.* (2015) COSMIC: exploring the world’s knowledge of somatic mutations in human cancer. *Nucleic Acids Res*, **43**(Database issue): D805–811.
- Fraser, A. (2004) Towards full employment: using RNAi to find roles for the redundant. *Oncogene*, **23**(51): 8346–52.
- Friedman, N., Linial, M., Nachman, I., and Pe’er, D. (2000) Using Bayesian networks to analyze expression data. *J Comput Biol*, **7**(3-4): 601–620.
- Fromental-Ramain, C., Warot, X., Lakkaraju, S., Favier, B., Haack, H., Birling, C., Dierich, A., Doll e, P., and Chambon, P. (1996) Specific and redundant functions of the paralogous Hoxa-9 and Hoxd-9 genes in forelimb and axial skeleton patterning. *Development*, **122**(2): 461–472.
- Futreal, P.A., Coin, L., Marshall, M., Down, T., Hubbard, T., Wooster, R., Rahman, N., and Stratton, M.R. (2004) A census of human cancer genes. *Nat Rev Cancer*, **4**(3): 177–183.
- Futreal, P.A., Kasprzyk, A., Birney, E., Mullikin, J.C., Wooster, R., and Stratton, M.R. (2001) Cancer and genomics. *Nature*, **409**(6822): 850–852.
- Gao, B. and Roux, P.P. (2015) Translational control by oncogenic signaling pathways. *Biochimica et Biophysica Acta*, **1849**(7): 753–65.
- Gatza, M.L., Kung, H.N., Blackwell, K.L., Dewhirst, M.W., Marks, J.R., and Chi, J.T. (2011) Analysis of tumor environmental response and oncogenic pathway activation identifies distinct basal and luminal features in HER2-related breast tumor subtypes. *Breast Cancer Res*, **13**(3): R62.
- Gatza, M.L., Lucas, J.E., Barry, W.T., Kim, J.W., Wang, Q., Crawford, M.D., Datto, M.B., Kelley, M., Mathey-Prevot, B., Potti, A., *et al.* (2010) A pathway-based classification of human breast cancer. *Proc Natl Acad Sci USA*, **107**(15): 6994–6999.

- Gatza, M.L., Silva, G.O., Parker, J.S., Fan, C., and Perou, C.M. (2014) An integrated genomics approach identifies drivers of proliferation in luminal-subtype human breast cancer. *Nat Genet*, **46**(10): 1051–1059.
- Gentleman, R.C., Carey, V.J., Bates, D.M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B., Gautier, L., Ge, Y., Gentry, J., *et al.* (2004) Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol*, **5**(10): R80.
- Genz, A. and Bretz, F. (2009) Computation of multivariate normal and t probabilities. In *Lecture Notes in Statistics*, volume 195. Springer-Verlag, Heidelberg.
- Genz, A., Bretz, F., Miwa, T., Mi, X., Leisch, F., Scheipl, F., and Hothorn, T. (2016) *mvtnorm: Multivariate Normal and t Distributions*. R package version 1.0-5. URL.
- Glaire, M.A., Brown, M., Church, D.N., and Tomlinson, I. (2017) Cancer predisposition syndromes: lessons for truly precision medicine. *J Pathol*, **241**(2): 226–235.
- Globus (Globus) (2017) Research data management simplified. <https://www.globus.org/>. Accessed: 25/03/2017.
- Goodwin, S., McPherson, J.D., and McCombie, W.R. (2016) Coming of age: ten years of next-generation sequencing technologies. *Nat Rev Genet*, **17**(6): 333–351.
- Grady, W.M., Willis, J., Guilford, P.J., Dunbier, A.K., Toro, T.T., Lynch, H., Wiesner, G., Ferguson, K., Eng, C., Park, J.G., *et al.* (2000) Methylation of the CDH1 promoter as the second genetic hit in hereditary diffuse gastric cancer. *Nat Genet*, **26**(1): 16–17.
- Graziano, F., Humar, B., and Guilford, P. (2003) The role of the E-cadherin gene (*CDH1*) in diffuse gastric cancer susceptibility: from the laboratory to clinical practice. *Annals of Oncology*, **14**(12): 1705–1713.
- Guaragnella, N., Palermo, V., Galli, A., Moro, L., Mazzoni, C., and Giannattasio, S. (2014) The expanding role of yeast in cancer research and diagnosis: insights into the function of the oncosuppressors p53 and BRCA1/2. *FEMS Yeast Res*, **14**(1): 2–16.
- Güell, O., Sagus, F., and Serrano, M. (2014) Essential plasticity and redundancy of metabolism unveiled by synthetic lethality analysis. *PLoS Comput Biol*, **10**(5): e1003637.

- Guilford, P., Hopkins, J., Harraway, J., McLeod, M., McLeod, N., Harawira, P., Taite, H., Scoular, R., Miller, A., and Reeve, A.E. (1998) E-cadherin germline mutations in familial gastric cancer. *Nature*, **392**(6674): 402–5.
- Guilford, P., Humar, B., and Blair, V. (2010) Hereditary diffuse gastric cancer: translation of *CDH1* germline mutations into clinical practice. *Gastric Cancer*, **13**(1): 1–10.
- Guilford, P.J., Hopkins, J.B., Grady, W.M., Markowitz, S.D., Willis, J., Lynch, H., Rajput, A., Wiesner, G.L., Lindor, N.M., Burgart, L.J., *et al.* (1999) E-cadherin germline mutations define an inherited cancer syndrome dominated by diffuse gastric cancer. *Hum Mutat*, **14**(3): 249–55.
- Guo, J., Liu, H., and Zheng, J. (2016) SynLethDB: synthetic lethality database toward discovery of selective and sensitive anticancer drug targets. *Nucleic Acids Res*, **44**(D1): D1011–1017.
- Hajian-Tilaki, K. (2013) Receiver Operating Characteristic (ROC) Curve Analysis for Medical Diagnostic Test Evaluation. *Caspian J Intern Med*, **4**(2): 627–635.
- Hall, M., Frank, E., Holmes, G., Pfahringer, B., Reutemann, P., and Witten, I.H. (2009) The weka data mining software: an update. *SIGKDD Explor Newsl*, **11**(1): 10–18.
- Hammerman, P.S., Lawrence, M.S., Voet, D., Jing, R., Cibulskis, K., Sivachenko, A., Stojanov, P., McKenna, A., Lander, E.S., Gabriel, S., *et al.* (2012) Comprehensive genomic characterization of squamous cell lung cancers. *Nature*, **489**(7417): 519–525.
- Hanahan, D. and Weinberg, R.A. (2000) The hallmarks of cancer. *Cell*, **100**(1): 57–70.
- Hanahan, D. and Weinberg, R.A. (2011) Hallmarks of cancer: the next generation. *Cell*, **144**(5): 646–674.
- Hanna, S. (2003) Cancer incidence in new zealand (2003-2007). In D. Forman, D. Bray F Brewster, C. Gombe Mbalawa, B. Kohler, M. Piñeros, E. Steliarova-Foucher, R. Swaminathan, and J. Ferlay (editors), *Cancer Incidence in Five Continents*, volume X, 902–907. International Agency for Research on Cancer, Lyon, France. Electronic version <http://ci5.iarc.fr> Accessed 22/03/2017.

- Hansford, S., Kaurah, P., Li-Chang, H., Woo, M., Senz, J., Pinheiro, H., Schrader, K.A., Schaeffer, D.F., Shumansky, K., Zogopoulos, G., *et al.* (2015) Hereditary Diffuse Gastric Cancer Syndrome: CDH1 Mutations and Beyond. *JAMA Oncol*, **1**(1): 23–32.
- Heiskanen, M.A. and Aittokallio, T. (2012) Mining high-throughput screens for cancer drug targets-lessons from yeast chemical-genomic profiling and synthetic lethality. *Wiley Interdisciplinary Reviews: Data Mining and Knowledge Discovery*, **2**(3): 263–272.
- Hell, P. (1976) Graphs with given neighbourhoods i. problèmes combinatoires at theorie des graphes. *Proc Coil Int CNRS, Orsay*, **260**: 219–223.
- Higgins, M.E., Claremont, M., Major, J.E., Sander, C., and Lash, A.E. (2007) CancerGenes: a gene selection resource for cancer genome projects. *Nucleic Acids Res*, **35**(Database issue): D721–726.
- Hillenmeyer, M.E. (2008) The chemical genomic portrait of yeast: uncovering a phenotype for all genes. *Science*, **320**: 362–365.
- Hoadley, K.A., Yau, C., Wolf, D.M., Cherniack, A.D., Tamborero, D., Ng, S., Leiserson, M.D., Niu, B., McLellan, M.D., Uzunangelov, V., *et al.* (2014) Multiplatform analysis of 12 cancer types reveals molecular classification within and across tissues of origin. *Cell*, **158**(4): 929–944.
- Hoehndorf, R., Hardy, N.W., Osumi-Sutherland, D., Tweedie, S., Schofield, P.N., and Gkoutos, G.V. (2013) Systematic analysis of experimental phenotype data reveals gene functions. *PLoS ONE*, **8**(4): e60847.
- Holm, S. (1979) A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, **6**(2): 65–70.
- Hopkins, A.L. (2008) Network pharmacology: the next paradigm in drug discovery. *Nat Chem Biol*, **4**(11): 682–690.
- Hu, Z., Fan, C., Oh, D.S., Marron, J.S., He, X., Qaqish, B.F., Livasy, C., Carey, L.A., Reynolds, E., Dressler, L., *et al.* (2006) The molecular portraits of breast tumors are conserved across microarray platforms. *BMC Genomics*, **7**: 96.

- Huang, E., Cheng, S., Dressman, H., Pittman, J., Tsou, M., Horng, C., Bild, A., Iversen, E., Liao, M., Chen, C., *et al.* (2003) Gene expression predictors of breast cancer outcomes. *Lancet*, **361**: 1590–1596.
- Hutchison, C.A., Chuang, R.Y., Noskov, V.N., Assad-Garcia, N., Deerinck, T.J., Ellisman, M.H., Gill, J., Kannan, K., Karas, B.J., Ma, L., *et al.* (2016) Design and synthesis of a minimal bacterial genome. *Science*, **351**(6280): aad6253.
- Imoto, S., Higuchi, T., Goto, T., Tashiro, K., Kuhara, S., and Miyano, S. (2004) Combining microarrays and biological knowledge for estimating gene networks via bayesian networks. *J Bioinform Comput Biol*, **2**(1): 77–98.
- International HapMap 3 Consortium (HapMap) (2003) The International HapMap Project. *Nature*, **426**(6968): 789–796.
- Jansen, R., Yu, H., Greenbaum, D., Kluger, Y., Krogan, N.J., Chung, S., Emili, A., Snyder, M., Greenblatt, J.F., and Gerstein, M. (2003) A Bayesian networks approach for predicting protein-protein interactions from genomic data. *Science*, **302**(5644): 449–453.
- Jeanes, A., Gottardi, C.J., and Yap, A.S. (2008) Cadherins and cancer: how does cadherin dysfunction promote tumor progression? *Oncogene*, **27**(55): 6920–6929.
- Jerby-Arnon, L., Pfetzer, N., Waldman, Y., McGarry, L., James, D., Shanks, E., Seashore-Ludlow, B., Weinstock, A., Geiger, T., Clemons, P., *et al.* (2014) Predicting cancer-specific vulnerability via data-driven detection of synthetic lethality. *Cell*, **158**(5): 1199–1209.
- Joachims, T. (1999) Making large-scale support vector machine learning practical. In S. Bernhard, I. Kropf, J.C.B. Christopher, and J.S. Alexander (editors), *Advances in kernel methods*, 169–184. MIT Press.
- Kaelin, Jr, W. (2005) The concept of synthetic lethality in the context of anticancer therapy. *Nat Rev Cancer*, **5**(9): 689–98.
- Kaelin, Jr, W. (2009) Synthetic lethality: a framework for the development of wiser cancer therapeutics. *Genome Med*, **1**: 99.
- Kamada, T. and Kawai, S. (1989) An algorithm for drawing general undirected graphs. *Information Processing Letters*, **31**(1): 7–15.

- Kawai, J., Shinagawa, A., Shibata, K., Yoshino, M., Itoh, M., Ishii, Y., Arakawa, T., Hara, A., Fukunishi, Y., Konno, H., *et al.* (2001) Functional annotation of a full-length mouse cDNA collection. *Nature*, **409**(6821): 685–690.
- Kelley, R. and Ideker, T. (2005) Systematic interpretation of genetic interactions using protein networks. *Nat Biotech*, **23**(5): 561–566.
- Kelly, S.T. (2013) *Statistical Predictions of Synthetic Lethal Interactions in Cancer*. Dissertation, University of Otago.
- Keshava Prasad, T.S., Goel, R., Kandasamy, K., Keerthikumar, S., Kumar, S., Mathivanan, S., Telikicherla, D., Raju, R., Shafreen, B., Venugopal, A., *et al.* (2009) Human Protein Reference Database–2009 update. *Nucleic Acids Res*, **37**(Database issue): D767–772.
- Kim, N.G., Koh, E., Chen, X., and Gumbiner, B.M. (2011) E-cadherin mediates contact inhibition of proliferation through Hippo signaling-pathway components. *Proc Natl Acad Sci USA*, **108**(29): 11930–11935.
- Koboldt, D.C., Fulton, R.S., McLellan, M.D., Schmidt, H., Kalicki-Veizer, J., McMichael, J.F., Fulton, L.L., Dooling, D.J., Ding, L., Mardis, E.R., *et al.* (2012) Comprehensive molecular portraits of human breast tumours. *Nature*, **490**(7418): 61–70.
- Kockel, L., Zeitlinger, J., Staszewski, L.M., Mlodzik, M., and Bohmann, D. (1997) Jun in drosophila development: redundant and nonredundant functions and regulation by two mapk signal transduction pathways. *Genes & Development*, **11**(13): 1748–1758.
- Kozlov, K.N., Gursky, V.V., Kulakovskiy, I.V., and Samsonova, M.G. (2015) Sequence-based model of gap gene regulation network. *BMC Genomics*, **15**(Suppl 12): S6.
- Kranthi, S., Rao, S., and Manimaran, P. (2013) Identification of synthetic lethal pairs in biological systems through network information centrality. *Mol BioSyst*, **9**(8): 2163–2167.
- Kroepil, F., Fluegen, G., Totikov, Z., Baldus, S.E., Vay, C., Schauer, M., Topp, S.A., Esch, J.S., Knoefel, W.T., and Stoecklein, N.H. (2012) Down-regulation of CDH1 is associated with expression of SNAIL in colorectal adenomas. *PLoS ONE*, **7**(9): e46665.

- Lander, E.S. (2011) Initial impact of the sequencing of the human genome. *Nature*, **470**(7333): 187–197.
- Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., *et al.* (2001) Initial sequencing and analysis of the human genome. *Nature*, **409**(6822): 860–921.
- Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009) Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol*, **10**(3): R25.
- Latora, V. and Marchiori, M. (2001) Efficient behavior of small-world networks. *Phys Rev Lett*, **87**: 198701.
- Laufer, C., Fischer, B., Billmann, M., Huber, W., and Boutros, M. (2013) Mapping genetic interactions in human cancer cells with RNAi and multiparametric phenotyping. *Nat Methods*, **10**(5): 427–31.
- Law, C.W., Chen, Y., Shi, W., and Smyth, G.K. (2014) voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol*, **15**(2): R29.
- Le Meur, N. and Gentleman, R. (2008) Modeling synthetic lethality. *Genome Biol*, **9**(9): R135.
- Le Meur, N., Jiang, Z., Liu, T., Mar, J., and Gentleman, R.C. (2014) Slgi: Synthetic lethal genetic interaction. r package version 1.26.0.
- Lee, A.Y., Perreault, R., Harel, S., Boulier, E.L., Suderman, M., Hallett, M., and Jenna, S. (2010a) Searching for signaling balance through the identification of genetic interactors of the rab guanine-nucleotide dissociation inhibitor gdi-1. *PLoS ONE*, **5**(5): e10624.
- Lee, I., Lehner, B., Vavouri, T., Shin, J., Fraser, A.G., and Marcotte, E.M. (2010b) Predicting genetic modifier loci using functional gene networks. *Genome Research*, **20**(8): 1143–1153.
- Lee, I. and Marcotte, E.M. (2009) Effects of functional bias on supervised learning of a gene network model. *Methods Mol Biol*, **541**: 463–75.

- Lee, M.J., Ye, A.S., Gardino, A.K., Heijink, A.M., Sorger, P.K., MacBeath, G., and Yaffe, M.B. (2012) Sequential application of anticancer drugs enhances cell death by rewiring apoptotic signaling networks. *Cell*, **149**(4): 780–94.
- Lehner, B., Crombie, C., Tischler, J., Fortunato, A., and Fraser, A.G. (2006) Systematic mapping of genetic interactions in *caenorhabditis elegans* identifies common modifiers of diverse signaling pathways. *Nat Genet*, **38**(8): 896–903.
- Li, B., Ruotti, V., Stewart, R.M., Thomson, J.A., and Dewey, C.N. (2010) RNA-Seq gene expression estimation with read mapping uncertainty. *Bioinformatics*, **26**(4): 493–500.
- Li, X.J., Mishra, S.K., Wu, M., Zhang, F., and Zheng, J. (2014) Syn-lethality: An integrative knowledge base of synthetic lethality towards discovery of selective anti-cancer therapies. *Biomed Res Int*, **2014**: 196034.
- Linehan, W.M., Spellman, P.T., Ricketts, C.J., Creighton, C.J., Fei, S.S., Davis, C., Wheeler, D.A., Murray, B.A., Schmidt, L., Vocke, C.D., *et al.* (2016) Comprehensive Molecular Characterization of Papillary Renal-Cell Carcinoma. *N Engl J Med*, **374**(2): 135–145.
- Lokody, I. (2014) Computational modelling: A computational crystal ball. *Nature Reviews Cancer*, **14**(10): 649–649.
- Lord, C.J., Tutt, A.N., and Ashworth, A. (2015) Synthetic lethality and cancer therapy: lessons learned from the development of PARP inhibitors. *Annu Rev Med*, **66**: 455–470.
- Lu, X., Kensche, P.R., Huynen, M.A., and Notebaart, R.A. (2013) Genome evolution predicts genetic interactions in protein complexes and reveals cancer drug targets. *Nat Commun*, **4**: 2124.
- Lu, X., Megchelenbrink, W., Notebaart, R.A., and Huynen, M.A. (2015) Predicting human genetic interactions from cancer genome evolution. *PLoS One*, **10**(5): e0125795.
- Lum, P.Y., Armour, C.D., Stepaniants, S.B., Cavet, G., Wolf, M.K., Butler, J.S., Hinshaw, J.C., Garnier, P., Prestwich, G.D., Leonardson, A., *et al.* (2004) Discovering modes of action for therapeutic compounds using a genome-wide screen of yeast heterozygotes. *Cell*, **116**(1): 121–137.

- Luo, J., Solimini, N.L., and Elledge, S.J. (2009) Principles of Cancer Therapy: Oncogene and Non-oncogene Addiction. *Cell*, **136**(5): 823–837.
- Machado, J., Olivera, C., Carvalh, R., Soares, P., Berx, G., Caldas, C., Sercuca, R., Carneiro, F., and Sorbrinho-Simoes, M. (2001) E-cadherin gene (*CDH1*) promoter methylation as the second hit in sporadic diffuse gastric carcinoma. *Oncogene*, **20**: 1525–1528.
- Markowetz, F. (2017) All biology is computational biology. *PLoS Biol*, **15**(3): e2002050.
- Masciari, S., Larsson, N., Senz, J., Boyd, N., Kaurah, P., Kandel, M.J., Harris, L.N., Pinheiro, H.C., Troussard, A., Miron, P., *et al.* (2007) Germline E-cadherin mutations in familial lobular breast cancer. *J Med Genet*, **44**(11): 726–31.
- Mattison, J., van der Weyden, L., Hubbard, T., and Adams, D.J. (2009) Cancer gene discovery in mouse and man. *Biochim Biophys Acta*, **1796**(2): 140–161.
- McLachlan, J., George, A., and Banerjee, S. (2016) The current status of parp inhibitors in ovarian cancer. *Tumori*, **102**(5): 433–440.
- McLendon, R., Friedman, A., Bigner, D., Van Meir, E.G., Brat, D.J., Mastrogianakis, G.M., Olson, J.J., Mikkelsen, T., Lehman, N., Aldape, K., *et al.* (2008) Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature*, **455**(7216): 1061–1068.
- Miles, D.W. (2001) Update on HER-2 as a target for cancer therapy: herceptin in the clinical setting. *Breast Cancer Res*, **3**(6): 380–384.
- Muzny, D.M., Bainbridge, M.N., Chang, K., Dinh, H.H., Drummond, J.A., Fowler, G., Kovar, C.L., Lewis, L.R., Morgan, M.B., Newsham, I.F., *et al.* (2012) Comprehensive molecular characterization of human colon and rectal cancer. *Nature*, **487**(7407): 330–337.
- Nagalla, S., Chou, J.W., Willingham, M.C., Ruiz, J., Vaughn, J.P., Dubey, P., Lash, T.L., Hamilton-Dutoit, S.J., Bergh, J., Sotiriou, C., *et al.* (2013) Interactions between immunity, proliferation and molecular subtype in breast cancer prognosis. *Genome Biol*, **14**(4): R34.
- Novomestky, F. (2012) *matrixcalc: Collection of functions for matrix calculations*. R package version 1.0-3.

- Nowak, M.A., Boerlijst, M.C., Cooke, J., and Smith, J.M. (1997) Evolution of genetic redundancy. *Nature*, **388**(6638): 167–171.
- Oliveira, C., Senz, J., Kaurah, P., Pinheiro, H., Sanges, R., Haegert, A., Corso, G., Schouten, J., Fitzgerald, R., Vogelsang, H., *et al.* (2009) Germline *CDH1* deletions in hereditary diffuse gastric cancer families. *Human Molecular Genetics*, **18**(9): 1545–1555.
- Oliveira, C., Seruca, R., Hoogerbrugge, N., Ligtenberg, M., and Carneiro, F. (2013) Clinical utility gene card for: Hereditary diffuse gastric cancer (HDGC). *Eur J Hum Genet*, **21**(8).
- Pandey, G., Zhang, B., Chang, A.N., Myers, C.L., Zhu, J., Kumar, V., and Schadt, E.E. (2010) An integrative multi-network and multi-classifier approach to predict genetic interactions. *PLoS Comput Biol*, **6**(9).
- Parker, J., Mullins, M., Cheung, M., Leung, S., Voduc, D., Vickery, T., Davies, S., Fauron, C., He, X., Hu, Z., *et al.* (2009) Supervised risk predictor of breast cancer based on intrinsic subtypes. *Journal of Clinical Oncology*, **27**(8): 1160–1167.
- Pereira, B., Chin, S.F., Rueda, O.M., Vollan, H.K., Provenzano, E., Bardwell, H.A., Pugh, M., Jones, L., Russell, R., Sammut, S.J., *et al.* (2016) Erratum: The somatic mutation profiles of 2,433 breast cancers refine their genomic and transcriptomic landscapes. *Nat Commun*, **7**: 11908.
- Perou, C.M., Sørlie, T., Eisen, M.B., van de Rijn, M., Jeffrey, S.S., Rees, C.A., Pollack, J.R., Ross, D.T., Johnsen, H., Akslen, L.A., *et al.* (2000) Molecular portraits of human breast tumours. *Nature*, **406**(6797): 747–752.
- R Core Team (2016) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. R version 3.3.2.
- Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research*, **43**(7): e47.
- Roguev, A., Bandyopadhyay, S., Zofall, M., Zhang, K., Fischer, T., Collins, S.R., Qu, H., Shales, M., Park, H.O., Hayles, J., *et al.* (2008) Conservation and rewiring of functional modules revealed by an epistasis map in fission yeast. *Science*, **322**(5900): 405–10.

- Roychowdhury, S. and Chinnaiyan, A.M. (2016) Translating cancer genomes and transcriptomes for precision oncology. *CA Cancer J Clin*, **66**(1): 75–88.
- Rung, J. and Brazma, A. (2013) Reuse of public genome-wide gene expression data. *Nat Rev Genet*, **14**(2): 89–99.
- Ryan, C., Lord, C., and Ashworth, A. (2014) Daisy: Picking synthetic lethals from cancer genomes. *Cancer Cell*, **26**(3): 306–308.
- Schena, M. (1996) Genome analysis with gene expression microarrays. *Bioessays*, **18**(5): 427–431.
- Scheuer, L., Kauff, N., Robson, M., Kelly, B., Barakat, R., Satagopan, J., Ellis, N., Hensley, M., Boyd, J., Borgen, P., *et al.* (2002) Outcome of preventive surgery and screening for breast and ovarian cancer in BRCA mutation carriers. *J Clin Oncol*, **20**(5): 1260–1268.
- Semb, H. and Christofori, G. (1998) The tumor-suppressor function of E-cadherin. *Am J Hum Genet*, **63**(6): 1588–93.
- Sing, T., Sander, O., Beerenwinkel, N., and Lengauer, T. (2005) Rocr: visualizing classifier performance in r. *Bioinformatics*, **21**(20): 7881.
- Slurm development team (Slurm) (2017) Slurm workload manager. <https://slurm.schedmd.com/>. Accessed: 25/03/2017.
- Sørlie, T., Perou, C.M., Tibshirani, R., Aas, T., Geisler, S., Johnsen, H., Hastie, T., Eisen, M.B., van de Rijn, M., Jeffrey, S.S., *et al.* (2001) Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci USA*, **98**(19): 10869–10874.
- Srihari, S., Singla, J., Wong, L., and Ragan, M.A. (2015) Inferring synthetic lethal interactions from mutual exclusivity of genetic events in cancer. *Biology Direct*, **10**(1): 57.
- Stajich, J.E. and Lapp, H. (2006) Open source tools and toolkits for bioinformatics: significance, and where are we? *Brief Bioinformatics*, **7**(3): 287–296.
- Stratton, M.R., Campbell, P.J., and Futreal, P.A. (2009) The cancer genome. *Nature*, **458**(7239): 719–724.

- Ström, C. and Helleday, T. (2012) Strategies for the use of poly(adenosine diphosphate ribose) polymerase (parp) inhibitors in cancer therapy. *Biomolecules*, **2**(4): 635–649.
- Tarazona, S., Garcia-Alcalde, F., Dopazo, J., Ferrer, A., and Conesa, A. (2011) Differential expression in RNA-seq: a matter of depth. *Genome Res*, **21**(12): 2213–2223.
- Telford, B.J., Chen, A., Beetham, H., Frick, J., Brew, T.P., Gould, C.M., Single, A., Godwin, T., Simpson, K.J., and Guilford, P. (2015) Synthetic lethal screens identify vulnerabilities in gpcr signalling and cytoskeletal organization in E-cadherin-deficient cells. *Mol Cancer Ther*, **14**(5): 1213–1223.
- The 1000 Genomes Project Consortium (1000 Genomes) (2010) A map of human genome variation from population-scale sequencing. *Nature*, **467**(7319): 1061–1073.
- The Cancer Genome Atlas Research Network (TCGA) (2017) The Cancer Genome Atlas Project. <https://cancergenome.nih.gov/>. Accessed: 26/03/2017.
- The Catalogue Of Somatic Mutations In Cancer (COSMIC) (2016) Cosmic: The catalogue of somatic mutations in cancer. <http://cancer.sanger.ac.uk/cosmic>. Release 79 (23/08/2016), Accessed: 05/02/2017.
- The Comprehensive R Archive Network (CRAN) (2017) Cran. <https://cran.r-project.org/>. Accessed: 24/03/2017.
- The ENCODE Project Consortium (ENCODE) (2004) The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science*, **306**(5696): 636–640.
- The National Cancer Institute (NCI) (2015) The genetics of cancer. <https://www.cancer.gov/about-cancer/causes-prevention/genetics>. Published: 22/04/2015, Accessed: 22/03/2017.
- The New Zealand eScience Infrastructure (NeSI) (2017) NeSI. <https://www.nesi.org.nz/>. Accessed: 25/03/2017.
- Tierney, L., Rossini, A.J., Li, N., and Sevcikova, H. (2015) *snow: Simple Network of Workstations*. R package version 0.4-2.
- Tiong, K.L., Chang, K.C., Yeh, K.T., Liu, T.Y., Wu, J.H., Hsieh, P.H., Lin, S.H., Lai, W.Y., Hsu, Y.C., Chen, J.Y., *et al.* (2014) Csnk1e/ctnnb1 are synthetic lethal to tp53 in colorectal cancer and are markers for prognosis. *Neoplasia*, **16**(5): 441–50.

- Tischler, J., Lehner, B., and Fraser, A.G. (2008) Evolutionary plasticity of genetic interaction networks. *Nat Genet*, **40**(4): 390–391.
- Tomasetti, C. and Vogelstein, B. (2015) Cancer etiology. Variation in cancer risk among tissues can be explained by the number of stem cell divisions. *Science*, **347**(6217): 78–81.
- Tong, A.H., Evangelista, M., Parsons, A.B., Xu, H., Bader, G.D., Page, N., Robinson, M., Raghibizadeh, S., Hogue, C.W., Bussey, H., *et al.* (2001) Systematic genetic analysis with ordered arrays of yeast deletion mutants. *Science*, **294**(5550): 2364–8.
- Tong, A.H., Lesage, G., Bader, G.D., Ding, H., Xu, H., Xin, X., Young, J., Berriz, G.F., Brost, R.L., Chang, M., *et al.* (2004) Global mapping of the yeast genetic interaction network. *Science*, **303**(5659): 808–13.
- Tran, B., Dancey, J.E., Kamel-Reid, S., McPherson, J.D., Bedard, P.L., Brown, A.M., Zhang, T., Shaw, P., Onetto, N., Stein, L., *et al.* (2012) Cancer genomics: technology, discovery, and translation. *J Clin Oncol*, **30**(6): 647–660.
- Travers, J. and Milgram, S. (1969) An experimental study of the small world problem. *Sociometry*, **32**(4): 425–443.
- Tunggal, J.A., Helfrich, I., Schmitz, A., Schwarz, H., Gunzel, D., Fromm, M., Kemler, R., Krieg, T., and Niessen, C.M. (2005) E-cadherin is essential for in vivo epidermal barrier function by regulating tight junctions. *EMBO J*, **24**(6): 1146–1156.
- Tutt, A., Robson, M., Garber, J.E., Domchek, S.M., Audeh, M.W., Weitzel, J.N., Friedlander, M., Arun, B., Loman, N., Schmutzler, R.K., *et al.* (2010) Oral poly(adp-ribose) polymerase inhibitor olaparib in patients with *BRCA1* or *BRCA2* mutations and advanced breast cancer: a proof-of-concept trial. *Lancet*, **376**(9737): 235–44.
- University of California, Santa Cruz (UCSC) (2012) Uscs cancer browser. Accessed 29/03/2012.
- van der Meer, R., Song, H.Y., Park, S.H., Abdulkadir, S.A., and Roh, M. (2014) RNAi screen identifies a synthetic lethal interaction between PIM1 overexpression and PLK1 inhibition. *Clinical Cancer Research*, **20**(12): 3211–3221.
- van der Post, R.S., Vogelaar, I.P., Carneiro, F., Guilford, P., Huntsman, D., Hoogerbrugge, N., Caldas, C., Schreiber, K.E., Hardwick, R.H., Ausems, M.G., *et al.* (2015)

- Hereditary diffuse gastric cancer: updated clinical guidelines with an emphasis on germline CDH1 mutation carriers. *J Med Genet*, **52**(6): 361–374.
- van Steen, K. (2012) Travelling the world of genegene interactions. *Briefings in Bioinformatics*, **13**(1): 1–19.
- van Steen, M. (2010) *Graph Theory and Complex Networks: An Introduction*. Maarten van Steen, VU Amsterdam.
- Vapnik, V.N. (1995) *The nature of statistical learning theory*. Springer-Verlag New York, Inc.
- Vizeacoumar, F.J., Arnold, R., Vizeacoumar, F.S., Chandrashekhar, M., Buzina, A., Young, J.T., Kwan, J.H., Sayad, A., Mero, P., Lawo, S., *et al.* (2013) A negative genetic interaction map in isogenic cancer cell lines reveals cancer cell vulnerabilities. *Mol Syst Biol*, **9**: 696.
- Vogelstein, B., Papadopoulos, N., Velculescu, V.E., Zhou, S., Diaz, L.A., and Kinzler, K.W. (2013) Cancer genome landscapes. *Science*, **339**(6127): 1546–1558.
- Vos, C.B., Cleton-Jansen, A.M., Berx, G., de Leeuw, W.J., ter Haar, N.T., van Roy, F., Cornelisse, C.J., Peterse, J.L., and van de Vijver, M.J. (1997) E-cadherin inactivation in lobular carcinoma in situ of the breast: an early event in tumorigenesis. *Br J Cancer*, **76**(9): 1131–3.
- Waldron, D. (2016) Cancer genomics: A multi-layer omics approach to cancer. *Nat Rev Genet*, **17**(8): 436–437.
- Wang, K., Singh, D., Zeng, Z., Coleman, S.J., Huang, Y., Savich, G.L., He, X., Mieczkowski, P., Grimm, S.A., Perou, C.M., *et al.* (2010) MapSplice: accurate mapping of RNA-seq reads for splice junction discovery. *Nucleic Acids Res*, **38**(18): e178.
- Wang, X. and Simon, R. (2013) Identification of potential synthetic lethal genes to p53 using a computational biology approach. *BMC Medical Genomics*, **6**(1): 30.
- Wappett, M. (2014) Bisep: Toolkit to identify candidate synthetic lethality. r package version 2.0.

- Wappett, M., Dulak, A., Yang, Z.R., Al-Watban, A., Bradford, J.R., and Dry, J.R. (2016) Multi-omic measurement of mutually exclusive loss-of-function enriches for candidate synthetic lethal gene pairs. *BMC Genomics*, **17**: 65.
- Warnes, G.R., Bolker, B., Bonebakker, L., Gentleman, R., Liaw, W.H.A., Lumley, T., Maechler, M., Magnusson, A., Moeller, S., Schwartz, M., *et al.* (2015) *gplots: Various R Programming Tools for Plotting Data*. R package version 2.17.0.
- Watts, D.J. and Strogatz, S.H. (1998) Collective dynamics of 'small-world' networks. *Nature*, **393**(6684): 440–2.
- Weinstein, I.B. (2000) Disorders in cell circuitry during multistage carcinogenesis: the role of homeostasis. *Carcinogenesis*, **21**(5): 857–864.
- Weinstein, J.N., Akbani, R., Broom, B.M., Wang, W., Verhaak, R.G., McConkey, D., Lerner, S., Morgan, M., Creighton, C.J., Smith, C., *et al.* (2014) Comprehensive molecular characterization of urothelial bladder carcinoma. *Nature*, **507**(7492): 315–322.
- Weinstein, J.N., Collisson, E.A., Mills, G.B., Shaw, K.R., Ozenberger, B.A., Ellrott, K., Shmulevich, I., Sander, C., Stuart, J.M., Chang, K., *et al.* (2013) The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet*, **45**(10): 1113–1120.
- Wickham, H. and Chang, W. (2016) *devtools: Tools to Make Developing R Packages Easier*. R package version 1.12.0.
- Wickham, H., Danenberg, P., and Eugster, M. (2017) *roxygen2: In-Line Documentation for R*. R package version 6.0.1.
- Wojtukiewicz, M.Z., Hempel, D., Sierko, E., Tucker, S.C., and Honn, K.V. (2016) Thrombin-unique coagulation system protein with multifaceted impacts on cancer and metastasis. *Cancer Metastasis Rev*, **35**(2): 213–233.
- Wong, S.L., Zhang, L.V., Tong, A.H.Y., Li, Z., Goldberg, D.S., King, O.D., Lesage, G., Vidal, M., Andrews, B., Bussey, H., *et al.* (2004) Combining biological networks to predict genetic interactions. *Proceedings of the National Academy of Sciences of the United States of America*, **101**(44): 15682–15687.
- World Health Organization (WHO) (2017) Fact sheet: Cancer. <http://www.who.int/mediacentre/factsheets/fs297/en/>. Updated February 2017, Accessed: 22/03/2017.

- Wu, M., Li, X., Zhang, F., Li, X., Kwoh, C.K., and Zheng, J. (2014) In silico prediction of synthetic lethality by meta-analysis of genetic interactions, functions, and pathways in yeast and human cancer. *Cancer Inform*, **13**(Suppl 3): 71–80.
- Yu, H. (2002) Rmpi: Parallel statistical computing in r. *R News*, **2**(2): 10–14.
- Zhang, F., Wu, M., Li, X.J., Li, X.L., Kwoh, C.K., and Zheng, J. (2015) Predicting essential genes and synthetic lethality via influence propagation in signaling pathways of cancer cell fates. *J Bioinform Comput Biol*, **13**(3): 1541002.
- Zhang, J., Baran, J., Cros, A., Guberman, J.M., Haider, S., Hsu, J., Liang, Y., Rivkin, E., Wang, J., Whitty, B., *et al.* (2011) International cancer genome consortium data portala one-stop shop for cancer genomics data. *Database: The Journal of Biological Databases and Curation*, **2011**: bar026.
- Zhong, W. and Sternberg, P.W. (2006) Genome-wide prediction of c. elegans genetic interactions. *Science*, **311**(5766): 1481–1484.
- Zweig, M.H. and Campbell, G. (1993) Receiver-operating characteristic (roc) plots: a fundamental evaluation tool in clinical medicine. *Clinical Chemistry*, **39**(4): 561–577.