

Literature Values Search

Thomas Locks

March 15, 2024

1 Escape Velocity from Dust Disk

This seems very difficult to find as a single equation. Making a very large assumption that the mass is evenly distributed and spherically distributed, we could just say that V_{esc} in the disk is given by Equation 1.

$$V_{esc} = \sqrt{\frac{2G(M_{star} + M_{disk})}{r_{star} + r_{disk}}} \quad (1)$$

2 Lifetime of Biological Material on Rocks in Space

Ginsburg et al. (2018) gives a link for the survival fraction of biological life to its lifetime:

$$f_{surv,bio}(t) = e^{-\frac{t}{\tau_{bio}}} \quad (2)$$

In Equation 2, τ_{bio} is the biological lifetime, and t is some timescale. Ginsburg et al. (2018) estimates $\tau_{bio} = 10^5$ yr for a polyextremophile, *D. radiodurans*.

Valtonen et al. (2008) gives a link between survival time and the radius of the body the biological material is in. This is seen in Equation 3.

$$t^2(My) \approx 75l^2 \quad (3)$$

Mileikowsky et al. (2000) gives a survival fraction that depends on many unknowns.

Adams & Napier (2022) looks promising. It gives a distribution of rocky objects by mass:

$$\frac{dN}{dm} = Am^{-p} \quad (4)$$

In Equation 4, m is the mass, N is the number of rocks, and p has values between 1 and 2. When the size-mass distribution is ‘determined by collisional processes’, $p \approx 1.8$. This can be linked to rock radius if we have an assumed density for these rocks. A is defined in Adams & Napier (2022).

https://astrostatistics.psu.edu/datasets/asteroid_dens.html#:~:text=However%2C%20the%20internal%20structure%20of,structures%20will%20have%20lower%20density. gives a density of asteroids (‘Oumuamua is said to have the same density – see the Wikipedia page) of 3–5 gcm⁻³.

Adams & Napier (2022) gives a minimum mass for the rocks to harbour biological life as 10 kg.

Adams & Napier (2022) also gives a minimum rock size for the rocks to harbour biological life as a few centimetres.

Linking density and the mass distribution, we can get the radius of these rocks and thus the survival time of the biological material.

Summary:

- Use Equation 3 to get a survival time for the biological material inside the rock. We still need a radius of the rock
- Use Equation 4 to get a distribution of rock masses and thus radius from typical values for asteroid densities.

2.1 Lifetime of These Rocks

If we have this, then we have t .

'Oumuamua is estimated to be in the age range of 0.20–0.45Gyr (Almeida-Fernandes & Rocha-Pinto 2018). Hsieh et al. (2021), however, gives 30Myr.

It would be better to have a time that depends on the star the rock has been ejected from.

References

- Adams, F. C. & Napier, K. J. (2022), 'Transfer of rocks between planetary systems: Panspermia revisited', *Astrobiology* **22**(12), 1429–1442.
- Almeida-Fernandes, F. & Rocha-Pinto, H. (2018), 'A kinematical age for the interstellar object 1i/'oumuamua', *Monthly Notices of the Royal Astronomical Society* **480**(4), 4903–4911.
- Ginsburg, I., Lingam, M. & Loeb, A. (2018), 'Galactic panspermia', *The Astrophysical Journal Letters* **868**(1), L12.
- Hsieh, C.-H., Laughlin, G. & Arce, H. G. (2021), 'Evidence suggesting that 'oumuamua is the 30 myr old product of a molecular cloud', *The Astrophysical Journal* **917**(1), 20.
- Mileikowsky, C., Cucinotta, F. A., Wilson, J. W., Gladman, B., Horneck, G., Lindegren, L., Melosh, J., Rickman, H., Valtonen, M. & Zheng, J. (2000), 'Risks threatening viable transfer of microbes between bodies in our solar system', *Planetary and Space Science* **48**(11), 1107–1115.
- Valtonen, M., Nurmi, P., Zheng, J.-Q., Cucinotta, F. A., Wilson, J. W., Horneck, G., Lindegren, L., Melosh, J., Rickman, H. & Mileikowsky, C. (2008), 'Natural transfer of viable microbes in space from planets in extra-solar systems to a planet in our solar system and vice versa', *The Astrophysical Journal* **690**(1), 210.