

Open Quantum System approach to in medium quarkonium

Tom Magorsch

QCD challenges from pp to AA collisions

03.09.2024





Quarkonium suppression

Propagation through QGP

T. Matsui, H. Satz, *Phys. Lett. B* 178 (1986) 416


$$T \approx O(100\text{MeV})$$



$$V(r) = -\frac{\alpha}{r}$$

↓ Debye-screening in medium

$$V(r) = -\frac{\alpha}{r}e^{-m_D r}$$



- Phenomenological predictions for **bottomonium** from **first principles**

Open Quantum Systems

- Quantum system not isolated
- Split into System S and Environment E


$$H = H_S \otimes I_E + I_S \otimes H_E + H_{\text{int}}$$

- Time evolution by Von-Neumann Equation

$$\frac{d}{dt}\rho = -i[H, \rho]$$

- Not interested in environmental d.o.f.: **Trace out!**

$$\rho_S = \text{Tr}_E[\rho]$$



Open Quantum Systems

- Time evolution by Von-Neumann Equation


$$\frac{d}{dt}\rho = -i[H, \rho]$$

- Environmental d.o.f. not needed Trace out!

$$\rho_S = \text{Tr}_E[\rho]$$

- “Master equation” for the System: **Lindblad Equation**

$$\frac{d\rho_S}{dt} = -i[H_S, \rho_S] + \sum_n \left(C_n \rho_S C_n^\dagger - \frac{1}{2} \{ C_n^\dagger C_n, \rho_S \} \right)$$



Open Quantum Systems

- Time evolution by Von-Neumann Equation


$$\frac{d}{dt}\rho = -i[H, \rho]$$

- Environmental d.o.f. not needed Trace out!

$$\rho_S = \text{Tr}_E[\rho]$$

- “Master equation” for the System: **Lindblad Equation**

$$\frac{d\rho_S}{dt} = -i[H_S, \rho_S] + \sum_n \left(C_n \rho_S C_n^\dagger - \frac{1}{2} \{ C_n^\dagger C_n, \rho_S \} \right)$$




Open Quantum Systems

- Time evolution by Von-Neumann Equation

$$\frac{d}{dt}\rho = -i[H, \rho]$$

- Environmental d.o.f. not needed Trace out!

$$\rho_S = \text{Tr}_E[\rho]$$




- “Master equation” for the System: **Lindblad Equation** *non-unitary*

$$\frac{d\rho_S}{dt} = -i[H_S, \rho_S] + \sum_n \left(C_n \rho_S C_n^\dagger - \frac{1}{2} \{ C_n^\dagger C_n, \rho_S \} \right)$$

OQS for quarkonium


- Quarkonium: System S
- QGP: Environment E

Aim to describe Quarkonium Suppression by
a master equation for encoding the
interaction with the QGP




$$\frac{d\rho_S}{dt} = -i[H_S, \rho_S] + \sum_n \left(C_n \rho_S C_n^\dagger - \frac{1}{2} \{ C_n^\dagger C_n, \rho_S \} \right)$$

Master equations from EFTs



Master equations from EFTs



- Degrees of Freedom: Heavy quarks

Lindblad eq. for *Heavy quarks*:


Akamatsu, Y. (2015). *Physical Review D*, 91(5), 056002.

Lindblad eq. for *Heavy quarkonium*:

Blaizot, J. P., & Escobedo, M. A. (2018). *Journal of High Energy Physics*, 2018(6), 1-57.

Blaizot, J. P., & Escobedo, M. A. (2018). *Physical Review D*, 98(7), 074007.

Master equations from EFTs



- Degrees of Freedom: Singlet and octet bound states

Lindblad eq. for *Heavy quarkonium*:

Brambilla, N., Escobedo, M. A., Soto, J., & Vairo, A. (2018).
Physical Review D, 97(7), 074009.

Brambilla, N., Escobedo, M. A., Soto, J., & Vairo, A. (2017).
Physical Review D, 96(3), 034021.

pNRQCD master equation

Brambilla, Escobedo, Soto, Vairo: Phys. Rev. D 97 (2018) 7, 074009

$$\frac{d\rho(t)}{dt} = -i [H, \rho(t)] + \sum_{nm} h_{nm} \left(L_i^n \rho(t) L_i^{m\dagger} - \frac{1}{2} \left\{ L_i^{m\dagger} L_i^n, \rho(t) \right\} \right)$$

$$h = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

$$L_i^0 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} r_i, \quad L_i^1 = \begin{pmatrix} 0 & 0 \\ 0 & \frac{N_c^2 - 4}{2(N_c^2 - 1)} A_i^{oo\dagger} \end{pmatrix}$$

$$L_i^2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} r_i, \quad L_i^3 = \begin{pmatrix} 0 & \frac{1}{N_c^2 - 1} A_i^{os\dagger} \\ A_i^{so\dagger} & 0 \end{pmatrix}$$

$$A_i^{uv} = \frac{g^2}{6N_c} \int_0^\infty ds e^{-ih_u s} r_i e^{ih_v s} \langle E_j^a(s) \Omega(s, 0)^{ab} E_j^b(0) \rangle$$

pNRQCD master equation

Brambilla, Escobedo, Soto, Vairo: Phys. Rev. D 97 (2018) 7, 074009

$$\frac{d\rho(t)}{dt} = -i [H, \rho(t)] + \sum_{nm} h_{nm} \left(L_i^n \rho(t) L_i^{m\dagger} - \frac{1}{2} \left\{ L_i^{m\dagger} L_i^n, \rho(t) \right\} \right)$$

- In general h_{nm} not completely positive:

Master equation *not necessarily* of Lindblad type

$$L_i^0 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} r_i, \quad L_i^1 = \begin{pmatrix} 0 & 0 \\ 0 & \frac{N_c^2 - 4}{2(N_c^2 - 1)} A_i^{oo\dagger} \end{pmatrix}$$

$$L_i^2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} r_i, \quad L_i^3 = \begin{pmatrix} 0 & \frac{1}{N_c^2 - 1} A_i^{os\dagger} \\ A_i^{so\dagger} & 0 \end{pmatrix}$$

$$h = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

E/T expansion

- Simplify using hierarchy of scales $\pi T \gg E$

$$A_i^{uv} = \frac{g^2}{6N_c} \int_0^\infty ds e^{-ih_us} r_i e^{ih_vs} \langle E_j^a(s) \Omega(s, 0)^{ab} E_j^b(0) \rangle$$

- Expand exponentials in $E/(\pi T)$
- At **LO** in $E/(\pi T)$ we get

$$\begin{aligned} A_i^{uv} &= \frac{g^2}{6N_c} \int_0^\infty ds r_i \langle E_j^a(s) \Omega(s, 0)^{ab} E_j^b(0) \rangle \\ &= \frac{r_i}{2} (\kappa - i\gamma) \end{aligned}$$

Transport coefficients

Viljamis Talk

pNRQCD LO Lindblad equation

$$A_i^{uv} = \frac{g^2}{6N_c} \int_0^\infty ds r_i \langle E_j^a(s) \Omega(s, 0)^{ab} E_j^b(0) \rangle$$

$$= \frac{r_i}{2} (\kappa - i\gamma)$$

Transport
coefficients

$$L_i^0 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} r_i, \quad L_i^1 \propto \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} r_i,$$

$$L_i^2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} r_i, \quad L_i^3 \propto \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} r_i,$$

Linearly dependent:
Lindblad possible!

pNRQCD LO Lindblad equation

$$\frac{d\rho(t)}{dt} = -i[H, \rho(t)] + \sum_n \left[C_i^n \rho(t) C_i^{n\dagger} - \frac{1}{2} \left\{ C_i^{n\dagger} C_i^n, \rho(t) \right\} \right],$$

$$H = \begin{pmatrix} h_s + \frac{r^2}{2}\gamma & 0 \\ 0 & h_o + \frac{N_c^2 - 2}{2(N_c^2 - 1)} \frac{r^2}{2}\gamma \end{pmatrix}$$

$$h_{s,o} = \vec{p}^2 / M + V_{s,o}$$

pNRQCD LO Lindblad equation

$$\frac{d\rho(t)}{dt} = -i[H, \rho(t)] + \sum_n \left[C_i^n \rho(t) C_i^{n\dagger} - \frac{1}{2} \left\{ C_i^{n\dagger} C_i^n, \rho(t) \right\} \right],$$

$$H = \begin{pmatrix} h_s + \frac{r^2}{2}\gamma & 0 \\ 0 & h_o + \frac{N_c^2 - 2}{2(N_c^2 - 1)} \frac{r^2}{2}\gamma \end{pmatrix}$$

$$h_{s,o} = \vec{p}^2 / M + V_{s,o}$$

Quarkonium
Potential

pNRQCD LO Lindblad equation

$$\frac{d\rho(t)}{dt} = -i[H, \rho(t)] + \sum_n \left[C_i^n \rho(t) C_i^{n\dagger} - \frac{1}{2} \left\{ C_i^{n\dagger} C_i^n, \rho(t) \right\} \right],$$

$$H = \begin{pmatrix} h_s + \frac{r^2}{2}\gamma & 0 \\ 0 & h_o + \frac{N_c^2 - 2}{2(N_c^2 - 1)} \frac{r^2}{2}\gamma \end{pmatrix} \quad C_i^0 = \sqrt{\frac{\kappa}{N_c^2 - 1}} r_i \begin{pmatrix} 0 & 1 \\ \sqrt{N_c^2 - 1} & 0 \end{pmatrix},$$

$$C_i^1 = \sqrt{\frac{\kappa(N_c^2 - 4)}{2(N_c^2 - 1)}} r_i \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

$$h_{s,o} = \vec{p}^2 / M + V_{s,o}$$

pNRQCD LO Lindblad equation

$$\frac{d\rho(t)}{dt} = -i[H, \rho(t)] + \sum_n \left[C_i^n \rho(t) C_i^{n\dagger} - \frac{1}{2} \left\{ C_i^{n\dagger} C_i^n, \rho(t) \right\} \right],$$


$$H = \begin{pmatrix} h_s + \frac{r^2}{2}\gamma & 0 \\ 0 & h_o + \frac{N_c^2 - 2}{2(N_c^2 - 1)} \frac{r^2}{2}\gamma \end{pmatrix} \quad C_i^0 = \sqrt{\frac{\kappa}{N_c^2 - 1}} r_i \begin{pmatrix} 0 & 1 \\ \sqrt{N_c^2 - 1} & 0 \end{pmatrix},$$

$$C_i^1 = \sqrt{\frac{\kappa(N_c^2 - 4)}{2(N_c^2 - 1)}} r_i \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

$$h_{s,o} = \vec{p}^2 / M + V_{s,o}$$

Transport
coefficients


pNRQCD NLO Lindblad equation

$$A_i^{uv} = \frac{g^2}{6N_c} \int_0^\infty ds e^{-ih_us} r_i e^{ih_vs} \langle E_j^a(s) \Omega(s, 0)^{ab} E_j^b(0) \rangle$$

$$1 + ih_vs$$

$$A_i^{uv} = \frac{r_i}{2} (\kappa - i\gamma) + \kappa \left(-\frac{ip_i}{2MT} + \frac{\Delta V_{uv}}{4T} r_i \right) + \dots$$

pNRQCD NLO Lindblad equation

$$A_i^{uv} = \frac{g^2}{6N_c} \int_0^\infty ds e^{-ih_us} r_i e^{ih_vs} \langle E_j^a(s) \Omega(s, 0)^{ab} E_j^b(0) \rangle$$



$$1 + ih_vs$$

$$A_i^{uv} = \frac{r_i}{2} (\kappa - i\gamma) + \kappa \left(-\frac{ip_i}{2MT} + \frac{\Delta V_{uv}}{4T} r_i \right) + \dots$$

$$L_i^0 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} r_i, \quad L_i^1 = \begin{pmatrix} 0 & 0 \\ 0 & \frac{N_c^2 - 4}{2(N_c^2 - 1)} A_i^{oo\dagger} \end{pmatrix} \quad \text{Not linearly dependent!}$$

But corrections of order $E^2 / (\pi T)^2$

pNRQCD LO Lindblad equation

$$\frac{d\rho(t)}{dt} = -i[H, \rho(t)] + \sum_n \left[C_i^n \rho(t) C_i^{n\dagger} - \frac{1}{2} \left\{ C_i^{n\dagger} C_i^n, \rho(t) \right\} \right],$$

$$H = \begin{pmatrix} h_s + \frac{r^2}{2}\gamma + \frac{\kappa}{4MT} \{r_i, p_i\} & 0 \\ 0 & h_o + \frac{N_c^2 - 2}{2(N_c^2 - 1)} \left(\frac{r^2}{2}\gamma + \frac{\kappa}{4MT} \{r_i, p_i\} \right) \end{pmatrix}$$

$$C_i^0 = \sqrt{\frac{\kappa}{N_c^2 - 1}} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \left(r_i + \frac{ip_i}{2MT} + \frac{\Delta V_{os}}{4T} r_i \right) \quad C_i^1 = \sqrt{\frac{\kappa(N_c^2 - 4)}{2(N_c^2 - 1)}} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \left(r_i + \frac{ip_i}{2MT} \right) \\ + \sqrt{\kappa} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \left(r_i + \frac{ip_i}{2MT} + \frac{\Delta V_{so}}{4T} r_i \right),$$

pNRQCD master equation

$$\frac{d\rho(t)}{dt} = -i[H, \rho(t)] + \sum_n \left[\cancel{C_i^n} \rho(t) \cancel{C_i^{n\dagger}} - \frac{1}{2} \left\{ \cancel{C_i^{n\dagger}} \cancel{C_i^n}, \rho(t) \right\} \right],$$

Projection on spherical harmonics



pNRQCD master equation

$$\frac{d\rho(t)}{dt} = -i[H, \rho(t)] + \sum_n \left[\cancel{C_i^n} \rho(t) \cancel{C_i^{n\dagger}} - \frac{1}{2} \left\{ \cancel{C_i^{n\dagger}} \cancel{C_i^n}, \rho(t) \right\} \right],$$

Hilbert space:

$$H_c \otimes H_l \otimes H_r$$

$\mathbf{c} = \mathbf{s}, \mathbf{0}$ $\mathbf{l} = 0, 1, 2, \dots$ Radial wavefunction
e.g. 2000 point lattice

Quantum trajectory algorithm

J. Dalibard, Y. Castin, and K. Mølmer, Wave-function approach to dissipative processes in quantum optics, Phys. Rev. Lett. 68 (1992), pp. 580–583.

- Idea:

1. Evolve individual trajectories $|\phi(t)\rangle$ stochastically


can evolve to arbitrary l

2. Calculate observables by averaging over trajectories $\overline{\langle \phi(t) | A | \phi(t) \rangle}$

Advantages:

- Evolve vector of size N_H instead N_H^2 density matrix
- Simulation of individual trajectories is **embarrassingly parallel**

Speedup



Qtraj!

Omar, H. B., Escobedo, M. Á., Islam, A., Strickland, M., Thapa, S., Vander Griend, P., & Weber, J. H. (2022). *Computer Physics Communications*, 273, 108266.

Connecting with pheno



- Initial state: Localized gaussian peak
- temperature evolution from hydrodynamics simulation

M. Alqahtani and M. Strickland, The European Physical Journal C 81 (2021)



$$\text{Survival Probability} = \frac{\langle \psi(t) | 1S \rangle}{\langle \psi(0) | 1S \rangle}$$

- Including Feed-down from PDG data

LO results





NLO results




NLO results

Michael Strickland, Sabin Thapa, PHYSICAL
REVIEW D 108, 014031 (2023)



New Potential





Spectrum:

	PDG	V_s^c	V_s^{3L}
$M(1S)/\text{GeV}$	9.445	9.445	9.445
$M(2S)/\text{GeV}$	10.017	9.635	10.042
$M(3S)/\text{GeV}$	10.355	9.670	10.395
$M(1P)/\text{GeV}$	9.888	9.635	9.887
$M(2P)/\text{GeV}$	10.251	9.670	10.279



Determination of transport coefficients

- Indirectly determine $\hat{\kappa}$ and $\hat{\gamma}$ from lattice measurements of the **in medium width** Γ and **mass shift** δm



Determination of transport coefficients

- Obtain $\hat{\kappa}$ from fits to $1S$ and $1P$ data and average




Coulomb: $\hat{\kappa} = 0.33 \pm 0.04$

New potential: $\hat{\kappa} = 1.88 \pm 0.16$



Nuclear modification factor results

- New potential can describe the experimental data
- Coulomb potential with $\hat{\kappa} = 0.33 \pm 0.04$ can not describe the data



Summary & Future challenges

- Progressing towards a more and more mature simulation framework
- Challenge assumptions:
 1. Markovian approximation
 2. Assumption of isotropy
 3. Estimate uncertainties from E/T expansion
- Charmonium ?



Backup slides

Quantum Trajectories

$$U(\Theta) = 1 - iH_{\text{eff}}\delta t$$

ψ_0
↓

1. Evolve state $|\psi(t)\rangle$ with $U(\Theta)$

$$|\psi(t + \delta t)\rangle = U(\Theta)|\psi(t)\rangle$$


2. Compute norm

$$\langle\psi(t + \delta t)|\psi(t + \delta t)\rangle = 1 - \delta p(\Theta) < 1$$

3. Apply jump operator $C(\Theta)$ with probability $\delta p(\Theta)$


$$|\psi(t + \delta t)\rangle = C(\Theta)|\psi(t)\rangle$$

4. Normalize $|\psi(t + \delta t)\rangle$



Non perturbative correction

N. Brambilla, M. A. Escobedo, M. Strickland,
A. Vairo, P. Vander Griend, and J. H. Weber,
JHEP 05, 136 (2021), 2012.01240



Heavy quark diffusion coefficient

$$V_s^{\text{non-pert}}(r) = -i \frac{g^2 T_F}{3N_c} r^2 \int_0^\infty dt \langle E^a(t) \Omega(t, 0)^{ab} E^b(0) \rangle$$

$$\gamma = \frac{g^2}{3N_c} \text{Im} \int_0^\infty dt \langle E^a(t) \Omega(t, 0)^{ab} E^b(0) \rangle$$

In medium width

- Width given by collapse operators

$$\Gamma = \sum_n C_n^\dagger C_n$$

- At LO in E/T

$$\Gamma = \hat{\kappa} T^3 r^2$$

Determination of transport coefficients

- Indirectly determine $\hat{\kappa}$ and $\hat{\gamma}$ from lattice measurements of the **in medium width** Γ and **mass shift** δm

$$\kappa = \hat{\kappa} T^3$$

no vacuum
part

$$\gamma = \gamma(T = 0) + \hat{\gamma} T^3$$

- Assume simple model for the vacuum part $\gamma(T = 0)$

$$\langle E^a(t) \Omega(t, 0)^{ab} E^b(0) \rangle = \langle E^2(0) \rangle e^{-i\Lambda_E t}$$

G. S. Bali and A. Pineda, Physical Review D 69 (2004)

$$\langle g^2 E^2(0) \rangle = -0.2 \text{ GeV}^4 \quad \Lambda_E = 1.25 \text{ GeV}$$

EFTs for Quarkonium Suppression

- Use NREFTs to exploit hierarchy of scales

$$M \gg 1/a_0 \gg \pi T \gg E$$

- Inverse radius:


$$1/a_0 \approx 1.2\text{GeV}$$

- Temperature regime:

$$250\text{MeV} < T < 425\text{MeV}$$

- Binding Energy:

$$E \sim 0.4\text{GeV}$$



New Potential

Kiyo, Y., Pineda, A., & Signer, A. (2010). *Nuclear Physics B*, 841(1-2), 231-256.

- Motivation: Implement a higher order potential with a more realistic spectrum

$$V_s^{3L}(r) = V_s^{\text{pert}}(r) + V_s^{\text{non-pert}}(r)$$

$$V_s^{\text{pert}}(\nu, \nu_r, r) = \begin{cases} \sum_{k=0}^3 V_{s,\text{RS}}^{(k)} \alpha_s^{k+1}(1/r) & \text{if } r < \nu_r^{-1} \\ \sum_{k=0}^3 V_{s,\text{RS}}^{(k)} \alpha_s^{k+1}(\nu) & \text{if } r > \nu_r^{-1} \end{cases}$$

three loop pNRQCD

$$\text{Re} \left(V_s^{\text{non-pert}}(r) \right) = \frac{\gamma}{2} r^2$$

leading non-perturbative correction