

icequake_example_time_domain

February 17, 2021

1 Example usage of SeisSrcMoment to calculate M_W in the time domain

This example is for an icequake at Vatnajokull, Iceland. The moment magnitude, M_W , is calculated in the time domain, i.e. the long period spectral level is calculated from the integral of the displacement through time.

1.1 1. Specify parameters to use:

```
[1]: import numpy as np
      from SeisSrcMoment import moment
```

```
[2]: # Specify variables:
stations_to_calculate_moment_for = ["SKR01", "SKR02", "SKR03", "SKR04",
    ↪ "SKR05", "SKR06", "SKR07"]
stations_not_to_process = ["SKG08", "SKG09", "SKG10", "SKG11", "SKG12",
    ↪ "SKG13", "GR01", "GR02", "GR03", "GR04", "BARD"]
mseed_filename = "data/mseed_data/20140629184210331.m"
inventory_fname = None
instruments_gain_filename = "data/instrument_gain_data.txt" # File with
    ↪ instrument name, instrument gains (Z,N,E) and digitaliser gains (Z,N,E)
NLLoc_event_hyp_filename = "data/NLLoc_data/loc.Tom_RunNLLoc000.20140629.
    ↪ 184210.grid0.loc.hyp"
window_before_after = [0.004, 0.196] # The time before and after the phase pick
    ↪ to use for calculating the magnitude within
filt_freqs = []
# MT_data_filename = "data/MT_data/20140629184210363MT.mat"
MT_six_tensor = np.array([1.,1.,1.,0.,0.,0.]) # Explosion in this example.
density = 917. # Density of medium, in kg/m3
Vp = 3630. # P-wave velocity in m/s
Q = 150. # Quality factor for the medium
use_full_spectral_method = False # If False, will use the time domain
    ↪ displacement integration method
verbosity_level = 1 # Verbosity level (1 for moment only) (2 for major
    ↪ parameters) (3 for plotting of traces)
```

1.2 Run moment calculation:

```
[3]: # Find seismic moment release:
av_M_0, std_err_seis_M_0, n_obs, event_obs_dict = moment.
    ↪ calc_moment(mseed_filename, NLLoc_event_hyp_filename,
    ↪ stations_to_calculate_moment_for, density, Vp,
    ↪ inventory_fname=inventory_fname,
    ↪ instruments_gain_filename=instruments_gain_filename, Q=Q,
    ↪ window_before_after=window_before_after, filt_freqs=filt_freqs,
    ↪ use_full_spectral_method=use_full_spectral_method,
    ↪ stations_not_to_process=stations_not_to_process,
    ↪ MT_six_tensor=MT_six_tensor, verbosity_level=verbosity_level)
print("Seismic moment release (Nm):", av_M_0)
```

```
Processing data for station: SKR01
Overall seismic moment (Nm): 7860258.2629
Processing data for station: SKR02
Overall seismic moment (Nm): 7596240.15988
Processing data for station: SKR03
Overall seismic moment (Nm): 5185040.49289
Processing data for station: SKR04
Overall seismic moment (Nm): 3427612.88854
Processing data for station: SKR05
Overall seismic moment (Nm): 4675845.68622
Processing data for station: SKR06
Overall seismic moment (Nm): 3582958.1162
Processing data for station: SKR07
Overall seismic moment (Nm): 7224947.64632
Average seismic moment for event: 5650414.75042 +/- 662869.178247
Seismic moment release (Nm): 5650414.75042
```

```
[4]: # And find corresponding moment magnitude, Mw (Hanks and Kanamori 1979):
M_w = (2./3.)*np.log10(av_M_0) - 6.0
print("Local moment magnitude, M:", M_w)
```

```
Local moment magnitude, M: -1.49861311535
```