

Mass Transfer Rates

Kavli Summer Program

TOM WAGG

Let's consider what the mass transfer rates for our stars should be. First let's just write some of the basic timescales. The nuclear timescale we can reduce assuming H core fusion and the mass-luminosity relation.

$$\tau_{\text{nuc}} \sim \frac{M f_M c^2 (\Delta m/m)}{L} \quad (1)$$

$$\sim 7.1 \times 10^{-4} \frac{M c^2}{L} \quad (2)$$

$$\sim 7.1 \times 10^{-4} \frac{M_{\odot} c^2}{1.4 L_{\odot}} \left(\frac{M}{M_{\odot}} \right)^{-2.5} \quad (3)$$

$$\boxed{\tau_{\text{nuc}} \sim 7.5 \times 10^9 \text{ yr} \left(\frac{M}{M_{\odot}} \right)^{-2.5}} \quad (4)$$

Then the thermal timescale is approximately:

$$\tau_{\text{therm}} \sim \frac{GM^2}{RL} \quad (5)$$

$$\sim \frac{G M_{\odot}^2}{1.4 R_{\odot} L_{\odot}} \left(\frac{R}{R_{\odot}} \right)^{-1} \left(\frac{M}{M_{\odot}} \right)^{-1.5} \quad (6)$$

$$\boxed{\tau_{\text{therm}} \sim 2.2 \times 10^7 \text{ yr} \left(\frac{R}{R_{\odot}} \right)^{-1} \left(\frac{M}{M_{\odot}} \right)^{-1.5}} \quad (7)$$

Now we can consider some different limits. Assuming that we accrete on a timescale of τ_{accrete} (and that $\tau_{\text{therm}} \ll \tau_{\text{nuc}}$):

- $\tau_{\text{accrete}} \gg \tau_{\text{nuc}}$: If the accretion is much slower than the nuclear timescale then the star will exhaust a significant fraction (or all!) of its central hydrogen before accretion can complete. This is not very realistic.
- $\tau_{\text{accrete}} \sim \tau_{\text{nuc}}$:
- $\tau_{\text{accrete}} \sim \tau_{\text{therm}}$:
- $\tau_{\text{accrete}} \ll \tau_{\text{therm}}$: