

COMPARISON OF STELLAR COMPANION EJECTION VELOCITIES FROM BINARIES EVOLVED IN DIFFERENT POPULATION SYNTHESIS CODES

TOM WAGG ¹, MATHIEU RENZO ², KATELYN BREIVIK ³, AND DAVID D. HENDRIKS ⁴

¹Department of Astronomy, University of Washington, Seattle, WA, 98195

²McWilliams Center for Cosmology and Astrophysics, Department of Physics, Carnegie Mellon University, Pittsburgh, PA 15213, USA

³University of Arizona, Department of Astronomy & Steward Observatory, 933 N. Cherry Ave., Tucson, AZ 85721, USA and

⁴Department of Physics, University of Surrey, Guildford GU2 7XH, Surrey, UK

Version March 29, 2025

ABSTRACT

We present a comparison of the predicted ejection velocities of disrupted binary companions in three different population synthesis codes: **COSMIC**, **COMPAS** and **binary_c**, which use two independent algorithms. We confirm that, despite the codes producing different pre-supernova orbital evolution from the same initial conditions, they each find the ejection velocities of secondary stars from disrupted binaries are narrowly distributed about their pre-supernova orbital velocity. We additionally include a correction to the derivation included in [Kiel & Hurley \(2009\)](#) that brings it into agreement with methods from other works for determining post-supernova binary orbital parameters. During this comparison we identified and resolved bugs in the kick prescriptions of *all three* codes we considered, highlighting how open-science practices and code comparisons are essential for addressing implementation issues.

1. INTRODUCTION

Observations of pulsars in the Milky Way necessitate that they have a high velocity dispersion and hence that these objects acquire high velocities upon formation (e.g., [Gunn & Ostriker 1970](#); [Hobbs et al. 2005](#); [Igoshev 2020](#)). For compact objects formed via isolated binary evolution, these velocities are expected to be imparted at the moment of core-collapse. Even for a symmetric explosion, the instantaneous mass loss from a supernova (SN) changes the post-SN orbital parameters ([Blaauw 1961](#); [Boersma 1961](#)). This is typically insufficient to disrupt the orbit, since the ejecta mass is often limited by loss during previous mass transfer phases, but in some cases ($\sim 16\%$) mass loss can unbind the binary and result in both binary constituents being ejected with their pre-SN orbital velocity ([Renzo et al. 2019](#)). Moreover, asymmetry in the explosion can impart a significant natal kick on the newly formed compact object (e.g., [Shklovskii 1970](#); [Lyne & Lorimer 1994](#); [Janka 2013, 2017](#)), which in the majority of cases is strong enough to disrupt a binary (e.g., [Renzo et al. 2019](#)). Similar asymmetry in neutrino emission can result in additional natal kicks, which can dominate in the case of full fallback (e.g., [Janka & Mueller 1994](#); [Burrows & Hayes 1996](#); [Scheck et al. 2006](#); [Wongwathanarat et al. 2013](#); [Coleman & Burrows 2022](#); [Vigna-Gómez et al. 2024](#)).

The impact of these SNe kicks on the post-SN orbital parameters of a binary system has been considered in detail by many previous works ([Flannery & van den Heuvel 1975](#); [Sutantyo 1978](#); [Hills 1983](#); [Dewey & Cordes 1987](#); [Wijers et al. 1992](#); [Brandt & Podsiadlowski 1995](#); [Kalogera 1996](#); [Tauris & Takens 1998](#); [Hurley et al. 2002](#); [Pfahl et al. 2002](#); [Belczynski et al. 2008](#); [Kiel & Hurley 2009](#)). These investigations account for a variety of effects, with the most extensive handling eccentric pre-SN

systems, including the effect of instantaneous symmetric mass loss, the impact of the natal kick on both the compact object and disruption of the system, the SN blast wave impulse on the secondary star and collisions between the compact object and star. For a detailed derivation, which explains how each of these effects impacts a binary in clear and compact vector notation, we refer the reader to Appendix B of [Pfahl et al. \(2002\)](#).

Binary stellar population synthesis codes apply different choices of these SN kick methodologies in their evolution. These difference have the potential to significantly change the subsequent evolution of binary systems after a SN. The post-SN eccentricity, semi-major axis and centre-of-mass velocity of bound systems, as well as rate of unbinding and ejection velocities of stars and compact objects, are sensitive to kick prescriptions. The impact on population synthesis predictions is therefore wide-ranging, for both bound (such as X-ray binaries and gravitational-wave sources) and unbound (such as runaway stars) binary products. In this paper we focus on how the ejections velocities of stellar companions compare for a binary with identical initial conditions evolved in different codes.

2. SECONDARY EJECTION VELOCITY COMPARISONS

In this Section we compare the ejection velocities for secondary stars as calculated by different population synthesis codes. The **COSMIC** population synthesis code ([Breivik et al. 2020a](#)) applies the method from [Pfahl et al. \(2002\)](#) (though previously followed [Kiel & Hurley \(2009\)](#), see Section 3). Similarly, **COMPAS** ([Team COMPAS: Riley, J. et al. 2022](#); [Compas et al. 2022](#)) uses the method of [Pfahl et al. \(2002\)](#). In contrast, **binary_c** ([Izzard et al. 2004, 2006, 2009, 2018](#); [Izzard & Jermyn 2023](#)) applies [Tauris & Takens \(1998\)](#). The combination of these codes therefore allows us to cross-validate




FIG. 1.— Secondary ejection velocities closely follow pre-SN orbital velocities, with extremely weak dependence on natal kick magnitudes, across different population synthesis codes. These plots compare a binary with identical initial conditions, evolved 50,000 times with different random natal kicks in three codes. For each code, the main panel shows the difference between the secondary ejection velocity and its pre-SN orbital velocity on the x -axis, while the y -axis shows the natal kick applied to the primary star. Each main panel shows a 2D kernel density estimator with a colourbar indicating the fraction of the population contained within each contour. Secondary panels show marginal histograms. Small black crosses and pluses indicate the mean and median of each distribution respectively. Panels are annotated with the population synthesis code used for evolution and information about the pre- and post-SN system.

three implementations of two different algorithms for determining secondary ejection velocities.

We evolve a representative mass-transferring massive binary with a range of random SN natal kicks in each code. This binary has the following initial conditions: primary mass, $m_1 = 20 M_{\odot}$, secondary mass $m_2 = 15 M_{\odot}$, orbital period, $P = 100$ days, eccentricity, $e = 0$, and metallicity, $Z = 0.02$. This system is representative of O-type stars that undergo case B mass transfer, which often produce ejected secondaries. We repeat the evolution 50,000 times, each with a different random natal kick magnitude and direction. For each code we draw natal kick magnitudes from a Maxwellian with $\sigma = 265 \text{ km s}^{-1}$ and apply the [Fryer et al. \(2012\)](#) delayed remnant mass

prescription, but otherwise use the code’s default settings.

The difference in binary physics settings between the codes results in moderately different pre-SN evolutionary states. This is likely in large part due to the treatment of wind mass loss and mass transfer. It may be possible to enforce agreement between codes through a detailed selection of binary physics settings but that is beyond the scope of this work.

The ejection velocities of the secondary star closely follow their pre-SN orbital velocities in each code, and have only a very weak dependence on the magnitude of natal kicks. In Figure 1, we show how the applied natal kick affects the difference between the pre- and post-SN ve-

locity of the secondary star in these simulations. The main panels show a 2D kernel density estimator for the disrupted binaries in the 50,000 realisations. Quantitatively, we find that, for each code, the ejection velocity of the secondary star is within $\sim 5 \text{ km s}^{-1}$ of its pre-SN orbital velocity for natal kicks spanning 3 orders of magnitude.

The independence of the secondary’s ejection velocity and the natal kick applied to the primary is expected. Due to the strength of the natal kicks, the secondary star experiences a near instantaneous disappearance of the compact object from its orbit, removing the centripetal acceleration and leading to its ejection at close to its pre-SN orbital velocity (see [Tauris & Takens 1998](#) for a more in-depth discussion). Following the same logic, stronger natal kicks more immediately remove the compact object from the system, therefore leading to a tighter relationship between the pre- and post-SN velocities as shown in Figure 1.

The agreement between the codes validates their implementation of the SN kick routines. Although each code produces a markedly different pre-SN evolution of the system (compare the annotations in each plot in Figure 1), the relationship between the pre- and post-SN velocities holds.

3. CORRECTION TO KIEL & HURLEY (2009)

Earlier versions of COSMIC ($< \text{v}3.5.0$) used the methodology described by [Kiel & Hurley \(2009\)](#) in calculating the post-SN orbital properties of binaries, as well as the ejection velocities of compact objects and their prior companions. However, when using this method, we found that the ejection velocities of companions were not strongly correlated with their pre-SN orbital velocities as one would expect. Instead, their ejection velocity showed an (unphysical) strong dependence on the natal kick received by the primary. Moreover, the ejection velocity of the compact object was often significantly slower than the natal kick that it received.

The reason for these effects is that, in [Kiel & Hurley \(2009\)](#), the z component of the final velocity of each binary constituent neglects the contributions from the velocity at infinity of the hyperbolic orbit, V_∞ . The derivation assumes that this velocity is directed entirely within the orbital plane of the binary, but does not fully account for the fact that the orbital plane is no longer aligned with the z axis after the SN. Therefore, the z component of the final velocity of the compact object (equation 17 from [Kiel & Hurley 2009](#)) should be as follows

$$V_{1z} = \frac{M_{\text{NS}}}{M'_b} V_{\text{kick}} \sin \phi + V_\infty \sin \nu \sin \gamma \frac{M_2}{M'_b}, \quad (1)$$

where the variables are defined in the same way as [Kiel & Hurley \(2009\)](#), such that M_{NS} is the mass of the resulting compact object, M'_b is the final total mass of the system, V_{kick} is the magnitude of natal kick, ϕ is the polar angle of the natal kick, V_∞ is the velocity at infinity of the hyperbolic orbit and ν and γ are angles accounting for the misalignment of the orbital plane after the SN (see Section 2.2.1 and Figure 2 of [Kiel & Hurley 2009](#)). Similarly the z component of the final velocity of the companion (equation 20 from [Kiel & Hurley 2009](#)) should be as fol-

lows

$$V_{2z} = \frac{M_{\text{NS}}}{M'_b} V_{\text{kick}} \sin \phi - V_\infty \sin \nu \sin \gamma \frac{M_{\text{NS}}}{M'_b}. \quad (2)$$

The latest versions of COSMIC ($\geq \text{v}3.5.0$) now use the [Pfahl et al. \(2002\)](#) method by default. We make the original [Kiel & Hurley \(2009\)](#) method accessible (via negative `kickflag` inputs) so that earlier COSMIC results can be reproduced. We have confirmed that the inclusion of the additional V_∞ terms in these equations brings the [Kiel & Hurley \(2009\)](#) method into agreement with that of [Pfahl et al. \(2002\)](#) and [Tauris & Takens \(1998\)](#). We note that this issue only affected the ejection velocities of stars and compact objects from *disrupted* systems, but the conditions for disruption and post-SN orbital parameters for bound systems were correct. Therefore prior work using COSMIC to explore bound systems remains unchanged.

4. CODE ISSUES IDENTIFIED AND RESOLVED AS A RESULT OF COMPARISON

During the course of this comparison we identified problems with either the kick prescription or implementation in each of the codes that we considered. As discussed above, COSMIC previously used an erroneous kick prescription, which has been resolved as of $\text{v}3.5.0^1$. The implementation in COMPAS had anomalous component velocities such that the ejection velocity of secondary stars was consistently overestimated by a factor of 2. This problem arose as a result of spurious vector indexing and incorrect rotation algebra in the code. This issue has been resolved as of COMPAS $\text{v}03.09.02^2$. In `binary_c`, the prior implementation did not sample natal kick magnitudes and azimuthal angles in an independent manner, which led to slightly lower ejection velocities on average. This correlated sampling occurred as a result of re-using a randomly generated number from the kick magnitude in the azimuthal angle sampling. This issue has been resolved as of `binary_c` **VERSIONNUMBER**³.

Despite the rigorous testing suites of each code that aim to address issues such as these, each bug was not uncovered until comparison with other open-source codes. This highlights the fact that open-science, and comparison projects such as these, are essential to identifying and resolving subtle issues in population synthesis.

ACKNOWLEDGEMENTS

TW and KB acknowledge support from NASA ATP grant 80NSSC24K0768.

Software: This work made use of the following software packages: `astropy` ([Astropy Collaboration et al. 2013, 2018, 2022](#)), `Jupyter` ([Perez & Granger 2007](#); [Kluyver et al. 2016](#)), `matplotlib` ([Hunter 2007](#)), `numpy` ([Harris et al. 2020](#)), `pandas` ([Wes McKinney 2010](#); [pandas development team 2024](#)), `python` ([Van Rossum & Drake 2009](#)), `scipy` ([Virtanen et al. 2020](#); [Gommers et al. 2024](#)), COSMIC ([Breivik et al. 2020b](#); [Coughlin](#)

¹<https://github.com/COSMIC-PopSynth/COSMIC/pull/678>

²<https://github.com/TeamCOMPAS/COMPAS/pull/1290>

³https://gitlab.com/binary_c/binary_c/-/commit/fba20687984e9cfb46bc38838d07655bfc99c6e5

et al. 2024), Cython (Behnel et al. 2011), h5py (Collette 2013; Collette et al. 2023), schwimmbad (Price-Whelan & Foreman-Mackey 2017), seaborn (Waskom 2021), and tqdm (da Costa-Luis et al. 2024). This research has made use of NASA’s Astrophysics Data System. Simulations in this paper made use of the COMPAS rapid binary population synthesis code (version 3.01.10), which

is freely available at <http://github.com/TeamCOMPAS/COMPAS> (Team COMPAS: Riley, J. et al. 2022). Software citation information aggregated using The Software Citation Station (Wagg & Broekgaarden 2024; Wagg et al. 2024).

REFERENCES

- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, *ApJ*, 935, 167, doi: [10.3847/1538-4357/ac7c74](https://doi.org/10.3847/1538-4357/ac7c74)
- Behnel, S., Bradshaw, R., Citro, C., et al. 2011, *Computing in Science Engineering*, 13, 31, doi: [10.1109/MCSE.2010.118](https://doi.org/10.1109/MCSE.2010.118)
- Belczynski, K., Kalogera, V., Rasio, F. A., et al. 2008, *ApJS*, 174, 223, doi: [10.1086/521026](https://doi.org/10.1086/521026)
- Blaauw, A. 1961, *Bull. Astron. Inst. Netherlands*, 15, 265
- Boersma, J. 1961, *Bull. Astron. Inst. Netherlands*, 15, 291
- Brandt, N., & Podsiadlowski, P. 1995, *MNRAS*, 274, 461, doi: [10.1093/mnras/274.2.461](https://doi.org/10.1093/mnras/274.2.461)
- Breivik, K., Coughlin, S., Zevin, M., et al. 2020a, *ApJ*, 898, 71, doi: [10.3847/1538-4357/ab9d85](https://doi.org/10.3847/1538-4357/ab9d85)
- . 2020b, *ApJ*, 898, 71, doi: [10.3847/1538-4357/ab9d85](https://doi.org/10.3847/1538-4357/ab9d85)
- Burrows, A., & Hayes, J. 1996, *Phys. Rev. Lett.*, 76, 352, doi: [10.1103/PhysRevLett.76.352](https://doi.org/10.1103/PhysRevLett.76.352)
- Coleman, M. S. B., & Burrows, A. 2022, *MNRAS*, 517, 3938, doi: [10.1093/mnras/stac2573](https://doi.org/10.1093/mnras/stac2573)
- Collette, A. 2013, *Python and HDF5* (O’Reilly)
- Collette, A., Kluyver, T., Caswell, T. A., et al. 2023, *h5py/h5py*: 3.8.0, 3.8.0, Zenodo, doi: [10.5281/zenodo.7560547](https://doi.org/10.5281/zenodo.7560547)
- Compas, T., Riley, J., Agrawal, P., et al. 2022, *The Journal of Open Source Software*, 7, 3838, doi: [10.21105/joss.03838](https://doi.org/10.21105/joss.03838)
- Coughlin, S., Breivik, K., Zevin, M., et al. 2024, *COSMIC-PopSynth/COSMIC: Hot fix for pypi install, v3.4.17*, Zenodo, doi: [10.5281/zenodo.13854578](https://doi.org/10.5281/zenodo.13854578)
- da Costa-Luis, C., Larroque, S. K., Altendorf, K., et al. 2024, *tqdm: A fast, Extensible Progress Bar for Python and CLI, v4.66.5*, Zenodo, doi: [10.5281/zenodo.13207611](https://doi.org/10.5281/zenodo.13207611)
- Dewey, R. J., & Cordes, J. M. 1987, *ApJ*, 321, 780, doi: [10.1086/165671](https://doi.org/10.1086/165671)
- Flannery, B. P., & van den Heuvel, E. P. J. 1975, *A&A*, 39, 61
- Fryer, C. L., Belczynski, K., Wiktorowicz, G., et al. 2012, *ApJ*, 749, 91, doi: [10.1088/0004-637X/749/1/91](https://doi.org/10.1088/0004-637X/749/1/91)
- Gommers, R., Virtanen, P., Haberland, M., et al. 2024, *scipy/scipy: SciPy 1.14.1, v1.14.1*, Zenodo, doi: [10.5281/zenodo.13352243](https://doi.org/10.5281/zenodo.13352243)
- Gunn, J. E., & Ostriker, J. P. 1970, *ApJ*, 160, 979, doi: [10.1086/150487](https://doi.org/10.1086/150487)
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, *Nature*, 585, 357, doi: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2)
- Hills, J. G. 1983, *ApJ*, 267, 322, doi: [10.1086/160871](https://doi.org/10.1086/160871)
- Hobbs, G., Lorimer, D. R., Lyne, A. G., & Kramer, M. 2005, *MNRAS*, 360, 974, doi: [10.1111/j.1365-2966.2005.09087.x](https://doi.org/10.1111/j.1365-2966.2005.09087.x)
- Hunter, J. D. 2007, *Computing in Science & Engineering*, 9, 90, doi: [10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)
- Hurley, J. R., Tout, C. A., & Pols, O. R. 2002, *MNRAS*, 329, 897, doi: [10.1046/j.1365-8711.2002.05038.x](https://doi.org/10.1046/j.1365-8711.2002.05038.x)
- Igoshev, A. P. 2020, *MNRAS*, 494, 3663, doi: [10.1093/mnras/staa958](https://doi.org/10.1093/mnras/staa958)
- Izzard, R. G., Dray, L. M., Karakas, A. I., Lugaro, M., & Tout, C. A. 2006, *A&A*, 460, 565, doi: [10.1051/0004-6361/20066129](https://doi.org/10.1051/0004-6361/20066129)
- Izzard, R. G., Glebbeek, E., Stancliffe, R. J., & Pols, O. R. 2009, *A&A*, 508, 1359, doi: [10.1051/0004-6361/200912827](https://doi.org/10.1051/0004-6361/200912827)
- Izzard, R. G., & Jermyn, A. S. 2023, *MNRAS*, 521, 35, doi: [10.1093/mnras/stac2899](https://doi.org/10.1093/mnras/stac2899)
- Izzard, R. G., Preece, H., Joffe, P., et al. 2018, *MNRAS*, 473, 2984, doi: [10.1093/mnras/stx2355](https://doi.org/10.1093/mnras/stx2355)
- Izzard, R. G., Tout, C. A., Karakas, A. I., & Pols, O. R. 2004, *MNRAS*, 350, 407, doi: [10.1111/j.1365-2966.2004.07446.x](https://doi.org/10.1111/j.1365-2966.2004.07446.x)
- Janka, H.-T. 2013, *MNRAS*, 434, 1355, doi: [10.1093/mnras/stt1106](https://doi.org/10.1093/mnras/stt1106)
- . 2017, *ApJ*, 837, 84, doi: [10.3847/1538-4357/aa618e](https://doi.org/10.3847/1538-4357/aa618e)
- Janka, H. T., & Mueller, E. 1994, *A&A*, 290, 496
- Kalogera, V. 1996, *ApJ*, 471, 352, doi: [10.1086/177974](https://doi.org/10.1086/177974)
- Kiel, P. D., & Hurley, J. R. 2009, *MNRAS*, 395, 2326, doi: [10.1111/j.1365-2966.2009.14711.x](https://doi.org/10.1111/j.1365-2966.2009.14711.x)
- Kluyver, T., Ragan-Kelley, B., Pérez, F., et al. 2016, in *ELPUB*, 87–90
- Lyne, A. G., & Lorimer, D. R. 1994, *Nature*, 369, 127, doi: [10.1038/369127a0](https://doi.org/10.1038/369127a0)
- pandas development team, T. 2024, *pandas-dev/pandas: Pandas, v2.2.3*, Zenodo, doi: [10.5281/zenodo.13819579](https://doi.org/10.5281/zenodo.13819579)
- Perez, F., & Granger, B. E. 2007, *Computing in Science and Engineering*, 9, 21, doi: [10.1109/MCSE.2007.53](https://doi.org/10.1109/MCSE.2007.53)
- Pfahl, E., Rappaport, S., Podsiadlowski, P., & Spruit, H. 2002, *ApJ*, 574, 364, doi: [10.1086/340794](https://doi.org/10.1086/340794)
- Price-Whelan, A. M., & Foreman-Mackey, D. 2017, *The Journal of Open Source Software*, 2, doi: [10.21105/joss.00357](https://doi.org/10.21105/joss.00357)
- Renzo, M., Zapartas, E., de Mink, S. E., et al. 2019, *A&A*, 624, A66, doi: [10.1051/0004-6361/201833297](https://doi.org/10.1051/0004-6361/201833297)
- Scheck, L., Kifonidis, K., Janka, H. T., & Müller, E. 2006, *A&A*, 457, 963, doi: [10.1051/0004-6361:20064855](https://doi.org/10.1051/0004-6361:20064855)
- Shklovskii, I. S. 1970, *Soviet Ast.*, 13, 562
- Sutantyo, W. 1978, *Ap&SS*, 54, 479, doi: [10.1007/BF00639450](https://doi.org/10.1007/BF00639450)
- Tauris, T. M., & Takens, R. J. 1998, *A&A*, 330, 1047
- Team COMPAS: Riley, J., Agrawal, P., Barrett, J. W., et al. 2022, *ApJS*, 258, 34, doi: [10.3847/1538-4365/ac416c](https://doi.org/10.3847/1538-4365/ac416c)
- Van Rossum, G., & Drake, F. L. 2009, *Python 3 Reference Manual* (Scotts Valley, CA: CreateSpace)
- Vigna-Gómez, A., Willcox, R., Tamborra, I., et al. 2024, *Phys. Rev. Lett.*, 132, 191403, doi: [10.1103/PhysRevLett.132.191403](https://doi.org/10.1103/PhysRevLett.132.191403)
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature Methods*, 17, 261, doi: [10.1038/s41592-019-0686-2](https://doi.org/10.1038/s41592-019-0686-2)
- Wagg, T., Broekgaarden, F., & Gültekin, K. 2024, *TomWagg/software-citation-station: v1.2, v1.2*, Zenodo, doi: [10.5281/zenodo.13225824](https://doi.org/10.5281/zenodo.13225824)
- Wagg, T., & Broekgaarden, F. S. 2024, *arXiv e-prints*, arXiv:2406.04405. <https://arxiv.org/abs/2406.04405>
- Waskom, M. L. 2021, *Journal of Open Source Software*, 6, 3021, doi: [10.21105/joss.03021](https://doi.org/10.21105/joss.03021)
- Wes McKinney. 2010, in *Proceedings of the 9th Python in Science Conference*, ed. Stéfan van der Walt & Jarrod Millman, 56–61, doi: [10.25080/Majora-92bf1922-00a](https://doi.org/10.25080/Majora-92bf1922-00a)
- Wijers, R. A. M. J., van Paradijs, J., & van den Heuvel, E. P. J. 1992, *A&A*, 261, 145
- Wongwathanarat, A., Janka, H. T., & Müller, E. 2013, *A&A*, 552, A126, doi: [10.1051/0004-6361/20120636](https://doi.org/10.1051/0004-6361/20120636)

provides fast and easy peer review for new papers in the **astro-ph** section of the arXiv, making the reviewing process simpler for authors and referees alike. Learn more at <http://astro.theoj.org>.

This paper was built using the Open Journal of Astrophysics L^AT_EX template. The OJA is a journal which