Ornithology, 2026, **143**, 1–14
https://doi.org/10.1093/ornithology/ukag037
Advance access publication 20 October 2026
Check List Supplement

Ornithology

OXFORD

Check List Supplement

Sixty-seventh Supplement to the American Ornithological Society's *Check-list of North American Birds*

R. Terry Chesser,[1,9,*] Shawn M. Billerman,[2] Kevin J. Burns,[3] Carla Cicero,[4] Jon L. Dunn,[5] Blanca E. Hernández-Baños,[6] Oscar Johnson,[7] Nicholas A. Mason,[8] and Pamela C. Rasmussen[2]

[1]National Museum of Natural History, Smithsonian Institution, Washington, D.C., USA
[2]Cornell Laboratory of Ornithology, Ithaca, New York, USA
[3]Department of Biology, San Diego State University, San Diego, California, USA
[4]Museum of Vertebrate Zoology, University of California, Berkeley, California, USA
[5]24 Idaho Street, Bishop, California, USA
[6]Departamento de Biología Evolutiva, Facultad de Ciencias, Museo de Zoología, Universidad Nacional Autónoma de México, Ciudad de México, México
[7]Department of Biological Sciences, Florida Gulf Coast University, Fort Myers, Florida, USA
[8]Museum of Natural Science and Dept. Biological Sciences, Louisiana State University, Baton Rouge, Louisiana, USA
[9]Authors are members of the Committee on Classification and Nomenclature—North and Middle America, of the American Ornithological Society (formerly American Ornithologists' Union), listed alphabetically after the Chairman.

*Corresponding author: chessert@si.edu

This is the 26th supplement since publication of the 7th edition of the *Check-list of North American Birds* (American Ornithologists' Union [AOU] 1998). It summarizes decisions made between 1 May 2025 and 10 August 2026 by the American Ornithological Society's (formerly American Ornithologists' Union) Committee on Classification and Nomenclature—North and Middle America. The Committee has continued to operate in the manner outlined in the 42nd Supplement (Banks et al. 2000). During the past year, Rosa Alicia Jiménez took a leave of absence from the committee.

Changes in this supplement include the following: (1) 6 species (*Apus affinis*, *Ichthyaetus audouinii*, *Ardeola ralloides*, *Circus cyaneus*, *Pernis ptilorhynchus*, and *Tachycineta albiventer*) are added to the main list on the basis of new distributional information, including 1 transferred from Appendix 1; (2) 7 species (*Numenius hudsonicus*, *Tringa inornata*, *Ardea egretta*, *Momotus castaneiceps*, *Mitrephanes aurantiiventris*, *Cistothorus paludicola*, and *Turdus ardosiaceus*) are added to the main list because of splits from species already on the list; (3) 1 species (*Setophaga aestiva*) is added to the main list because of a split from a species already on the list, and the English name of that species (*S. petechia*) is transferred to *S. aestiva*; (4) 3 species names are changed (to *Pterodroma deserta*, *Camptostoma flaviventre*, and *Deconychura typica*) because of splits from extralimital species; (5) the English names and distributional statements of 3 species (*Onychorhynchus coronatus, Formicarius nigricapillus*, and *Tunchiornis ochraceiceps*) are changed because of splits from extralimital species; (6) the distributional statements of 3 species (*Bubo virginianus, Myiopsitta monachus*, and *Fringilla coelebs*) are changed because of splits from extralimital species; (7) 1 genus (*Nannus*) is added due to a split from another genus, resulting in changes to 2 scientific names (*N. pacificus* and *N. hiemalis*); (8) 1 genus (*Thryorchilus*) is lost due to merger with another genus, resulting in a change to 1 scientific name *(Troglodytes browni*); (9) the English group names of 2 genera (*Tunchiornis* and *Myioborus*) are changed, resulting in changes to the English names of 4 species; (10) 5 species (*Apus pallidus*, *Zapornia pusilla*, *Pelecanus onocrotalus*, *Botaurus stellaris*, and *Parus cinereus*) are added to Appendix 1; (11) 1 species (*Apus nipalensis*) is deleted from Appendix 1 due to merger with a species added to the main list; and (12) 1 species (*Margarops fuscatus*) is added to the list of species known to occur in the United States.

A new linear sequence is adopted for species and genera in the superfamily Certhioidea (families Sittidae, Certhiidae, Polioptilidae, and Troglodytidae) due to new phylogenetic data.

Literature that provides the basis for the Committee's decisions is cited at the end of this supplement, and citations not already in the Literature Cited of the 7th edition (with supplements) become additions to it. A list of the bird species known from the AOS *Check-list* area can be found at http://checklist.americanornithology.org/taxa, and proposals that form the basis for this supplement can be found at https://americanornithology.org/about/committees/nacc/current-prior-proposals/2026-proposals/.

The following changes to the 7th edition (page numbers refer thereto) and its supplements result from the Committee's actions:

pp. xvii–liv. Increase the number in the title of the list of species to 2,226. Insert the following names in the proper position as indicated by the text of this supplement:

Submission Date: August 7, 2026; Editorial Acceptance Date: August 7, 2026
© American Ornithological Society 2026. All rights reserved. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

Apus affinis Little Swift. (A)
Numenius phaeopus Eurasian Whimbrel.
Numenius hudsonicus Hudsonian Whimbrel.
Tringa inornata Prairie Willet.
Tringa semipalmata Saltmarsh Willet.
Ichthyaetus audouinii Audouin's Gull. (A)
Pterodroma deserta Desertas Petrel. (N)
Ardeola ralloides Squacco Heron. (A)
Ardea alba Great White Egret. (A)
Ardea egretta American Egret.
Circus cyaneus Hen Harrier. (A)
Pernis ptilorhynchus Oriental Honey-buzzard. (A)
Momotus mexicanus Mexican Motmot.
Momotus castaneiceps Guatemalan Motmot.
Onychorhynchus coronatus Tropical Royal-Flycatcher.
Camptostoma flaviventre Panama Beardless-Tyrannulet.
Mitrephanes phaeocercus Tawny Tufted-Flycatcher.
Mitrephanes aurantiiventris Lemon-bellied Tufted-Flycatcher.
Formicarius nigricapillus Black-capped Antthrush.
Deconychura typica Piping Long-tailed Woodcreeper.
Tunchiornis ochraceiceps Ochre-crowned Brownlet.
Tachycineta albiventer White-winged Swallow.
Troglodytes browni Timberline Wren.
Nannus pacificus Pacific Wren.
Nannus hiemalis Winter Wren.
Cistothorus paludicola Western Marsh-Wren.
Cistothorus palustris Eastern Marsh-Wren.
Turdus plumbeus Western Red-legged Thrush.
Turdus ardosiaceus Eastern Red-legged Thrush.
Setophaga aestiva Yellow Warbler.
Setophaga petechia Mangrove Warbler.
Myioborus pictus Painted Whitestart.
Myioborus miniatus Slate-throated Whitestart.
Myioborus torquatus Collared Whitestart.

Delete the following names:

Numenius phaeopus Whimbrel.
Tringa semipalmata Willet.
Pterodroma feae Fea's Petrel. (N)
Ardea alba Great Egret.
Momotus mexicanus Russet-crowned Motmot.
Onychorhynchus coronatus Royal Flycatcher.
Camptostoma obsoletum Southern Beardless-Tyrannulet.
Mitrephanes phaeocercus Tufted Flycatcher.
Formicarius nigricapillus Black-headed Antthrush.
Deconychura longicauda Long-tailed Woodcreeper.
Tunchiornis ochraceiceps Tawny-crowned Greenlet.
Troglodytes pacificus Pacific Wren.
Troglodytes hiemalis Winter Wren.
Cistothorus palustris Marsh Wren.
Thryorchilus browni Timberline Wren.
Turdus plumbeus Red-legged Thrush.
Setophaga petechia Yellow Warbler.
Myioborus pictus Painted Redstart.
Myioborus miniatus Slate-throated Redstart.
Myioborus torquatus Collared Redstart.

Adopt the following linear sequence for species and genera in the families Sittidae, Certhiidae, Polioptilidae, and Troglodytidae:

Sittidae
Sitta carolinensis
Sitta pygmaea
Sitta pusilla
Sitta insularis
Sitta canadensis
Certhiidae
Certhia americana
Polioptilidae
Ramphocaenus melanurus
Microbates cinereiventris
Polioptila schistaceigula
Polioptila lembeyei
Polioptila albiventris
Polioptila bilineata
Polioptila caerulea
Polioptila nigriceps
Polioptila albiloris
Polioptila melanura
Polioptila californica
Troglodytidae
Salpinctes obsoletus
Microcerculus philomela
Microcerculus marginatus
Catherpes mexicanus
Hylorchilus sumichrasti
Hylorchilus navai
Thryomanes bewickii
Thryothorus ludovicianus
Campylorhynchus brunneicapillus
Campylorhynchus albobrunneus
Campylorhynchus zonatus
Campylorhynchus megalopterus
Campylorhynchus yucatanicus
Campylorhynchus jocosus
Campylorhynchus chiapensis
Campylorhynchus griseus
Campylorhynchus gularis
Campylorhynchus capistratus
Campylorhynchus humilis
Campylorhynchus rufinucha
Ferminia cerverai
Nannus pacificus
Nannus hiemalis
Cistothorus paludicola
Cistothorus palustris
Cistothorus stellaris
Cistothorus platensis
Troglodytes browni
Troglodytes rufociliatus
Troglodytes ochraceus
Troglodytes tanneri
Troglodytes sissonii
Troglodytes aedon
Troglodytes musculus
Troglodytes beani
Troglodytes martinicensis
Troglodytes mesoleucus
Troglodytes musicus
Troglodytes grenadensis
Pheugopedius atrogularis

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

Pheugopedius spadix
Pheugopedius maculipectus
Pheugopedius rutilus
Pheugopedius felix
Pheugopedius fasciatoventris
Cantorchilus thoracicus
Cantorchilus leucopogon
Cantorchilus modestus
Cantorchilus zeledoni
Cantorchilus elutus
Cantorchilus nigricapillus
Cantorchilus semibadius
Cantorchilus leucotis
Henicorhina leucosticta
Henicorhina leucophrys
Cyphorhinus phaeocephalus
Uropsila leucogastra
Thryophilus sinaloa
Thryophilus pleurostictus
Thryophilus rufalbus

Note: *The entries below follow the current linear sequence as established in this and previous supplements, although entries continue to be cross-referenced to page numbers in* AOU (1998).

1. [pp. 280, 694] *Apus nipalensis*, currently in Appendix 1, is treated as conspecific with *A. affinis*, and a record for *A. affinis* in the Checklist area is recognized. Remove the account for *A. nipalensis* from the Appendix, and insert the following new species account in the main list after the account for *Apus pacificus*:

Apus affinis (J. E. Gray). Little Swift.

Cypselus affinis J. E. Gray, 1830, in Gray and Hardwicke's Illustrated Indian Zoology 1, pt. 2, pl. 35, f. 2. (no locality = Ganges.)

Habitat.—Most frequently in cities and towns where it nests on buildings, but occurs in most terrestrial habitats, less so in very arid deserts. Also nests on cliffs.

Distribution.—*Breeds* [*affinis* group] from southern Spain, North Africa, sub-Saharan Africa, eastern Türkiye, and the Middle East locally east to Central Asia in Tajikistan and Uzbekistan and south through the Indian Subcontinent to Sri Lanka; [*nipalensis* group] from the Himalayas east to eastern China, central Japan, Taiwan, the northern Philippines, and in eastern Indonesia east to southern Sulawesi. Both groups mostly resident, but [*affinis* group] populations in the Palearctic and South Africa are partially to entirely migratory, and [*nipalensis* group] northern populations believed to be migratory, and records away from resident range may indicate migratory movements.

Infrequently noted [*affinis* group] in migration (Israel, Saudi Arabia, and Chad).

Casual or accidental [*affinis* group] to northern Europe as far as Sweden and the British Isles, southern Europe, Madeira, the Canary Islands, Madagascar, the Maldives, and [*nipalensis* group] northwestern Australia.

Accidental to Barbados (3 June 2023; 24–25 June 2023; photos; https://ebird.org/checklist/S140353653, https://ebird.org/checklist/S142526956). The origin of an individual of the *nipalensis* group found dead at a container terminal in southwestern British Columbia is considered questionable (Chesser et al. 2019).

Notes.—Groups: *A. affinis* [Little Swift] and *A. nipalensis* (Hodgson, 1836) [House Swift]. *Apus affinis* and *A. nipalensis* were formerly (e.g., Chesser et al. 2019) considered separate species, but are treated as conspecific based on the lack of diagnostic morphological characters (Abdulali 1966, Brooke 1971, Inskipp and Praveen 2026) and vocalizations (Chantler et al. 2026), and paraphyly in mitochondrial DNA (Päckert et al. 2012).

2. [pp. 160-161] *Numenius hudsonicus* is treated as a species separate from *N. phaeopus*. In the species account for *N. phaeopus*, change the English name to Eurasian Whimbrel, and change the distributional statement and existing Notes to the following:

Distribution.—*Breeds* in coastal eastern Greenland (Jameson Land; Boertmann et al. 1985) and in Eurasia from Iceland, the Faroe Islands, northern Scandinavia, northern Russia, and northern Siberia south to the Orkney and Shetland islands, southern Scandinavia, central Russia, central Siberia, and east to Anadyrland and the Sea of Okhotsk.

Winters in the Old World from the Atlantic Islands and Mediterranean region (occasionally the British Isles), east through the Arabian Peninsula to southeastern China, and south to southern Africa, Madagascar, Indian Ocean islands, South and Southeast Asia, Australasia, and western Pacific islands.

Migrates through the western Aleutians (Near Islands) and rarely to the Pribilof and St. Lawrence islands (specimens are of subspecies *variegatus;* Gibson and Withrow 2015).

Casual in the Hawaiian Islands, western Alaska, along the Pacific coast of North America south to California, and along the Atlantic coast of North America from southern Labrador, Newfoundland, and Nova Scotia south to Georgia; also in Bermuda and the Lesser Antilles. Accidental in Yukon Territory, interior California, Ontario, Quebec, Michigan, Ohio, Texas, and Cuba.

Notes.—See comments under *N. hudsonicus*.

Insert the following new species account after the account for *N. phaeopus*:

Numenius hudsonicus Latham. Hudsonian Whimbrel.

Numenius hudsonicus Latham, 1790, Index Ornithologicus 2: 712. (Hudson Bay.)

Habitat.—Tundra (breeding); beaches, rocky coasts, tidal mudflats, marshes, estuaries, flooded fields, and pastures (nonbreeding).

Distribution.—[same as for the *hudsonicus* group in the account for *Numenius phaeopus* in AOU (1998), except delete "(except Cuba)" from the *Winters* paragraph and replace the Casual paragraph with the following: "Casual or accidental in Greenland, the Hawaiian Islands, Clipperton Island, Rapa Nui, Bolivia, the Azores, Oman, China, Taiwan, New Guinea, Australia, and New Zealand. Rare in western Europe."]

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *N. phaeopus*, but separated based on differences in plumage (Cramp and Simmons 1983), displays (Williamson 1946,

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

Cramp and Simmons 1983, Skeel 1978, Skeel et al. 2025), and genomics (Tan et al. 2019, 2023; McLaughlin et al. 2020).

3. [p. 156] *Tringa inornata* is treated as a species separate from *T. semipalmata*. In the species account for *T. semipalmata*, change the English name to Saltmarsh Willet and change the habitat and distributional statements to the following:

Habitat.—Salt marshes and beaches (breeding); marshes, tidal mudflats, beaches, and margins of shallow coastal lagoons (nonbreeding).

Distribution.—*Breeds* locally along the Atlantic-Gulf coast from eastern Quebec (Magdalen Islands), southern New Brunswick, Prince Edward Island, and Nova Scotia south to southern Florida and west to southern Texas and Tamaulipas; in the West Indies, on the Bahamas, Antilles (Cuba, Beata Island off Hispaniola, Anegada and St. Croix in the Virgin Islands, Antigua, and Guadeloupe, possibly also to Barbuda, St. Martin, Anguilla, and St. Kitts), and in the Cayman Islands; and on Los Roques, off northern Venezuela. Nonbreeding individuals occur sporadically in summer as far south as Argentina.

Winters from the Caribbean and Atlantic coast of South America to Uruguay, southern Brazil, and Argentina (Buenos Aires province). Winter status in southern Mexico, both coasts of Central America, and the Pacific coast of South America uncertain, but tracking data indicate that some winter on the Pacific coast of southern Central America and Ecuador (Heath et al. 2021).

Migrates primarily through coastal areas.

Accidental to Ontario, Vermont, Pennsylvania, central North Carolina, Tennessee, and Norway (14 October 1992–23 March 1993; photos; Gustad 1994).

Insert the following sentence at the end of the existing Notes for this species: See comments under *T. inornata*.

Insert the following new species account before the account for *T. semipalmata*:

Tringa inornata (Brewster). Prairie Willet.

Symphemia semipalmata inornata Brewster, 1887, Auk 4: 145. (Larimer County, Colorado.)

Habitat.—Marshy lake and pond margins and wet meadows (breeding); marshes, tidal mudflats, beaches, and saline lake margins (nonbreeding).

Distribution.—*Breeds* locally from eastern Oregon, southern Idaho, central Alberta, southern Saskatchewan, and southern Manitoba south to northeastern and east-central California, western Nevada, central Utah, northern Colorado, western and northern Nebraska, and eastern South Dakota, formerly in western Minnesota and northwestern Iowa. Nonbreeding individuals occur sporadically in summer as far south as Argentina.

Winters from western Washington (casually from southwestern British Columbia; also regularly inland in southeastern California at the Salton Sea and Owens Lake), the Gulf coast, and New Jersey south along both coasts of Middle America (including offshore islands), the West Indies, and both coasts of South America to the Galapagos Islands, central Chile, and east at least to French Guiana. Winters in small numbers in coastal southern Brazil, Uruguay, and Argentina (Buenos Aires province).

Migrates primarily through coastal areas but also irregularly through most of the interior western and central United States, regularly east to the Great Lakes and on the east and Gulf coasts, rarely elsewhere in the eastern United States.

Casual north to northeastern British Columbia, southern Hudson Bay, southern Quebec, and Newfoundland. Accidental in the Hawaiian Islands (Maui), Alaska (Kenai Peninsula; sight report from Minto Lakes), Bermuda, Finland, France, the Azores, and Cabo Verde. Although the text in Pyle and Pyle (2017) identifies the Maui record as *semipalmata*, the photographs indicate *inornata*; other records from Oahu, Kauai, and Hawaii are not separable from *semipalmata* (Pyle and Pyle 2017), but are very likely to be *inornata*.

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *T. semipalmata*, but separated based on differences in vocalizations (Sordahl 1979, Lowther et al. 2020), including differential response to playback (Douglas 1998), genomics (Oswald et al. 2016), morphology (O'Brien 2006), and ecology.

4. [p. 185] After the account for *Ichthyaetus ichthyaetus*, insert the following new species account:

Ichthyaetus audouinii (Payraudeau). Audouin's Gull.

Larus audouinii Payraudeau, 1826, Annales Des Sciences Naturelles 8: 462. (Sardinia and Corsica.)

Habitat.—Sandy or rocky beaches, saltpans, coastal fields, and at sea; seldom scavenges. Nests on rocky islands or sandy soils with short vegetation.

Distribution.—*Breeds* locally throughout much of the Mediterranean region, mostly in the western Mediterranean (in particular the Ebro Delta, Spain, with 20,000+ pairs, some two-thirds of the population). Breeding range recently extended to southern Portugal (Algarve).

Winters mainly on the Atlantic coast north to Portugal, Mediterranean coasts, and coastal Morocco; fewer south to the Canary Islands, Mauritania, Senegal, and The Gambia.

Migrates regularly in spring and fall past Gibraltar.

Casual or accidental to the Azores, western France, the British Isles, the Netherlands, Scandinavia, Germany, Switzerland, Bulgaria, Cabo Verde, the Sea of Marmora, and Jordan. Casual recently to Trinidad and northern South America with documented records (photos) from Trinidad, Suriname, and French Guiana.

Accidental to Barbados, one present from 20 June to at least 12 October 2024 (photos; https://ebird.org/checklist/S182610062).

5. [p. 14] *Pterodroma deserta* is treated as a species separate from the now extralimital *P. feae*. Remove the account for *P. feae* and replace it with the following new account:

Pterodroma deserta Mathews. Desertas Petrel.

Pterodroma mollis deserta Mathews, 1934, Bulletin of the British Ornithologists' Club 54: 179. (Desertas Islands.)

Habitat.—Pelagic waters; nests in burrows or crevices on islands.

Distribution.—*Breeds* on Bugio Island in the Desertas Islands, Madeira Archipelago.

Ranges at sea in the eastern North Atlantic, at least casually north to the British Isles.

Rare but annual in western Atlantic waters off North America; most records are in late spring and are from off North Carolina, but documented north to Nova Scotia and reported south to Florida (Dunn et al. 2012, Kratter 2018).

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

Accidental inland in Virginia following Hurricane Fran (September 1996; Howell 2012). Data from geolocators indicated that one individual (of 17 tracked) wintered off the coast of Georgia and Florida in 2007–2008 (Ramírez et al. 2013).

Notes.—Formerly (Chesser et al. 2013) considered conspecific with *P. feae* Salvadori, 1899 [Fea's Petrel] but separated based on differences in nesting phenology and vocalizations (Robb and Mullarney 2008). Similarity of plumage makes distinguishing *P. deserta* from *P. feae* difficult, but available information suggests that records from the western Atlantic are of *P. deserta*.

6. [p. 43] Before the account for *Ardeola bacchus*, insert the following new species account:

Ardeola ralloides (Scopoli). Squacco Heron.

Ardea ralloides Scopoli, 1769, Annus I Historico-Naturalis, p. 88. ("In Carniolica" = Carniola.)

Habitat.—Shallow aquatic habitats, typically fresh or brackish water with plenty of reedy cover; also flooded rice fields. Nests in bushes or short trees within this habitat, often with other herons, *Microcarbo pygmaeus* (Pallas, 1873) [Pygmy Cormorant], and *Plegadis falcinellus*.

Distribution.—*Breeds* from Morocco and locally across southern Europe, north to about the Danube and east to southern Ukraine, southern Russia, southwestern Kazakhstan, eastern Uzbekistan, and eastern Turkmenistan, and south locally through Egypt and the Middle East to Iran.

Resident in Africa south of the Sahara and on Madagascar.

Winters mostly in Africa south of the Sahara; a few winter in the southern Mediterranean region (especially Nile Valley), Iraq, and the Persian Gulf.

Migrates over the Sahara and the Arabian Peninsula where recorded mostly from oases.

Rare or casual to northern Europe north to the British Isles and Fennoscandia, and to the Canary Islands, Madeira, Azores, and Cabo Verde. A sight report for northeastern Afghanistan. Accidental in Iceland.

Casual to the Leeward Islands in the West Indies, where recorded on Martinique (4 April 2021; photos, alternate plumage and therefore a definitive identification; https://ebird.org/checklist/S269658823), Guadeloupe (10 November 2018, basic plumage; photos; https://ebird.org/checklist/S49838084), and Barbados (26 November 2021, basic plumage; photos; https://ebird.org/checklist/S98087082).

7. [p. 40-41] *Ardea egretta* is treated as a species separate from *A. alba*. In the species account for *A. alba*, change the English name to Great White Egret, remove the New World parts of the distribution from the *Breeds*, *Winters*, and Wanders paragraphs of the distributional statement, and replace the Accidental paragraph with the following:

Casual in Alaska (central and western Aleutians; photos and two specimens, both of subspecies *modesta*; Gibson and Byrd 2007) and accidental south to British Columbia and Washington. Accidental in Chincoteague, Virginia (tentatively identified as subspecies *melanorhynchos*; May 2008–April 2026; photos, eBird records).

Change the existing Notes to:

Notes.—Formerly placed in the genus *Casmerodius*; placed by some authors (e.g., Blake 1977) in *Egretta*. See comments under *A. egretta*.

Insert the following new species account after the account for *A. alba*:

Ardea egretta Gmelin. American Egret.

Ardea Egretta Gmelin, 1789, Systema Naturae 1: 629; based chiefly on "La Grande Aigrette" of Buffon, 1780, Histoire Naturelle Générale 22, Histoire Naturelle des Oiseaux 7: 377. (Santo Domingo, Falkland Islands, South America to Louisiana.)

Habitat.—[same as the habitat statement in account for *Ardea alba* in AOU (1998)]

Distribution.— [same as the New World parts of the *Breeds*, *Winters*, and Wanders paragraphs of the account for *Ardea alba* in AOU (1998)]

Casual to southeastern and south-central Alaska. Accidental in interior and northern Alaska (Gibson and Withrow 2015), the Hawaiian Islands (Oahu, Maui, Kauai; Pyle and Pyle 2017), the Azores, and the Canary Islands (Reyt et al. 2025).

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *A. alba* [as Great Egret], but separated based on differences in vocalizations (Boesman 2016) and coloration of bare parts (Pratt 2011, Reyt et al. 2025). See comments under *A. alba*.

8. [p. 92] Before the account for *Circus hudsonius*, insert the following new species account:

Circus cyaneus (Linnaeus). Hen Harrier.

Falco cyaneus Linnaeus, 1766, Systema Naturae, ed. 12, vol. 1, p. 126; based on "The Blue Hawk" of Edwards, 1756, Gleanings of Natural History, p. 33, pl. 225. (Europe, Africa = region of London *ex* Edwards.)

Habitat.—Open ground such as moors, heaths, bogs, and fields. In N Europe nests also in open woods on swampy ground and in the British Isles in reforestation plantations. More catholic in habitat preferences in migration and winter but still prefers open habitats.

Distribution.—*Breeds* from the Scandinavian peninsula east to Kamchatka, south to eastern Hungary, northern Romania and central Ukraine, east through Russia, far northern Kazakhstan, Mongolia, and northeastern China. Also resident in the British Isles and Western Europe from Denmark south to northern Spain and east to western Poland.

Winters from the British Isles and western Europe east through southern Europe, Türkiye, the northern Middle East, and locally through western Asia, northern India east to the southern Korean Peninsula, Japan, southeastern China, northern Vietnam, and Laos; and in northwestern Africa from Morocco south to western Mauritania, Tunisia, and locally in the southern Arabian Peninsula and southern South and Southeast Asia.

Accidental in Alaska (Attu Island, early June 1999; partial wing identified using mtDNA; Withrow et al. 2024; and Adak Island, January 2025; photos; J. J. Withrow, in litt.).

Notes.—Included in AOU (1983, 1998) as *C. cyaneus sensu lato*, prior to separation from *C. hudsonius* (Chesser et al. 2017). See comments under *C. hudsonius*.

9. [p. 88] After the account for *Chondrohierax wilsonii*, insert the following new heading, citation, and species account:

Genus ***PERNIS*** Cuvier

Pernis Cuvier, 1817, Règne Animal 1 (1816): 322. Type, by monotypy, *Falco apivorus* Linnaeus.

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

Pernis ptilorhynchus (Temminck). Oriental Honey-buzzard.

Falco ptilorhyncus [sic] Temminck, 1821, Nouveau Recueil de Planches Coloriées d'Oiseaux, livr. 8, pl. 44. (Java.)

Habitat.—Resident, tropical populations in rainforest to more open wooded country. In the Eastern Palearctic, breeds in broadleaf and mixed forest and other well-timbered lowland and hill areas, often with open areas interspersed. In winter also in wooded to partly wooded areas. Often migrates over wooded ridges.

Distribution.—*Resident* in South and Southeast Asia, from far eastern Pakistan and India east to extreme southeastern China, Taiwan, and Vietnam, and south through the Malay Peninsula, Sumatra, western Java, Borneo, and the Philippine Islands.

Breeds from south-central Russia east to Sakhalin, and south to northern Mongolia, northeastern China, the Korean Peninsula, and Japan.

Winters locally around the Persian Gulf, Pakistan, Taiwan, Hainan, and the Lesser Sundas.

Migrates from Japan in fall over East China Sea to reach China and thence southward to wintering grounds, but in spring migrates over land through the Korean Peninsula en route to Japan. In Central Asia may join migrating flocks of *Pernis apivorus* (Linnaeus, 1758) [European Honey-buzzard].

Rare migrant and winterer in the Caucasus (Georgia), Middle East (Israel), and East Africa (Kenya). Rare in Australia. Accidental to Europe.

Accidental in Alaska (Shemya Island, 27 May 2024; photos; Pyle et al. 2025, Pohlen et al. 2026).

10. [pp. 257-258] Extralimital species *Bubo nigrescens, B. nacurutu*, and *B. magellanicus* are treated as species separate from *B. virginianus*. In the species account for *B. virginianus*, replace "south to Honduras, rarely in Guatemala and El Salvador, and perhaps in north-central Nicaragua, and from northern Colombia and Venezuela south to Tierra del Fuego (except for most of Amazonia). Absent from the West Indies, most other islands, and apparently much of Central America" in the distributional statement (as modified in Banks et al. 2000) with "south to Honduras, rarely in Guatemala and El Salvador, and in north-central Nicaragua. Absent from the West Indies.", and insert the following paragraph at the end of the distributional statement:

Introduced to Hiva Oa, in the southern Marquesas.

Add the following Notes to the end of the species account:

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *B. nigrescens* Berlepsch, 1884 [Paramo Horned Owl], *B. nacurutu* (Vieillot, 1817) [Tropical Horned Owl], and *B. magellanicus* (Lesson, 1828) [Magellanic Horned Owl], but separated, following Remsen et al. (2026), based on differences in vocalizations (López-Lanús 2015), genetics (Ostrow et al. 2023), and morphology (Traylor 1958, König et al. 1996).

11. [pp. 320-321] *Momotus castaneiceps* is treated as a species separate from *M. mexicanus*. In the species account for *M. mexicanus*, change the English name to Mexican Motmot, change "Guatemala (upper Motagua Valley)" to "far western Guatemala" in the distributional statement, and insert the following at the end of the account:

Notes.—See comments under *M. castaneiceps*.

Insert the following new species account after the account for *M. mexicanus*:

Momotus castaneiceps Gould. Guatemalan Motmot.

Momotus castaneiceps Gould, 1854, Proceedings of the Zoological Society of London, pt. 22 (1855), p. 154. (Guatemala.)

Habitat.—Tropical Deciduous Forest, Gallery Forest, Secondary Forest (0-1900 m; Tropical and lower Subtropical zones).

Distribution.—*Resident* in the interior of central and southern Guatemala (upper Motagua Valley).

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *M. mexicanus*, but separated based on differences in morphology and vocalizations commensurate with differences between other closely related species of *Momotus* (Howell and Juárez 2025).

12. [p. 233] Extralimital species *Myiopsitta luchsi* is treated as a species separate from *M. monachus*. In the species account for *M. monachus*, change "central Bolivia" to "southeastern Bolivia" in the distributional statement and replace the existing "Introduced and established" paragraph with the following:

Widely introduced into urban centers, with established populations mainly in Illinois, New York, and the southern United States from California locally to Florida, Mexican urban areas, Puerto Rico, Grand Cayman, central Chile, southern Europe, the Canary Islands, Israel, and Singapore.

Add the following Notes to the end of the species account:

Notes.—Formerly considered conspecific with *M. luchsi* (Finsch, 1868) [Cliff Parakeet], but separated, following Remsen et al. (2026), based on differences in vocalizations (Boesman 2017) and breeding behavior.

13. [p. 386] Extralimital species *Onychorhynchus swainsoni* is treated as a species separate from *O. coronatus*. In the species account for *O. coronatus*, change the English name to Tropical Royal-Flycatcher, remove "; and [*swainsoni* group] in southeastern Brazil" from the distributional statement, and replace the existing Notes with:

Notes.—Formerly considered conspecific with *O. swainsoni* (Pelzeln, 1858) [Atlantic Royal-Flycatcher], but separated, following Remsen et al. (2026), based on differences in vocalizations (Kirwan et al. 2024) consistent with genomic differences (Harvey et al. 2020, Reyes et al. 2023). The remaining groups may also represent separate species: *O. mexicanus* (Sclater, 1857) [Northern Royal-Flycatcher], *O. occidentalis* (Sclater, 1860) [Western Royal-Flycatcher], and *O. coronatus* [Amazonian Royal-Flycatcher].

14. [p. 374] *Camptostoma flaviventre* is treated as a species separate from the now extralimital species *C. obsoletum*. Remove the species account for *C. obsoletum* and insert the following new species account:

Camptostoma flaviventre Sclater and Salvin. Panama Beardless-Tyrannulet.

Camptostoma flaviventre Sclater and Salvin, 1864, Proceedings of the Zoological Society of London, p. 358. (Panama; presumably from near Lion Hill, Canal Zone.)

Habitat.—Secondary Forest, Gallery Forest, Riparian Thickets, Second-growth Scrub (0-1200 m; Tropical and lower Subtropical zones).

Distribution.—*Resident* in southwestern Costa Rica (Pacific slope north to Tárcoles) and Panama (Pacific slope throughout,

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

including Coiba, Cébaco, and the Pearl islands; locally on the Caribbean slope in Colón, Canal area, and San Blas).

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *C. obsoletum* (Temminck, 1824) [Southeastern Beardless-Tyrannulet], but separated, along with *C. caucae* Chapman, 1914 [Cauca Beardless-Tyrannulet], *C. pusillum* (Cabanis and Heine, 1859) [Venezuelan Beardless-Tyrannulet], *C. sclateri* (Berlepsch and Taczanowski, 1883) [Pacific Beardless-Tyrannulet], and *C. napaeum* (Ridgway, 1888) [Amazonian Beardless-Tyrannulet], following Remsen et al. (2026), based on differences in vocalizations (Lima and Vaz 2024).

15. [pp. 389-390] *Mitrephanes aurantiiventris* is treated as a species separate from *M. phaeocercus*. In the species account for *M. phaeocercus*, change the English name to Tawny Tufted-Flycatcher and change the distributional statement and existing Notes to the following:

Distribution.—*Resident* in the highlands from northeastern Sonora, western Chihuahua, Sinaloa, Durango, western Zacatecas, southeastern San Luis Potosí, and southern Tamaulipas south through Mexico, Guatemala, El Salvador, and Honduras to north-central Nicaragua.

Casual to southeastern Arizona. Accidental to central and northern Arizona, southern New Mexico, and western and southeastern Texas (Kleberg County).

Notes.—See comments under *M. aurantiiventris*.

Insert the following new species account after the account for *M. phaeocercus*:

Mitrephanes aurantiiventris (Lawrence). Lemon-bellied Tufted-Flycatcher.

Mitrephorus aurantiiventris Lawrence, 1865, Annals of the Lyceum of Natural History of New York 8: 173. (Tabacales, Costa Rica.)

Habitat.—Montane Evergreen Forest, Pine-Oak Forest, Montane Evergreen Forest Edge (600-3600 m; upper Tropical to lower Temperate zones).

Distribution.—*Resident* [*aurantiiventris* group] from both slopes of Costa Rica south locally through Panama (Bocas del Toro, Chiriquí, Veraguas, Coclé); and [*berlepschi* group] extreme eastern Panama (eastern Darién) and western Colombia to northwestern Ecuador.

Notes.— Formerly (AOU 1983, 1998) considered conspecific with the extralimital *M. olivaceus* Berlepsch and Stolzmann, 1894 [Olive Tufted-Flycatcher], then, following the separation of *M. phaeocercus* from *M. olivaceus* (Chesser et al. 2018), treated as conspecific with *M. phaeocercus*. Now separated from *M. phaeocercus* based on differences in vocalizations (Dyer and Howell 2023, Iliff and Rasmussen 2026), genomics (Harvey et al. 2020), and plumage (Webster 1968, Farnsworth and Lebbin 2020). Groups: *M. aurantiiventris* [Costa Rican Tufted-Flycatcher] and *M. berlepschi* Hartert, 1902 [Choco Tufted-Flycatcher]. The population in the Cerro Tacarcuna range, Panama, may represent an undescribed taxon.

16. [p. 370] Extralimital species *Formicarius destructus* is treated as a species separate from *F. nigricapillus*. In the species account for *F. nigricapillus*, change the English name to Black-capped Antthrush, delete ", and along the Pacific slope of northwestern South America south to Ecuador" from the distributional statement, and insert the following at the end of the account:

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *F. destructus* Hartert, 1898 [Black-hooded Antthrush], but separated based on differences in vocalizations (Areta and Benítez Saldívar 2025).

17. [p. 356] *Deconychura typica* is treated as a species separate from the now extralimital *D. longicauda*. Remove the species account for *D. longicauda* and insert the following new species account:

Deconychura typica Cherrie. Piping Long-tailed Woodcreeper.

Deconychura typica Cherrie, 1891, Proceedings of the United States National Museum 14: 339. (Pozo Azul de Pirrís, southwestern Costa Rica.)

Habitat.—Tropical Lowland Evergreen Forest (0-1300 m; Tropical and lower Subtropical zones).

Distribution.—Locally in southeastern Honduras (Olancho), Costa Rica (Caribbean slope of Cordillera Central, and northwestern region from the Gulf of Nicoya southward) to Panama (Chiriquí, and from eastern Panamá province east through Darién) and northern Colombia.

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *D. longicauda* (Pelzeln, 1868) [Whistling Long-tailed Woodcreeper], but separated, along with *D. pallida* Zimmer, 1929 [Mournful Long-tailed Woodcreeper], following Remsen et al. (2026), based on differences in vocalizations (Marantz et al. 2003, Boesman 2016a).

18. [p. 439] Extralimital species *Tunchiornis ferrugineifrons*, *T. luteifrons*, and *T. rubrifrons* are treated as species separate from *T. ochraceiceps*, and the English group name for *Tunchiornis* is changed from Greenlet to Brownlet.

After the heading and citation for genus *Tunchiornis*, insert the following Notes:

Notes.—English group name changed from Greenlet to Brownlet, following Remsen et al. (2026), to better reflect the plumage coloration of species of this genus and their phylogenetic distance from greenlets of the genera *Hylophilus* and *Pachysylvia*.

In the species account for *T. ochraceiceps*, change the English name to Ochre-crowned Brownlet and replace the distributional statement and Notes with the following:

Distribution.—*Resident* from northeastern Oaxaca and southern Veracruz south on the Gulf-Caribbean slope (except Yucatán) to Nicaragua, and on both slopes of Costa Rica (absent from the drier parts of the northwest) and Panama (mainly on the Caribbean slope, but both slopes in the southeast).

Notes.—Formerly considered conspecific with *T. ferrugineifrons* (Sclater, 1862) [Rufous-fronted Brownlet], *T. luteifrons* (Sclater, 1862) [Guianan Brownlet], and *T. rubrifrons* (Sclater and Salvin, 1867) [Para Brownlet], but separated, following Remsen et al. (2026), based on deep divergence in genomics and average differences in song, plumage, and iris color (Buainain et al. 2021). Formerly placed in the genus *Hylophilus*, but genetic data (Slager et al. 2014) indicate that *Hylophilus sensu lato* is paraphyletic and that *Tunchiornis* is not closely related to true *Hylophilus*.

19. [p. 457] After the account for *Tachycineta bicolor*, insert the following new species account:

Tachycineta albiventer (Boddaert). White-winged Swallow.

Hirundo albiventer Boddaert, 1783, Table des Planches Enluminéez d'Histoire Naturelle de M. Daubenton, p. 32. (Cayenne, *ex* Daubenton, pl. 546, fig. 2.)

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

Habitat.— Freshwater Lakes and Ponds, Rivers, Pastures/Agricultural Lands (0-1500 m; Tropical and lower Subtropical zones).

Distribution.—*Resident* in lowlands from northern Colombia and Venezuela south through the Guianas, Brazil, Ecuador, and Peru east of the Andes, to northern Bolivia, eastern Paraguay, and extreme northeastern Argentina. Recent arrival to Grenada, where it has bred locally (based on the presence of juvenile birds; https://ebird.org/checklist/S71545684).

Accidental in the Grenadines. Sight reports from Martinique and Panama.

20. [pp. 468-486, 491-494] Phylogenetic analyses of nuclear and mitochondrial DNA sequences (Imfeld et al. 2024) have shown that our current linear sequence of species and genera in the Sittidae, Certhiidae, Polioptilidae, and Troglodytidae does not reflect their evolutionary relationships. These findings result in the following changes:

Rearrange the linear sequence of species and genera as follows (note that this species listing incorporates changes from items 21 and 22 below):

Sittidae
- *Sitta carolinensis*
- *Sitta pygmaea*
- *Sitta pusilla*
- *Sitta insularis*
- *Sitta canadensis*

Certhiidae
- *Certhia americana*

Polioptilidae
- *Ramphocaenus melanurus*
- *Microbates cinereiventris*
- *Polioptila schistaceigula*
- *Polioptila lembeyei*
- *Polioptila albiventris*
- *Polioptila bilineata*
- *Polioptila caerulea*
- *Polioptila nigriceps*
- *Polioptila albiloris*
- *Polioptila melanura*
- *Polioptila californica*

Troglodytidae
- *Salpinctes obsoletus*
- *Microcerculus philomela*
- *Microcerculus marginatus*
- *Catherpes mexicanus*
- *Hylorchilus sumichrasti*
- *Hylorchilus navai*
- *Thryomanes bewickii*
- *Thryothorus ludovicianus*
- *Campylorhynchus brunneicapillus*
- *Campylorhynchus albobrunneus*
- *Campylorhynchus zonatus*
- *Campylorhynchus megalopterus*
- *Campylorhynchus yucatanicus*
- *Campylorhynchus jocosus*
- *Campylorhynchus chiapensis*
- *Campylorhynchus griseus*
- *Campylorhynchus gularis*
- *Campylorhynchus capistratus*
- *Campylorhynchus humilis*
- *Campylorhynchus rufinucha*
- *Ferminia cerverai*
- *Nannus pacificus*
- *Nannus hiemalis*
- *Cistothorus paludicola*
- *Cistothorus palustris*
- *Cistothorus stellaris*
- *Cistothorus platensis*
- *Troglodytes browni*
- *Troglodytes rufociliatus*
- *Troglodytes ochraceus*
- *Troglodytes tanneri*
- *Troglodytes sissonii*
- *Troglodytes aedon*
- *Troglodytes musculus*
- *Troglodytes beani*
- *Troglodytes martinicensis*
- *Troglodytes mesoleucus*
- *Troglodytes musicus*
- *Troglodytes grenadensis*
- *Pheugopedius atrogularis*
- *Pheugopedius spadix*
- *Pheugopedius maculipectus*
- *Pheugopedius rutilus*
- *Pheugopedius felix*
- *Pheugopedius fasciatoventris*
- *Cantorchilus thoracicus*
- *Cantorchilus leucopogon*
- *Cantorchilus modestus*
- *Cantorchilus zeledoni*
- *Cantorchilus elutus*
- *Cantorchilus nigricapillus*
- *Cantorchilus semibadius*
- *Cantorchilus leucotis*
- *Henicorhina leucosticta*
- *Henicorhina leucophrys*
- *Cyphorhinus phaeocephalus*
- *Uropsila leucogastra*
- *Thryophilus sinaloa*
- *Thryophilus pleurostictus*
- *Thryophilus rufalbus*

Replace the existing Notes after the heading Family **SITTIDAE:** Nuthatches with the following:

Notes.—Linear sequence of species follows Imfeld et al. (2024).

Replace the existing Notes after the heading Family **POLIOPTILIDAE:** Gnatcatchers with the following:

Notes.—Linear sequence of species and genera follows Imfeld et al. (2024).

Replace the existing Notes after the heading Family **TROGLODYTIDAE:** Wrens with the following:

Notes.—Linear sequence of species and genera follows Imfeld et al. (2024).

Delete the existing Notes under Genus ***CAMPYLORHYNCHUS*** Spix.

21. [pp. 480-484] Phylogenetic analyses of nuclear and mitochondrial DNA sequences (Rice et al. 1999, Barker 2017, Imfeld et al. 2024) have shown that the genus *Troglodytes* is paraphyletic with respect to *Cistothorus*, *Ferminia*, and *Thryorchilus*, and that *Thryorchilus browni* is either very closely related to or nested within *Troglodytes sensu stricto*. These findings result in the following changes:

Remove the citation for *Nannus* from the synonymy of *Troglodytes*.

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

After the species account for *Ferminia cerverai*, insert the following new heading, citation, and Notes:

Genus ***NANNUS*** Billberg

Nannus Billberg, 1828, Synopsis Faunae Scandinaviae (ed. 2) 1(2): 57, tab. A. Type, by monotypy, *Motacilla troglodytes* Linnaeus.

Notes.—Formerly included in *Troglodytes*, but genetic data (Rice et al. 1999, Barker 2017, Imfeld et al. 2024) indicate that *Troglodytes sensu lato* is paraphyletic.

Change ***Troglodytes pacificus*** Baird to ***Nannus pacificus*** (Baird) and ***Troglodytes hiemalis*** Vieillot to ***Nannus hiemalis*** (Vieillot), make the appropriate changes to generic abbreviations in the Notes, move the accounts for these species to follow the heading, citation, and Notes for *Nannus*, and add the following to the end of the existing Notes for both species: See comments under *Nannus*.

Change ***Thryorchilus browni*** (Bangs) to ***Troglodytes browni*** Bangs, remove the heading for *Thryorchilus*, move the citation for *Thryorchilus* into the synonymy of *Troglodytes*, move the species account for *T. browni* to follow the heading and citation for *Troglodytes*, and replace the existing Notes with the following:

Notes.—Formerly placed in *Thryorchilus* based on larger feet and fewer rectrices than in other species of *Troglodytes*, but genetic data (Rice et al. 1999, Barker 2017, Imfeld et al. 2024) indicate that *T. browni* is very closely related to or nested within *Troglodytes*.

22. [pp. 482-483] *Cistothorus paludicola* is treated as a species separate from *C. palustris*. In the species account for *C. palustris*, change the English name to Eastern Marsh-Wren, remove the *paludicola* group from the distributional statement, remove the term "[*palustris* group]" from the distributional statement, delete "; also locally in Mexico (Río Toluca)" from the distributional statement, and change the existing Notes to:

Notes.—See comments under *C. paludicola*.

Insert the following new species account before the account for *C. palustris*:

Cistothorus paludicola Baird. Western Marsh-Wren.

Cistothorus palustris var. *paludicola* Baird, 1864, Review of American Birds in the Museum of the Smithsonian Institution, Part 1 (1864-1873): 148. (Shoalwater Bay, Washington.)

Habitat.—Freshwater and brackish marshes in cattails, tule, bulrush, and reeds; in winter and migration, also in rice fields, low dense growth bordering wetlands, and salt marsh.

Distribution.—[same as for the *paludicola* group in the account for *C. palustris* in AOU (1998), except add "also southern Mexican Plateau" to the end of the *Breeds* paragraph in the distributional statement]

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *C. palustris*, but separated based on differences in vocalizations and genomics maintained in the contact zone between the taxa (Kroodsma and Canady 1985, Kroodsma 1988, Kroodsma et al. 2026). Subspecies *tolucensis* of the southern Mexican Plateau placed with *C. paludicola* based on vocal similarities (Kroodsma et al. 2026), but may represent a separate species.

23. [p. 522] A record of Pearly-eyed Thrasher *Margarops fuscatus* in the United States is recognized. Replace the Accidental paragraph in the distributional statement with the following:

Accidental in Florida (Key West, March-April 2024; photos; Pyle et al. 2025) and on Barbados (where possibly formerly resident).

24. [p. 512] *Turdus ardosiaceus* is treated as a species separate from *T. plumbeus*. In the species account for *T. plumbeus*, change the English name to Western Red-legged Thrush, remove the *ardosiaceus* group from the distributional statement, remove the term "[*plumbeus* group]" from the distributional statement, and insert the following paragraph at the end of the distributional statement:

Accidental in Florida (four records of Bahamas subspecies *plumbeus* from Brevard County to Miami-Dade County; photos; Anderson and Ponce 2010, Kratter 2020, Johnson and Lamoreaux 2025), and one record of Cuban subspecies *rubripes* from Key West (October 2020–April 2021; photos; Kratter 2022).

Change the existing Notes to:

Notes.—See comments under *T. ardosiaceus*.

Replace the existing Notes in the species account for *Turdus ravidus* with:

Notes.—*Turdus ravidus*, *T. plumbeus*, and *T. ardosiaceus* were sometimes formerly placed in the genus *Mimocichla*.

Insert the following new species account after the account for *T. plumbeus*:

Turdus ardosiaceus Vieillot. Eastern Red-legged Thrush.

Turdus ardosiaceus Vieillot, 1823, Tableau encyclopédique et méthodique des trois régnes de la nature. Ornithologie. 2, livr. 91, p. 646. (Santo Domingo.)

Habitat.—Tropical Deciduous Forest, Tropical Lowland Evergreen Forest, Montane Evergreen Forest, Secondary Forest (0-1200 m).

Distribution.—*Resident* in Hispaniola (including Gonâve, Tortue, and Saona islands), Puerto Rico, and Dominica.

Notes.—Formerly (AOU 1983, 1998) considered conspecific with *T. plumbeus*, but separated based on differences in vocalizations (Boesman 2016b), plumage and soft part colors, and genetics (Ricklefs and Bermingham 2008, Batista et al. 2020).

25. [p. 568] Extralimital species *Fringilla spodiogenys*, *F. moreletti*, *F. maderensis*, and *F. canariensis* are treated as species separate from *F. coelebs*. In the species account for *F. coelebs*, delete "the eastern Atlantic islands," from the distributional statement and insert the following at the end of the existing Notes: Formerly (AOU 1983, 1998) considered conspecific with *F. spodiogenys* Bonaparte, 1841 [African Chaffinch], *F. moreletti* Pucheran, 1859 [Azores Chaffinch], *F. maderensis* Sharpe, 1888 [Madeira Chaffinch], and *F. canariensis* Vieillot, 1817 [Canary Islands Chaffinch], but separated based on morphological and genomic differences (Recuerda et al. 2021a, b).

26. [p. 539] *Setophaga aestiva* is treated as a species separate from *S. petechia*. In the species account for *S. petechia*, change the English name to Mangrove Warbler, remove the *aestiva* group from the distributional and habitat statements, replace

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

"southern Tamaulipas south along both coasts of Middle America (including the Bay Islands off Honduras" in the distributional statement for the *erithachorides* group with "southern Texas south along both coasts of Middle America (including Cozumel Island, the Bay Islands off Honduras,", replace the "Casual or accidental" statement for the *erithachorides* group with "in southern California (San Diego, Salton Sea), southern Arizona (Roosevelt Lake), the Revillagigedo Islands (Socorro; sight reports), and southern Texas (Cameron County, Rockport).", and replace the existing Notes with the following:

Notes.—Also known as Mangrove Yellow Warbler. Groups: *S. petechia* [Golden Warbler] and *S. erithachorides* (Baird, 1858) [Mangrove Warbler]. A new population on Cozumel Island is phenotypically intermediate between the resident subspecies of the *petechia* group and the subspecies of the *erithachorides* group on the nearby mainland, suggesting that the two groups hybridize (Machkour-M'Rabet et al. 2023). Formerly placed in the genus *Dendroica*. See comments under *Setophaga* and *S. aestiva*.

Insert the following new species account before the account for *S. petechia*:

Setophaga aestiva (Gmelin). Yellow Warbler.

Motacilla aestiva Gmelin, 1789, Systema Naturae 1(2): 996; based on "Figuier de Canada" of Daubenton 1765-1780, Planches Enluminéez, pl. 58, fig. 2. (Guiana, summers in Canada; restricted to City of Quebec, Canada, by American Ornithologists' Union 1957, Check-list of North American Birds, ed. 5, p. 487.)

Habitat.—[same as the habitat statement for the *aestiva* group in the account for *S. petechia* in AOU (1998)]

Distribution.—[same as the distributional statement for the *aestiva* group in the account for *S. petechia* in AOU (1998)]

Notes.—Also known as Northern Yellow Warbler. Formerly considered conspecific with *S. petechia*, but separated based on differences in genetics (Klein and Brown 1994, Chaves et al. 2012), vocalizations (Mennill 2001, Morton and Stutchbury 2012), morphology (Wiedenfeld 1991, Luther and Greenberg 2014), and ecology (Salgado-Ortíz et al. 2008). See comments under *Setophaga* and *S. petechia*.

27. [pp. 564-565] The group name of species placed in the genus *Myioborus* is changed from Redstart to Whitestart. Insert the following Notes under the heading and citation for Genus ***MYIOBORUS*** Baird:

Notes.—Members of the genus *Myioborus* were formerly (AOU 1983, 1998) referred to under the group name Redstart, but this was changed to Whitestart due to the increasingly widespread use of this phenotypically more appropriate name in recent literature (e.g., Ridgely and Greenfield 2001, Hilty 2003, del Hoyo et al. 2016, Gill et al. 2025), and to reflect the phylogenetic separation of *Setophaga ruticilla* [American Redstart] from the genus *Myioborus*.

28. [p. 694] Add the following species account to Part 1 of the Appendix:

Apus pallidus (Shelley). Pallid Swift.

Cypselus pallidus Shelley, 1870, Ibis 1870: 445. (Egypt.)

An emaciated swift, originally thought to be a Chimney Swift *Chaetura pelagica*, was found on a street in Montreal, Quebec, on 7 or 8 June 2007, rehabilitated, and released later in June. Photographs of the bird indicate that it was actually a juvenile swift of the genus *Apus*, likely *A. pallidus*. The date would seem to preclude a juvenile *A. apus*, but difficulty in definitively separating juveniles of *A. pallidus* from those of *A. apus* makes the identification uncertain (Pyle et al. 2025).

29. [p. 691] Add the following species account to Part 1 of the Appendix:

Zapornia pusilla (Pallas). Baillon's Crake.

Rallus pusillus Pallas, 1776, Reise durch verschiedene Provinzen des Russischen Reichs (1768-74), vol. 3, p. 700. (Dauria.)

An individual seen in flight on Attu Island, Alaska, on 20-21 September 2000, was likely this species but could also have been *Z. parva* or perhaps a juvenile *Porzana carolina*. Without a specimen or photographic documentation this record is considered unsubstantiated and the identification remains uncertain (J. J. Withrow/D. D. Gibson, in litt.; Pyle et al. 2025).

30. [p. 688] Add the following species account to Part 1 of the Appendix:

Pelecanus onocrotalus Linnaeus. Great White Pelican.

Pelecanus Onocrotalus Linnaeus, 1758, Systema Naturae, ed. 10, p. 132; based on "The Pelican" of Edwards, 1747, Nat. Hist. Birds, p. 92, pl. 92. (Africa, Asia: Caspian Sea, Russia, suggested as restricted type locality by Grant and Mackworth-Praed, 1933, Bulletin of the British Ornithologists' Club 53: 189.)

An individual of this species was seen at various localities in Florida from February 2016 (Kratter 2019) through at least January 2021. Although known as a vagrant within Eurasia outside its normal breeding and wintering range, more evidence of longer-distance movement would be needed before treating this individual as a natural vagrant (Pyle et al. 2025).

31. [p. 689] Add the following species account to Part 1 of the Appendix:

Botaurus stellaris (Linnaeus). Eurasian Bittern.

Ardea stellaris Linnaeus, 1758, Systema Naturae, ed. 10, p. 144. (Europe; restricted to Sweden by Linnaeus, 1761, Fauna Svevica, ed. 2, p. 58)

Two poor photographs of a bird seen on Buldir Island, Alaska, on 25 June 2012 (probably of the same individual seen on 13 June) suggest that the individual was this species, but are inadequate to rule out *B. lentiginosus* (J. J. Withrow/D. D. Gibson, in litt.; Pyle et al. 2025).

32. [p. 696] Delete the species account for *Tachycineta albiventer* from Part 1 of the Appendix.

33. [p. 696] Add the following species account to Part 1 of the Appendix:

Parus cinereus Vieillot. Asian Tit.

Parus cinereus Vieillot, 1818, Nouveau Dictionnaire d'Histoire Naturelle, nouv. éd. 20: 316. (Batavia, Java.)

A bird seen on Little Diomede Island, Alaska, on 2 September 1988 was likely this species, although it could have been *P. major*. Without a specimen or photographic documentation this record is considered unsubstantiated, and the identification remains uncertain (J. J. Withrow/D. D. Gibson, in litt.; Pyle et al. 2025).

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

34. [pp. 705 ff.] Make the following changes to the list of French names of North American birds:

Insert the following names in the proper position as indicated by the text of this and previous supplements:

Apus affinis	Martinet des maisons
Numenius hudsonicus	Courlis hudsonien
Tringa inornata	Chevalier des Prairies
Ichthyaetus audouinii	Goéland d'Audouin
Pterodroma deserta	Pétrel des Desertas
Sula brewsteri	Fou de Cocos
Ardeola ralloides	Crabier chevelu
Ardeola bacchus	Crabier de Chine
Ardea egretta	Aigrette d'Amérique
Circus cyaneus	Busard Saint-Martin
Pernis ptilorhynchus	Bondrée orientale
Surnia ulula	Épervière boréale
Momotus mexicanus	Motmot du Mexique
Momotus castaneiceps	Motmot du Guatemala
Camptostoma flaviventre	Tyranneau isthmien
Mitrephanes aurantiiventris	Moucherolle du Costa Rica
Deconychura typica	Grimpar de Cherrie
Pseudocolaptes lawrencii	Pseudocolapte chamois
Tachycineta albiventer	Hirondelle à ailes blanches
Nannus pacificus	Troglodyte de Baird
Nannus hiemalis	Troglodyte des forêts
Cistothorus paludicola	Troglodyte paludicole
Troglodytes browni	Troglodyte des volcans
Troglodytes musculus	Troglodyte des maisons
Troglodytes martinicensis	Troglodyte des Kalinagos
Turdus ardosiaceus	Merle dominicain
Tarsiger cyanurus	Robin à flancs roux
Setophaga aestiva	Paruline jaune
Setophaga petechia	Paruline des mangroves
Setophaga adelaidae	Paruline d'Adelaide
Chlorothraupis carmioli in APPENDIX (Part 1)	Habia de Carmiol
Apus pallidus	Martinet pâle
Zapornia pusilla	Marouette de Baillon
Pterodroma defilippiana	Pétrel de Filippi
Pelecanus onocrotalus	Pélican blanc
Botaurus stellaris	Butor étoilé
Parus cinereus	Mésange d'Asie

Delete the following names:

Pterodroma feae	Pétrel gongon
Sula brewsteri	Fou de Brewster
Ardeola bacchus	Crabier chinois
Surnia ulula	Chouette épervière
Camptostoma obsoletum	Tyranneau passegris
Deconychura longicauda	Grimpar à longue queue
Pseudocolaptes lawrencii	Anabate chamois
Troglodytes musculus	Troglodyte austral
Troglodytes martinicensis	Troglodyte de la Martinique
Troglodytes pacificus	Troglodyte de Baird
Troglodytes hiemalis	Troglodyte des forêts
Thryorchilus browni	Troglodyte des volcans
Tarsiger cyanurus	Rossignol à flancs roux
Setophaga petechia	Paruline jaune
Setophaga adelaidae	Paruline d'Adélaïde
Chlorothraupis carmioli in APPENDIX (Part 1)	Habia olive
Apus nipalensis	Martinet malais
Pterodroma defilippiana	Pétrel de De Filippi
Tachycineta albiventer	Hirondelle à ailes blanches

Change the sequence of species and genera in the family SITTIDAE, POLIOPTILIDAE, and TROGLODYTIDAE as indicated by the text of this supplement.

Proposals considered but not accepted by the Committee include changing the English group name of species of *Tympanuchus* from Prairie-Chicken to Prairie-Grouse, separating *Melanerpes caymanensis* and *M. nyeanus* from West Indian Woodpecker *M. superciliaris*, changing the linear sequence of species of *Sialis*, separating *Loxia bifasciata* from White-winged Crossbill *L. leucoptera*, separating *Leiothlypis ridgwayi* from Nashville Warbler *L. ruficapilla*, and subsuming *Leiothlypis* into *Oreothlypis*. Action has been postponed on a proposal to separate *Vireo caymanensis* from Yucatan Vireo *V. magister*.

ACKNOWLEDGMENTS

Michel Gosselin is the Committee's authority for French names, Doug Stotz provides information on Neotropical species, and Andrew W. Kratter updates the master files for the Checklist. Jacob C. Cooper, Max T. Kirsch, Diego Ocampo, David Vander Pluym, and Nicholas Vinciguerra serve on the Committee's Early Professional Systematics Group. We thank Fernanda Araujo, Juan I. Areta, Gustav Asplund, M. Juliana Benítez Saldívar, Erling Jirle, Chad David Ferguson, Martin Freeland, Bruce Hallett, Peggy M. Huss, Marshall Iliff, Tim Inskipp, Praveen Jayadevan, Roselvy Juárez, Markus Lagerqvist, Daniel F. Lane, Rafael D. Lima, Nils Navarro Pacheco, Michael O'Brien, J. V. Remsen, Jr., Andrew Spencer, Martin Stervander, Brian Sullivan, and Javier Torres López for writing or contributing to proposals for consideration by the Committee; and Juan I. Areta, Kimball Garrett, Daniel D. Gibson, Michel Gosselin, Andrew W. Kratter, Alan P. Peterson, Jack J. Withrow, the South American Classification Committee (SACC), and the Palearctic Regional Advisory Group, the Tropical Asian Regional Advisory Group, and the Taxonomic Committee of AviList for assistance, suggestions, and comments.

LITERATURE CITED

Abdulali, H. (1966). Notes on Indian birds 5 – the races of *Apus affinis* (J. E. Gray) in the Indian region. *Journal of the Bombay Natural History Society* 62:520–528.

American Ornithologists' Union (AOU) (1983). *Check-list of North American Birds*, 6th ed. American Ornithologists' Union, Lawrence, KS, USA.

American Ornithologists' Union (AOU) (1998). *Check-list of North American Birds*, 7th ed. American Ornithologists' Union, Washington, DC, USA.

Anderson, B. H., and M. S. Ponce (2010). First record of Red-legged Thrush (*Turdus plumbeus*) for Florida and the North American mainland. *North American Birds* 64:364–367.

Areta, J. I., and M. J. Benítez Saldívar (2025). Species limits in the Black-headed Antthrush (*Formicarius nigricapillus*). *Journal of Ornithology* 166:841–851.

Banks, R. C., C. Cicero, J. L. Dunn, A. W. Kratter, H. Ouellet, P. C. Rasmussen, J. V. , Remsen, J. D. Rising, and D. F. Stotz (2000). Forty-second supplement to the American Ornithologists' Union *Check-list of North American Birds*. *The Auk* 117:847–858.

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

Barker, F. K. (2017). Molecular phylogenetics of the wrens and allies (Passeriformes: Certhioidea), with comments on the relationships of *Ferminia*. *American Museum Novitates* 3887:1–28.

Batista, R., U. Olsson, T. Andermann, A. Aleixo, C. Ribas, and A. Antonelli (2020). Phylogenomics and biogeography of the world's thrushes (Aves: *Turdus*): New evidence for a more parsimonious evolutionary history. *Proceedings of the Royal Society B* 287:20192400.

Blake, E. R. (1977). *Manual of Neotropical Birds. Volume 1. Spheniscidae (Penguins) to Laridae (Gulls and Allies)*. University of Chicago Press, Chicago, IL, USA.

Boertmann, D., J. Madsen, and C. E. Mortensen (1985). Rare birds in Jameson Land, East Greenland, 1982-1984. *Dansk Ornitologisk. Forenings Tidsskrift* 79:151–152 (Danish, with English summary).

Boesman, P. (2016a). Notes on the vocalizations of the Long-tailed Woodcreeper (*Deconychura longicauda*). HBW Alive Ornithological Note 78. In *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona, Spain.

Boesman, P. (2016b). Notes on the vocalizations of Red-legged Thrush (*Turdus plumbeus*). HBW Alive Ornithological Note 310. In *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona, Spain.

Boesman, P. (2017). Notes on the vocalizations of Monk Parakeet (*Myiopsitta monachus*) and Cliff Parakeet (*Myiopsitta luchsi*). HBW Alive Ornithological Note 452. In *Handbook of the Birds of the World Alive*. Lynx Edicions, Barcelona, Spain.

Brooke, R. K. (1971). Geographical variation in the little swift *Apus affinis* (Aves: Apodidae). *Durban Museum Novitates* 9:93–103.

Buainain, N., M. F. A. Maximiano, M. Ferreira, A. Aleixo, B. C. Faircloth, R. T. Brumfield, J. Cracraft, and C. C. Ribas (2021). Multiple species and deep genomic divergences despite little phenotypic differentiation in an ancient Neotropical songbird, *Tunchiornis ochraceiceps* (Sclater, 1860) (Aves: Vireonidae). *Molecular Phylogenetics and Evolution* 162:107206.

Chantler, P., P. F. D. Boesman, P. Pyle, and C. Hansasuta (2026). House Swift (*Apus nipalensis*), version 1.1. In *Birds of the World* (J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, E. de Juana, and B. K. Keeney, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.houswi1.01.1

Chaves, J. A., P. G. Parker, and T. B. Smith (2012). Origin and population history of a recent colonizer, the Yellow Warbler in Galapagos and Cocos Islands. *Journal of Evolutionary Biology* 25:509–521.

Chesser, R. T., R. C. Banks, F. Keith Barker, C. Cicero, J. L. Dunn, A. W. Kratter, I. J. Lovette, P. C. Rasmussen, J. V. , Remsen, J. D. Rising, D. F. Stotz, and K. Winker (2013). Fifty-fourth supplement to the American Ornithologists' Union *Check-list of North American Birds*. *The Auk* 130:558–571.

Chesser, R. T., K. J. Burns, C. Cicero, J. L. Dunn, A. W. Kratter, I. J. Lovette, P. C. Rasmussen, J. V. , Remsen, J. D. Rising, D. F. Stotz, and K. Winker (2017). Fifty-eighth supplement to the American Ornithological Society's *Check-list of North American Birds*. *The Auk: Ornithological Advances* 134:751–773.

Chesser, R. T., K. J. Burns, C. Cicero, J. L. Dunn, A. W. Kratter, I. J. Lovette, P. C. Rasmussen, J. V. Remsen, Jr., D. F. Stotz, B. M. Winger, and K. Winker (2018). Fifty-ninth supplement to the American Ornithological Society's *Check-list of North American Birds*. *The Auk: Ornithological Advances* 135:798–813.

Chesser, R. T., K. J. Burns, C. Cicero, J. L. Dunn, A. W. Kratter, I. J. Lovette, P. C. Rasmussen, J. V. , Remsen, D. F. Stotz, and K. Winker (2019). Sixtieth supplement to the American Ornithological Society's *Check-list of North American Birds*. *The Auk: Ornithological Advances* 136:ukz042.

Cramp, S., and K. E. L. Simmons (1983). *The Birds of the Western Palearctic. Volume 3: Waders to Gulls*. Oxford University Press, Oxford, UK.

del Hoyo, J., N. J. Collar, D. A. Christie, A. Elliott, L. D. C. Fishpool, P. Boesman, and G. M. Kirwan (2016). *HBW and BirdLife International Illustrated Checklist of the Birds of the World. Volume 2: Passerines*. Lynx Edicions and BirdLife International, Barcelona, Spain and Cambridge, UK.

Douglas, H. D. (1998). Response of Eastern Willets (*Catoptrophorus s. semipalmatus*) to vocalizations of Eastern and Western (*C. s. inornatus*) willets. *The Auk* 115:514–518.

Dunn, J. L., D. D. Gibson, K. L. Garrett, M. J. Iliff, M. Lockwood, R. Pittaway, D. Sibley, and K. J. Zimmer (2012). 23rd report of the ABA Checklist Committee. *Birding* 44:28–33.

Dyer, D., and S. N. G. Howell (2023). *Birds of Costa Rica*. Princeton Field Guides. Princeton University Press, Princeton, NJ, USA.

Farnsworth, A., and D. J. Lebbin (2020). Tufted Flycatcher (*Mitrephanes phaeocercus*), version 1.0. In *Birds of the World* (J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, and E. de Juana, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA.

Gibson, D. D and G. V. Byrd (2007). *Birds of the Aleutian Islands, Alaska*. Series in Ornithology 1. Nuttall Ornithological Club and American Ornithologists' Union, Cambridge, MA, and Washington, D.C., USA.

Gibson, D. D., and J. Withrow (2015). Inventory of the species and subspecies of Alaska birds, second edition. *Western Birds* 46:94–185.

Gill, F., D. Donsker, and P. C. Rasmussen (Editors) (2025). IOC World Bird List (v 15.1). https://doi.org/10.14344/IOC.ML.15.1. http://www.worldorg/

Gustad, J. R. (1994). Sjeldne fugler i Norge i 1992. Rapport fra Norsk sjeldenhetskomite for fugl (NSKF). *Vår Fuglefauna* 17: 259–278.

Harvey, M. G., G. A. Bravo, S. Claramunt, *et al.* (2020). The evolution of a tropical biodiversity hotspot. *Science* 370:1343–1348.

Heath, S. A., J. K. Wilson, J. Smith, and N. W. Cooper (2021). Migration and wintering locations of breeding eastern Willet (*Tringa semipalmata semipalmata*) in the western Gulf of Mexico. *Waterbirds* 44:483–491.

Hilty, S. L. (2003). *Birds of Venezuela*. Christopher Helm, London, UK.

Howell, S. N. G. (2012). *Petrels, Albatrosses, and Storm-Petrels of North America: A Photographic Guide*. Princeton University Press, Princeton, NJ, USA.

Howell, S. N. G., and R. Juárez (2025). A tale of two motmots. *Neotropical Birding* 37:19–23.

Iliff, M., and P. C. Rasmussen (2026). Treat Tufted Flycatcher *Mitrephanes phaeocercus* as two or three species. Proposal 2026-B-9 to N&MA Classification Committee, American Ornithological Society. https://americanornithology.org/about/committees/nacc/current-priorproposals/2026-proposals/

Imfeld, T. S., F. K. Barker, H. Vázquez-Miranda, J. A. Chaves, P. Escalante, G. M. Spellman, and J. Klicka (2024). Diversification and dispersal in the Americas revealed by new phylogenies of the wrens and allies (Passeriformes: Certhioidea). *Ornithology* 141:ukae007.

Inskipp, T., and J. Praveen (2026) Treat Little Swift *Apus affinis* and House Swift *A. nipalensis* as a single species. Proposal 2026-C-9 to N&MA Classification Committee, American Ornithological Society. https://americanornithology.org/about/committees/nacc/current-prior-proposals/2026-proposals/

Johnson, W., and A. Lamoreaux (2025). Red-legged Thrush and other Bahamian vagrants in Florida, Spring 2025. *North American Birds* 76(1).

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

Kirwan, G. M., R. Sample, B. Shackelford, R. Kannan, and P. F. D. Boesman (2024). Atlantic Royal Flycatcher (*Onychorhynchus swainsoni*), version 1.1. In *Birds of the World* (T. S. Schulenberg and B. K. Keeney, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.royfly5.01.1

Klein, N. K., and W. M. Brown (1994). Intraspecific molecular phylogeny in the Yellow Warbler (*Dendroica petechia*) and implications for avian biogeography in the West Indies. *Evolution* 48:1914–1932.

König, C., P. Heidrich, and M. Wink (1996). Zur Taxonomie der Uhus (*Bubo* ssp.) im südlichen Südamerika. *Stuttgarter Beiträge zur Naturkunde, Series A*, 540.

Kratter, A. W. (2018). Twenty-sixth report of the Florida Ornithological Society Records Committee: 2016. *Florida Field Naturalist* 46:8–28.

Kratter, A. W. (2019). Twenty-eighth report of the Florida Ornithological Society Records Committee: 2018. *Florida Field Naturalist* 47:60–81.

Kratter, A. W. (2020). Twenty-ninth report of the Florida Ornithological Society Records Committee: 2019. *Florida Field Naturalist* 48:105–129.

Kratter, A. W. (2022). Thirtieth report of the Florida Ornithological Society Records Committee: 2019–2021. *Florida Field Naturalist* 50:100–138.

Kroodsma, D. E., and R. A. Canady (1985). Differences in repertoire size, singing behavior, and associated neuroanatomy among Marsh Wren populations have a genetic basis. *The Auk* 102:439–446.

Kroodsma, D. E. (1988). Two species of Marsh Wren (*Cistothorus palustris*) in Nebraska? *Nebraska Bird Review* 56:40–42.

Kroodsma, D. E., D. A. DeRaad, B. E. Byers, D. E. Winslow, and F. K. Barker (2026). Genetic analysis of *Cistothorus palustris* (Marsh Wren) across a broad transition zone in Southern Saskatchewan reveals deep divergence and little hybridization between two cultural phenotypes. *Ornithology* 143:ukaf056.

Lima, R. D., and R. V. Vaz (2024). Divergence in vocalizations indicates cryptic speciation in *Camptostoma* tyrannulets. *Ornithology* 142:ukae058.

López-Lanús, B. (2015). Análisis comparativo de las vocalizaciones de distintos taxa del género *Bubo* en América. *Hornero* 30:69–88.

Lowther, P. E., H. D. Douglas, and C. L. Gratto-Trevor (2020). Willet (*Tringa semipalmata*), version 1.0. In *Birds of the World* (A. F. Poole and F. B. Gill, Eds.). Cornell Lab of Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.willet1.01

Luther, D., and R. Greenberg (2014). Habitat type and ambient temperature contribute to bill morphology. *Ecology and Evolution* 4:699–705.

Machkour-M'Rabet, S., W. Santamaría-Rivero, A. Dzib-Chay, L. T. Cristiani, and B. MacKinnon-Haskins (2023). Multi-character approach reveals a new mangrove population of the Yellow Warbler complex, *Setophaga petechia*, on Cozumel Island, Mexico. PLoS One18:e0287425.

Marantz, C. A., A. Aleixo, L. R. Bevier, and M. A. Patten (2003). Family Dendrocolaptidae (woodcreepers). In *Handbook of the Birds of the World, Vol. 8. Broadbills to Tapaculos*. (J. del Hoyo *et al.*, Editors). Lynx Edicions, Barcelona, Spain. pp. 358–447.

McLaughlin, J. F., B. C. Faircloth, T. C. Glenn, and K. Winker (2020). Divergence, gene flow, and speciation in eight lineages of trans-Beringian birds. *Molecular Ecology* 29:3526–3542.

Mennill, D. J. (2001). Song characteristics and singing behavior of the Mangrove Warbler (*Dendroica petechia bryanti*). *Journal of Field Ornithology* 72:327–337.

Morton, E. S., and B. J. M. Stutchbury (2012). Vocal communication in androgynous territorial defense by migratory birds. *International Scholarly Research Notices Zoology* 729301. https://onlinelibrary.wiley.com/doi/10.5402/2012/729307

O'Brien, M. (2006). Subspecific identification of the Willet (*Catoptrophorus semipalmatus*). *Birding* 38(3):41–47.

Ostrow, E. N., L. H. DeCicco, and R. G. Moyle (2023). Range-wide phylogenomics of the Great Horned Owl (*Bubo virginianus*) reveals deep north-south divergence in northern Peru. *PeerJ* 11:e15787.

Oswald, J. A., M. G. Harvey, R. C. Remsen, D. U. Foxworth, S. Cardiff, D. L. Dittmann, L. C. Megna, M. D. Carling, and R. T. Brumfield (2016). Willet be one species or two? A genomic view of the evolutionary history of *Tringa semipalmata*. *The Auk* 133:593–614.

Päckert, M., J. Martens, J. M. Wink, A. Feigl, and D. T. Tietze. (2012). Molecular phylogeny of Old World swifts (Aves: Apodiformes, Apodidae, *Apus* and *Tachymarptis*) based on mitochondrial and nuclear markers. *Molecular Phylogenetics and Evolution* 63:606–616.

Pohlen, Z. M., J. A. Johnson, R. B. Benter, and J. Helm (2026). First North American record of the Oriental Honey-Buzzard (*Pernis ptilorhynchus*). *Western Birds* 57:64–67.

Pyle, P., L. DeCicco, D. Gochfeld, M. Hafting, L. Harter, M. Heindel, A. Jaramillo, A. W. Kratter, M. Lockwood, M. Mutchler, and D. Sibley (2025). 36th report of the ABA Checklist Committee 2025. *North American Birds* 76:26–35.

Pyle, R.L., and P. Pyle (2017). *The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status*. B.P. Bishop Museum, Honolulu, HI, USA. Version 2 (1 January 2017). http://hbs.bishopmuseum.org/birds/rlp-monograph/

Ramírez, I., V. H. Paiva, D. Menezes, I. Silva, R. A. Phillips, J. A. Ramos, and S. Garthe (2013). Year-round distribution and habitat preferences of the Bugio Petrel. *Marine Ecology Progress Series* 476:269–284.

Recuerda, M., J. C. Illera, G. Blanco, R. Zardoya, and B. Milá (2021a). Sequential colonization of oceanic archipelagos led to a species-level radiation in the common chaffinch complex (Aves: *Fringilla coelebs*). *Molecular Phylogenetics and Evolution* 164:107291.

Recuerda, M., J. Vizueta, C. Cuevas-Caballé, G. Blanco, J. Rozas, and B. Milá (2021b). Chromosome-level genome assembly of the Common Chaffinch (Aves: *Fringilla coelebs*): A valuable resource for evolutionary biology. *Genome Biology and Evolution* 13:evab034.

Remsen, , J. V.Jr., J. I. Areta, E. Bonaccorso, S. Claramunt, D. F. Lane, L. N. Naka, M. B. Robbins, F. G. Stiles, and K. J. Zimmer (Version 25 June 2026). *A Classification of the Bird Species of South America*. Museum of Natural Science, Louisiana State University, LA, USA. http://www.museum.lsu.edu/Remsen/SACCBaseline.htm

Reyes, P., J. M. Bates, L. N. Naka, M. J. Miller, I. Caballero, C. Gonzalez-Quevedo, J. L. Parra, H. F. Rivera-Gutierrez, E. Bonaccorso, and J. G. Tello (2023). Phylogenetic relationships and biogeography of the ancient genus *Onychorhynchus* (Aves: Onychorhynchidae) suggest cryptic Amazonian diversity. *Journal of Avian Biology* 2023:e03159.

Reyt, S., P.-A. Crochet, S. Reeber, and P. Dufour (2025). Identification of 'Great Egret' taxa in winter plumage. *Dutch Birding* 47:73–97.

Rice, N. H., A. T. Peterson, and G. Escalona-Segura (1999). Phylogenetic patterns in montane *Troglodytes* wrens. *The Condor* 101:446–451.

Ricklefs, R. E., and E. Bermingham (2008). Likely human introduction of the Red-legged Thrush (*Turdus plumbeus*) to Dominica, West Indies. *The Auk* 125:299–303.

Ridgely, R. S., and P. J. Greenfield (2001). *The Birds of Ecuador, Volumes 1–2*. Cornell University Press, Ithaca, NY, USA.

Robb, M., and K. Mullarney (2008) *Petrels Night and Day*. Sound Approach, Dorset, UK.

Salgado-Ortíz, J., P. P. Marra, T. S. Sillett, and R. J. Robertson (2008). Breeding ecology of the Mangrove Warbler (*Dendroica petechia*

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

bryanti) and comparative life history of the Yellow Warbler species complex. *The Auk* 125:402–410.

Skeel, M. A. (1978). Vocalizations of the Whimbrel on its breeding grounds. *The Condor* 80:194–202.

Skeel, M. A., E. P. Mallory, and P. Pyle (2025). Hudsonian Whimbrel (*Numenius hudsonicus*), version 1.0. In *Birds of the World* (S. M. Billerman and M. G. Smith, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA. https://doi.org/10.2173/bow.whimbr3.01

Slager, D. L., C. J. Battey, R. W. Bryson, G. Voelker, and J. Klicka (2014). A multilocus phylogeny of a major New World avian radiation: The Vireonidae. *Molecular Phylogenetics and Evolution* 80:95–104.

Sordahl, T. A. (1979). Vocalizations and behavior of the Willet. *The Wilson Bulletin* 91:551–574.

Tan, H. Z., E. Y. X. Ng, Q. Tang, G. A. Allport, J. J. F. J. Jansen, P. S. Tomkovich, and F. E. Rheindt (2019). Population genomics of two congeneric Palaearctic shorebirds reveals differential impacts of Quaternary climate oscillations across habitat types. *Scientific Reports* 9:18172.

Tan, H. Z., J. J. F. J. Jansen, G. A. Allport, K. M. Garg, B. Chattopadhyay, M. Irestedt, S. E. Pang, G. Chilton, C. Y. Gwee, and F. E. Rheindt (2023). Megafaunal extinctions, not climate change, may explain Holocene genetic diversity declines in *Numenius* shorebirds. *eLife* 12:e85422.

Traylor, M. A. (1958). Variation in South American Great Horned Owls. *The Auk* 75:143–149.

Webster, J. D. (1968). A revision of the tufted flycatchers of the genus *Mitrephanes*. *The Auk* 85:9.

Wiedenfeld, D. A. (1991). Geographical morphology of male Yellow Warblers. *The Condor* 93:712–723.

Williamson, K. (1946). Field-notes on the breeding-biology of the Whimbrel, *Numenius phaeopus phaeopus* (Linnaeus). *The North Western Naturalist* 21:167–184.

Withrow, J. J., A. M. Spicer, and D. W. Sonneborn (2024). Mitochondrial DNA demonstrates that a Hen Harrier (*Circus cyaneus*) reached Attu Island, Alaska. *Western Birds* 55:280–292.

Downloaded from academic.oup.com/auk/article/143/4/1/8934017 by guest on 31 October 2026

© American Ornithological Society 2026. All rights reserved. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

Ornithology 2026 143(4) 1–14
https://doi.org/10.1093/ornithology/ukag037
Check List Supplement