



universität
wien

Φnm

FWF

Der Wissenschaftsfonds.

erc

Open-Source 4D-STEM Phase Reconstructions of Focused-Probe Data with Near-Ideal Direct-Electron Detection


Toma Susi, Jani Kotakoski, Clemens Mangler (U. Vienna)
Niklas Dellby, Russ Hayner, Tracy Lovejoy, Andreas Mittelberger, Benjamin Plotkin-Swing (Nion Co. R&D / Bruker AXS)

Christoph Hofer, Timothy Pennycook (U. Antwerp)



Open-source software

Language




Python
(3.11.9)

Data Acquisition




Nion Swift
(0.16.10)

Phase Reconstructions



PyPtychoSTEM
(0.14.16)



py4DSTEM
(0.14.16)



Stephanie
Ribet



Georgios
Varnavides

Reconstruction Algorithms:

single-sideband (**SSB**), Wigner distribution deconvolution (**WDD**),
(iterative) differential phase contrast (**DPC**),
parallax imaging ie. tilt-corrected bright-field STEM (**parallax**),
interative gradient descent ptychography (**GD**)



Near-ideal direct-electron 4D-STEM

Number of pixels (W x H)	192 × 192
Active area (W x H) [mm²]	20 × 20
Pixel size (W x H) [μm²]	100 × 100
Sensor material	Silicon (Si) or high-Z
Energy range [keV]	30–300
Frame rate (max.) [Hz]	120,000
Count rate (max.) [el/s/pixel]	10⁸
Detective Quantum Efficiency, DQE(0)	0.82 @ 80 keV 0.75 @ 200 keV, 0.75 @ 300 keV
Detector mounting	Retractable



Near-ideal 4D-STEM performance at 60 keV

**Installation of production unit #001
in Vienna at the start of 2024**



4D-STEM data volumes


Data volumes collected during typical 4D-STEM experiments (assuming a 512×512 px scan).

Mode	Pixels	Bit depth	Bits / CBED	Frame rate	MB / s	GB/scan
Saved file (hdf5)	192×192	12	442368	20000	1106	
Numpy (fp16)	192×192	16	589824	-	-	39
Reduced file (hdf5)	96×96	8	73728	120000	1106	
Numpy (fp16)	96×96	16	147456	-	-	9.7


- Huge data volumes can be collected in seconds (challenge for storage, processing...)

Ptychography can correct aberrations post-acquisition

PACBED




Spatial frequencies




$A(K)^* A(K-Q)$


'Exp'






SVD



corrected



Theory



SVD = singular value decomposition



Timothy
Pennycook




Christoph
Hofer


3rd order aberration correction might not be sufficient

Experimental data of monolayer graphene, 35 mrad (Vienna)


uncorrected



3rd order corrected



5th order corrected






Timothy
Pennycook



Christoph
Hofer

Site-specific phase via iterative optimization



Maximize correlation between experimental and simulated image

$$R = \frac{\sum_{i=1}^N \left((\mu^{sim} - I_i^{sim}) (\mu^{exp} - I_i^{exp}) \right)}{\sigma^{sim} \sigma^{exp} (N - 1)}$$

I^{exp} and I^{sim} = image intensities

μ^{exp} and μ^{sim} = mean values

σ^{sim} and σ^{exp} = standard deviations

- Can account for:
- aberrations
 - incoherence
 - scan distortions
 - sample drift
 - sample tilt
 - etc...

Phase analysis




STEM_Optimization*
(commit 6c549ba9)

Defective WS_2

Can account for:

- aberrations
- incoherence
- scan distortions
- sample drift
- sample tilt
- etc...





Timothy
Pennycook




Christoph
Hofer

Ptychography for charge transfer in defective WS₂




Graphene: ideal uniform phase object

- ✓ ∞ radiation hardness <80 keV \Rightarrow no dose limitation
- ✓ low- Z , one-atom-thick \Rightarrow perfect weak phase object
- ✓ only one element \Rightarrow each atomic site identical






Nion + ARINA 4D-STEM data collection (original)



HAADF (80–300 mrad)


Nion + ARINA 4D-STEM data collection (original)

512 × 512 px scan (**Rpix**), 96 × 96 px CBED (**Qpix**; binned by 4 for analysis), 100 us dwell time




- 34 mrad semi-convergence angle, ~70 mrad max collection angle, ~8 pA
- counting every electron, corrected dose per pixel can be normalized

Nion + ARINA 4D-STEM data collection (final)




$\sim 200 \text{ pA current!}$

Comparing phase reconstruction algorithms (Q binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 192×192 Qpix
1× binning in Q

Comparing phase reconstruction algorithms (Q binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 192×192 Qpix
2× binning in Q

Comparing phase reconstruction algorithms (Q binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 192×192 Qpix
4× binning in Q

Comparing phase reconstruction algorithms (Q binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 192×192 Qpix
8× binning in Q

Comparing phase reconstruction algorithms (Q binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 192×192 Qpix
16× binning in Q

Comparing phase reconstruction algorithms (Q binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 192×192 Qpix
32× binning in Q

Comparing phase reconstruction algorithms (R binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 96×96 Qpix (HW binQ by 2)
1× binning in R

Comparing phase reconstruction algorithms (R binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 96×96 Qpix (HW binQ by 2)
2× binning in R

Comparing phase reconstruction algorithms (R binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 96×96 Qpix (HW binQ by 2)
4× binning in R

Comparing phase reconstruction algorithms (R binning)



36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 96×96 Qpix (HW binQ by 2)
8× binning in R


Comparing phase reconstruction algorithms (R binning)




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 96×96 Qpix (HW binQ by 2)
16x binning in R

Atomic phase assignment


Optimization of atomic phases for reconstructions



Phase variation over graphene C atoms




Quantification of phase variation




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 192×192 Qpix (**binQ**)

Quantification of phase variation




36 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 192×192 Qpix (**binR**)

Quantification of phase variation



60 mrad maximum scattering angle, 33.3 us dwell time, 512×512 Rpix, 192×192 Qpix

Quantification of phase variation




109 mrad maximum scattering angle, 100 us dwell time, 256×256 Rpix, 192×192 Qpix

Computational cost

Scan (px)	binQ	CBED (Qpix)	Array (MB)	DPC (s)	Parallax (s)	Iter. GD* (s)	SSB (s)	SSB AC (s)	WDD (s)	WDD AC (s)
512×512	1	192×192	38656	16	–	578	236	18097		–
	2	96×96	9664	6.1	855	517	59	888	1938	2375
	4	48×48	2416	6.9	728	195	34	272	714	784
	8	24×24	604	5.4	68	146	13	87	306	317
	16	12×12	151	5.0	24	126	7.2	41	163	165
	32	6×6	38	4.4	22	122	7.1	35	144	147

Future prospects



- Defocused-probe ptychography with iterative reconstructions
- Variable camera lengths (for super-resolution up to $\sim 4 \times \alpha$)
- Live virtual detectors / (semi-)live single-sideband ptychography?
- Structure & charge transfer in various materials... *and much more*

Upcoming Japan stay (Sep 2024 – Feb 2025)




Nagashima (TUS)




Suenaga (Osaka)



Shibata (Tokyo)



you? (???)



Thank you!

Questions, collaborations: *toma.susi@univie.ac.at*



Open code & talk slides:
<https://github.com/TomaSusi/arina-ptycho>

Open datasets:
<https://phaidra.univie.ac.at/o:2081765>

